Nanoparticle-delivered resiquimod induces brain tumor regression in medulloblastoma and diffuse midline glioma models by interrupting paracrine growth support and activating myeloid immune signaling and phagocytosis

Leon McSwain[1], Kyongtea Kim[2], Duhyeong Hwang[2,3], Chaemin Lim[2,4], Cynthia Winham[1], Jon Jacques[1], Elias Rosen[5], Sudhir Kasturi[6], Arpan Pradhan[6] Andrey Tikunov[1], Alexander V. Kabanov[2], Jessica Raper[1,6], Timothy R. Gershon[1,7], Marina Sokolsky[2,7]

[1] Department of Pediatrics, Emory University, Atlanta, GA 30323, USA
[2] Center for Nanotechnology in Drug Delivery and Division of Pharmacoengineering and Molecular Pharmaceutics, Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, NC 27599, USA.
[3] College of Pharmacy, Keimyung University, 1095 Dalgubeol-daero, Dalseo-gu, Daegu 42601, Republic of Korea
[4] College of Pharmacy, CHA University, 335 PangyoPangyo-ro, Bundang-gu, Seongnam-si, Gyeonggi-do 13488, Republic of Korea.
[5] Eshelman School of Pharmacy, University of North Carolina, Chapel Hill, North Carolina, United States.
[6] Please fill in
[7] These authors contributed equally.

ABSTRACT

We studied the effect of stimulating innate immune function in tumor-associated myeloid cells (TAMs) in medulloblastoma (MB) and diffuse midline glioma (DMG), using a polyoxazoline nanoparticle formulation of the TLR7/8 agonist resiquimod (ResiPOx). Children with MB and DMG need novel therapeutic strategies to improve outcomes and reduce recurrence. We investigated the effect of systemically administered ResiPOx on TAMs in MB and DMG using endogenous MB and DMG models in immune-competent mice and identified multiple mechanisms of anti-tumor effect.

We packaged resiquimod into polyoxazoline micelles to generate ResiPOx. We studied ResiPOx efficacy as a single agent or paired with radiation therapy (RT). We determined ResiPOx pharmacokinetics (PK) using tritium-labeled resiquimod and mass spectroscopy imaging (MSI). We determined ResiPOx pharmacodynamics (PD) using flow cytometry immunohistochemistry, bulk and single-cell RNA-seq and immunoblotting. We then studied ResiPOx safety and PD in a non-human primate model using rhesus macaques.

ResiPOx formulation improved the blood-brain barrier penetration and anti-tumor efficacy of resiquimod. ResiPOx treatment extended progression-free survival (PFS) in mice with MB and DMG. In both tumor types, ResiPOx expanded TAM populations and reprogrammed TAMs toward anti-tumoral states, characterized by activation of IFNβ and extrinsic apoptosis pathway signaling, antigen presentation, and T cell activation signatures. In rhesus macaques, systemic ResiPOx administration was well tolerated and induced brain transcriptional responses that resembled ResiPOx responses in DMG and MB mouse models, indicating common effects across species from mice to non-human primates, and highlighting potential for similar effects in patients.

ResiPOx is a brain-penetrant immunomodulatory therapeutic that reshapes the immune-privileged brain tumor microenvironment. Systemic administration activates myeloid-driven anti-tumoral immunity mediated by microglial and macrophage TAMs, and improves survival in preclinical models of DMG and MB.

INTRODUCTION

Growth-supportive interactions between brain tumors and tumor-associated myeloid cells (TAMs) present opportunities for therapeutic intervention. Myeloid cells, including microglia and bone marrow-derived macrophages (BMDMs), are the most numerous immune cells in the CNS. These brain myeloid cells shift between pro-inflammatory and anti-inflammatory phenotypes and modulate cytokine and growth factor secretion in order to coordinate immunosurveillance, response to injury and developmental proliferation [1-4]. In brain tumors, however, including both medulloblastoma (MB) and high-grade glioma, tumor-associated myeloid cells (TAMs) predominantly show immunosuppressive phenotypes and support tumor growth through paracrine signaling [5-9]. Here we examined the potential for pathogen receptor activation to shift TAMs from tumor-supportive to tumor-suppressive states and to produce therapeutic benefits in primary mouse models of pediatric brain tumors, including Sonic Hedgehog (SHH)-driven MB and diffuse midline glioma (DMG).

TAMs play critical roles in SHH MB and DMG. TAMs in SHH MBs secrete IGF1 ligand, which acts as a paracrine growth factor that promotes tumor progression [9], by activation IGF receptor (IGF1R) on MB cells, which potentiates SHH signaling [10, 11]. Reducing TAM populations by treatment with CSF1R inhibitors slows SHH MB progression in mouse models [12]. However, TAMs in SHH MB are heterogenous [13] and while specific TAM populations are associated with tumor growth [9, 14], other populations are tumor suppressive [15, 16]. For example, meningeal macrophages secrete the chemokine CXCL4, which inhibits tumor-promoting CXCL12/CXCR4 signaling, with the effect of slowing SHH MB tumorigenesis [16]. In DMGs, TAMs secrete growth factors that promote tumor progression [17, 18] and adopt immunosuppressive phenotypes [19, 20], but retain a capacity for anti-tumor effects through phagocytosis of tumor cells [21]. Therapies that can enable control of TAM phenotypes may provide new approaches broadly applicable to both MB and DMG.

TAMs express pathogen-associated molecular pattern (PAMP) receptors, including Toll-like receptors 7 and 8 (TLR7/8) [22, 23], which mediate dynamic reactions to changes in the tumor microenvironment (TME) [24, 25, 20]. TLR7/8 activation has received significant attention as a promising immunomodulatory approach in diverse cancers [26-30]. Resiquimod (R848) is a TLR7/8 agonist that has shown activity with systemic administration in trials for melanoma and other solid malignancies, but has been limited by side effects associated with immune hyper-activation, including cerebral edema [31-34]. Incorporating resiquimod into a nanoparticle delivery system can mitigate the challenges of poor solubility and both local and systemic toxicities [30, 35-37]. In brain tumor therapy, resiquimod has shown promise in glioma models when delivered in systemic nanoparticles [38] or locally in slow-releasing solid matrix formulation [39]. These studies showed potent antitumor effects when used to treat mice with implanted syngeneic gliomas, in which the tumor and host derive from different individual animals of the same species.

To advance resiquimod as a potential therapy for multiple types of pediatric brain tumors, we formulated Resiquimod-loaded polyoxazoline nanoparticles (ResiPOx), and investigated ResiPOx in mice engineered to develop primary SHH MB or DMG. These models feature endogenous tumors, with intact blood brain barrier (BBB) [40] and without potentially immunostimulatory tumor-host mismatch, thus modeling the typically immunologically "cold" phenotype of MBs and DMGs that occur in patients [20, 41]. Polyoxazoline nanoparticle formulations have shown improved CNS delivery of diverse small molecule agents [42, 43] suggesting the potential for ResiPOx to deliver resiquimod across the blood brain barrier directly to the brain tumor TAMs. To investigate this potential, we administered ResiPOx to mice with SHH MB, determined PK and efficacy, and defined pharmacodynamic (PD) effects with cellular and molecular detail. We then compared efficacy in mice with DMG and tolerability and CNS PD in Rhesus macaques, as a model closer in evolution to humans. Our data show that ResiPOx delivers active resiquimod to brain tumors, acts directly on TAMs, synergizes with radiotherapy, and produces similar effects on brain myeloid cells in multiple species, supporting translational potential.

RESULTS

ResiPOx enhances delivery across the blood brain barrier and achieves bioactive exposure.

We used tritium-labeled resiquimod to compare the PK of ResiPOx or free drug after systemic administration. We incorporated tritium-labeled resiquimod into ResiPOx and generated mice with medulloblastoma by breeding *hGFAP-Cre* and *SmoM2* mouse lines to generate *hGFAP-Cre/SmoM2* (*G-Smo*) progeny [44, 45]. We administered tritium-labeled resiquimod as ResiPOx or free drug to replicate P10 *G-Smo* mice by IP injection. We then harvested the mice at successive time points and analyzed resiquimod concentrations in plasma and tissue samples by scintillation counting, processed with R permutation analysis.

ResiPOx showed higher peak drug concentrations (Cmax) in plasma, forebrain, and tumor (**Fig. 1A**), and higher systemic exposure, determined by area under the curve (**AUC**). Plasma, forebrain, and tumor showed slower decay of ResiPOx and increased AUC (ΔAUC = 4773.5 p < 0.001 (Plasma), 1523.0 p < 0.0001 (Forebrain), 2265.6 p < 0.0001 (Tumor)) compared to free resiquimod. We also tracked PK in liver, kidney and spleen (**Supplementary Fig. 1**). For orthogonal validation of ResiPOx delivery into the tumor and adjacent brain, we used spectroscopy imaging (MSI) effected by matrix-assisted laser desorption electrospray ionization (IR-MALDESI). This technique applies an infrared laser to frozen tissue sections in an ice matrix and generates concentration maps for all detected metabolites within a present molecular weight range. We mapped the neuronal marker N-acetylaspartic acid (NAA) to distinguish brain from tumor, and mapped creatine to distinguish the edges of the tissue. MSI confirmed detectable accumulation of resiquimod in the tumor and surrounding brain 3 hours after ResiPOx injection, which was no longer detectable by 24 hours (**Fig. 1B**); Together, these data show that ResiPOx enhanced resiquimod PK and successfully delivered the drug into the brain and TME.

To determine the potential clinical relevance of ResiPOx-enhanced PK, we compared progression-free survival (PFS) time in *G-Smo* mice treated with ResiPOx or free resiquimod to saline-treated *G-Smo* controls. We randomized *G-Smo* mice to ResiPOx ResiPOx or free resiquimod, administered in three 5 mg/kg doses at P10, P12 and P14. We then followed mouse weight and neurologic status and determined PFS as the time to the humane endpoint of symptomatic tumor progression, detected by sustained weight loss >10% of body weight or onset of neurologic symptoms. ResiPOx-treated mice showed significantly increased PFS compared to either saline or free resiquimod, which were similar (**Fig. 1C**).

ResiPOx was not curative as a single agent in any mice, as all mice showed eventual tumor progression. Because medulloblastoma chemotherapy is typically combined with radiation therapy (RT), we evaluated the efficacy of ResiPOx combined with concurrent RT as a more clinically-relevant approach. Our prior studies showed that *G-Smo* tumors are relatively radiation resistant compared to other genetic mouse MB models [46] but show a modest survival benefit with an optimal xRT dose of three fractions of 0.5Gy [45]. We randomized mice to ResiPOx or saline on P10, P12 and P14, and treated all mice with 0.5 Gy xRT on P11, 13 and 15. We compared survival between these two groups, and to *G-Smo* mice treated with ResiPOx alone shown in **Fig. 1C**. *G-Smo* mice treated with ResiPOx plus RT showed improved PFS compared to either RT alone or ResiPOx alone (**Fig. 1D**), demonstrating mutual enhancement.

ResiPOx induces changes in TAM and tumor populations

To gain insight into the mechanisms of ResiPOx anti-tumor efficacy, we analyzed ResiPOx effects on TAM populations in *G-Smo* MBs. We randomly assigned *G-Smo* mice to treatment with ResiPOx at P10, P12 and P14 or to no treatment. We then harvested tumors on P15, 24 hours after the last ResiPOx dose, subjected dissociated tumors to flow cytometry and compared treated versus control replicates. Immunostaining for CD45 and CD11b identified discrete TAM subsets in *G-Smo* MBs, including BMDM TAMs, characterized by high CD45 and CD11b expression (**CD45^{hi}/CD11B^{hi}**) and microglial TAMs with intermediate expression of both markers (**CD45^{int}/CD11B^{int}**). The microglial TAM compartment comprised two subpopulations, one with relatively higher CD45 expression (MG^{CD45}) and the other with higher CD11b (MG^{CD11b}) as shown in **Fig. 2A**. ResiPOx treatment increased the fractions of both BMDM and microglial TAMs, relative to the total cell population (**Fig. 2B**). Within the microglial compartment, the MG^{CD11b} subpopulation increased relative to the total cell population whereas the MG^{CD45} subpopulation showed no significant change (**Fig. 2C**). ResiPOx thus increased TAM populations and showed evidence of change in TAM phenotype.

To determine the role of proliferation in the ResiPOx-induced increase in TAMs, we compared *in vivo* EdU uptake in TAMs of ResiPOx-treated and control *G-Smo* mice. We treated one group of *G-Smo* mice with ResiPOx on P10, 12 and P14, administered EdU IP 24 hours later on P15, and another group of *G-Smo* mice with ResiPOx on P10 and 12, administered EdU IP 24 hours later on P13. Tumors from all mice in both groups were harvested 1 hour after EdU injection and compared to age matched untreated *G-Smo* controls, injected IP with EdU IP at P15 or P13. Harvested tumors were dissociated and subjected to flow cytometry. We

quantified EdU+ cells in the CD45+/CD11b+ TA population. This approach also enabled assessment of ResiPOx effects on MB cell proliferation by quantifying EdU+ cells in the CD45-/CD11b- population, which consisted predominantly of MB cells.

Within the TAM population, ResiPOx induced proliferation in both microglial and BMDM subsets at P13, 24 hours after two doses. In BMDMs, this proliferation persisted at P15, 24 hours after 3 doses of ResiPOx (**Fig 2D**). In contrast, in the larger CD45-/CD11b- population that primarily consisted of MB cells, ResiPOx decreased proliferation (**Fig. 2E**).

As an orthogonal approach to quantify TAM populations, we subjected tumors to IHC for the pan-myeloid marker IB1A1. We treated sets of replicate *G-Smo* mice with a single IP dose of ResiPOx or saline at P10, harvesting replicates at 6h or 24h after administration. Additionally we treated sets of replicate *G-Smo* mice with a 3 dose series of ResiPOx or saline at P10, P12 and P14 and harvested tumors 24h later. We then quantified IBA1+ cells in sections of replicate tumors in each condition. We found that ResiPOx increased IBA1+ fractions as early as 6 hours after the first dose and that IBA1+ fractions continued to increase through P16 (**Fig. 2F**). The initial increase at 6h was too rapid to include the effect of proliferation, and most likely reflected increased migration into tumors, while the continued increase included the contribution from increased TAM proliferation.

ResiPOx induces an acute immune response in MBs and suppresses TAM-to-tumor IGF1 signaling

To identify molecular events induced by ResiPOx, we subjected *G-Smo* MBs from ResiPOx-treated and control mice to transcriptomic analysis. To define acute and chronic transcriptomic changes after ResiPOx without age variation as a potentially confounding variable, we sampled all mice at P14-15. For acute ResiPOx studies (6h and 24h time points), we administered one dose of ResiPOx to 8 P14 *G-Smo* mice, harvested MBs from 4 replicate mice at 6h or 24h later and compared to 4 *G-Smo* controls injected at P14 with POx-vehicle and harvested 24 hours later. For chronic ResiPOx studies (5D time point), we administered ResiPOx to 4 *G-Smo* mice at P10, P12 and P14 and then harvested MBs at P15 and compared to 4 *G-Smo* controls injected with POx-vehicle at P10, P12 and P14 and harvested at P15. We subjected all MB samples to RNA-seq and compared normalized transcript counts in ResiPOx-treated MBs versus the set of corresponding controls to identify differentially expressed genes (DEGs) with adjusted *p* value < 0.05.

ResiPOx induced maximal transcriptomic changes at 6 hours that decreased with time post-administration. We identified 472 ResiPOx-affected DEGs in MBs at the 6h time point, 313 at 24h, and 131 at 5D (**Fig. 3A–C**). The sets of DEGs at each time point were highly similar, with 240 DEGs commonly affected at 6h and 24h after ResiPOx (p=2e-227 by hypergeometric test) and 55 commonly affected at 24h and 5D (p=2.2e-142 by hypergeometric test) (**Supplementary Fig. 2A,B**). GSEA analysis of DEGs at each time point identified common biologic functions indicating pathogen recognition and innate immune activation (**Fig. 3D and Supplementary Fig 2C,D**). The pattern of DEGs in isolated BMDMs treated *in vitro* closely resembled the

ResiPOx-affected DEGs identified by RNA-seq in *G-Smo* tumors (**Supplementary Fig. 2E,F**), suggesting that the transcriptomic effects induced by ResiPOx in tumors originated from TAMs.

We used IHC for individual gene products to validate the ResiPOx-induced transcriptomic changes and to identify cell types driving these changes in gene expression. We selected MARCO for protein expression studies as it was consistently up-regulated in ResiPOx-treated tumors at 6h, 24h and 5D, was undetectably low in control tumors and showed the highest fold change in the RNA-seq data 5D timepoint. While selected on informatic grounds, MARCO was also of interest as a TAM receptor implicated in tumor control in other cancers [47]. In sagittal sections containing tumor and adjacent brain, MARCO was undetectable by IHC in control samples, but robustly detected in cells with myeloid morphology; co-localization with IBA1 confirmed that ResiPOx induced MARCO specifically in myeloid cells, including both TAMs within the MBs and microglia in normal brain regions (**Fig. 3E**). MARCO protein expression began as early as 6 hours after ResiPOx, increased by 24 hours and persisted over 6 days (**Fig. 3F**). These MARCO studies show that TAMs, which comprise a fraction of the tumor population, were the predominant source of a transcript that was strongly detected in ResiPOx-treated MBs in the bulk RNA-seq dataset.

Along with MARCO, which we selected as an example of DEG that increased with ResiPOx, we studied the protein expression of IGF1, as an example DEG that decreased with ResiPOx (adj p=0.0042; log2FC=-0.49; Supplementary Data 1). IGF1 is of particular interest as a previous study showed that SHH MBs depend on TAM-derived, IGF1 for paracrine growth factor signaling [9]. Dual IHC for IGF1 and IBA1 confirmed TAMs as the source of IGF1 in the control *G-Smo* tumors and showed that ResiPOx decreased TAM IGF1 protein expression (**Fig. 3E**). To probe the biologic significance of decreased IGF1, we analyzed phosphorylation of the IGF receptor beta subunit (IGF1Rβ) in whole MB lysates, using immunoblot. In MB samples from P15 *G-Smo* mice treated with of ResiPOx as in the 5D timepoint, the total IGF1Rβ and the fraction of phosphorylated IGF1Rβ were significantly lower compared to controls (**Fig. 3G and Supplementary. Fig. 3A additional loading controls)**, indicating a decrease in IGF1R activation corresponding with the ResiPOx-suppressed IGF1 ligand. These protein studies show that ResiPOx-induced changes in TAM gene expression could be detected as positive (MARCO) and negative (IGF1) signals in bulk transcriptomic assays, and that produced biologically relevant changes in TAM protein expression and TAM-tumor signaling.

ResiPOx induces cytokine secretion and IFNβ signaling

We noted that ResiPOx up-regulated mRNAs coding for cytokines and interferon signaling markers, and that these effects showed variable temporal patterns. We identified 28 cytokines that were up-regulated at 6h and 15 cytokines up-regulated at 24h, of which 13 cytokines were up-regulated at both 6 and 24 hours**,** and specific cytokines, including *Ccl5*, *Ccl12* and *Ccl19* remained elevated at 24 hours after 3 doses (**Fig 4A**). Similarly, ResiPOx induced multiple interferon response genes (**Fig. 4A**).

Protein studies using cytokine arrays and immunoblot provided orthogonal validation of the RNA-seq data. We compared MB lysates from P10 *G-Smo* mice treated with ResiPOx and harvested at 6h or 24h and P15 *G-*

Smo mice treated with ResiPOx at P10, P12 P14 versus a control set of P10 *G-Smo* control mice treated with POx vehicle and harvested at 24 hours. A cytokine array that included proteins corresponding to 22 of 28 cytokine DEGs, detected IL12p40 and CCL12 (MCP5) as induced by ResiPOx with maximal signal at 6 hours and waning over time, matching the temporal pattern of *Il12b* and *Ccl12* mRNA (**Fig. 4B, Supplementary Fig. 3B**). On a different array that included an overlapping set of 14 of the 28 cytokine DEGs, CCL9 and CXCL10 were similarly increased (**Supplementary. Fig. 3C**). These protein data were consistent with prior studies showing resiquimod up-regulated IL12p40 [38, 48] and confirm that ResiPOx induced dynamic expression of specific cytokines detected by RNA-seq.

Protein immunoblot studies confirmed that ResiPox up-regulated expression of multiple interferon signaling proteins, including STAT1, IFIT3 and IRF9, and also showed that ResiPOx induced STAT1 phosphorylation (**Fig. 4C, Supplementary. Fig. 3D additional loading controls**). Interferon signaling peaked at 24h after the first ResiPOx dose and remained elevated at 5D compared to untreated controls, at lower levels compared to peak IFNb ligand was also elevated at 5D post ResiPOx, varying inversely with anti-inflammatory TAM marker IBA1, linking persistent interferon signaling to reduced anti-inflammatory TAM phenotype (**Fig. 4D, Supplementary. Fig. 3A additional loading controls**). Together, these PD studies demonstrate ResiPOx responses with different temporal patterns, including a rapid, transient induction of specific cytokines, a slower activation of interferon signaling and a sustained increase in MARCO expression.

TAMs are the predominant driver of the transcriptomic response to ResiPOx

To resolve ResiPOx-induced changes with greater cellular resolution, we subjected *G-Smo* mice treated with ResiPOx to scRNA-seq. We compared MBs from 4 *G-Smo* mice treated with ResiPOx at P10, P12 and P14 versus MBs from 4 age-matched, saline-injected *G-Smo* controls. All tumors were dissociated and processed for split-seq-based scRNA-seq using the Evercode Whole Transcriptome (WT) kit v1 (Parse Biotechnologies). Control replicates showed fewer cells sequenced, compared to ResiPOx-treated replicates and to facilitate comparisons we randomly down-sampled the cells from each replicate to match the numbers of cells in the replicate with the smallest number of cells, in keeping with best practices [45]. After down-sampling, 3865 cells in each replicate passed filtering and were analyzed.

After filtering to remove doublets, unsupervised clustering defined 18 clusters, including 9 discrete clusters and 9 clusters that mapped to a large, multi-cluster group (**Fig. 5A, Supplementary Fig. 4A**). Cluster-specific DEGs and representative markers identified the discrete clusters as astrocytes, oligodendrocytes, myeloid cells, gabaergic neurons and other non-tumor cells expected in the CNS (**Fig. 5A, Table 1**), and identified the multicluster group as MB cells in a range of states that paralleled the developmental trajectory of cerebellar granule neuron progenitors (**Table 1**), as in our prior studies in this model [42, 45, 46, 49, 50]. We used the scRNA-seq data to identify the cell types that expressed the genes that were most strongly induced by ResiPOx in the bulk RNA-seq data (adjusted p< 0.05 and log2 FC > 2). There were 43 genes meeting these criteria in the bulk RNA-seq dataset, and 36 of these 43 genes were detected in the scRNA-seq dataset. The scRNA-seq data

showed that all of these 36 genes that were identified as ResiPOx-induced DEGs in the bulk RNA-seq data were specifically expressed by TME cells. While 30/36 ResiPOx-induced DEGs were specific to the myeloid cluster (**Fig. 5B**), 6/36 DEGs, including MHC and interferon response genes were also expressed by endothelial and choroid plexus cells. TAMs were thus the major drivers of the transcriptomic response to ResiPOx, and vascular-facing cells showed an interferon-driven response. TAMs were the only cells to express TLR7, the resiquimod target (**Fig. 5C**), supporting the interpretation that direct drug effects operated through TAMs, and effects on other cell types proceeded indirectly through changes in TAM-mediated signaling.

ResiPOx alters the proportions of TAMs with specific phenotypes

To investigate the effect of ResiPOx on TAM phenotypes, we isolated the myeloid cluster *in silico* and resolved discrete elements within the TAM population by repeating the PCA and Louvain clustering on the isolated TAMs. This second round of PCA analysis generated a UMAP with three distinct regions, while cluster analysis identified 3 clusters that predominantly correlated with UMAP localization (**Fig. 5D**). Supervised analysis using genes known to mark brain myeloid subsets showed that 2 of the clusters expressed the microglial marker *Cx3cr1* [51-53] (**Fig. 5E**), *P2ry12* and *Fcrls* (**Supplementary Fig. 4B,C**) [13]. One of the two microglial clusters expressed the anti-inflammatory marker *Mrc1+* [54] (MG^{Mrc1+}; **Fig. 5E**). We further characterized the clusters by determining cluster-specific DEGs (Table 2, Supplementary Data X), which showed that the *Mrc1-* microglial cluster uniquely expressed the ramified microglial marker *Sparc* [55] (MG^R) and MG^{Mrc1+} expressed anti-inflammatory marker *Cd36* [56], supporting an anti-inflammatory role. The *Cx3cr1-* cluster expressed the BMDM markers *Ccr2* [53] (**Fig. 5F**) and *Cytip* (**Supplementary Fig. 4D**) [13]. The MG^{Mrc1+} cluster was the predominant source of *Igf1* (**Fig. 5F**).

To determine the interaction between ResiPOx and TAM heterogeneity, we determined how each TAM subset contributed to the overall transcriptomic response to ResiPOx, and how ResiPOx affected the relative proportions of TAM subclusters. The MG^R, and MG^{Mrc1+} and BMDM subclusters each expressed different subset of the ResiPOx-induced DEGs identified in the bulk RNA-seq (**Fig. 5G**). ResiPOx shifted TAM populations, enriching the MG^R (p=6.199 x 10^{-6} Chi square test) at the same time as increasing the overall TAM population relative to all other TME cells (**Fig. 5H**). Across the overall TAM population, *Mrc1* and *Igf1* were both decreased by ResiPOx (**Supplementary Fig. 4E,F**). Consistent with this change in TAM phenotypes, IHC studies of MRC1 expression, co-localized with pan-myeloid marker IBA1, showed that ResiPOx decreased MRC1+ TAMs (**Fig. 5I**). Together, these scRNA-seq data show that ResiPOx most strongly altered the gene expression in TAMs, induced different, specific patterns of DEGs from different types of TAMs, and shifted TAM phenotypes, enriching MG^R TAMs and depleting the *Mrc1+* TAMs that secrete IGF1.

scRNA-seq identifies ResiPOx effects on tumor cell gene expression obscured in bulk RNA-seq

To determine the effect of ResiPOx on MB cells, we isolated the combined set of 9 clusters of MB cells *in silico*. We note that ResiPOx increased the population at the differentiating/differentiated pole in the UMAP (**Fig. 5J**). To facilitate statistical comparisons, we grouped the 9 tumor clusters into three groups according to

their position along the developmental trajectory from proliferative, undifferentiated cells to early and late differentiating cells, based on gene expression patterns. Representative markers confirmed the phenotypic characterization of each of these groups, including proliferation markers *Cyclins A1, B1, D1 and E1*, the SHH-driven transcription factor *Gli2*, neural developmental genes *Cntn2*, *Neurod1*, *Neurod2* and the glutamatergic neuron marker *Slc17a7* (aka *Vglut1*; **Fig. 5K**). ResiPOx-treated MBs showed significantly reduced proliferative MB populations and significantly increased early and late differentiating MB populations (**Fig. 5L**), confirming that ResiPOx induced cell cycle exit and differentiation.

We compared gene expression of the combined set of 9 MB clusters in ResiPOx-treated versus control tumors (Supp. Data X). We noted that ResiPOx increased expression of genes associated with cerebellar granule neuron differentiation, including *Adamts18* [57] and *Unc5* [58]. Consistent with inducing a shift in the MB populations toward differentiation, ResiPOx decreased expression of the proliferation marker Mki67 (**Fig. 5M, Supplementary Fig. 4G**). GO Annotation analysis of the set of 253 DEGs with adjusted p<0.05 and log2 FC>1 showed ResiPOx treatment enriched DEGs associated with oxidative phosphorylation (p=8.83E-08) including multiple subunits of Cytochrome Oxidase and ATP Synthase. In addition, we noted that ResiPOx induced expression of the immunomodulatory gene *Il31ra*, and the phagocytosis inhibitory signal Cd47 (**Fig. 5M**). Analysis of scRNA-seq thus identified ResiPOx-induced changes in MB cells that were obscured by the signal from TME cells in the bulk RNA-seq.

ResiPOx increased phagocytosis of tumor cells by TAMs

To investigate ResiPOx-induced changes in TAM function, we analyzed effects of ResiPOx on genes related to phagocytosis and antigen presentation. ResiPOx induced genes relevant to both innate and adaptive immune responses including non-opsonic receptors, opsonic Fc receptors, complement cascade components, MHC antigen presentation, and extrinsic apoptosis-inducing ligands and receptors (**Fig. 6A**). Non-opsonic receptors mediate innate recognition of PAMPs and DAMPs. Additionally, opsonic Fc receptors, antigen presentation mechanisms and complement components were also increased, suggesting effects that may link TAMs to humoral responses while upregulation of extrinsic apoptosis pathway components suggests potential activation of TRAIL- and FAS-mediated apoptosis, which may increase phagocytosis of newly killed tumor cells. GSEA confirmed enrichment of transcriptomic patterns consistent with phagocytosis and antigen presentation (**Fig. 6B**).

To test directly for an effect on phagocytosis, we compared uptake of MB cells by TAM in tumors from mice treated with ResiPOx versus controls. For this purpose, we bred our tumor models with *Ai6* mice, which feature Cre conditional expression of ZsGreen. We bred *Ai6* mice with either *Math1-Cre* mice that express Cre in the cerebellar progenitors that are cells of origin for SHH MB [59], or with *Gfap-Cre* mice that express Cre in stem cells that give rise to neurons and glia throughout the brain [60]. We bred the resulting *Math1-Cre/Ai6* and *Gfap-Cre/Ai6* mice with *SmoM2* mice, to generate pups that developed ZsGreen-expressing MBs. All mice with *Math1-Cre/Ai6/SmoM2* (*M-Smo^Ai6^*) and *Gfap-Cre/Ai6/SmoM2* (*G-Smo^Ai6^*) genotypes developed MBs as expected. *M-Smo^Ai6^* mice developed MBs with ZsGreen in specifically in tumor cells, while *G-Smo^Ai6^* mice expressed ZsGreen in both MBs and in glial cells throughout the brain. In both genotypes, myeloid cells were

outside the lineages that inherited Cre-driven recombination, so that TAMs would only be expected to show ZsGreen fluorescence if they phagocytosed ZsGreen+ cells. IHC for IBA1 and ZsGreen showed ZsGreen+ vacuoles in IBA1+ TAMs (**Fig. 6C**), confirming detection of TAM phagocytosis of MB cells.

To determine the effect of ResiPOx on TAM phagocytosis of MB cells, we treated M-Smo^{Ai6} and G-Smo^{Ai6} mice with ResiPOx or saline at P10,P12, P14 and then harvested mice at P15, dissociated MBs and used flow cytometry to quantify ZsGreen+ cells in the $CD45^{int}/CD11B^{int}$ microglia and $CD45^{high}/CD11B^{high}$ BMDMs (**Fig. 6D**). Representative flow dot plots indicate gating strategy for the myeloid compartment and distribution of ZsGreen+ cells (**Fig. 6 D,E**). Comparison of replicate mice in each genotype and condition showed that ResiPOx increased both total fractions of microglial and BMDM TAMs, and the ZsGreen+ fractions of microglial and BMDM TAMs, relative to the total cell population (**Fig. 6F,G**). ResiPOx thus increased TAM phagocytosis of MB cells, proportionally with the increase in TAM populations.

ResiPOx shows similar effects in mice with DMG

Considering the common importance of TAMs in MB and in malignant gliomas [5, 7, 17, 53, 61, 62], we analyzed ResiPOx efficacy in an immunocompetent DMG model [63]. We generated mice with DMGs by transducing *Pdgfb*, *Cre*, *H3K27M-Gfp* fusion and *Luciferase* into P2 *Nestin-tva/Tp53^{f/f}* mice by intracranial injection of RCAS viruses. We then monitored bioluminescence imaging (BLI) twice weekly to determine the onset of DMG growth. At the first BLI measurement detecting tumor, we initiated 3 IP doses of ResiPOx or POx vehicle, 48 hours apart, as in our MB studies. The glioma model was more slowly progressive than the *G-Smo* MB model, and to address the potential for subjectivity in the identification of glioma symptoms, we randomized mice to ResiPOx or vehicle with the animal handler blinded as to which mice received which treatment. We then continued twice weekly BLI and monitored mice for the onset of symptoms indicating tumor progression, which we defined as the humane endpoint. After 12 mice per group died of progressive tumor, we unblinded the study and analyzed the results.

ResiPOx-treated mice reached the humane endpoint after significantly longer PFS (**Fig. 7A**, $p = 0.0141$). Analysis of BLI measurements over time showed significantly delayed DMG progression in ResiPOx-treated (**Fig. 7B**, $p < 0.02$, determined by a generalized linear model). Individual ResiPOx-treated mice with RCAS-DMG showed transient tumor regression (**Fig. 7C**) that was never seen in controls. Together these data demonstrate ResiPOx efficacy in DMG, comparable to SHH MB.

As in SHH MB, ResiPOx-treated mice with RCAS DMG showed increased TAM populations and increased MARCO+ TAMs (**Fig. 7D**). We used transcriptomic analysis to compare ResiPOx-induced PD in DMG versus SHH MB. For this study, we generated 4 replicate ResiPOx-treated mice and 3 replicate POx-vehicle controls with RCAS-DMG. We harvested brains from all mice at 24 hours after IP injection and dissected out tumors, guided by GFP expression and processed tumor samples for RNA-seq analysis. We compared tumors from ResiPOx-treated versus control DMGs and identified 193 DEGs (**Supp. Fig. 7A**). These ResiPOx-induced DEGs in DMGs showed high concordance with the DEGs up-regulated in MB 24 hours after ResiPOx

(p=1.7903 x10^{-43} by hypergeometric test), with common induction of *Marco*, *Cd72*, *Ccl5* and *Irf7* (**Fig. 7E**). Together, these results indicate that ResiPOx induces a shared immune-activating program across brain tumors with distinctly different genetics and ontogeny, supporting a common mechanism of action.

Systemic ResiPOx is well tolerated and shows similar PD in the brain in Rhesus macaques

The efficacy of ResiPOx in multiple mouse brain tumor models supports the clinical potential of ResiPOx for adjuvant brain tumor therapy. To determine ResiPOx safety and PD in the brain in a primate species, we therefore analyzed ResiPOx in adult rhesus macaques at the Emory National Primate Research Center that had pre-existing clinical conditions that warranted a plan for euthanasia, but were able to contribute to this short study prior to euthanasia.

Considering the CNS penetration of ResiPOx in mice, we confirmed safety in non-human primates through dose escalation studies. Using doses of 0.2 mg/kg, 0.4 mg/kg and 0.8 mg/kg in individual animals, we administered 2 IV ResiPOx doses 24 hours apart, monitored animal vital signs and compared complete blood cell counts prior to ResiPOx and 2 hours after the first and second dose. In all 3 animals, all doses were well-tolerated, produced mild, transient temperature elevation. Noting no acute toxicity, we then treated 4 replicate animals, including 2 males and 2 females with 0.8 mg/kg ResiPOx using 2 doses at 24 hours and 2 hours prior to euthanasia. We compared 4 age and sex-matched controls treated with saline. We sampled blood at baseline, and at several time points after each injection. To analyze CNS effects, we humanely euthanized animals, transcardially perfused with saline and then rapidly harvested brain tissue from animals at 24h after the first ResiPOx or saline injection, which was also 2h after the second injection. Brain samples from hippocampus, parietal cortex and corpus callosum were dissected from similar regions in each animal. We isolated RNA from each brain sample and pooled RNA from each region to generate pooled-region samples from each individual animal. We then sequenced and analyzed the RNA as in our mouse studies, mapping to the Rhesus macaque genome.

RNA-seq studies showed that ResiPOx induced significant PD effects in the brain. We identified 1344 DEGs in ResiPOx-treated brain samples (Supp data. 7X). These DEGs closely matched the ResiPOx induced DEGs in *G-Smo* MBs (p=3.4056 x10^{-121} by hypergeometric test) with 216 DEGs commonly induced (**Fig. 7F**). Analysis of blood samples showed that ResiPOx induced changes in circulating myeloid cells, with decreased circulating PBMCs and increased fractions of band neutrophils (**Fig. 7G.H**), consistent with systemic immune response and increased passage of myeloid cells from blood to tissues. These data show that ResiPOx was tolerable in non-human primates, crossed the BBB and induced a set of genes that was common across divergent species, indicating engagement of common mechanisms and supporting the potential for similar PD effects in patients.

Discussion

These studies shows that ResiPOx delivers resiquimod across the BBB, where it acts on TAMs in MB and DMG, enriching pro-inflammatory TAM phenotypes and inducing cytokine secretion, interferon signaling and phagocytosis of tumor cells, ultimately slowing tumor progression. Immunotherapy for MB and DMG face challenges posed by the BBB, by immunosuppressive myeloid populations that also secrete tumor-supportive growth factors, and by the paucity of immune effectors such as T cells and NK cells [20, 41, 64]. Our data shows ResiPOx addresses these specific challenges.

ResiPOx showed ResiPOx showed superior PK and efficacy compared with free resiquimod in *G-Smo* MBs, a genetically-engineered mouse model known to have intact BBB [40]. ResiPOx induced consistent PD effects within the CNS. In MBs these PD effects included acute induction of cytokine secretion, sustained activation of interferon signaling to vascular cells, increased TAM proliferation and altered TAM phenotypes. ResiPOx induced similar TAM phenotypic changes in DMG, accompanied by comparable anti-tumor efficacy. In rhesus macaques, IV ResiPOx induced transcriptomic changes in normal brain tissue that closely resembled those following IP ResiPOx in mouse brain tumors, indicating consistent BBB penetrance across species and conserved patterns of drug response in normal microglia and in TAMs.

In repolarizing TAMs in *G-Smo* MBs, ResiPOx interrupted TAM-tumor paracrine growth support mediated by IGF1 signaling. Our scRNA-seq data localized tumor-supportive IGF1 signaling specifically to the MRC1+ fraction of TAMs, consistent with reports localizing IGF1 to MRC1+ myeloid cells outside of the CNS [65]. ResiPOx decreased MRC1+ TAMs and increased pro-inflammatory TAM populations expressing IL12-p40, matching resiquimod-induced changes in myeloid cell polarization previously described in macrophages *in vitro* [66]. Concurrent with depleting MRC1+ TAMs in *G-Smo* MBs, ResiPOx decreased TAM IGF1 expression, producing a significant decrease in IGF1R phosphorylation in MBs. As microglial TAMs, unlike BMDMs do not circulate outside the CNS, these changes in microglial TAMs indicate local PD effects, in which ResiPOx acted within the CNS compartment to shift TAMs from immunosuppressive phenotypes and to interrupt TAM support for tumor growth.

Specific ResiPOx effects on BMDM TAMs indicate systemic effects that were concurrent with local effects in *G-Smo* MBs. The scRNA-seq analysis of *G-Smo* MBs identified discrete patterns of ResiPOx response in BMDM TAMs that differed significantly from microglial TAMs. Moreover, ResiPOx increased proliferation and phagocytosis by BMDM TAMs as well as by microglial TAMs. In our Rhesus macaque studies, which included peripheral blood studies at sequential time points, ResiPOx altered circulating immune populations, markedly reducing PBMCs and transiently increasing band neutrophils. The reduced circulating PBMCs in the Rhesus macaques together with the increased BMDMs in *G-Smo* MBs, suggest migration of circulating immune cells into peripheral organs, as well have BMDM proliferation, may have increased BMDM populations, consistent with prior reports of TLR7-activated myeloid cell migration [67, 68]. These peripheral effects of ResiPOx on circulating immune cells are distinct from, and co-incident with the local ResiPOx effects on microglial TAMs.

A prior study using β-cyclodextrin nanoparticle-formulated resiquimod was the first to show efficacy in glioma [38]. In this study, nanoparticle resiquimod administered peripherally to mice with syngeneic, implanted gliomas striking blocked tumor growth through effects on circulating myeloid cells that trafficked to the tumors [38]. In two more recent studies, local administration of matrix-bound resiquimod showed locally-mediated brain tumor suppression [39, 69]. These studies show the potential for local resiquimod to stimulate anti-tumor immune mechanisms. However, these papers modeled gliomas using implanted cell lines, and while these cell lines were syngenic, and orthotopic, they did not derive from the host animals. The T cell responses seen in the studies [39, 69] suggest stronger immune responses than the endogenous tumor models that we studied, which more closely model the immunologically cold brain tumors of patients, where T cells are typically rare.

Our data show ResiPOx efficacy in endogenous brain tumors that form in immunocompetent mouse hosts in our MB and DMG models. In both of these endogenous brain tumor models models, ResiPOx acted both peripherally and locally. ResiPOx achieved local delivery of resiquimod throughout the brain and without surgery, which may be advantageous for patients with tumors not amenable to resection, including recurrent MB and DMG and for patients with wider dissemination of tumor than the volume of drug distribution that can be achieved with a wafer approach[70]. The simultaneous engagement of both local and systemic immune function by ResiPOx may contribute to the anti-tumor effect.

While ResiPOx did not detectably increase T cells or NK cells in mouse brain tumors, in a prior study, resiquimod loaded in β-cyclodextrin nanoparticles demonstrated anti-glioma efficacy in mice with global T cell depletion [38]. In that study, resiquimod increased myeloid cell ROS, suggesting the potential for direct cell killing as an anti-tumor mechanism. The ResiPOx-induced increase in TAM phagocytosis of MB cells may also be a direct anti-tumor mechanism, or may reflect increased tumor cell killing rapidly followed by phagocytosis. Resiquimod therapy thus addresses the lack of immune effectors in brain tumors by recruiting TAMs to this function.

In summary, we have developed a nanoparticle platform with distinct advantages for delivery of resiquimod, allowing for systemic administration with safety and tolerability in both mouse models and non-human primates and enabling sustained activation of myeloid cells both systemically and within the TME. Our data are consistent with recent studies showing anti-brain tumor effects of resiquimod delivered systemically in β-cyclodextrin nanoparticles [38] or locally in solid matrices [39, 69]. We now show the efficacy of our agent genetically-engineered primary models of pediatric brain tumors MB and DMG, where endogenous tumors have less potential for tumor-host mismatch. In these models, which are rapidly progressive and highly refractory to therapy, brain tumors arise through a process of tumorigenesis *in situ* and recapitulate intratumoral heterogeneity, TME and treatment-refractory nature of recurrent MB [46] and DMG [63]. Through integrated transcriptomic, protein and cellular analyses in our models, we localized specific features of the resiquimod-induced response to different microglial and BMDM subsets of TAMs and define responses of

tumor cells to the disruption caused by ResiPOx. In these models, ResiPOx combined effectively with RT in MB, and ResiPOx produced similar effects in the brain in non-human primates, without apparent toxicity. ResiPOx delivery may be ideal for pediatric brain tumor patients not undergoing resection, including DMGs, which are not resectable, and recurrent MBs, which are typically multifocal and not resected. We conclude that ResiPOx is a promising approach for safe, systemic immunotherapy for pediatric brain tumors that synergizes with RT that is integral to the current standard of treatment.

METHODS

Materials

We purchased 2-methyl-2-oxazoline (MeOx, 98%), 2-ethyl-2-oxazoline (EtOx, ≥99%), 2-n-butyl-2-oxazoline (BuOx, 99.3%), anhydrous acetonitrile (ACN, for synthesis, 99.8%), and piperidine (≥99.5%) from Millipore Sigma (Burlington, MA, USA). Methyl trifluoromethanesulfonate (MeOTf) (>98.0%) was obtained from Tokyo Chemical Industry (Tokyo, Japan). We acquired resiquimod (Res, >98%) and 3H-Resiquimod (≥96%, 1 mCi mL-1 in EtOH) from MedKoo Biosciences (Durham, NC, USA) and Moravek (Brea, CA, USA), respectively. We procured Ultima gold liquid scintillation cocktail and soluene-350 from Revvity (Waltham, MA, USA). Spectra/Por 3 dialysis membrane tubing (Molecular Weight Cut-Off 3.5 kDa) was purchased from Repligen (Waltham, MA, USA). All antibodies used for IHC, western blots or flow cytometry studies are described in Supplementary Data XXX Materials Table.

Ethical compliance

This research complies with all relevant ethical regulations. All mouse experiments at UNC were performed with the approval of the UNC IACUC under protocols 19-098 and 21-011. All mouse experiments at Emory University were performed with the approval of the Emory IACUC under protocol 202200148. These protocols all require that mice with brain tumors must be euthanized at the first emergence of symptoms, which is considered the maximum allowable tumor burden. This maximum tumor burden was not exceeded

Synthesis of Poly(2-oxazoline) block copolymer

Amphiphilic poly(2-oxazoline) block copolymer P(MeOx)$_{35}$-b-P(BuOx)$_{24}$-b-P(MeOx)$_{35}$ (POx) was synthesized by the living cationic ring-opening polymerization as previously described [71, 72]. The polymer synthesis was initiated by mixing 1 equivalent (eq) of MeOTf with 35 eq of MeOx in ACN. The polymerization proceeded at 60 °C for 2 days and the degree of polymerization was monitored by nuclear magnetic resonance spectroscopy (NMR). The second and third blocks of the polymer sequentially propagated with 24 eq of BuOx and 35 eq of MeOx in the stated order. To terminate polymerization, we added excess piperidine (> 3 eq) into the reactant and carried out the reaction at 40 °C for 4 hr. Upon completion, the polymer was dissolved in acetone and filtered through a nylon membrane filter (pore size 0.2 μm). The filtrate was dialyzed against deionized water (molecular weight cut-off 3.4 kDa) for a week with semi-daily replacement of water. The dialyzed polymer was

lyophilized and stored at 4 °C with desiccant until use. The molecular weight of the polymer (*ca.* 9.2 kDa) was determined by NMR.

Preparation of ResiPOx

The formulation was prepared by the thin-film method [73]. Resiquimod and POx were dissolved in 200 proof EtOH to a concentration of 10 mg mL^{-1} and subsequently sonicated at room temperature for 5 min. For ResiPOx preparation, 100 µL of POx solution was mixed with 40 µL of Res solution in a 1.5 mL centrifugation tube (10:4 w/w). The mixture was subjected to drying at 60 °C under nitrogen flow until the solvent completely evaporated. To induce ResiPOx formation, we added 200 µL of saline to the dried thin film. All formulations were freshly prepared from dried thin films immediately prior to administration.

Determination of ResiPOx pharmacokinetics

To evaluate the PK profile of the formulation in model animals, tritium-labeled ResiPOx (^{3}H-ResiPOx) was prepared *via* the thin-film method. (*vide supra*) An EtOH solution of unlabeled Res was mixed with the tritium-labeled Res (^{3}H-Res) to yield a total Res concentration of 10 mg mL^{-1} and a radioactivity of 25 µCi mL^{-1} in the ^{3}H-ResiPOx. Prior to organ and blood collection at each time point, 100 µL of 1% heparin diluted in saline was subcutaneously administered. Whole blood was collected by cardiac puncture and was centrifuged at 2000 × g for 10 min at 4 °C to separate plasma from whole blood. The plasma fraction from the supernatant was mixed with 1 mL of the 1:1 (v/v) mixture of Soluen-350 and isopropyl alcohol. Harvested organs were completely dissolved in 1 mL of Solene-350 for 3 days. Immediately prior to scintillation counting, 15 mL of Ultima gold liquid scintillation cocktail was added onto each specimen, followed by through vortexting. The level of scintillation from each specimen was determined by Tri-Carb 3110 TR liquid scintillation analyzer (PerkinElmer, Waltham, MA, USA).

Immunoblotting and Chemokine Array

Whole tumor tissue was isolated into DPBS containing 6g/L glucose and resuspended in 500-1000uL of RIPA + 1x protease/phosphatase inhibitors (RIPA cat CST #9806S, Ptase inhibitors cat CST #5872). Samples were sonicated thoroughly prior to centrifugation and protein quantification using BCA assay. 15-30ug of protein was loaded into each lane of a Tris-Glycine gel and transferred onto 0.45um PVDF immobilon membrane. Membranes were blocked with 5% milk in TBST and probed with respective antibodies in TBST with 5% BSA. For chemokine array (Ray Biotech Cat#AAM-CYT-3-2), 500ug of protein lysate was diluted in array buffer and supplemented with additional protease/phosphatase inhibitors. Lysate was incubated overnight on membranes and washed according to manufacturer's instructions. Secondary antibodies were then applied overnight followed by wash and development according to manufacturer's instructions.

Flow Cytometry

Tumors were first triturated into a single cell suspension using Worthington papain kit (cat#) and subject to RBC lysis. For surface staining, samples were incubated with zombie viability die at 1:500 for 25min, blocked with mouse Fc block for 10 min in FACS buffer, and stained with antibodies to CD45 or CD11b. For surface and intracellular staining, surface proteins were stained as described and subsequently fixed and permed using eBiosciences fix/perm kit (cat# 88-8824-00) according to manufacturer's instructions and finally intracellularly stained. For cell cycle analysis cells were stained with Ki67 and 7AAD prior to EdU azide reaction using thermo EdU plus flow kit (cat#). All samples were run on Cytek Aurora equipped with 5 laser lines.

Immunohistochemistry

Brain samples containing MB or DMG were fixed in 4% formaldehyde for 24-48 hours, embedded in paraffin and sections were cut at 4um for MB tissue and 5um for DIPG tissue. MB sections were stained imaged and quantified as previously described [45, 74]. For DMG, samples were incubated at 60C for 1h followed by de-paraffinization and rehydration and antigen retrieval in sodium citrate buffer in an instant pot at high pressure for 15 minutes. Slide were then photobleached to quench RBCs and background using 3% H2O2 and 20mM NaOH in H2O under activated fluorescent lamps 2x 30min with quench solution change. Samples were blocked with serum and incubated with antibody overnight at 4C followed by HRP-secondary antibody incubation and tyramide amplification reaction. For multiplexing, samples were re-boiled in sodium citrate buffer prior to re-blocking and staining. Slide images were acquired using an Akoya slide scanner with HALO imaging software used for quantification.

Bulk RNA Sequencing

Tumor samples were harvested in Trizol and flash frozen. After thaw, RNA was purified from Trizol according to manufacturers instructions and treated with DNase prior to paired-end sequencing with 50 million reads. Following sequencing, samples were checked using FastQC, trimmed using TrimGalore, and aligned using STAR genome aligner. Gene reads were normalized and differential genes analysis performed using DEseq in R. For gene set enrichment analysis of DEGs, the fgsea package was used (Korotkevich G et al. 2019). Correlation analysis, including DEGs and GSEA NES values, was performed using Pearson correlation with gene overlap assessed using a hypergeometric test. Principal component analysis of the DMG dataset revealed two samples affected by harvesting-related artifacts. These outliers were excluded prior to subsequent analyses.

scRNA sequencing

Split-seq based barcoding and library construction was performed using the Parse Fixation and barcoding kits (Cell Fixation Kit v1, Evercode WT (100K) v1) and sequenced on an Illumina Novaseq6000, using an S2 flow cell, as previously described [45]. The scRNA-seq data were analyzed using Parse software to identify cells by bar code, and to map transcripts to the mouse transcriptome. Data were then analyzed using our previously reported pipeline [46, 50].

MSI

MSI was accomplished using IR-MALDESI, as previously described [43]. In brief, an IR laser was applied to 10 µm frozen sections of mouse brain with tumor, to desorb adjacent 50µm diameter sampling locations, and an electrospray (50/50 mixture of methanol/water (v/v) with 0.2% formic acid) ionized the desorbed neutral molecules. The resulting ions were sampled into a high resolving power Thermo Fisher Scientific Q Exactive Plus (Bremen, Germany) mass spectrometer for synchronized analysis. The mass spectrometer was operated in positive ion mode from m/z 100 to 400, with resolving power of 140,000FWHM at m/z 200. Resiquimod, creatinine and NAA were identified by m/z through reference to the Human Metabolome Database (hmdb.ca, Canada).

Acknowledgements

Research reported in this publication was supported in part by the national Institutes of Health (NIH). MS was supported by the National Institute for Neurologic Disease and Stroke (NINDS) (R01NS125073). TRG, AT and LM were supported by the NINDS (R01NS088219, R01NS102627, R01NS106227). AVK was supported by the National Cancer Institute (NCI) (CA264488). Research reported in this publication was supported in part by the Cancer Tissue and Pathology Shared Resource of Winship Cancer Institute of Emory University and NIH/NCI under award number P30CA138292. This research was also supported in part by the National Institutes of Health (NIH) Office of the Director (P51OD011132). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Histopathology/Digital Pathology was also performed by the Pathology Services Core at the University of North Carolina-Chapel Hill. We thank Nicholas Pankow and Edison Floyd in the Pathology Services Core for expert technical assistance with Histopathology and Digital Pathology. The PSC is supported in part by an NCI Center Core Support Grant (P30CA016086) and NC Biotech Impact Innovation Grant (2024-IIG-0015). We thank Kaitlyn Love, Alex van Schoor, Rebecca Richardson and veterinarian, Dr. Fawn Stroud, at the Emory National Primate Research Center for their assistance with data collection on rhesus macaques.

Author Contributions

Competing Interests

A.V.K. declares a conflict of interest due to issued and pending patents related to this work and his role as co-founder, shareholder, and licensor of technology to DelAQUA Pharmaceuticals, which develops poly(2-oxazoline)–based drug formulations. M.S. declares a conflict of interest as the spouse of A.V.K. The other authors report no competing interests.

Antibody	Species	Dilution
CD11b	Rat, APC or AF647 BD cat#553312	Flow: 1:75
CD45	Rat, BUV805 BD cat#748370	Flow: 1:125
IBA1	Rabbit, CST cat#17198	IHC: 1:750
CD206	Rabbit, ProteinTech cat#18704-1-AP	WB: 1:1000, IHC: 1:750
zsGreen	Mouse, Takara Bio cat #632598	IHC: 1:500
pSTAT1 S727	Rabbit, CST cat#9167	1:1000
Total STAT1	Rabbit, CST cat#9172	1:1000
IFIT3	Rabbit, ProteinTech cat#15201-1-AP	1:1000
IRF9	Rabbit, ProteinTech cat#14167-1-AP	1:1000
MARCO	Rabbit, Abcam cat#ab259264	WB: 1:1000, IHC: 1:750
β-Tubulin	Mouse, CST cat#2146	1:1000
pIGF1Rβ Y1316	Rabbit, CST cat#28897	1:1000
pIGF1Rβ Y1135	Rabbit, CST cat#3918	1:1000
IGF1Rβ	Rabbit, CST cat#9750	1:1000
IFNβ	Rabbit, ProteinTech cat#27506-1-AP	1:1000

Main Tables

Table 1. Cell type designations for each cluster in the initial scRNA-seq analysis including both tumor cells and stromal cells. Cluster numbers were assigned in size order, with cluster 0 the most populous.

Cluster	designation	best markers
0	early differentiating MB cells	*Obscn, Sema7a*
1	proliferating MB cells	*Rrn2, E2f8*
2	mitotic MB cells	*Pif1, Ccnb1, Kif11*
3	differentiating MB cells	*L1cam, Grin2b*
4	neural progenitor-like, non-proliferative MB cells	*Angpt1, Sox6,Ttc23*
5	early differentiating MB cells	*Obscn*
6	differentiating, cholinergic MB cells	*Chrna3*
7	differentiating MB cells	*Gabra2, Grin1, Unc5b*
8	late differentiating MB cells	*Grin2c, Gabra6, Gabra1*
9	endothelial cells	*Cldn5, Ctla2*
10	oligodendrocytes	*Sox10*

11	myeloid cells	*Ptprc, Tlr7*
12	myelinating oligodendrocytes	*Mog, Mag, Sox10*
13	astrocytes	*Aqp4, Aldh1l1*
14	fibroblasts	*Dcn, Cl1a1*
15	interneurons	*Pax2, Gad2, Slc6a5*
16	pericytes	*Cysltr2, Notch3*
17	choroid plexus	*Folr1*

Table 2. Cell type designations for the myeloid subclustering of the scRNA-seq data. Cluster numbers were assigned in size order, with cluster 0 the most populous.

Cluster	designation	best markers
0	MG- ramified	*Sparc, Sall1*
1	MG- MRC1+	*Mrc1, Cd36*
2	BMDMs	*Cytip*

References

1.	Rusin D, Vahl Becirovic L, Lyszczarz G, Krueger M, Benmamar-Badel A, Vad Mathiesen C, Sigurethardottir Schioth E, Lykke Lambertsen K, Wlodarczyk A. Microglia-Derived Insulin-like Growth Factor 1 Is Critical for Neurodevelopment. Cells. 2024;13(2). Epub 20240118. doi: 10.3390/cells13020184. PubMed PMID: 38247874; PMCID: PMC10813844.

2.	Yu D, Jain S, Wangzhou A, Zhu B, Shao W, Coley-O'Rourke EJ, De Florencio S, Kim J, Choi JJ, Paredes MF, Nowakowski TJ, Huang EJ, Piao X. Microglia regulate GABAergic neurogenesis in prenatal human brain through IGF1. Nature. 2025;646(8085):676-86. Epub 20250806. doi: 10.1038/s41586-025-09362-8. PubMed PMID: 40770097; PMCID: PMC12527950.

3.	Rodriguez-Gomez JA, Kavanagh E, Engskog-Vlachos P, Engskog MKR, Herrera AJ, Espinosa-Oliva AM, Joseph B, Hajji N, Venero JL, Burguillos MA. Microglia: Agents of the CNS Pro-Inflammatory Response. Cells. 2020;9(7). Epub 20200717. doi: 10.3390/cells9071717. PubMed PMID: 32709045; PMCID: PMC7407646.

4.	Colonna M, Butovsky O. Microglia Function in the Central Nervous System During Health and Neurodegeneration. Annu Rev Immunol. 2017;35:441-68. Epub 20170209. doi: 10.1146/annurev-immunol-051116-052358. PubMed PMID: 28226226; PMCID: PMC8167938.

5.	Read RD, Tapp ZM, Rajappa P, Hambardzumyan D. Glioblastoma microenvironment-from biology to therapy. Genes Dev. 2024;38(9-10):360-79. Epub 20240625. doi: 10.1101/gad.351427.123. PubMed PMID: 38811170; PMCID: PMC11216181.

6.	Persson ML, Douglas AM, Alvaro F, Faridi P, Larsen MR, Alonso MM, Vitanza NA, Dun MD. The intrinsic and microenvironmental features of diffuse midline glioma: Implications for the development of effective immunotherapeutic treatment strategies. Neuro Oncol. 2022;24(9):1408-22. doi: 10.1093/neuonc/noac117. PubMed PMID: 35481923; PMCID: PMC9435509.

7.	Chen Z, Giotti B, Kaluzova M, Vallcorba MP, Rawat K, Price G, Herting CJ, Pinero G, Cristea S, Ross JL, Ackley J, Maximov V, Szulzewsky F, Thomason W, Marquez-Ropero M, Angione A, Nichols N, Tsankova NM, Michor F, Shayakhmetov DM, Gutmann DH, Tsankov AM, Hambardzumyan D. A paracrine circuit of IL-1beta/IL-1R1 between myeloid and tumor cells drives genotype-dependent glioblastoma progression. J Clin Invest. 2023;133(22). Epub 20231115. doi: 10.1172/JCI163802. PubMed PMID: 37733448; PMCID: PMC10645395.

8.	Zhu C, Mustafa D, Zheng PP, van der Weiden M, Sacchetti A, Brandt M, Chrifi I, Tempel D, Leenen PJM, Duncker DJ, Cheng C, Kros JM. Activation of CECR1 in M2-like TAMs promotes paracrine stimulation-mediated glial tumor progression. Neuro Oncol. 2017;19(5):648-59. doi: 10.1093/neuonc/now251. PubMed PMID: 28453746; PMCID: PMC5464467.

9.	Yao M, Ventura PB, Jiang Y, Rodriguez FJ, Wang L, Perry JSA, Yang Y, Wahl K, Crittenden RB, Bennett ML, Qi L, Gong CC, Li XN, Barres BA, Bender TP, Ravichandran KS, Janes KA, Eberhart CG, Zong H. Astrocytic trans-Differentiation Completes a Multicellular Paracrine Feedback Loop Required for Medulloblastoma Tumor Growth. Cell. 2020;180(3):502-20 e19. Epub 2020/01/28. doi: 10.1016/j.cell.2019.12.024. PubMed PMID: 31983537; PMCID: PMC7259679.

10.   Fernandez C, Tatard VM, Bertrand N, Dahmane N. Differential modulation of Sonic-hedgehog-induced cerebellar granule cell precursor proliferation by the IGF signaling network. Dev Neurosci. 2010;32(1):59-70. Epub 20100325. doi: 10.1159/000274458. PubMed PMID: 20389077; PMCID: PMC2866582.

11.   Rao G, Pedone CA, Del Valle L, Reiss K, Holland EC, Fults DW. Sonic hedgehog and insulin-like growth factor signaling synergize to induce medulloblastoma formation from nestin-expressing neural progenitors in mice. Oncogene. 2004;23(36):6156-62. doi: 10.1038/sj.onc.1207818. PubMed PMID: 15195141.

12.   Tan IL, Arifa RDN, Rallapalli H, Kana V, Lao Z, Sanghrajka RM, Sumru Bayin N, Tanne A, Wojcinski A, Korshunov A, Bhardwaj N, Merad M, Turnbull DH, Lafaille JJ, Joyner AL. CSF1R inhibition depletes tumor-associated macrophages and attenuates tumor progression in a mouse sonic Hedgehog-Medulloblastoma model. Oncogene. 2021;40(2):396-407. Epub 20201106. doi: 10.1038/s41388-020-01536-0. PubMed PMID: 33159168; PMCID: PMC7855734.

13.   Dang MT, Gonzalez MV, Gaonkar KS, Rathi KS, Young P, Arif S, Zhai L, Alam Z, Devalaraja S, To TKJ, Folkert IW, Raman P, Rokita JL, Martinez D, Taroni JN, Shapiro JA, Greene CS, Savonen C, Mafra F, Hakonarson H, Curran T, Haldar M. Macrophages in SHH subgroup medulloblastoma display dynamic heterogeneity that varies with treatment modality. Cell Reports. 2021;34(13):108917. doi: https://doi.org/10.1016/j.celrep.2021.108917.

14.   Lee C, Lee J, Choi SA, Kim SK, Wang KC, Park SH, Kim SH, Lee JY, Phi JH. M1 macrophage recruitment correlates with worse outcome in SHH Medulloblastomas. BMC Cancer. 2018;18(1):535. Epub 2018/05/10. doi: 10.1186/s12885-018-4457-8. PubMed PMID: 29739450; PMCID: PMC5941618.

15.   Maximov V, Chen Z, Wei Y, Robinson MH, Herting CJ, Shanmugam NS, Rudneva VA, Goldsmith KC, MacDonald TJ, Northcott PA, Hambardzumyan D, Kenney AM. Tumour-associated macrophages exhibit anti-tumoural properties in Sonic Hedgehog medulloblastoma. Nat Commun. 2019;10(1):2410. Epub 2019/06/05. doi: 10.1038/s41467-019-10458-9. PubMed PMID: 31160587; PMCID: PMC6546707.

16.   Pokrajac NT, Tokarew NJA, Gurdita A, Ortin-Martinez A, Wallace VA. Meningeal macrophages inhibit chemokine signaling in pre-tumor cells to suppress mouse medulloblastoma initiation. Dev Cell. 2023;58(20):2015-31 e8. Epub 20230928. doi: 10.1016/j.devcel.2023.08.033. PubMed PMID: 37774709.

17.   Ross JL, Chen Z, Herting CJ, Grabovska Y, Szulzewsky F, Puigdelloses M, Monterroza L, Switchenko J, Wadhwani NR, Cimino PJ, Mackay A, Jones C, Read RD, MacDonald TJ, Schniederjan M, Becher OJ, Hambardzumyan D. Platelet-derived growth factor beta is a potent inflammatory driver in paediatric high-grade glioma. Brain. 2021;144(1):53-69. doi: 10.1093/brain/awaa382. PubMed PMID: 33300045; PMCID: PMC7954387.

18.   Daginakatte GC, Gutmann DH. Neurofibromatosis-1 (Nf1) heterozygous brain microglia elaborate paracrine factors that promote Nf1-deficient astrocyte and glioma growth. Hum Mol Genet. 2007;16(9):1098-112. Epub 20070330. doi: 10.1093/hmg/ddm059. PubMed PMID: 17400655.

19.   Ausejo-Mauleon I, Labiano S, de la Nava D, Laspidea V, Zalacain M, Marrodan L, Garcia-Moure M, Gonzalez-Huarriz M, Hervas-Corpion I, Dhandapani L, Vicent S, Collantes M, Penuelas I, Becher OJ, Filbin MG, Jiang L, Labelle J, de Biagi-Junior CAO, Nazarian J, Laternser S, Phoenix TN, van der Lugt J, Kranendonk M, Hoogendijk R, Mueller S, De Andrea C, Anderson AC, Guruceaga E, Koschmann C, Yadav VN, Gallego Perez-Larraya J, Patino-Garcia A, Pastor F, Alonso MM. TIM-3 blockade in diffuse intrinsic pontine glioma models promotes tumor regression and antitumor immune memory. Cancer Cell. 2023;41(11):1911-26 e8. Epub 20231005. doi: 10.1016/j.ccell.2023.09.001. PubMed PMID: 37802053; PMCID: PMC10644900.

20.   Lin GL, Nagaraja S, Filbin MG, Suva ML, Vogel H, Monje M. Non-inflammatory tumor microenvironment of diffuse intrinsic pontine glioma. Acta Neuropathol Commun. 2018;6(1):51. Epub 20180628. doi: 10.1186/s40478-018-0553-x. PubMed PMID: 29954445; PMCID: PMC6022714.

21.   Pachocki CJ, Hol EM. Current perspectives on diffuse midline glioma and a different role for the immune microenvironment compared to glioblastoma. J Neuroinflammation. 2022;19(1):276. Epub 20221119. doi: 10.1186/s12974-022-02630-8. PubMed PMID: 36403059; PMCID: PMC9675250.

22.   Lu H, Dietsch GN, Matthews MA, Yang Y, Ghanekar S, Inokuma M, Suni M, Maino VC, Henderson KE, Howbert JJ, Disis ML, Hershberg RM. VTX-2337 is a novel TLR8 agonist that activates NK cells and augments ADCC. Clin Cancer Res. 2012;18(2):499-509. Epub 20111129. doi: 10.1158/1078-0432.CCR-11-1625. PubMed PMID: 22128302.

23.   Peng J, Tsang JY, Li D, Niu N, Ho DH, Lau KF, Lui VC, Lamb JR, Chen Y, Tam PK. Inhibition of TGF-beta signaling in combination with TLR7 ligation re-programs a tumoricidal phenotype in tumor-associated macrophages. Cancer Lett. 2013;331(2):239-49. Epub 20130111. doi: 10.1016/j.canlet.2013.01.001. PubMed PMID: 23318200.

24.   Kawai T, Ikegawa M, Ori D, Akira S. Decoding Toll-like receptors: Recent insights and perspectives in innate immunity. Immunity. 2024;57(4):649-73. doi: 10.1016/j.immuni.2024.03.004. PubMed PMID: 38599164.

25.   Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. Nat Immunol. 2010;11(5):373-84. Epub 20100420. doi: 10.1038/ni.1863. PubMed PMID: 20404851.

26.   Isvoranu G, Surcel M, Enciu AM, Munteanu AN, Neagu M, Niculae AM, Chiritoiu G, Munteanu CVA, Chiritoiu-Butnaru M. Toll-like Receptor 7/8 Agonists Exert Antitumor Effect in a Mouse Melanoma Model. Medicina (Kaunas). 2026;62(1). Epub 20260109. doi: 10.3390/medicina62010141. PubMed PMID: 41597427; PMCID: PMC12843957.

27.   Yazdimamaghani M, Kolupaev OV, Lim C, Hwang D, Laurie SJ, Perou CM, Kabanov AV, Serody JS. Tumor microenvironment immunomodulation by nanoformulated TLR 7/8 agonist and PI3k delta inhibitor enhances therapeutic benefits of radiotherapy. Biomaterials. 2025;312:122750. Epub 20240808. doi: 10.1016/j.biomaterials.2024.122750. PubMed PMID: 39126779; PMCID: PMC11401478.

28.   Frega G, Wu Q, Le Naour J, Vacchelli E, Galluzzi L, Kroemer G, Kepp O. Trial Watch: experimental TLR7/TLR8 agonists for oncological indications. Oncoimmunology. 2020;9(1):1796002. Epub 20200721. doi: 10.1080/2162402X.2020.1796002. PubMed PMID: 32934889; PMCID: PMC7466852.

29.   Lesniak M, Lipniarska J, Majka P, Kopyt W, Lejman M, Zawitkowska J. The Role of TRL7/8 Agonists in Cancer Therapy, with Special Emphasis on Hematologic Malignancies. Vaccines (Basel). 2023;11(2). Epub 20230128. doi: 10.3390/vaccines11020277. PubMed PMID: 36851155; PMCID: PMC9967151.

30.   Vinod N, Hwang D, Azam SH, Van Swearingen AED, Wayne E, Fussell SC, Sokolsky-Papkov M, Pecot CV, Kabanov AV. High-capacity poly(2-oxazoline) formulation of TLR 7/8 agonist extends survival in a chemo-insensitive, metastatic model of lung adenocarcinoma. Sci Adv. 2020;6(25):eaba5542. Epub 20200617. doi: 10.1126/sciadv.aba5542. PubMed PMID: 32596460; PMCID: PMC7299629.

31.   Bhagchandani SH, Vohidov F, Milling LE, Tong EY, Brown CM, Ramseier ML, Liu B, Fessenden TB, Nguyen HV-T, Kiel GR, Won L, Langer RS, Spranger S, Shalek AK, Irvine DJ, Johnson JA. Engineering kinetics of

TLR7/8 agonist release from bottlebrush prodrugs enables tumor-focused immune stimulation. Science Advances. 2023;9(16):eadg2239. doi: doi:10.1126/sciadv.adg2239.

32.   Dudek AZ, Yunis C, Harrison LI, Kumar S, Hawkinson R, Cooley S, Vasilakos JP, Gorski KS, Miller JS. First in human phase I trial of 852A, a novel systemic toll-like receptor 7 agonist, to activate innate immune responses in patients with advanced cancer. Clin Cancer Res. 2007;13(23):7119-25. doi: 10.1158/1078-0432.CCR-07-1443. PubMed PMID: 18056192.

33.   Harrison LI, Astry C, Kumar S, Yunis C. Pharmacokinetics of 852A, an imidazoquinoline Toll-like receptor 7-specific agonist, following intravenous, subcutaneous, and oral administrations in humans. J Clin Pharmacol. 2007;47(8):962-9. doi: 10.1177/0091270007303766. PubMed PMID: 17660481.

34.   Zahr NM, Zhao Q, Goodcase R, Pfefferbaum A. Systemic Administration of the TLR7/8 Agonist Resiquimod (R848) to Mice Is Associated with Transient, In Vivo-Detectable Brain Swelling. Biology (Basel). 2022;11(2). Epub 20220210. doi: 10.3390/biology11020274. PubMed PMID: 35205140; PMCID: PMC8869423.

35.   Shang H, Xia D, Geng R, Wu J, Deng W, Tong Y, Ba X, Zhong Z, He Y, Huang Q, Ye T, Yang X, Jiang K, Peng E, Zhu J, Liu Y, Chen Z, Tang K. Thermosensitive Resiquimod-Loaded Lipid Nanoparticles Promote the Polarization of Tumor-Associated Macrophages to Enhance Bladder Cancer Immunotherapy. ACS Nano. 2025;19(21):19599-621. Epub 20250517. doi: 10.1021/acsnano.4c17444. PubMed PMID: 40380939; PMCID: PMC12139615.

36.   Geng R, Shang H, Chen Z, Tang K, Liu Y, Zhu J. ROS-responsive resiquimod prodrug nanoparticles for macrophage reprogramming and enhanced immune checkpoint inhibitor therapy in bladder cancer. Biomaterials. 2026;326:123677. Epub 20250902. doi: 10.1016/j.biomaterials.2025.123677. PubMed PMID: 40907393.

37.   Li JX, Shu N, Zhang YJ, Tong QS, Wang L, Zhang JY, Du JZ. Self-Assembled Nanoparticles from the Amphiphilic Prodrug of Resiquimod for Improved Cancer Immunotherapy. ACS Appl Mater Interfaces. 2024;16(20):25665-75. Epub 20240512. doi: 10.1021/acsami.4c01563. PubMed PMID: 38735053.

38.   Turco V, Pfleiderer K, Hunger J, Horvat NK, Karimian-Jazi K, Schregel K, Fischer M, Brugnara G, Jahne K, Sturm V, Streibel Y, Nguyen D, Altamura S, Agardy DA, Soni SS, Alsasa A, Bunse T, Schlesner M, Muckenthaler MU, Weissleder R, Wick W, Heiland S, Vollmuth P, Bendszus M, Rodell CB, Breckwoldt MO, Platten M. T cell-independent eradication of experimental glioma by intravenous TLR7/8-agonist-loaded nanoparticles. Nat Commun. 2023;14(1):771. Epub 20230211. doi: 10.1038/s41467-023-36321-6. PubMed PMID: 36774352; PMCID: PMC9922247.

39.   Graham-Gurysh EG, Woodring RN, Simpson SR, Mendell SE, Lukesh NR, Pena ES, Moore KM, Ontiveros-Padilla LA, Hendricksen AT, Lopez AM, Williamson GL, Murphy CT, Genito CJ, Hipp KA, Singh G, Zamboni WC, Hingtgen SD, Fecci PE, Bachelder EM, Ainslie KM. Post-resection delivery of a TLR7/8 agonist from a biodegradable scaffold achieves immune-mediated glioblastoma clearance and protection against tumor challenge in mice. Nat Commun. 2025;16(1):8603. Epub 20250929. doi: 10.1038/s41467-025-63692-9. PubMed PMID: 41022761; PMCID: PMC12479796.

40.   Phoenix TN, Patmore DM, Boop S, Boulos N, Jacus MO, Patel YT, Roussel MF, Finkelstein D, Goumnerova L, Perreault S, Wadhwa E, Cho YJ, Stewart CF, Gilbertson RJ. Medulloblastoma Genotype Dictates Blood Brain Barrier Phenotype. Cancer Cell. 2016;29(4):508-22. Epub 2016/04/07. doi: 10.1016/j.ccell.2016.03.002. PubMed PMID: 27050100; PMCID: PMC4829447.

41.   Eisemann T, Wechsler-Reya RJ. Coming in from the cold: overcoming the hostile immune microenvironment of medulloblastoma. Genes Dev. 2022;36(9-10):514-32. doi: 10.1101/gad.349538.122. PubMed PMID: 35680424; PMCID: PMC9186392.

42.   Lim C, Dismuke T, Malawsky D, Ramsey JD, Hwang D, Godfrey VL, Kabanov AV, Gershon TR, Sokolsky-Papkov M. Enhancing CDK4/6 inhibitor therapy for medulloblastoma using nanoparticle delivery and scRNA-seq-guided combination with sapanisertib. Sci Adv. 2022;8(4):eabl5838. Epub 20220126. doi: 10.1126/sciadv.abl5838. PubMed PMID: 35080986; PMCID: PMC8791615.

43.   Hwang D, Dismuke T, Tikunov A, Rosen EP, Kagel JR, Ramsey JD, Lim C, Zamboni W, Kabanov AV, Gershon TR, Sokolsky-Papkov Ph DM. Poly(2-oxazoline) nanoparticle delivery enhances the therapeutic potential of vismodegib for medulloblastoma by improving CNS pharmacokinetics and reducing systemic toxicity. Nanomedicine. 2020;32:102345. Epub 2020/12/02. doi: 10.1016/j.nano.2020.102345. PubMed PMID: 33259959.

44.   Schuller U, Heine VM, Mao J, Kho AT, Dillon AK, Han YG, Huillard E, Sun T, Ligon AH, Qian Y, Ma Q, Alvarez-Buylla A, McMahon AP, Rowitch DH, Ligon KL. Acquisition of granule neuron precursor identity is a critical determinant of progenitor cell competence to form Shh-induced medulloblastoma. Cancer Cell. 2008;14(2):123-34. doi: 10.1016/j.ccr.2008.07.005. PubMed PMID: 18691547; PMCID: PMC2597270.

45.   Li Y, Lim C, Dismuke T, Malawsky DS, Oasa S, Bruce ZC, Offenhauser C, Baumgartner U, D'Souza RCJ, Edwards SL, French JD, Ock LSH, Nair S, Sivakumaran H, Harris L, Tikunov AP, Hwang D, Alicea Pauneto CDM, Maybury M, Hassall T, Wainwright B, Kesari S, Stein G, Piper M, Johns TG, Sokolsky-Papkov M, Terenius L, Vukojevic V, McSwain LF, Gershon TR, Day BW. Suppressing recurrence in Sonic Hedgehog subgroup medulloblastoma using the OLIG2 inhibitor CT-179. Nat Commun. 2025;16(1):1091. Epub 20250204. doi: 10.1038/s41467-024-54861-3. PubMed PMID: 39904981; PMCID: PMC11794477.

46.   Malawsky DS, Weir SJ, Ocasio JK, Babcock B, Dismuke T, Cleveland AH, Donson AM, Vibhakar R, Wilhelmsen K, Gershon TR. Cryptic developmental events determine medulloblastoma radiosensitivity and cellular heterogeneity without altering transcriptomic profile. Commun Biol. 2021;4(1):616. Epub 2021/05/23. doi: 10.1038/s42003-021-02099-w. PubMed PMID: 34021242; PMCID: PMC8139976.

47.   Xing Q, Feng Y, Sun H, Yang S, Sun T, Guo X, Ji F, Wu B, Zhou D. Scavenger receptor MARCO contributes to macrophage phagocytosis and clearance of tumor cells. Exp Cell Res. 2021;408(2):112862. Epub 20211006. doi: 10.1016/j.yexcr.2021.112862. PubMed PMID: 34626585.

48.   Heil F, Hemmi H, Hochrein H, Ampenberger F, Kirschning C, Akira S, Lipford G, Wagner H, Bauer S. Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8. Science. 2004;303(5663):1526-9. Epub 20040219. doi: 10.1126/science.1093620. PubMed PMID: 14976262.

49.   Malawsky DS, Dismuke T, Liu H, Castellino E, Brenman J, Dasgupta B, Tikunov A, Gershon TR. Chronic AMPK inactivation slows SHH medulloblastoma progression by inhibiting mTORC1 signaling and depleting tumor stem cells. iScience. 2023;26(12):108443. Epub 20231114. doi: 10.1016/j.isci.2023.108443. PubMed PMID: 38094249; PMCID: PMC10716552.

50.   Ocasio J, Babcock B, Malawsky D, Weir SJ, Loo L, Simon JM, Zylka MJ, Hwang D, Dismuke T, Sokolsky M, Rosen EP, Vibhakar R, Zhang J, Saulnier O, Vladoiu M, El-Hamamy I, Stein LD, Taylor MD, Smith KS, Northcott PA, Colaneri A, Wilhelmsen K, Gershon TR. scRNA-seq in medulloblastoma shows cellular heterogeneity and lineage expansion support resistance to SHH inhibitor therapy. Nat Commun. 2019;10(1):5829. Epub 2019/12/22. doi: 10.1038/s41467-019-13657-6. PubMed PMID: 31863004; PMCID: PMC6925218.

51.   Zhao XF, Alam MM, Liao Y, Huang T, Mathur R, Zhu X, Huang Y. Targeting Microglia Using Cx3cr1-Cre Lines: Revisiting the Specificity. eNeuro. 2019;6(4). Epub 20190710. doi: 10.1523/ENEURO.0114-19.2019. PubMed PMID: 31201215; PMCID: PMC6620394.

52.   Jung S, Aliberti J, Graemmel P, Sunshine MJ, Kreutzberg GW, Sher A, Littman DR. Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green fluorescent protein reporter gene insertion. Mol Cell Biol. 2000;20(11):4106-14. doi: 10.1128/MCB.20.11.4106-4114.2000. PubMed PMID: 10805752; PMCID: PMC85780.

53.   Chen Z, Feng X, Herting CJ, Garcia VA, Nie K, Pong WW, Rasmussen R, Dwivedi B, Seby S, Wolf SA, Gutmann DH, Hambardzumyan D. Cellular and Molecular Identity of Tumor-Associated Macrophages in Glioblastoma. Cancer Res. 2017;77(9):2266-78. Epub 2017/02/27. doi: 10.1158/0008-5472.CAN-16-2310. PubMed PMID: 28235764; PMCID: PMC5741820.

54.   Stein M, Keshav S, Harris N, Gordon S. Interleukin 4 potently enhances murine macrophage mannose receptor activity: a marker of alternative immunologic macrophage activation. J Exp Med. 1992;176(1):287-92. doi: 10.1084/jem.176.1.287. PubMed PMID: 1613462; PMCID: PMC2119288.

55.   Lloyd-Burton SM, York EM, Anwar MA, Vincent AJ, Roskams AJ. SPARC regulates microgliosis and functional recovery following cortical ischemia. J Neurosci. 2013;33(10):4468-81. Epub 2013/03/08. doi: 10.1523/JNEUROSCI.3585-12.2013. PubMed PMID: 23467362; PMCID: PMC6704956.

56.   Hou X, Qu X, Chen W, Sang X, Ye Y, Wang C, Guo Y, Shi H, Yang C, Zhu K, Zhang Y, Xu H, Lv L, Zhang D, Hou L. CD36 deletion prevents white matter injury by modulating microglia polarization through the Traf5-MAPK signal pathway. J Neuroinflammation. 2024;21(1):148. Epub 20240605. doi: 10.1186/s12974-024-03143-2. PubMed PMID: 38840180; PMCID: PMC11155181.

57.   Zhu R, Pan YH, Sun L, Zhang T, Wang C, Ye S, Yang N, Lu T, Wisniewski T, Dang S, Zhang W. ADAMTS18 Deficiency Affects Neuronal Morphogenesis and Reduces the Levels of Depression-like Behaviors in Mice. Neuroscience. 2019;399:53-64. Epub 20181221. doi: 10.1016/j.neuroscience.2018.12.025. PubMed PMID: 30579834; PMCID: PMC8006808.

58.   Talbot J, Fombonne J, Torrejon J, Babcock BR, McSwain LF, Rama N, Lospinoso Severini L, Bonerandi E, Marsaud V, Bernardi F, Gharsalli T, Guix C, Ducarouge B, Neururer V, Basili I, Mercier AL, Yu H, Forget A, Indersie E, Leboucher S, Souphron J, Okonechnikov K, Wang W, Kawauchi D, Wainwright BJ, Frappaz D, Varlet P, Dufour C, Beccaria K, Blauwblomme T, Martignetti L, Di Marcotullio L, Puget S, Doz F, Bourdeaut F, Masliah-Planchon J, Gershon TR, Mehlen P, Ayrault O. Sonic hedgehog medulloblastomas are dependent on Netrin-1 for survival. Nat Commun. 2025;16(1):5137. Epub 20250603. doi: 10.1038/s41467-025-59612-6. PubMed PMID: 40461501; PMCID: PMC12134228.

59.   Machold R, Fishell G. Math1 is expressed in temporally discrete pools of cerebellar rhombic-lip neural progenitors. Neuron. 2005;48(1):17-24. doi: 10.1016/j.neuron.2005.08.028. PubMed PMID: 16202705.

60.   Zhuo L, Theis M, Alvarez-Maya I, Brenner M, Willecke K, Messing A. hGFAP-cre transgenic mice for manipulation of glial and neuronal function in vivo. Genesis. 2001;31(2):85-94. Epub 2001/10/23. PubMed PMID: 11668683.

61.   Ross JL, Puigdelloses-Vallcorba M, Pinero G, Soni N, Thomason W, DeSisto J, Angione A, Tsankova NM, Castro MG, Schniederjan M, Wadhwani NR, Raju GP, Morgenstern P, Becher OJ, Green AL, Tsankov AM, Hambardzumyan D. Microglia and monocyte-derived macrophages drive progression of pediatric high-grade

gliomas and are transcriptionally shaped by histone mutations. Immunity. 2024;57(11):2669-87 e6. Epub 20241011. doi: 10.1016/j.immuni.2024.09.007. PubMed PMID: 39395421; PMCID: PMC11578068.

62.   Bhatia B, Northcott PA, Hambardzumyan D, Govindarajan B, Brat DJ, Arbiser JL, Holland EC, Taylor MD, Kenney AM. Tuberous sclerosis complex suppression in cerebellar development and medulloblastoma: separate regulation of mammalian target of rapamycin activity and p27 Kip1 localization. Cancer Res. 2009;69(18):7224-34. Epub 2009/09/10. doi: 0008-5472.CAN-09-1299 [pii]

10.1158/0008-5472.CAN-09-1299. PubMed PMID: 19738049; PMCID: 2745891.

63.   Cordero FJ, Huang Z, Grenier C, He X, Hu G, McLendon RE, Murphy SK, Hashizume R, Becher OJ. Histone H3.3K27M Represses p16 to Accelerate Gliomagenesis in a Murine Model of DIPG. Mol Cancer Res. 2017;15(9):1243-54. Epub 20170518. doi: 10.1158/1541-7786.MCR-16-0389. PubMed PMID: 28522693; PMCID: PMC5581686.

64.   Price G, Bouras A, Hambardzumyan D, Hadjipanayis CG. Current knowledge on the immune microenvironment and emerging immunotherapies in diffuse midline glioma. EBioMedicine. 2021;69:103453. Epub 20210619. doi: 10.1016/j.ebiom.2021.103453. PubMed PMID: 34157482; PMCID: PMC8220552.

65.   Huang L, Chen W, Tan Z, Huang Y, Gu X, Liu L, Zhang H, Shi Y, Ding J, Zheng C, Guo Z, Yu B. Mrc1(+) macrophage-derived IGF1 mitigates crystal nephropathy by promoting renal tubule cell proliferation via the AKT/Rb signaling pathway. Theranostics. 2024;14(4):1764-80. Epub 20240217. doi: 10.7150/thno.89174. PubMed PMID: 38389846; PMCID: PMC10879870.

66.   Rodell CB, Arlauckas SP, Cuccarese MF, Garris CS, Li R, Ahmed MS, Kohler RH, Pittet MJ, Weissleder R. TLR7/8-agonist-loaded nanoparticles promote the polarization of tumour-associated macrophages to enhance cancer immunotherapy. Nat Biomed Eng. 2018;2(8):578-88. Epub 20180521. doi: 10.1038/s41551-018-0236-8. PubMed PMID: 31015631; PMCID: PMC6192054.

67.   Wu N, Morsey BM, Emanuel KM, Fox HS. Sequence-specific extracellular microRNAs activate TLR7 and induce cytokine secretion and leukocyte migration. Mol Cell Biochem. 2021;476(11):4139-51. Epub 20210727. doi: 10.1007/s11010-021-04220-3. PubMed PMID: 34313894.

68.   Tanaka R, Murakami Y, Ellis D, Seita J, Yinga W, Kakuta S, Kumano K, Fukui R, Miyake K. TLR7 responses in glomerular macrophages accelerate the progression of glomerulonephritis in NZBWF1 mice. Int Immunol. 2025;37(6):339-53. Epub 20250127. doi: 10.1093/intimm/dxaf005. PubMed PMID: 40401698.

69.   Kaiser Y, Garris CS, Marinari E, Kim HS, Oh J, Pedard M, Halabi EA, Choi M, Parvanian S, Kohler R, Migliorini D, Weissleder R. Targeting immunosuppressive myeloid cells via implant-mediated slow release of small molecules to prevent glioblastoma recurrence. Nat Biomed Eng. 2025. Epub 20251022. doi: 10.1038/s41551-025-01533-2. PubMed PMID: 41125869.

70.   Wolak DJ, Thorne RG. Diffusion of macromolecules in the brain: implications for drug delivery. Mol Pharm. 2013;10(5):1492-504. Epub 20130131. doi: 10.1021/mp300495e. PubMed PMID: 23298378; PMCID: PMC3646902.

71.   Luxenhofer R, Schulz A, Roques C, Li S, Bronich TK, Batrakova EV, Jordan R, Kabanov AV. Doubly amphiphilic poly(2-oxazoline)s as high-capacity delivery systems for hydrophobic drugs. Biomaterials. 2010;31(18):4972-9. Epub 2010/03/30. doi: 10.1016/j.biomaterials.2010.02.057. PubMed PMID: 20346493; PMCID: PMC2884201.

72.   Sahn M, Weber C, Schubert US. Poly(2-oxazoline)-Containing Triblock Copolymers: Synthesis and Applications. Polymer Reviews. 2019;59(2):240-79. doi: 10.1080/15583724.2018.1496930.

73.   Cho S, Rasoulianboroujeni M, Kang RH, Kwon GS. From Conventional to Next-Generation Strategies: Recent Advances in Polymeric Micelle Preparation for Drug Delivery. Pharmaceutics. 2025;17(10):1360. PubMed PMID: doi:10.3390/pharmaceutics17101360.

74.   De la Cruz G, Nikolaishvili-Feinberg N, Gershon TR. Automated Immunofluorescence Staining for Analysis of DNA Damage and Apoptosis in Brain Sections. Methods Mol Biol. 2023;2583:55-61. doi: 10.1007/978-1-0716-2752-5_6. PubMed PMID: 36418725.

Figure 1: ResiPOx enhances delivery across the blood brain barrier and achieves bioactive exposure.

Supplementary Fig. 5: ResiPOx induces BTK expression in *G-Smo* mice

A.

BTK ResiPOx

Supplementary Fig. 5: (A) Quantification of BTK immunostaining of *G-Smo* following ResiPOx treatment.

Supplementary Fig. 6: ResiPOx induces transcriptional changes in DMG

A.

RCAS DMG 24h ResiPOx

Supplementary Fig. 6: (A) Volcano plot of RCAS DMG DEGs 24h after a single dose of ResiPOx.

Supplementary Fig 4. ResiPOx induces changes to tumor and myeloid populations in *G-Smo* mice

Supplementary Fig. 4: (A) Seurat Cluster UMAP of Control and ResiPOx treated G-Smo tumors. **(B)** Violin plot of P2ry12 expression in 3 myeloid sub-clusters demonstrating Microglial enrichment. **(C)** Violin plot of Fcrls expression in 3 myeloid sub-clusters indicating Microglial enrichment. **(D)** Violin plot of Cytip in 3 myeloid sub-clusters indicating BMDM enrichment. **(E)** Comparison of Mrc1 expression in myeloid cluseter 11 between control and ResiPOx treated G-Smo, wilcox t-test. **(F)** Comparison of Igf1 expression in myeloid cluseter 11 between control and ResiPOx treated G-Smo, wilcox t-test. **(G)** Representative pRb immunostaining of control or ResiPOx treated G-Smo.

Supplementary Fig 3. ResiPOx induces changes to IFNβ and myeloid signaling at the protein level

Supplementary Fig. 3. ResiPOx induces an IFNβ Response and Reduction in IGF signaling. (A) Additional β-Tubulin controls for Fig. 3G and 4D. **(B)** Additional biological replicates for chemokine array Fig. 4B.. **(C)** Additional chemokokine arrays indicating induction of CXCL10 and CCL9 24h after a single dose of ResiPOx (n=2 per timepoint). **(D)** Additional β-Tubulin controls for Fig. 4C.

Supplementary Fig. 1. ResiPOx enhances delivery across the blood brain barrier and achieves bioactive exposure.

Supplementary Fig. 1. Polyoxazoline Nano-particle Resiquimod demonstrates greater tissue penetrance. In Liver, Kidney, and Spleen, POx formulation improves tissue penetration and persistence as indicated by differences in area under the curve (AUC) and statistically significant differences within time points (permutation test (AUC) or students t-test within time point *** p < 0.0001, ** p < 0.001, * p < 0.05).

Supplementary Fig. 2. ResiPOx induces changes in TAM and tumor populations.

Supplementary Fig. 2. ResiPOx induces changes in TAM and tumor populations. (A) Correlation of *G-Smo* DEGs 6h and 24h after a single ResiPOx injection. **(B)** Correlation of *G-Smo* DEGs 24h after a single ResiPOx injection or 24h after 3 EOD ResiPOx injections. **(C)** Correlation of *G-Smo* NES 6h and 24h after a single ResiPOx injection. **(D)** Correlation of *G-Smo* NES 24h after a single ResiPOx injection or 24h after 3 EOD ResiPOx injections. **(E)** Volcano plot of BMDM RNAseq 24h following a 2h pulse of ResiPOx. **(F)** Correlation *G-Smo* and BMDM DEG 24h after a single ResiPOx injection or 24h following a 2h pulse of ResiPOx, respectively.

Figure 7. ResiPOx shows similar efficacy in mice with DMG

Figure 6. ResiPOx increased phagocytosis of tumor cells by TAMs

Figure 2. ResiPOx induces changes in TAM and tumor populations.