Functional characterization of bat limb regulatory elements

Aki Ushiki[1,2,3*], Guy Kelman[4], Rory Sheng[1,2], Elizabeth Murray[1,2], Walter Eckalbar[1,2],
Yichi Zhang[1,2], Mai Nobuhara[1,2], Riya Rajani[1,2], Katja Friess[1,2], Vladyslav Barskyi[1,2],
Kelly Ngo[1,2] , Shiori Kinoshita[3], Stephen A Schlebusch[5], Mandy Mason[6], Shijun Zhan[7],
Maggie Liang[1,2], Sarah Fong[1,2], Md Yasin Haider[3], Varda Singhal[1,2], Tony Schountz[7],
Dorit Hockman[6,8,9], Nicola Illing[6], Tommy Kaplan[10,11], Nadav Ahituv[1,2*]

[1]Department of Bioengineering and Therapeutic Sciences, University of California San Francisco, San Francisco, CA, USA.
[2]Institute for Human Genetics, University of California San Francisco, San Francisco, CA, USA.
[3]Department of Genetics, Washington University School of Medicine, St. Louis, MO, USA
[4]The Jerusalem Center for Personalized Computational Medicine, Hebrew University of Jerusalem, Jerusalem, Israel.
[5]Department of Zoology, Faculty of Science, Charles University, Prague, Czech Republic.
[6]Department of Molecular and Cell Biology, University of Cape Town, Rondebosch 7701, South Africa.
[7]Department of Microbiology, Immunology and Pathology, Colorado State University, Fort Collins, CO, USA.
[8]Division of Cell Biology, Department of Human Biology, University of Cape Town, Rondebosch 7701, South Africa.
[9]Neuroscience Institute, University of Cape Town, Cape Town, South Africa.
[10]School of Computer Science and Engineering, The Hebrew University of Jerusalem, Jerusalem, Israel.
[11]Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel

*Corresponding authors: akiu@wustl.edu, nadav.ahituv@ucsf.edu

Abstract

Bats are the only mammals capable of powered flight and roost head-down. However, the molecular changes shaping bat limbs remain largely unknown. Here, we used comparative functional genomics coupled with mouse-bat sequence swaps to identify key regulatory elements important in bat limb development. We generated and compared bat and mouse forelimb and hindlimb genomic datasets at key wing developmental timepoints, followed by mouse enhancer assays to characterize sequences showing differences between species. We then swapped six mouse enhancer sequences with their corresponding bat sequences, obtaining a variety of bat limb associated phenotypes, including ossification delay, longer digits, thicker skin and symmetrical hindlimb digits. Our work provides a genomic catalog of genes and regulatory elements involved in bat limb development and through extensive characterization in mice shows how changes in regulatory elements lead to small phenotypic changes that together contribute to bat limb development.

Main

Bats are the only mammal capable of powered flight and roost head-down. To achieve this, they have undergone a variety of morphological innovations, that include not only the wing itself but also unique muscles that power it along with higher metabolic rates, flexible bones, the lack of claws on digits II to V in the forelimb, and craniofacial changes such as having a smaller premaxilla and changes in ear placement to reduce drag[1]. In terms of the wing, during development the bat forelimb autopods retain extended embryonic growth of the metacarpals and proximal and intermediate phalanges of digits II to V[1-4]. In addition, developing bat autopods retain their interdigital tissue which forms part of the wing[1,5]. In contrast, the bat hindlimbs have relatively short, free, symmetrical digits. The hindlimb is diminutive in comparison, rotated by 180°, with short symmetrical clawed digits that allows bats to lock themselves into a roosting position with a tight grip[6,7].

Examining gene expression differences between bat forelimb and hindlimb at key embryonic developmental stages has found numerous genes to be associated with bat wing development. This was initially carried out in a 'gene-by-gene' manner, finding genes such as *Tbx3, Brinp3, Meis2, HoxD* genes and members of the *Shh-Fgf* signaling pathway to be associated with bat-specific changes in forelimb development[5,8-11]. Later studies used genome-wide assays, such as RNA-seq, on developing bat forelimbs and hindlimbs at key wing developmental stages finding a multitude of gene expression changes between forelimbs and hindlimbs, primarily in known limb developmental associated genes[12,13]. These include multiple ribosomal proteins and known limb patterning signaling pathways, including FGF, WNT, and BMP signaling. Recent work that used single-cell RNA-seq on the bat wing found a specific fibroblast cell population in the bat wing membrane that expresses *MEIS2* and *TBX3* that is thought to be important in tissue retention[14,15] and a unique bat forelimb mesenchymal progenitor cell population with high *PDGFD* expression that is thought to promote bone proliferation[15]. Combined, these studies suggest that changes in regulatory elements that differentially activate these genes in the forelimb are major drivers in the evolution of the bat wing.

Gene regulatory elements, such as promoters and enhancers, control spatial temporal and levels of gene expression. Several individual enhancers associated with bat wing development have been investigated on a 'one-by-one' basis. One such example is the *Prx1* limb enhancer. The replacement of this enhancer with the homologous bat *Prx1* sequence resulted in mouse embryonic day (E) 18.5 embryos having 6% longer forelimbs[16]. Using comparative genomics to annotate bat accelerated regions (BARs), sequences that are conserved in mammals but significantly changed in bats, identified hundreds of candidate regions[13,17,18]. Mouse enhancer assays using the bat sequence of five of these BARs, chosen due to their proximity to known limb genes, found all five to be limb enhancers and three to have differential enhancer activity compared to their homologous mouse sequence[17]. In addition, using chromatin immunoprecipitation followed by sequencing (ChIP-seq) for active [H3K27ac;[19,20]] and repressive

[H3K27me3;[21]] marks on autopods dissected from bat forelimbs and hindlimbs at key
wing developmental stages (CS15, CS16 and CS17) from the Natal long-fingered bat
(*Miniopterus natalensis*), identified thousands of differential active regulatory
elements[13]. Combined, these studies have identified thousands of candidate genes and
regulatory elements involved in this morphological adaptation, making it difficult to
pinpoint the sequences driving this adaptation.
98
We took a variety of approaches to narrow down the number of candidate genes and
regulatory elements specifically involved in bat wing development for further studies
including: 1) expression/activity differences from genomic datasets; 2) focusing on
biologically relevant genes; 3) BAR selection. We first generated genomic datasets
(RNA-seq, H3K27ac & H3K27me3 ChIP-seq) for mouse forelimb and hindlimb for
stages of development that match bat embryonic stages of development (CS15-
CS17)[22]. In addition, we generated ATAC-seq datasets for developing bat forelimb and
hindlimb samples at CS17and embryonic day (E) 13.5 embryos. Utilizing these
datasets, we were able to tease out genes and candidate regulatory sequences that
specifically changed in the bat forelimbs during these key stages of wing development.
We next carried out mouse enhancer assays for eleven candidate bat enhancer
sequences, finding several functional limb enhancers that show differential activity
compared to their homologous mouse sequences. To characterize their functional
importance, we next swapped six sequences in the mouse with the homologous bat
sequence, finding a multitude of phenotypes associated with bat limb development,
including delayed bone ossification, longer digits, thicker skin and symmetrical hindlimb
digits. Combined, our work highlights specific genes and regulatory regions important
for bat wing development. In addition, by functionally linking these elements to different
phenotypic outcomes through extensive characterization in mice, we show how
changes in regulatory elements lead to small phenotypic changes that together
contribute to bat limb development in line with recent hypotheses that large evolutionary
changes are caused by multiple factors[23].
121
122

Results

Transcriptomic comparison of embryonic bat and mouse limbs

Previous work by our labs identified over 7,000 genes to be differentially expressed between forelimb (FL) and hindlimb (HL) in *M. natalensis* during Carnegie Stages 15-17 (CS15-CS17), three key stages of bat wing development[13]. As there are inherent differences in gene expression in FL compared to HL autopods in flightless mammals, we carried out RNA-seq on mouse FL and HL at matching developmental changes, embryonic day (E) 12.0, 13.0 and 13.5 matching CS15, CS16 and CS17 respectively (**Fig. 1a**). We then compared FL and HL differentially expressed (DE) genes in bats and mice, highlighting specific gene expression changes that are unique to bat. We identified a total of 1,715 DE genes between FL and HL. Amongst them, 123 genes (7%) were shared between bats and mice and 1,234 (72%) and 358 (21%) were unique to bat and mouse respectively (**Fig. 1b**). For the bat FL, these included *HoxD* genes (*Hoxd10-13*) and *Tbx3* that showed higher expression in the bat FL compared to mouse (**Fig. 1c-d**). In the bat HL, we observed *HoxC* genes (*Hoxc8-10*) and *Tbx4* to have higher expression compared to mouse (**Fig. 1c-d**). Gene ontology (GO) enrichment analyses using DAVID[24] for the bat differential genes (N=1,234) found enrichment for terms that are involved in development (**Extended Data Fig. 1a**, **Supplementary Data 1**). Comparison of GeneORGANizer[25] results for bat only (N=1,234) versus mouse only (N=358) DE genes found an enrichment for bat DE genes known to be expressed in the skeleton (**Extended Data Fig. 1b**), including ones specific to the digits (**Extended Fig. 1c**).

Comparison of embryonic bat and mouse limb regulomes

We previously used ChIP-seq to characterize differentially active (H3K27ac) and silenced (H3K27me3) regulatory elements in the three *M. natalensis* developmental stages, CS15, CS16, and CS17[13]. We identified 4,553 and 19,352 differentially enriched regions for H3K27ac and H3K27me3 respectively and 2,475 regions to be differentially enriched between FL and HL for both H3K27ac and H3K27me3[13]. To reduce the number of potential regulatory element candidates involved in wing development, we carried out a similar ChIP-seq for H3K27ac and H3K27me3 in matching mouse FL and HL developmental timepoints (E12.0, E13.0, E13.5) (**Fig. 1a**). To directly compare ChIP-seq peaks between bat and mouse, we aligned our bat peaks to the mouse genome, using only sequences that aligned to a similar location in the mouse genome (see Methods). We then characterized differentially active (DA) peaks between bat and mouse in FL and HL at each developmental time point for either chromatin mark (H3K27ac or H3K27me3), identifying hundreds of DA peaks for each condition (**Fig. 1e**).

We next investigated differentially active (DA) peaks between FL and HL and bat and mouse. Using the Genomic Regions Enrichment of Annotations Tool (GREAT[26]), we analyzed each time point (CS15, CS16, CS17), tissue (FL or HL), and mark (H3K27ac or H3K27me3) individually for enriched phenotype terms. We found enrichment for

167 terms such as small hindlimb buds, abnormal tarsal bone morphology, abnormal
168 appendicular skeleton morphology, thin apical ectodermal ridge, abnormal limb bud
169 morphology, abnormal skeleton morphology, and abnormal tendon morphology (**Fig. 1f**,
170 **Extended Data Fig. 1d**).
171
To refine the length of candidate sequences, as histone ChIP-seq peaks are quite
lengthy, we also carried out ATAC-seq, which provides narrower defined peaks. For bat,
this was done on CS17 FL and HL as we only had available *M. natalensis* embryos for
this time point and for mouse at E13.5 to match that developmental time point. We
identified thousands of peaks in each condition and annotated those that overlap
H3K27ac ChIP-seq at a similar time point and that overlap between mouse and bat
(**Fig. 1g**). We used our ATAC-seq data as an additional layer to choose candidates for
our subsequent mouse assays. We next analyzed individual bat FL DA peaks. We first
searched for clusters of DA peaks near FL and HL DE genes. We found gene regulatory
DA peak clusters near the *HoxD* genes (*Hoxd10-12*)(**Fig. 1h**) and *Tbx3* for FL and
HoxC genes and *Tbx4* for HL (**Fig. 1i**), suggesting that these loci are important regions
for bat wing and hindlimb development respectively. *Tbx3* and *HoxC* loci were analyzed
further in the next results section.
185

186
Fig. 1: Comparison of RNA-seq and ChIP-seq data between mouse and bat.
a, Experimental Design. RNA-seq and ChIP-seq (H3K27ac and H3K27me3) were carried out on autopods from
forelimb (FL) and *hindlimb* (HL) at three developmental stages (E12.0, E13.0, and E13.5 in mouse and the
corresponding CS15, CS16 and CS17 in bat). ATAC-seq was conducted at E13.5 and CS17. **b**, Venn diagram
showing the number of differentially expressed genes between FL and HL in bats and mice. **c**, Comparison of FL vs.
HL differentially expressed genes in mice (left panel) and bats (right panel). The X and Y axes are normalized read
counts. The blue lines reference a log2-fold change of 1, i.e. deviation twice away from what would be expected

under the assumption that FL and HL expression are identical. **d**, MA plot of differential gene expression between
bat and mouse, considering the tissue (FL vs. HL) and stage (E12 through E13.5 and CS15 through C17). The X-axis
provides the mean normalized counts, placed against the model's log2-fold change on the Y-axis. The blue dots mark
a statistical significance of this differential gene expression across the species, tissue, and time stages. Similar to
panel C, the light blue horizontal lines reference a log2 fold-change of 1. Selected genes are marked by orange
circles and their gene symbols are given. Transcripts in the *Hoxc* and *Hoxd* clusters are within the green and blue
shades respectively. **e**, FL and HL differentially active (DA) ChIP-seq peak numbers for H3K27ac and H3K27me3 at
three stages of development for mouse compared to bat. The stage designation in the headers represent mouse and
bat, such that E12.0 is the equivalent bat developmental stage CS15. **f**, GREAT[26] analysis of H3K27ac and
H3K27me3 ChIP-seq DA peaks for CS15 FL and HL autopods, compared to the read count pattern of the
corresponding region in the mouse. No limb-related associated terms were found for CS15 HL H3K27me3 and hence
not shown. **g**-**h**, DA peaks in the *HoxD* (**g**) and *Tbx4* (**h**) locus with the following tracks: RNA-seq (pink), ATAC-seq
(blue), H3K27ac ChIP-seq (green) for E13.5 mouse and CS17 bat. The blue highlighted regions indicate DA peaks.
The Y-axis is similar for FL and HL in all tracks. **i**, ATAC-seq peaks in bats (CS17) or mice (E13.5) and their overlap
with H3K27ac ChIP-seq and each other.
209
Functional validation of candidate regions from computational analysis

We next selected three top ranking DA sequences between bat and mouse FL from our
genomic assays to characterize for mouse enhancer activity. We chose the following
three regions: 1) A previously characterized mouse E11.5 limb enhancer (mm1417,
VISTA Enhancer Browser[27]) in the homeobox C cluster (*HoxC*) which has high
H3K27ac and ATAC-seq peaks in the bat HL (**Fig. 2a**); 2) A sequence downstream to
T-box transcription factor 5 (*Tbx5*) which showed strong H3K27ac and ATAC-seq peaks
in the bat FL; 3) A previously characterized mouse E11.5 limb enhancer (hs483, VISTA
Enhancer Browser[27]) in the *Tbx3* locus that showed strong H3K27ac activity in the bat
FL.
220
All three *M. natalensis* bat sequences were cloned into a mouse enhancer assay vector
in front of the *Hsp68* minimal promoter and LacZ reporter gene. In addition, we also
cloned the corresponding mouse sequences for the *HoxC*, *Tbx5* and *Tbx3* regions. We
observed differences in limb enhancer activity for all three bat sequences, compared to
their corresponding mouse sequences at the specified embryonic stage (E12.5 to
E13.5) (**Fig. 2a-c**, **Extended Data Fig. 2a-c**). For *HoxC*, we observed limb enhancer
activity at E12.5 in the proximal posterior side in both FL and HL for both mouse and bat
sequences (**Fig. 2a**). For *Tbx5,* in some embryos we found that the bat sequence had
strong enhancer activity in the developing digits while the mouse sequence showed limb
enhancer activity in the base of the zeugopod (**Fig. 2b top**); however most embryos had
weak enhancer activity in the zeugopod for both bat and mouse sequences (**Fig. 2b
bottom**). For *Tbx3*, we observed weak enhancer activity for the mouse sequence while
the bat sequence showed strong enhancer activity in the anterior side of the autopod
(**Fig. 2c**).
235
Due to the strong enhancer activity differences between the bat and mouse *Tbx3*
sequences and the known role of this gene in early limb development[28], we swapped
the mouse sequence with the bat sequence using a modified protocol for the Precise
Integration into Target Chromosome (PITCh) system[29] that uses longer homology arms
(1 kb) and two gRNAs targeting the mouse homologous sequence and the bat enhancer
donor plasmid (**Fig. 2d**; see Methods for more detail). We next checked the

endogenous *Tbx3* gene expression level at E12.0 limb buds, as this time point is similar to the one we observed enhancer activity differences between the bat and mouse sequence. We observed no significant *Tbx3* expression differences between wild type (WT) and *bat-Tbx3 enhancer* homozygous replacement mice($bTbx3^{E/E}$) (**Fig. 2e**). We also checked for skeletal differences at E18.5 using alcian blue and alizarin red staining, finding no apparent differences (**Fig. 2f**). In summary, despite the strong DA peak differences and transgenic enhancer differences between bat and mouse for this sequence, we did not observe any apparent *Tbx3* expression changes at E12.0 or skeletal differences at E18.5.

Fig 2: Functional characterization of the bat *HoxC*, *Tbx5* and *Tbx3* enhancers.
a-c, Genomic landscape and lacZ enhancer transgenic assay. **Left:** RNA-seq (pink), ATAC-seq (blue), and H3K27ac ChIP-seq (green) data from bat or mouse FL and HL at E13.5 (mouse) or CS17 (bat). **Right**: LacZ enhancer

transgenic assay result. The number below the representative embryo picture indicates the number of embryos
showing similar enhancer activity. Bars below denote 1mm. The *HoxC* locus (**a**) shows mm1417 (red bar in genome
snapshot), a validated limb enhancer from the VISTA enhancer browser [30]. For *Tbx5* (**b**) two distinct staining patterns
were observed, with the one above showing strong limb staining and below the black arrows point to the weaker limb
staining pattern. For *Tbx3* (**c**) the genome browser shows hs483 (red bar in genome snapshot), a validated limb
enhancer from the VISTA enhancer browser [30]. **d**, The strategy used for bat enhancer replacement. Two gRNAs were
designed to target the mouse homologous region (orange) and the donor plasmid is targeted by one of these gRNA
(green). **e**, Gene expression levels of *Tbx3* in wild-type (WT) and *bTbx3*[E/E] homozygous knockin mice in FL and HL.
Each value represents the ratio of *Tbx3* gene expression to that of *b-Actin* (n=11–13 biological replicates), and
values are mean ± standard deviation. The expression value of wild type (WT) FL was arbitrarily set at 1.0. Each dot
represents one embryo, and statistical differences were determined using unpaired t test (N.S., not significant).
f, Alcian blue and alizarin red staining for wild-type (WT) and *bTbx3*[E/E] homozygous knockin mice. Bars below denote
1mm.
268
Functional validation of candidate regions from biologically relevant genes
We selected additional enhancer candidates for mouse enhancer assays, based on
their known involvement in wing development and/or webbing. These included a
sequence near the receptor interacting serine/threonine kinase 4 (*Ripk4*) gene that had
strong bat ATAC-seq and H3K27ac peaks in the FL and HL compared to mouse (**Fig.**
3a). Mutations in *RIPK4* cause Bartsocas-Papas type 1 popliteal pterygium syndrome
(BPS1; OMIM #263650), that is characterized by cleft lip and/or palate, webbing of the
skin in the knees, elbows and neck, syndactyly and genital and digit abnormalities [31]. In
addition, *Ripk4* homozygous mouse knockouts die at birth due to epidermal
differentiation abnormalities, including reduced skin folds, fusion of the tail and hindlimb
to the body and fusion of the oral cavity, esophagus and external orifices and a more
thick epidermis[32]. *Ripk4* was also found to be differentially expressed in the developing
bat plagiopatagium (membrane connecting the bat FL and HL) between an
insectivorous bat and omnivorous bat suggesting that it is important in shaping wing
structure[33]. We also selected a sequence near the gremlin (*Grem1*) gene, which is
known to play an important role in limb development[34,35] and interdigital membrane
retention in several mammals including bats[5]. *Grem1* showed high expression levels in
the bat FL compared to HL and the chosen sequence showed a FL specific ATAC-seq
peak (**Fig. 3b**). This sequence did not overlap any of the previously characterized
Grem1 limb enhancers[36].
289
We cloned both sequences into a mouse enhancer assay vector and analyzed their
LacZ expression at E13.5, as this was the time point that showed strong DA in the
ATAC-seq data. We observed strong reproducible expression in the digits and arm for
the bat *Ripk4* associated sequence while the mouse sequence only provided dim arm
expression (**Fig. 3c**, **Extended Data Fig. 3a**). For the bat *Grem1* associated sequence,
we observed enhancer activity in the limb proximal ventral region, while the mouse
sequence was negative for enhancer activity (**Fig. 3d**, **Extended Data Fig. 3b**).
297
298 Due to the inherent bat and mouse enhancer expression differences, we next swapped
299 both sequences in the mouse using the modified PITCh protocol[29]. We measured
300 mRNA expression for either gene at E13.0, finding a significant increase in *Ripk4*
301 expression in the FL of the *bRipk4*[E/E] mice in the whole limb bud (**Fig. 3e**). We analyzed
302 the *Grem1* expression in autopod (hand) and stylopod/zeugopod (arm) separately.

Grem1 was significantly upregulated in stylopod/zeugopod; however there was no significant difference in the autopod between the *Grem1* homozygous knockin (*bGrem1E/E*) and wild-type mice (**Fig. 3f**, **Extended Data Fig. 3c**). We next characterized bone development at E18.5 for both mouse lines using alcian blue and alizarin red staining, observing no observable differences between *bGrem1E/E* and wild-type mice (**Extended Data Fig. 3d**). In *bRipk4E/E* mice, we observed a complete absence of ossification in the intermediate phalanges compared to wild-type mice (**Fig. 3g-h**). In bats, delayed ossification in the FL was observed as a means to allow longer digit length[2–4].

As both *Ripk4* and *Grem1* have important roles in skin development, we next analyzed the ventral and dorsal skin region of the limb of homozygous knockin and wild type mice at P0 using hematoxylin and eosin (H&E) staining. We observed no significant differences in skin epidermis thickness for *bGrem1E/E* mice compared to wild-type mice (**Extended Data Fig. 3e**). In contrast, *bRipk4E/E* mice had a significantly thicker epidermal skin layer in the FL and HL ventral region and in the FL dorsal region (**Fig. 3i-j**).

To further characterize the molecular mechanisms driving this epidermal skin layer length difference, we conducted RNA-seq on *bRipk4E/E* and WT P0 FL ventral skin samples (**Fig. 3k**). Gene Ontology (GO)[37,38] term enrichment for DE genes between *bRipk4E/E* and WT mice found significant enrichment for "skin development" and "positive regulation of epidermis development". One of the top DE genes was the Fos proto-oncogene (*Fos*) that showed 6.7 fold higher expression in *bRipk4E/E* mice (**Extended Data Fig. 4**). Inducible expression of *c-fos* in the epidermis of adult mice is known to promote inflammation-mediated epidermal hyperplasia[39]. Other DE genes included the ERBB receptor feedback inhibitor 1 (*Errfi1*, also known as *Mig6i)*, a protein that regulates the epidermal growth factor receptor (EGFR) pathway and is involved in skin development[40] and the Transcription Factor AP-2 Alpha (*Tfap2a*, also known as *AP-2α*) whose loss leads to altered epidermal morphology[41]. We also found the fibroblast growth factor 2 (*Fgf2*) which binds to the fibroblast growth factor receptor 2 (Fgfr2) in keratinocytes to trigger a signaling cascade that stimulates cell division, promoting epidermal growth and thickness[42] to be DE. Combined, our RNA-seq results identified numerous DE genes which could contribute to a thickened epidermis layer in *bRipk4E/E.*

340
Fig 3: Mouse characterization of bat *Ripk4* and *Grem1* enhancers.
a-f, Genomic landscape and lacZ enhancer transgenic assay. (**a-b**) RNA-seq (pink), ATAC-seq (blue), and H3K27ac
ChIP-seq (green) data from bat or mouse FL and HL at E13.5 (mouse) or CS17 (bat). (**c-d**) LacZ enhancer
transgenic assay result. The number below the representative embryo picture indicates the number of embryos
showing similar enhancer activity. Bars below denote 1mm. (**e-f**) Gene expression levels of *Ripk4* and *Grem1* in wild-
type (WT) and bat enhancer homozygous knockin mice in FL and HL as measured by qRT-PCR. Each value
represents the ratio of *Ripk4* and *Grem1* gene expression to that of *b-Actin* (*n*=6-10 biological replicates), and values
are mean ± standard deviation. The expression value of wild type (WT) FL was arbitrarily set at 1.0. Each dot
represents one embryo. **g,** Alcian blue and alizarin red staining for wild-type (WT) and *bRipk4^E/E* homozygous

replacement mice. Bars below denote 0.5mm. **h,** Ossification zone length of 4th digit intermediate phalanx at E18.5.
Each dot represents one embryo. Biological replicate numbers are shown beneath the graph. **i**, Epidermal skin layer
thickness in the FL and HL, measured in both ventral and dorsal regions at P0. Each dot represents one mouse. **j,**
H&E staining of FL ventral side skin. Black arrow indicates the epidermis layer. Bars above denote 50μm. **k,** Heat
map showing gene expression differences between WT and *bRipk4^E/E^* homozygous replacement mice at P0, based
on RNA-seq. Differentially expressed genes associated with the Gene Ontology terms 'skin development' or 'positive
regulation of epidermis development' are shown. All statistical differences were determined using unpaired t test (*$p <$
0.05, **$p < 0.01$, ***$p < 0.005$, ****$p < 0.001$, N.S., not significant).

Functional validation of bat accelerated regions

We next characterized BARs near genes involved in limb development. These included:
1) BAR116, a sequence in the *Hoxd* locus (**Extended Data Fig. 5a**) that we previously
showed to have strong FL expression in the entire autopod at E12.5 while the mouse
sequence was negative for enhancer activity[17]. It is also worth noting, that the
homologous BAR116 chicken sequence showed increased FL expression compared to
the HL in mouse enhancer assays at E12.5[43]. 2) The sonic hedgehog (*Shh*) limb
enhancer (BAR61), also called the zone of polarizing activity (ZPA) regulatory sequence
(ZRS) or the mammal-fish-conserved-sequence 1 (MFCS1)[44], did not show significant
expression differences between the *M.natalensis* bat and mouse sequences in a
transgenic mouse enhancer assay at E12.5[17] but showed extended activity at the
anterior edge of the limb for the *Pteropus Vampyrus* (megabat) sequence at E11.5
compared to mouse[45]. The ZRS is associated with a variety of limb malformations
including polydactyly, triphalangeal thumb and syndactyly in humans[46], limb loss in
snakes[45] and limb malformations in chickens[47]. 3) Two adjacent BARs near the msh
homeobox 2 (*Msx2*) gene[18] that along with *Msx1* plays an important role in
mesenchyme interdigit cell death during limb development[48,49] and both genes were
found to have significantly lower expression in the embryonic bat FL compared to the
HL[13].

As mouse enhancer assays for the bat and mouse sequence were previously carried
out for BAR116 and ZRS (BAR61), we swapped these two sequences. For BAR116, we
did not observe any gene expression differences via qRT-PCR in FL for *Hoxd9, 10, 11,*
12, and *13* at E12.5, the time point where the bat and mouse showed differential
enhancer activity[17], between the homozygous knockin and wild-type mice, but observed
some differences in the HL for *Hoxd9, Hoxd10,* and *Hoxd12* (**Extended Data Fig. 5b**).
However, we did not observe any apparent skeletal differences in the FL and HL at P1
using alcian blue and alizarin red staining (**Extended Data Fig. 5c**). In contrast, for
*bZRS^E/E^*, we observed polydactyly in the HL in heterozygous mice that became more
symmetrical and 100% penetrant in homozygous mice, similar to a bat HL (**Fig. 4a**). We
also observed a shortened tibia (**Fig. 4b**). In the FL, we observed lower levels of mild
polydactyly (55% of the mice) (**Fig. 4b**). qRT-PCR on the E11.5 limb showed *Shh* to be
significantly upregulated only in the HL (**Fig. 4e**). Whole-mount *in situ* hybridization
(WISH) for *Shh* and its receptor patched 1 (*Ptch1*) found both to show ectopic
expression in the anterior HL (**Fig. 4c-d**). We also analyzed previous *Shh* WISH on bat
M. natalensis embryos[8] at CS14E (equivalent to E11.0 in mice) observing similar
ectopic expression in the anterior part of the HL in two different embryos (**Fig. 4f**,
Extended Data Fig. 5d).

We next carried out RNA-seq on both WT and *bZRS^{E/E}* E11.5 whole limb buds finding

numerous genes involved in embryonic limb morphogenesis and anterior/posterior

pattern specification to be differentially expressed (**Fig. 4g-h**). We observed

upregulation of the Meis Homeobox 2 (*Meis2)* and *Grem1* genes that are known to be

upregulated in the bat FL[50,51]. We also saw the ALX Homeobox 4 (*Alx4*) gene which is a

downstream factor of Shh signalling [52] to be upregulated in *bZRS^{E/E}*. The zinc finger and

BTB domain-containing protein 16 (*Zbtb16*, also known as *Plzf*), a transcription factor

involved in the development of the anterior-posterior axis in limbs[53] was also found to be

upregulated. *Dlx5;Dlx6* double mutants exhibit hindlimb ectrodactyly [54] and both these

genes were found to be downregulated in *bZRS^{E/E}*. In summary, we found several limb

developmental associated genes to be differentially expressed in *bZRS^{E/E}* E11.5

embryos.

410
Fig 4: Mouse characterization of bat ZRS.
a, Hindlimb phenotype in *bZRS^E/E*. Hindlimb morphology in WT, *bZRS^E/W*, *bZRS^E/E*, and *Artibeus jamaicensis*.
Polydactylous digits are indicated as poly. **b,** Alcian blue and alizarin red staining for wild-type (WT) and *bZRS^E/E*
homozygous replacement mice. Bars below denote 0.5 mm. Percentage noted on the *bZRS^E/E* panel is the
penetrance of this phenotype. **c-d,** Whole-mount *in situ* hybridization for *Shh* (**c**) and *Ptch1* (**d**) of E11.5 mouse
embryos. Black bar represents 1 mm (whole embryo) and 0.5 mm (limb bud). **e,** Gene expression levels of *Shh* in
wild-type (WT) and bat enhancer homozygous replacement mice in FL and HL as measured by qRT-PCR. Each
value represents the ratio of *Shh* gene expression to that of *b-Actin* (*n*=3 biological replicates), and values are mean
± standard deviation. The expression value of wild type (WT) FL as arbitrarily set at 1.0. Each dot represents one
embryo, and statistical differences were determined using unpaired t test (**p* < 0.01, N.S., not significant). **f,** Whole-
mount *in situ* hybridization for *Shh* of CS14 early *Miniopterus natalensis* embryo. Black bar represents 1 mm (whole
embryo) and 0.2 mm (limb bud). Pink arrowhead indicates the *Shh* gene expression in HL. **g,** Volcano plots showing
the global transcriptional changes for the indicated groups. Each circle represents one gene. The log2 fold change in

the indicated genotype is represented on the x axis. The y axis shows the p value. A p value of 0.05 and a fold
change of 2 are indicated by lines. **h,** Heat map showing gene expression differences between WT and *bZRS^{E/E}*
homozygous knock-in mice at E11.5, based on RNA-seq. Differentially expressed genes associated with the Gene
Ontology terms 'Embryonic limb morphogenesis' or 'Anterior/posterior pattern specification' are shown.

For the two adjacent *Msx2* BARs (**Fig. 5a**), we first carried out a bat and mouse
enhancer assay, cloning both of them together as they were adjacent to each other,
finding that the mouse *Msx2* sequence has stronger enhancer activity at E13.5 in the
distal autopod while the bat sequence showed reduced expression in this region (**Fig.**
5b, **Extended Data Fig. 6a**). We next swapped this sequence in mice with the bat
sequence using the modified PITCh protocol[29]. We found *BAR-Msx2* homozygous mice
(*bMsx2^{E/E}*) to have significantly reduced expression of *Msx2* in the FL and HL compared
to WT mice, with the FL showing a stronger difference (**Fig. 5c**). Analysis of bone
ossification using alcian blue/alizarin red staining at E18.5 showed reduced ossification
of the 4th intermediate phalanx for *bMsx2^{E/E}* compared to wild-type embryos; however,
this was not a statistically significant reduction (**Fig. 6a-b**).

Limb malformations are only observed when both *Msx1* and *Msx2* genes are altered but
not individually[55], likely due to functional redundancy. Deleting *Msx2* conditionally in the
limb using a *Prx1* Cre that drives expression in the limb mesenchyme along with *Msx1*
knockout mice showed impairment of the apical ectodermal ridge (AER) and
subsequent mesenchyme along with digit abnormalities including preaxial polydactyly[56].
We thus carried out mouse transgenic enhancer assays for five putative *cis*-regulatory
elements bat sequences spread broadly over 100 kb of intergenic regions in the bat
Msx1 locus (**Extended Data Fig. 7a**). We did not observe any differences in activity
between FL and HL (**Extended Data Fig. 7b**). We thus set out to test whether placing
bMsx2^{E/E} mice on an *Msx1* heterozygous gene knockout background instead of
enhancer replacement will exacerbate the limb phenotype. Using the *i*-GONAD
protocol[57], we generated *Msx1* heterozygous gene knockout mice (*Msx1^{+/-}*). *Msx1*
homozygous mice are known to die at birth[58]. We then crossed *Msx1^{+/-}* gene knockout
mice to *bMsx2^{E/E}* mice obtaining *bMsx2^{E/E}; Msx1^{-/-}* compound mice. As *Msx1*
homozygous knockout mice die at birth, we carried out alcian blue/alizarin red staining
at E18.5 finding these mice to have significantly reduced ossification in the 4th
intermediate phalanx compared to *bMsx2^{E/E}* or *Msx1^{-/-}* mice (**Fig. 6a-b**). These results
suggest that lower *Msx1* expression levels leads to reduced compensation for
differential expression of *Msx2*, leading to a stronger limb ossification phenotype.
460

461
Fig 5: Mouse characterization of the bat *Msx2* enhancer.
a, Genomic landscape of *Msx2* locus. RNA-seq (pink), ATAC-seq (blue), and H3K27ac ChIP-seq (green) data from
bat or mouse FL and HL at E13.5 (mouse) or CS17 (bat). **b,** LacZ enhancer transgenic assay at E13.5 result. The
number below the representative embryo picture indicates the number of embryos showing similar enhancer activity.
Bars below denote 1mm. **c,** Gene expression levels of *Msx2* in WT and bat enhancer homozygous knockin mice in
FL and HL as measured by qRT-PCR. Each value represents the ratio of *Msx2* gene expression to that of *b-Actin*
(n=7-11 biological replicates), and values are mean ± standard deviation. The expression value of wild type (WT) FL
was arbitrarily set at 1.0. Each dot represents one embryo, and statistical differences were determined using unpaired
t test (***$p < 0.005$, *$p < 0.05$, N.S., not significant).

Compound mice show bat associated phenotypes

We next set out to test whether combining bat-mouse swapped alleles could have an
additive effect on phenotype. As both *bRipk4[E/E]* and *bMsx2[E/E]; Msx1[-/-]* mice showed
delayed bone ossification in the digits, we mated these lines to one another. We
observed that *bRipk4[E/E]; bMsx2[E/E]; Msx1[-/-]* mice (triple compound mice) had significantly
reduced ossification at E18.5 (**Fig. 6a-b**). In bats, delayed ossification in the FL was
observed as a means to allow for longer digit length[2,3]. Additionally, *Msx2* knockout
mice showed defective chondrogenesis and endochondral ossification[59,60]. We thus
examined the length of the 4th intermediate phalanx in WT and triple compound mice
FL and HL from E18.5 to P6 using alcian blue and alizarin red staining of bone
specimens, as this is one of the bones showing significant differences during the
neonatal period. As *Msx1* homozygous knockout mice are lethal at birth we used *Msx1*
heterozygous mice (*Msx1[+/-]*) for these analyses. Forelimb bone length was comparable
or slightly shorter from E18.5 to P1, increased at P2 and P4, and showed no difference
by P6 (**Fig. 6c**). The HL showed a similar trend, but no significant difference was
observed from E18.5 to P4 (**Fig. 6c**). We measured all bone lengths at P2 and P4 in
bRipk4[E/E] and triple compound mice, and compared with WT (**Fig. 6d**). Although both
bRipk4[E/E] and triple compound mice showed delayed bone ossification in the digits,
bone length differences were more pronounced in the triple compound mice (**Fig. 6d-e**).
bRipk4[E/E] mice had shorter distal phalanges compared to WT at P2 and P4. In contrast,

triple compound mice showed increased bone length at P2 in multiple bones, including
the phalanges, humerus, femur, and fibula (**Fig. 6d-e**). Four forelimb phalanges and
one hindlimb phalanx remained longer at P4 (**Fig. 6d-e**).
495
To identify genes involved in these differences, RNA-seq was carried out on wild-type,
bRipk4^{E/E} and *bRipk4^{E/E}; bMsx2^{E/E}; Msx1^{+/-}* mice from P4 digits (**Fig. 6f-g**). In *bRipk4^{E/E}*,
we observed upregulation of *Krtap 5-1, 2, 3* and *4* genes that are major components of
hair and nails[61,62] (**Fig. 6f**). We also saw downregulation of ossification related genes,
such as *Osr1* and *Osr2*. Dominant negative *Osr2* transgenic mice, in which the function
of both *Osr1* and *Osr2* genes is inhibited, show delayed mineralization in calvarial and
tibial bone tissues[63]. *Smad3* and *Samd6* were slightly down- and up-regulated
respectively. TGF-β/Smad3 signals are known to repress chondrocyte hypertrophic
differentiation[64] and overexpression of *Smad6* significantly delays chondrocyte
hypertrophy, which can lead to postnatal dwarfism with osteopenia[65]. *Col10a1* is a
marker gene of hypertrophic chondrocytes[66,67] that was also found to be upregulated.
These results suggest there are perturbations of the endochondral ossification process
in digits.
509
In the triple compound mice, we observed upregulation of skeletal development-related
genes, including *Hoxa7, Cnmd, Foxo3* and *Eya1*, as well as extracellular matrix (ECM)
genes enriched in growth plate chondrocytes, such as *Col9a1*, *Col9a3, Col10a1* and
Matn3[67,68], along with structural and matrix-organizing ECM components *Col11a1* and
Hapln1(**Fig. 6g**). In the context of endochondral ossification, *Cnmd* (ChM-1;
Chondromodulin-I) is a negative regulator of cartilage vascularization, maintaining
avascular cartilage and thereby controlling the timing of ossification[69]. *Foxo3* inhibits
Wnt signaling and delays ossification[70]. Overall, our transcriptomic analyses show
inhibited Wnt signaling and delayed ossification, marked by retained hypertrophic
chondrocytes, upregulation of collagen genes, and a potential role in prolonged bone
elongation in triple compound mice.
521

523 **Fig 6: Characterization of compound mice carrying multiple bat enhancers**

524 **a,** Alcian blue and alizarin red staining for WT, *bMsx2^E/E^*, *Msx1^-/-^*, *bMsx2^E/E^; Msx1^-/-^* mice. Bars below denote 0.5mm.

525 Arrowheads indicate regions of delayed ossification. **b,** Ossification zone length of 4th digit intermediate phalanx at

526 E18.5. Each dot represents one embryo. Biological replicate numbers are shown beneath the graph. **c,** Transition of

527 the 4^th^ intermediate phalanx length from E18.5 to P6. **d,** Schematic representation of each bone length difference.

Statistical significant P2 and P4 bone length differences between WT and *bRipk4$^{E/E}$* or WT and *bRipk4$^{E/E}$; bMsx2$^{E/E}$;*
Msx1$^{+/-}$ are shown. **e,** Bone lengths in P2 or P4 WT, *bRipk4$^{E/E}$* and *bRipk4$^{E/E}$; bMsx2$^{E/E}$; Msx1$^{+/-}$*. Each dot represents
one embryo (*n* = 5-9 biological replicates). **f** (wild-type vs *bRipk4$^{E/E}$*) **-g** (wild-type vs *bRipk4$^{E/E}$; bMsx2$^{E/E}$; Msx1$^{+/-}$*
mice)**,** Left: Volcano plots showing the global transcriptional changes for the indicated groups digits at P4. Each circle
represents one gene. The log2 fold change in the indicated genotype is represented on the x axis. The y axis shows
the *p* value. A *p* value of 0.05 and a fold change of 2 are indicated by lines. Right: Heat map showing gene
expression differences between WT and bat enhancer replacement mice, based on RNA-seq. All statistical
differences were determined using an unpaired t test (*$p < 0.05$, **$p < 0.01$, ***$p < 0.005$, ns, not significant, nd, not
determined).
537

Discussion

The bat wing is a unique organ that underwent numerous morphological changes during evolution to allow for powered flight. Here, we used bat and mouse embryos to identify in a genome-wide manner numerous genes and regulatory elements that show unique activity in the developing bat FL. Using mouse transgenic enhancer assays, we identified several sequences that show differential limb enhancer activity between the bat and mouse sequence. We then swapped six sequences in the mouse genome with the corresponding bat sequence finding relevant morphological changes associated with both growth and patterning aspects of the bat limb phenotype, including delayed bone ossification, slightly longer digits, increased epidermal layer thickness and symmetrical hindlimbs. Combining swapped lines that affected bone ossification had a small but significant additive effect on the phenotype within a specific neonatal time window and led to longer phalanges, humerus, femur, and fibula.

Our genomic datasets (RNA-seq, ChIP-seq, and ATAC-seq) provide useful catalogs for bat limb development. By generating matching developmental time points for mouse FL and HL these datasets can be scanned for differential gene expression and regulation across these two species. We focused our analyses on differential activity of homologous sequences between the mouse and bat genome. However, in the future it will be intriguing to also characterize sequences that are either unique to bats or missing in bats but appear in other mammals, which could also be associated with limb development. In addition, further analyses of noncoding RNAs, such as miRNAs and lncRNAs, that are associated with bat wing development would also be of interest.

The bat *Ripk4* regulatory sequence led to the strongest phenotypic effect from an individual swapped sequence. This included both a delay in bone ossification, mild changes in bone length and a thicker epidermal layer. Wnt signaling has been shown to play an important role in limb organogenesis[71]. Our previous bat FL and HL genomic characterization[13] showed an important role for this pathway in wing development with higher levels of canonical Wnt pathway antagonists and an overall suppression of the canonical Wnt/β-catenin pathway in the bat FL compared to HL and an increased expression of canonical Wnt receptors in the bat HL[13]. Ripk4 phosphorylates Dishevelled (Dvl) which leads to an increase in β-catenin accumulation and promotes the activation of the canonical Wnt signaling[72]. We observed increased activity for the bat *Ripk4* enhancer in the developing digits of both FL and HL, likely leading to increased Wnt activity. Mouse knockouts for *Ripk4* show shorter limbs, fusion of digits and reduced skin folds[31]. Thus, increased activity of *Ripk4* and subsequently the Wnt pathway will likely lead to an opposite phenotype of longer limbs and thicker skin folds, as we observed in *bRipk4$^{E/E}$* mice. While we characterized the limb bud , *Ripk4* has also been shown to be highly expressed in the bat patagium[33] and could also have an important role in shaping this structure. Our genomic data showed additional regions to be differentially active between bats and mice in this locus (**Fig. 3a**), suggesting that

581 additional regulatory elements might be involved in various wing associated
582 phenotypes.
583
The triple combined line, *bRipk4^{E/E}; bMsx2^{E/E}; Msx1^{+/-}* had small additive effects on
phenotype with mice exhibiting a more pronounced phenotype than *bRipk4^{E/E}* by itself at
P2 and P4. These included delayed bone ossification in the digits and increased bone
length including the phalanges, humerus, femur, and fibula (**Fig. 6d-e**). Bats are known
to have delayed ossification in the forelimb[3]. RNA-seq analyses of these triple
compound mice showed increased inhibition of the Wnt pathway, which is known to
contribute to delayed ossification[73,74]. Endochondral ossification is a key process in
longitudinal bone elongation, characterized by an increase in chondrocyte number and
size, followed by matrix calcification and subsequent replacement of cartilage with bone,
ultimately forming trabecular bone. Our results suggest that multiple regulatory factors
involved in the attenuation of Wnt signaling might be responsible for delayed ossification
in bats. In terms of longitudinal bone elongation, not only delayed ossification but also
increasing both the size and number of chondrocytes are essential[75,76]. Further work
could explore the factors involved in chondrocyte phenotype to drive bone elongation.
598
Msx2 is known to be involved in endochondral ossification and stimulate the maturation
of chondrocytes into the prehypertrophic and hypertrophic stages[59,60]. However, in our
study, we focused on earlier developmental stages, and indeed, our *bMsx2^{E/E}* mice
showed gene expression changes at E13.5 but not at P4, during the endochondral
ossification stage. This suggests that other putative regulatory elements may be
involved in the stage-specific regulation of *Msx2*. Based on our genomics dataset, we
did not detect differential regulatory regions associated with key limb development
genes such as *Fgf*, *Bmp*, *Pthlh*, *Sox9*, or *Ihh*. There are several potential reasons for
this: 1) Our targeted developmental stages may be too early to capture *Pthlh*, *Sox9*, or
Ihh activity; 2) We used bulk RNA-seq, which may not identify expression changes in
smaller cell populations, such as localized expression of *Fgf8* or *Shh*. It is also worth
noting that a recent study that carried out single-cell RNA-seq on developing mouse and
bat (*Carollia perspicillata*) FL and HL at E11.5, E12.5 and E13.5 in mice and CS15 and
CS17 in bats did not observe differential expression in *Fgf* or *Bmp*[51]. From our results
we did have an observable limb phenotype in *bZRS^{E/E}* mice, while this sequence did not
show any significant differences in our genomics data; and 3) gene expression changes
during early development may lead to phenotypic differences at later stages. For
example, in the triple compound mice, we did not detect differential expression of *Msx2*
or *Ripk4* at P4, despite clear differences in ossification and bone length at that stage.
This highlights the complexity and importance of capturing dynamic gene regulation
within specific developmental time windows, which remain a challenge but also an
opportunity for future temporal-resolution genomics.
621
Interestingly, swapping the mouse sequence with a bat accelerated region, BAR116,
that showed strong FL enhancer activity for the bat sequence[17] and chicken sequence[43]
but was negative for the mouse sequence did not lead to an observable phenotype. The

HoxD locus contains a cluster of nine *HoxD* genes that are regulated in a
spatiotemporal manner by multiple enhancers whose chromatin interactions are thought
to be regulated by two distinct topologically associating domains (TADs). When a distal
Hoxd limb enhancer is inserted into the proximal limb TAD, its distal limb-specific
activity is suppressed in this ectopic context[77], suggesting that the native chromatin
environment can mask the activity of exogenously inserted enhancers even in the same
Hoxd locus. To overcome this limitation and more accurately recapitulate endogenous
cis-regulatory function in bats, strategies such as TAD-level structural reconstruction or
the use of longer transgenes (e.g., BAC/YAC constructs) might be needed.
634
For the ZRS, we primarily obtained a bat-HL phenotype. This sequence was chosen
due to it being a BAR. However, we did not observe strong differences between bat and
mouse ZRS enhancer activity and also between bat FL and HL in our genomic analyses
(**Extended Data Fig. 8a**). However, we reasoned that due to the strong phenotypic
effects of point mutations in this enhancer that lead to a multitude of limb developmental
phenotypes in humans, mice and other mammals[46,78,79], we would likely observe a limb
development associated phenotype. Interestingly, while we did not obtain a strongly
penetrant FL-associated phenotype, we did observe a HL phenotype in *bZRS* mice that
included polydactyly and the formation of symmetrical HL digits.
644
A key region that differs between bats and mice in the ZRS is a known binding site for
the ETS translocation variants 4 and 5 (Etv4/5) proteins. Posterior *Shh* gene expression
in the limb bud is regulated by a balance between transcriptional activators of the E26
transformation-specific (ETS) family and the repressors Etv4/5[80]. The ZRS contains
multiple binding sites for both ETS and ETV factors (**Extended Data Fig. 8b**). *ETS1/2*
are predominantly expressed in the posterior region of the limb bud, while *Etv4/5* are
expressed throughout the distal limb bud, encompassing both anterior and posterior
regions. The spatial expression patterns of these factors define the posterior-specific
expression of *Shh*. Mutations in ETV binding sites can disrupt this balance, leading to
ectopic activation of *Shh* in the anterior limb bud[80]. For example, conditional knockout of
Etv5 in mesoderm-derived cells using *Brachyury* (*T*)-*Cre* leads to anterior reactivation of
Shh and results in polydactyly specifically in the hindlimbs, mirroring the phenotype
observed in *bZRS* mice[81]. In the *bZRS* sequence, we identified sequence differences
within the ETV-A site and its surrounding region, which overlap with variants previously
linked to triphalangeal thumb polydactyly[82–85] (**Extended Data Fig. 8b-c**). Notably, the
rs606231152 variant (295T>C, represented as A>G in the figure due to reverse
complement orientation) is highly dominant over the wild-type allele in families with
triphalangeal thumb polydactyly. Intriguingly, most bat species examined also carry the
A>G substitution at this site (**Extended Data Fig. 8b-c**). Testing the human ZRS
sequence with the rs606231152 A>G change in a mouse enhancer assay showed that it
leads to anterior enhancer activity[85]. In addition, it was shown that mutating the ETV-A
site alone is insufficient to drive anterior enhancer activity in a LacZ transgenic assay[80],
suggesting that multiple mutations within the ZRS, including variants like rs606231152,
are likely required to induce robust anterior enhancer function. Interestingly, a recent

study highlighted the importance of a low-affinity ETS binding site (ETS-A)[86], which we identified as partially overlapping with the aforementioned ETV-A site[80] (**Extended Data Fig. 8b**). High-affinity mutations in ETS-A lead to anterior reactivation of *Shh* and result in a similar hindlimb polydactyly phenotype as observed in our *bZRS* mice. It is possible that ETS binding to ETS-A interferes with ETV-A binding, suggesting that competitive binding and the balance between ETS and ETV factors, particularly in overlapping ETS-A/ETV-A sites, could play a crucial role in this phenotype.

Our results suggest that the bat wing arose from numerous small molecular changes, in line with past work showing that morphological novelty in evolution likely arose from numerous changes[23]. In bats, comparative evolutionary genomics is particularly difficult as there are no intermediate species from the fossil records[87] thus restricting the ability to decipher the morphological skeletal evolution that leads to bats. Recent work comparing bird and bat limb evolution by analyzing hundreds of skeletal structures suggests that the bat FL and HL evolved in unison in contrast to birds where they evolved independently thus restricting each other's development and morphology[6]. Thus, it could be that analyzing changes driving both FL and HL phenotypes could identify sequences that are more likely to drive wing development. As mentioned above, our top phenotype altering sequence, *bRipk4*, showed increased enhancer activity in both FL and HL.

Our study had several limitations. We focused our genomics work on three bat developmental time points, thus potentially missing earlier or later time points that could also be important in wing development. We selected sequences primarily in a biased manner, focusing on those that are thought to regulate known genes involved in limb development and malformations. We only changed the targeted mouse genomic sequences to one specific bat species, *M. natalensis*. Other bat sequences could have a different effect, due to the large diversity of wing morphology in bats. To circumvent this, we could have predicted the last common ancestor sequence of bats and swapped this sequence in, but it may not have represented a current existing bat. As bat genome editing is currently not possible, we modeled the hypothesised genomic changes in mice whose chromatin architecture or 'trans' environment could have a differential effect on limb development and might not be a good model to assay for limb elongation, as to our knowledge, there are very limited mouse mutants that have an elongated limb phenotype. Finally, we focused our work on individual alleles and only generated one combined allele.

In summary, our work provides a detailed genomic catalog of candidate genes and putative *cis*-regulatory elements involved in bat FL and HL development, compared with corresponding stages in mice. Through extensive *in vivo* functional analyses, we demonstrate the phenotypic relevance of several of these elements, highlighting the role of enhancers in bat wing morphogenesis and suggesting that multiple regulatory changes contributed to the emergence of the bat wing, one of the most eminent evolutionary adaptations in mammals.

Methods

RNA-seq

RNA was extracted from paired forelimbs and hindlimbs from three C57BL/6J (Jackson labs, 000664) embryos (biological replicates) at three developmental stages (E12.0, E13.0, E13.5) using the RNeasy Midi (Qiagen) kit. Total RNA samples were enriched for poly-A containing transcripts using the Oligotex mRNA Mini kit (Qiagen) and strand-specific RNA-seq libraries were generated using PrepX RNA library preparation kits (IntegenX) following the manufacturer's protocol. After clean up with AMPure XP beads (Beckman Coulter) and amplification with Phusion High-Fidelity polymerase (NEB), RNA libraries were sequenced on a HiSeq 2500 to a depth of at least 30 M reads. All mouse work (other than ATAC-seq which was approved by the University of Cape Town) was approved by the UCSF Institutional Animal Care and Use Committee (IACUC) protocol number AN197608. Sequence alignment was performed using STAR[88]. Mappings were restricted to those that were uniquely assigned to the mouse genome; unique read alignments were used to quantify expression and aggregated on a per-gene basis using the Ensembl GRCm37.67 annotation. We analysed these raw data with DESeq2[89] to assess variance and differential expression.

ChIP-seq

C57BL/6J (Jackson labs, 000664) mouse forelimbs and hindlimbs dissected from E12.0, E13.0, E13.5 were cross-linked with 1% formaldehyde for 10 minutes, quenched with glycine and flash frozen. Cross-linked limbs were then combined into pools of 4-7 pairs per stage for chromatin sheering using a Covaris S2 sonicator. Sheared chromatin was then used for chromatin immunoprecipitation (ChIP) with antibodies against active (anti-H3K27ac; Abcam ab4729) or repressed (anti-H3K27me3; Millipore 07-449) chromatin marks using the Diagenode LowCell# ChIP kit following the manufacture's protocol. Libraries were prepared using the Rubicon ThruPLEX-FD Prep Kit following the manufacturer's protocol and sequenced on an Illumina HiSeq2500 using single end 50 bp reads to a sequencing depth of at least 25 M. Raw sequence files were aligned from fastq to mm9 or mnat_v1 bam format using samtools[90] and a pre-built annotation index of the respective assembly. ChIP-Seq files include an input channel for each species, stage, and tissue. This allowed us to obtain increased accuracy when calling the peaks. Peak calling was done two-fold. First, using MACS v2[91,92] we called the peaks on each of the channels separately, then with elaborate background processing. In the first step, MACS[91], we used the following parameters: *macs2 -n file-name -t file-name.sort.bam -c input-file.sort.bam -g mm --nomodel --pvalue=5e-5 --slocal=5000 --llocal=20000 --shiftsize=300* for H3K27ac. We used *macs2 -n file-name -t file-name.sort.bam -c input-file.sort.bam -g mm --nomodel --pvalue=1e-4 --slocal=10000 --llocal=50000 --shiftsize=500* for H3K27me3. In the bat we used the same parameters for either antibody: *--nomodel --pvalue=1e-5 --slocal=5000 --llocal=20000 --shiftsize=30*0. Later, we added background processing from[93] to improve the ability to align the called peaks across the channels. For example, peaks called on E12.0FL and E12.0HL are expected to misalign. Using the background analysis of Diaz et al and the

background profiling provided by deepTools[94], we were able to orient and identify the
same called peaks. After peak calling for each channel we aligned the peaks across the
species. Misaligned peaks could be missing from the other species but as we do not
having any prior information about the nature of these regions, we did not use them in a
differential analysis context. To characterize regions across the species, we used
lifOver[95,96] allowing multiple targets to be listed in the output. After obtaining all the
lifted-over targets, we employed a search algorithm for synteny anchors on either side
of each peak. To match the best candidate peak across the two species, the algorithm
was required to observe two synteny blocks in the flanks of each peak (see **Extended**
Data Fig. 9). We then used the following statistical analysis to list the differential
chromatin (Differentially Active) regions across bat and mouse. We created a design
that accounts for chromatin activity levels in all the possible combinations of species (2 -
mouse vs. bat), tissue (2 - FL vs. HL), and stage (3 - E12.0, E13.0, or E13.5). We used
the R package DESeq2[89], which we customized to use counts estimated from FPKM
using the normalization method due to[93], then used these numbers only on aligned
regions. The count data were averaged across the replicates, and we then performed
Maximum Likelihood Estimation (MLE) given two models; a simple model where the
main effects considered are *Species + Stage*, and an elaborate model whose main
effects are further enhanced by interactions: *Species + Stage + Tissue + Tissue:Stage*
+ Tissue:Species. The MLE between the two models that was marked after FDR-
correction as statistically significant, gave the differential chromatin (DA) regions listed
in the supplementary master_features_v3_pvalSort_20230806.xlsx available in Zenodo.
779
ATAC-seq
Mouse embryos for ATAC-seq (C57BL/6 strain UCT3) were supplied by the Animal
Research Facility, University of Cape Town. Ethical approval for their use was granted
by the Health Science Animal Ethics Committee (FHS AEC014/07). *Miniopterus*
natalensis embryos were collected from wild-caught, pregnant females in 2016 and
2017 from De Hoop Nature Reserve, Western Cape Province, South Africa (Western
Cape Nature Conservation Board permit number: AAA007-00133-0056). Approval for
the use of the bat embryos was granted by the Science Faculty Animal Ethics
Committee (2012/V41/NI and 2012/V39/NI). Limbs were dissected from embryos and
forelimbs and hindlimbs from the same embryo were pooled separately and placed in
solutions of 0.25%Trypsin/0.1M EDTA:HBSS at 1:1 or 1:3 dilution and incubated at 4°C
overnight. For all E13.5 mouse limbs, limbs from 4 embryos were pooled. Samples were
centrifuged at 250 x g for 3 minutes, all but 1 ml of solution removed, and incubated at
37°C for 25 minutes. A further 1 ml of 0.25% trypsin/0.1M EDTA was added, followed by
incubation at 37°C for 100 minutes. Limbs were dissociated by gentle pipetting and
centrifuged at 250 x g for 5 minutes to pellet the cells. The solution was removed and
pellets were washed in Hanks followed by a further centrifugation at 250 x g for 5
minutes. Cells were resuspended in DMEM with 10% fetal calf serum and quantified
using a haemocytometer. 50,000 cells were centrifuged at 500 x g for 5 minutes at 4°C,
washed twice with ice cold 1x PBS and centrifuged for a further 5 minutes. The cell
pellet was resuspended in cold lysis buffer (10 mM Tris-HCl, pH7.4; 10 mM NaCl; 3 mM

MgCl2; 0.1% IGEPAL CA-630), and centrifuged for a further 10 minutes at 500 x *g* at 4°C. The supernatant was discarded and the cell nuclei resuspended in a transposition reaction mix [25 µl 2x TD Buffer (C-121-1030, Illumina); 2.5 µl Tn5 Transposase (FC-121-1030, Illumina); 22.5 µl Nuclease Free H2O]. The reaction was incubated at 37°C for 30 minutes. To stop tagmentation, EDTA was added to a final concentration of 50 nM and the reaction was incubated at 50°C for 30 minutes. Tagmented DNA was cleaned up using Qiagen PCR purification MinElute Kit (28004, Qiagen) and amplified using KAPA HiFi Hotstart ReadyMix (TAQHSKB, Roche) for 12-14 cycles. The amplified DNA was purified using a Qiagen PCR purification MinElute kit (28004), and KAPA Pure Beads (KK8000, Roche). Tagmentation efficiency was assessed using Agilent BioAnalyzer and libraries quantified using a Qubit Fluorometer (Thermo Fisher) at the Central Analytical Facility (University of Stellenbosch). The concentration of ATAC library pools was assessed with the KAPA Library Quantification Kit (KK4835, Roche). CS17 bat and E13.5 mouse libraries were sequenced by BGI (Hong Kong) using paired-end 50 bp runs on the Illumina HiSeq4000 platform. We used a three-stage MACS v2[91,92] to eliminate background. First, we called peaks assuming a flat model: *macs2 callpeak -t bedGraph-file-name -B --nomodel --pvalue=5e-5 --slocal=5000 --llocal=20000 --SPMR -n file-name*. We then performed an enrichment estimation by creating a treatment vs. control statistical apparatus: *macs2 bdgcmp -t treatment_pileup.bdg -c control_lambda.bdg -o FE.bdg -m FE*. We then subtracted the noise residual of the two models: *macs2 bdgcmp -t treatment_pileup.bdg -c control_lambda.bdg -o sub.bdg -m subtract*. The ATAC-Seq channels were utilized in order to home in on the exact locations of the chromatin action in regions of differential chromatin (DA).

LacZ enhancer transgenic assays
Mouse work was approved by the UCSF Institutional Animal Care and Use Committee (IACUC), protocol number AN197608, and was conducted in accordance with AALAC and NIH guidelines. The bat and mouse enhancer regions were amplified from mouse or bat (*Miniopterus natalensis*) genomic DNA by PCR, cloned into the Hsp68-LacZ vector[97] and sequence verified. Cloned regiones are shown in **Data S2**. All LacZ transgenic mice were generated by Cyagen Biosciences using standard procedures[98], and harvested and stained for LacZ expression as previously described[99] at specific time points noted on figures. Pictures were obtained using a M165FC stereo microscope (Leica).

Generation of bat enhancer replacement mice
We used the Precise Integration into Target Chromosome (PITCh) system[29] with minor modifications. Briefly, homology between bat and mouse sequences was characterized using CLUSTALW[100]. For donor plasmid construction, the bat enhancer and 1 kb mouse homology arms (5′ and 3′) were amplified by PCR. The cloned regions are listed in **Data S2.** These three fragments were cloned into the backbone plasmid pLSODN-3 (Biodynamics Laboratory, DS620) or pGEM-T easy (Promega, A1360) using NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs, E2621L). Circular donor

plasmids were dialyzed with a membrane (Millipore, VSWP04700) and used for
pronuclear injection. Two gRNAs were designed to target the 5′ and 3′ ends of the
mouse endogenous homology region (light blue in **Fig. 2d**) using the gRNA design tool
on the Integrated DNA Technologies (IDT) website and were selected based on low off-
target and high on-target scores. One of the gRNA target sites was placed at the 5′ and
3′ end of the homology in the donor plasmid (green in **Fig. 2d**). gRNAs are listed in
Supplementary Table 1. A mixture of crRNA (IDT), tracrRNA (IDT), Cas9 protein (IDT),
and donor plasmid was used for pronuclear injection. FVB/NJ mouse stain (Jackson
Laboratory; 001800) was used for all bat enhancer replacement mice generation.
Injections were performed at the Gladstone Transgenic Core. Founder mice were
screened by PCR and sequencing.
856
Alcian blue and alizarin red staining
Embryos or neonate mice limbs were harvested at E18.5 and limbs were dissected and
used for staining. Alcian blue/Alizarin red staining was performed according to standard
procedures for late-gestation stage embryos[101]. Briefly, following skin removal, limb
tissues were fixed in 95% ethanol overnight at room temperature, then treated with
acetone overnight at room temperature. Samples were stained with Alcian blue
overnight at room temperature and destained with sequential quick washes in 70% and
95% ethanol, followed by overnight incubation at room temperature. Tissues were then
pre-cleared in 1% KOH for 1 hour at room temperature, stained with Alizarin red for 3–4
hours at room temperature, and incubated in a 1:1 solution of 50% glycerol and 1%
KOH to remove excess red stain. Ossification zone and phalanx bone lengths were
measured using M165FC stereo microscope (Leica).
869
Histology
Whole limb tissues were fixed in 4% PFA at 4 °C overnight. The dorsal and ventral skin
from the arm was then dissected and further fixed in 4% PFA for 3 hours. Skin tissues
were embedded in paraffin, sectioned at 10 μm using a rotary microtome, and used for
standard Hematoxylin and Eosin (H&E) staining[102]. Epidermis thickness was measured
using Fiji software[103].
876
RNA extraction and qPCR
Total RNA was collected from tissue using TRIzol (Thermo Fisher Scientific; 15596026)
and converted to cDNA using ReverTra Ace qPCR-RT master mix with genomic DNA
(gDNA) remover (Toyobo; FSQ-301). qPCR was performed using SsoFast EvaGreen
supermix (Bio Rad; 1725205). Primer sequences used for qPCR are shown in
Supplementary Table 2.
883
RNA-seq for bat enhancer replacement mice
RNA-seq was conducted by Novogene. The library was generated with NEBNext Ultra II
RNA Library Prep Kit for Illumina (NEB; 7770), and sequencing was done using the
NovaSeq 6000 S4 platform with PE150. The data were analyzed using Partek Flow
(Version 10.0). After primary quality assessment was performed, bases and reads with

low quality were filtered out and the reads were aligned to the mouse reference genome
(mm10) using STAR (Version 2.7.8a)[88]. The final BAM files were quantified using the
Partek E/M algorithm[104]. Normalization of read count was performed by the total number
of counts (count per million) plus 0.0001, and all genes with less than ten normalized
read counts were excluded from subsequent analyses. Differentially expressed genes
were identified using DEseq2[89] in Partek Flow. Gene ontology[37] and Morpheus (RRID:
SCR_017386, https://software.broadinstitute.org/morpheus) were used for data
analysis.
897
Whole-mount *in situ* hybridization
For bat embryos, whole-mount in situ hybridization was carried out as described in [8].
For mouse, E11.5 embryos were fixed in 4% paraformaldehyde. A plasmid containing
mouse *Shh* and *Ptch1* cDNA (GenScript; Shh_OMu22903D and Ptch1_OMu22749D)
were used as templates for DIG-labeled probes. Mouse whole-mount in situ
hybridization was performed using standard procedures[105].
904

References

1. Sadier, A. *et al.* Making a bat: The developmental basis of bat evolution. *Genet. Mol. Biol.* **43**, e20190146 (2021).
2. Hockman, D., Mason, M. K., Jacobs, D. S. & Illing, N. The role of early development in mammalian limb diversification: a descriptive comparison of early limb development between the Natal long-fingered bat (Miniopterus natalensis) and the mouse (Mus musculus). *Dev. Dyn.* **238**, 965–979 (2009).
3. Xie, M. *et al.* Secondary ossification center induces and protects growth plate structure. *Elife* **9**, e55212 (2020).
4. Koyabu, D. & Son, N. T. Patterns of postcranial ossification and sequence heterochrony in bats: life histories and developmental trade-offs. *J. Exp. Zool. B Mol. Dev. Evol.* **322**, 607–618 (2014).
5. Weatherbee, S. D., Behringer, R. R., Rasweiler, J. J. th & Niswander, L. A. Interdigital webbing retention in bat wings illustrates genetic changes underlying amniote limb diversification. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 15103–15107 (2006).
6. Orkney, A., Boerma, D. B. & Hedrick, B. P. Evolutionary integration of forelimb and hindlimb proportions within the bat wing membrane inhibits ecological adaptation. *Nat. Ecol. Evol.* 1–13 (2024).
7. Reyes-Amaya, N., Jerez, A. & Flores, D. Morphology and postnatal development of lower hindlimbs in Desmodus rotundus (Chiroptera: Phyllostomidae): A comparative study. *Anat. Rec. (Hoboken)* **300**, 2150–2165 (2017).
8. Hockman, D. *et al.* A second wave of Sonic hedgehog expression during the development of the bat limb. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 16982–16987 (2008).
9. Wang, Z. *et al.* Unique expression patterns of multiple key genes associated with the evolution of mammalian flight. *Proc. Biol. Sci.* **281**, 20133133 (2014).
10. Mason, M. K. *et al.* Retinoic acid-independent expression of Meis2 during autopod patterning in the developing bat and mouse limb. *Evodevo* **6:6.**, 10.1186/s13227-015-0001-y. eCollection 2015. (2015).
11. Wang, Z. *et al.* Digital gene expression tag profiling of bat digits provides robust candidates contributing to wing formation. *BMC Genomics* **11:619.**, 10.1186/1471-2164-11-619. (2010).
12. Maier, J. A. *et al.* Transcriptomic insights into the genetic basis of mammalian limb diversity. *BMC Evol. Biol.* **17**, 86 (2017).
13. Eckalbar, W. L. *et al.* Transcriptomic and epigenomic characterization of the developing bat wing. *Nat. Genet.* **48**, 528–36. doi: 10.1038/ng.3537. Epub 2016 Mar 28. (2016).
14. Schindler, M. *et al.* Comparative single-cell analyses reveal evolutionary repurposing of a conserved gene programme in bat wing development. *Nat. Ecol. Evol.* **9**, 1626–1642 (2025).
15. Lyu, X. *et al.* Single-cell expression profiling of bat wing development. *Nat. Commun.* **16**, 6612 (2025).

16. Cretekos, C. J. *et al.* Regulatory divergence modifies limb length between
mammals. *Genes Dev.* **22**, 141–51. (2008).
17. Booker, B. M. *et al.* Bat Accelerated Regions Identify a Bat Forelimb Specific
Enhancer in the HoxD Locus. *PLoS Genet.* **12**, e1005738 (2016).
18. Ferris, E., Abegglen, L. M., Schiffman, J. D. & Gregg, C. Accelerated evolution in
distinctive species reveals candidate elements for clinically relevant traits, including
mutation and cancer resistance. *Cell Rep.* **22**, 2742–2755 (2018).
19. Creyghton, M. P. *et al.* Histone H3K27ac separates active from poised enhancers
and predicts developmental state. *Proc* **107**, 21931–6. (2010).
20. Rada-Iglesias, A. *et al.* A unique chromatin signature uncovers early developmental
enhancers in humans. *Nature* **470**, 279–83. (2011).
21. Barski, A. *et al.* High-resolution profiling of histone methylations in the human
genome. *Cell* **129**, 823–837 (2007).
22. Hockman, D., Mason, M. K., Jacobs, D. S. & Illing, N. The role of early development
in mammalian limb diversification: A descriptive comparison of early limb
development between the natal long-fingered bat (*Miniopterus natalensis*) and the
mouse (*Mus musculus*). *Dev. Dyn.* **238**, spcone-spcone (2009).
23. Cooper, K. L. The case against simplistic genetic explanations of evolution.
Development **151**, dev203077 (2024).
24. Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and
functional annotation of gene lists (2021 update). *Nucleic Acids Res.* **50**, W216–
W221 (2022).
25. Gokhman, D. *et al.* Gene ORGANizer: linking genes to the organs they affect.
Nucleic Acids Res. **45**, W138–W145 (2017).
26. McLean, C. Y. *et al.* GREAT improves functional interpretation of cis-regulatory
regions. *Nat. Biotechnol.* **28**, 495–501 (2010).
27. Kosicki, M. *et al.* VISTA Enhancer browser: an updated database of tissue-specific
developmental enhancers. *Nucleic Acids Res.* (2024) doi:10.1093/nar/gkae940.
28. Khan, S. F. *et al.* The roles and regulation of TBX3 in development and disease.
Gene **726**, 144223 (2020).
29. Sakuma, T., Nakade, S., Sakane, Y., Suzuki, K.-I. T. & Yamamoto, T. MMEJ-
assisted gene knock-in using TALENs and CRISPR-Cas9 with the PITCh systems.
Nat. Protoc. **11**, 118–133 (2016).
30. Visel, A., Minovitsky, S., Dubchak, I. & Pennacchio, L. A. VISTA Enhancer Browser-
-a database of tissue-specific human enhancers. *Nucleic Acids Res.* **35**, D88-92
(2007).
31. Kalay, E. *et al.* Mutations in RIPK4 cause the autosomal-recessive form of popliteal
pterygium syndrome. *Am. J. Hum. Genet.* **90**, 76–85 (2012).
32. Rountree, R. B. *et al.* RIP4 regulates epidermal differentiation and cutaneous
inflammation. *J. Invest. Dermatol.* **130**, 102–112 (2010).
33. Anthwal, N. *et al.* Insights into the formation and diversification of a novel
chiropteran wing membrane from embryonic development. *BMC Biol.* **21**, 101
(2023).

34. Zuniga, A. Next generation limb development and evolution: old questions, new
perspectives. *Development* **142**, 3810–3820 (2015).
35. Khokha, M. K., Hsu, D., Brunet, L. J., Dionne, M. S. & Harland, R. M. Gremlin is the
BMP antagonist required for maintenance of Shh and Fgf signals during limb
patterning. *Nat. Genet.* **34**, 303–307 (2003).
36. Malkmus, J. *et al.* Spatial regulation by multiple Gremlin1 enhancers provides digit
development with cis-regulatory robustness and evolutionary plasticity. *Nat.*
Commun. **12**, 5557 (2021).
37. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene
Ontology Consortium. *Nat. Genet.* **25**, 25–29 (2000).
38. Gene Ontology Consortium *et al.* The Gene Ontology knowledgebase in 2023.
Genetics **224**, iyad031 (2023).
39. Briso, E. M. *et al.* Inflammation-mediated skin tumorigenesis induced by epidermal
c-Fos. *Genes Dev.* **27**, 1959–1973 (2013).
40. Ferby, I. *et al.* Mig6 is a negative regulator of EGF receptor-mediated skin
morphogenesis and tumor formation. *Nat. Med.* **12**, 568–573 (2006).
41. Smits, J. P. H. *et al.* The aryl hydrocarbon receptor regulates epidermal
differentiation through transient activation of TFAP2A. *bioRxivorg* (2023)
doi:10.1101/2023.06.07.544032.
42. Katoh, M. FGFR2 abnormalities underlie a spectrum of bone, skin, and cancer
pathologies. *J. Invest. Dermatol.* **129**, 1861–1867 (2009).
43. Yakushiji-Kaminatsui, N. *et al.* Similarities and differences in the regulation of HoxD
genes during chick and mouse limb development. *PLoS Biol.* **16**, e3000004 (2018).
44. Sagai, T., Hosoya, M., Mizushina, Y., Tamura, M. & Shiroishi, T. Elimination of a
long-range cis-regulatory module causes complete loss of limb-specific Shh
expression and truncation of the mouse limb. *Development* **132**, 797–803 (2005).
45. Kvon, E. Z. *et al.* Progressive loss of function in a limb enhancer during snake
evolution. *Cell* **167**, 633-642.e11 (2016).
46. VanderMeer, J. E. & Ahituv, N. Cis-regulatory mutations are a genetic cause of
human limb malformations. *Dev. Dyn.* **240**, 920–930 (2011).
47. Maas, S. A., Suzuki, T. & Fallon, J. F. Identification of spontaneous mutations within
the long-range limb-specific Sonic hedgehog enhancer (ZRS) that alter Sonic
hedgehog expression in the chicken limb mutants oligozeugodactyly and silkie
breed. *Dev. Dyn.* **240**, 1212–1222 (2011).
48. Lallemand, Y., Bensoussan, V., Cloment, C. S. & Robert, B. Msx genes are
important apoptosis effectors downstream of the Shh/Gli3 pathway in the limb. *Dev.*
Biol. **331**, 189–198 (2009).
49. Montero, J. A. *et al.* Role of FGFs in the control of programmed cell death during
limb development. *Development* **128**, 2075–2084 (2001).
50. Dai, M. *et al.* Differential expression of Meis2, Mab21l2 and Tbx3 during limb
development associated with diversification of limb morphology in mammals. *PLoS*
One **9**, e106100 (2014).

51. Schindler, M. *et al.* Comparative single-cell analyses reveal evolutionary repurposing of a conserved gene program in bat wing development. *Evolutionary Biology* (2024).

52. Kuijper, S. *et al.* Function and regulation of Alx4 in limb development: complex genetic interactions with Gli3 and Shh. *Dev. Biol.* **285**, 533–544 (2005).

53. Barna, M., Hawe, N., Niswander, L. & Pandolfi, P. P. Plzf regulates limb and axial skeletal patterning. *Nat. Genet.* **25**, 166–172 (2000).

54. Vieux-Rochas, M. *et al.* BMP-mediated functional cooperation between Dlx5;Dlx6 and Msx1;Msx2 during mammalian limb development. *PLoS One* **8**, e51700 (2013).

55. Lallemand, Y. *et al.* Analysis of Msx1; Msx2 double mutants reveals multiple roles for Msx genes in limb development. *Development* **132**, 3003–3014 (2005).

56. Bensoussan-Trigano, V., Lallemand, Y., Saint Cloment, C. & Robert, B. Msx1 and Msx2 in limb mesenchyme modulate digit number and identity. *Dev. Dyn.* **240**, 1190–1202 (2011).

57. Gurumurthy, C. B. *et al.* Creation of CRISPR-based germline-genome-engineered mice without ex vivo handling of zygotes by i-GONAD. *Nat. Protoc.* **14**, 2452–2482 (2019).

58. Satokata, I. & Maas, R. Msx1 deficient mice exhibit cleft palate and abnormalities of craniofacial and tooth development. *Nat. Genet.* **6**, 348–356 (1994).

59. Satokata, I. *et al.* Msx2 deficiency in mice causes pleiotropic defects in bone growth and ectodermal organ formation. *Nat. Genet.* **24**, 391–395 (2000).

60. Amano, K. *et al.* MSX2 stimulates chondrocyte maturation by controlling Ihh expression. *J. Biol. Chem.* **283**, 29513–29521 (2008).

61. Holthaus, K. B., Steinbinder, J., Sachslehner, A. P. & Eckhart, L. Skin appendage proteins of tetrapods: Building blocks of claws, feathers, hair and other cornified epithelial structures. *Animals (Basel)* **15**, 457 (2025).

62. Fujikawa, H., Fujimoto, A., Farooq, M., Ito, M. & Shimomura, Y. Characterization of the human hair keratin-associated protein 2 (KRTAP2) gene family. *J. Invest. Dermatol.* **132**, 1806–1813 (2012).

63. Kawai, S., Yamauchi, M., Wakisaka, S., Ooshima, T. & Amano, A. Zinc-finger transcription factor odd-skipped related 2 is one of the regulators in osteoblast proliferation and bone formation. *J. Bone Miner. Res.* **22**, 1362–1372 (2007).

64. Yang, X. *et al.* TGF-beta/Smad3 signals repress chondrocyte hypertrophic differentiation and are required for maintaining articular cartilage. *J. Cell Biol.* **153**, 35–46 (2001).

65. Horiki, M. *et al.* Smad6/Smurf1 overexpression in cartilage delays chondrocyte hypertrophy and causes dwarfism with osteopenia. *J. Cell Biol.* **165**, 433–445 (2004).

66. Kong, R. Y. *et al.* Intron-exon structure, alternative use of promoter and expression of the mouse collagen X gene, Col10a-1. *Eur. J. Biochem.* **213**, 99–111 (1993).

67. Gómez-Picos, P. & Eames, B. F. On the evolutionary relationship between chondrocytes and osteoblasts. *Front. Genet.* **6**, 297 (2015).

68. Bell, P. A. *et al.* Loss of matrilin 1 does not exacerbate the skeletal phenotype in a mouse model of multiple epiphyseal dysplasia caused by a Matn3 V194D mutation. *Arthritis Rheum.* **64**, 1529–1539 (2012).

69. Shukunami, C. & Hiraki, Y. Role of cartilage-derived anti-angiogenic factor, chondromodulin-I, during endochondral bone formation. *Osteoarthritis Cartilage* **9 Suppl A**, S91-101 (2001).

70. Almeida, M., Han, L., Martin-Millan, M., O'Brien, C. A. & Manolagas, S. C. Oxidative stress antagonizes Wnt signaling in osteoblast precursors by diverting beta-catenin from T cell factor- to forkhead box O-mediated transcription. *J. Biol. Chem.* **282**, 27298–27305 (2007).

71. Geetha-Loganathan, P., Nimmagadda, S. & Scaal, M. Wnt signaling in limb organogenesis. *Organogenesis* **4**, 109–115 (2008).

72. Huang, X. *et al.* Phosphorylation of Dishevelled by protein kinase RIPK4 regulates Wnt signaling. *Science* **339**, 1441–1445 (2013).

73. Hu, L., Chen, W., Qian, A. & Li, Y.-P. Wnt/β-catenin signaling components and mechanisms in bone formation, homeostasis, and disease. *Bone Res.* **12**, 39 (2024).

74. Regard, J. B., Zhong, Z., Williams, B. O. & Yang, Y. Wnt signaling in bone development and disease: making stronger bone with Wnts. *Cold Spring Harb. Perspect. Biol.* **4**, a007997–a007997 (2012).

75. Farnum, C. E., Tinsley, M. & Hermanson, J. W. Forelimb versus hindlimb skeletal development in the big brown bat, Eptesicus fuscus: functional divergence is reflected in chondrocytic performance in Autopodial growth plates. *Cells Tissues Organs* **187**, 35–47 (2008).

76. Saxena, A. *et al.* Interspecies transcriptomics identify genes that underlie disproportionate foot growth in jerboas. *Curr. Biol.* **32**, 289-303.e6 (2022).

77. Bolt, C. C. *et al.* Context-dependent enhancer function revealed by targeted inter-TAD relocation. *Nat. Commun.* **13**, 3488 (2022).

78. Tickle, C. & Towers, M. Sonic hedgehog signaling in limb development. *Front. Cell Dev. Biol.* **5**, 14 (2017).

79. Lettice, L. A., Devenney, P., De Angelis, C. & Hill, R. E. The conserved sonic hedgehog limb enhancer consists of discrete functional elements that regulate precise spatial expression. *Cell Rep.* **20**, 1396–1408 (2017).

80. Lettice, L. A. *et al.* Opposing functions of the ETS factor family define Shh spatial expression in limb buds and underlie polydactyly. *Dev. Cell* **22**, 459–467 (2012).

81. Zhang, Z., Verheyden, J. M., Hassell, J. A. & Sun, X. FGF-regulated Etv genes are essential for repressing Shh expression in mouse limb buds. *Dev. Cell* **16**, 607–613 (2009).

82. Kvon, E. Z. *et al.* Comprehensive in vivo interrogation reveals phenotypic impact of human enhancer variants. *Cell* **180**, 1262-1271.e15 (2020).

83. Lettice, L. A. *et al.* A long-range Shh enhancer regulates expression in the developing limb and fin and is associated with preaxial polydactyly. *Hum. Mol. Genet.* **12**, 1725–1735 (2003).

84. Albuisson, J. *et al.* Identification of two novel mutations in Shh long-range regulator associated with familial pre-axial polydactyly. *Clin. Genet.* **79**, 371–377 (2011).

85. Furniss, D. *et al.* A variant in the sonic hedgehog regulatory sequence (ZRS) is associated with triphalangeal thumb and deregulates expression in the developing limb. *Hum. Mol. Genet.* **17**, 2417–2423 (2008).

86. Lim, F. *et al.* Affinity-optimizing enhancer variants disrupt development. *Nature* **626**, 151–159 (2024).

87. Teeling, E. C. *et al.* A molecular phylogeny for bats illuminates biogeography and the fossil record. *Science* **307**, 580–584 (2005).

88. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

89. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

90. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

91. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

92. Feng, J., Liu, T., Qin, B., Zhang, Y. & Liu, X. S. Identifying ChIP-seq enrichment using MACS. *Nat. Protoc.* **7**, 1728–1740 (2012).

93. Diaz, A., Park, K., Lim, D. A. & Song, J. S. Normalization, bias correction, and peak calling for ChIP-seq. *Stat. Appl. Genet. Mol. Biol.* **11**, Article 9 (2012).

94. Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. deepTools: a flexible platform for exploring deep-sequencing data. *Nucleic Acids Res.* **42**, W187-91 (2014).

95. Karolchik, D. *et al.* The UCSC Genome Browser Database. *Nucleic Acids Res.* **31**, 51–4. (2003).

96. Kuhn, R. M., Haussler, D. & Kent, W. J. The UCSC genome browser and associated tools. *Brief. Bioinform.* **14**, 144–161 (2013).

97. Kothary, R. *et al.* A transgene containing lacZ inserted into the dystonia locus is expressed in neural tube. *Nature* **335**, 435–437 (1988).

98. Pu, X.-A., Young, A. P. & Kubisch, H. M. Production of transgenic mice by pronuclear microinjection. *Methods Mol. Biol.* **1874**, 17–41 (2019).

99. Pennacchio, L. A. *et al.* In vivo enhancer analysis of human conserved non-coding sequences. *Nature* **444**, 499–502 (2006).

100. Thompson, J. D., Higgins, D. G. & Gibson, T. J. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673–4680 (1994).

101. Rigueur, D. & Lyons, K. M. Whole-mount skeletal staining. *Methods Mol. Biol.* **1130**, 113–121 (2014).

102. Fischer, A. H., Jacobson, K. A., Rose, J. & Zeller, R. Hematoxylin and eosin staining of tissue and cell sections. *CSH Protoc.* **2008**, db.prot4986 (2008).

103. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).

104. Xing, Y. *et al.* An expectation-maximization algorithm for probabilistic reconstructions of full-length isoforms from splice graphs. *Nucleic Acids Res.* **34**, 3150–3160 (2006).

105. Hargrave, M., Bowles, J. & Koopman, P. In situ hybridization of whole-mount embryos. *Methods Mol. Biol.* **326**, 103–113 (2006).

Acknowledgements

We want to thank Dr. Hai P. Nguyen for discussion, advice and encouragement and the Ahituv lab for helpful advice. This work was funded in part by the National Human Genome Research Institute grant numbers R01HG012396 (NA) and K99HG012576 (AU) and R00HG012576 (AU), Japan Society for the Promotion of Science (JSPS) postdoctoral fellowships for research abroad (AU), Uehara Memorial Foundation postdoctoral fellowship (AU) and the Sydney Brenner Fellowship from the Academy of Science of South Africa (DH). US-Israel Binational Science Foundation (no. 2015307, NA and TK) and the Bergmann Memorial Research Award (TK).

Author information

Contributions

Conceptualization: AU, GK, DH, NI, TK, NA

Methodology: AU, GK, RS, EM, WE, YZ, MN, RR, KF, VB, KN, SK, SAS, MM, SZ, ML, SF, VS, TS, DH

Investigation: AU, GK, RS, EM, YZ, MN, RR, KF, VB, KN, SK, SAS, MM, SZ, ML, SF, MYH, VS, TS, DH, NI, TK, NA

Visualization: AU, GK, RS, EM, MN, KF, VB, SZ, ML, MYH, DH, NI, TK, NA

Funding acquisition: AU, DH, NI, TK, NA

Project administration: AU, GK, DH, NI, TK, NA

Supervision: AU, NI, TK, NA

Writing – original draft: AU, GK, DH, NI, TK, NA

Writing – review & editing: AU, GK, DH, NI, TK, NA

Ethics declarations

Competing interests

NA is a cofounder and on the scientific advisory board of Regel Therapeutics Inc. NA received funding from BioMarin Pharmaceutical Incorporate. The remaining authors declare no competing interests.

Supplementary information

Extended Data Figures 1-9

Supplementary Tables 1-2

Supplementary Data 1-2: Excel file containing additional data too large to fit in a PDF