1   **Title: *Myrmecocystus* honeypot ants have species specific resident gut microbiome**
2

3   Authors: DiemQuynh V. Nguyen[1], Charlotte B. Francoeur[1], Bianca R. Nogueira[1], Indira Sawh[1],
4   Michele Lanan[2], Lily Khadempour[1]
5
6   **Affiliations**
7   1. Department of Earth and Environmental Sciences, Rutgers University, Newark, NJ, USA
8   07102
9   2. Department of Entomology, University of Arizona, Tucson, AZ, USA 85721
10
11   **Corresponding author** Lily Khadempour, lily.khadempour@rutgers.edu

12
Abstract

Myrmecocystus honeypot ants rely on specialized workers, repletes, to store dissolved
carbohydrates in their crops long term. The repletes store this liquid, which does not spoil in
their crops, for many months at a time. When resources are scarce, repletes redistribute the
stored nutrients to their colony members via trophallaxis. While we suspect that the gut
microbiome of honeypot ants may aid in spoilage prevention, before we can investigate this, we
must first characterize it. Here, we used 16S rRNA gene sequencing to determine the microbial
community composition across six *Myrmecocystus* honeypot ant species, sampling multiple
colonies, castes, and organs. We found that microbiome community composition was strongly
shaped by species, with variation between colonies in *M. arenarius*, *M. depilis*, and *M.
mexicanus*. Organ level differences were observed in the crop and midgut in *M. mexicanus*.
Caste differences were observed in *M. flaviceps* and *M. mexicanus.* Replete crops of *M.
mexicanus* and *M. depilis* were enriched in *Fructilactobacillus,* other lactic acid bacteria, and
acetic acid bacteria, whereas halophiles were more prominent in the gut of species such as *M.
flaviceps* and *M. wheeleri.* In this study we demonstrate that *Myrmecocystus* ants host species-
specific gut microbiomes and identify an association between lactic acid bacteria, acetic acid
bacteria, and halophiles within replete crops. While much work remains in understanding the
roles of the microbes in the symbiosis with their host ants, the dominance of these particular
taxonomic groups suggests an association with a high sugar environment and a potential
microbial role in preventing spoilage of the crop contents.

33
34   **Keywords**
35   Replete, crop, lactic acid bacteria*,* halophiles
36

Introduction

Many animals, particularly those with specialized diets, rely on their microbiomes to supplement
essential nutrients and provide beneficial services to the host (Engel & Moran, 2013). Compared
to vertebrates, insects provide more tractable systems for studying host-microbiome
relationships because they tend to have simpler gut morphology and gut microbiome
communities. Host-associated microbiomes can contribute to a range of tasks, including
nitrogen fixation and recycling (e.g., *Cephalotes* turtle ants), amino acid production (e.g.,

Onthophagus taurus dung beetles), toxin remediation (e.g., *Apis mellifera* honeybees), and the
breakdown of complex plant polysaccharides the host cannot digest independently (e.g.,
Nasuititermes termites) (Brune, 2014; Auer et al., 2017; Hu et al., 2018; Flynn et al., 2021;
Motta et al., 2022; Jones et al., 2025), However, not all insects require a resident gut
microbiome. For example, despite their herbivorous diet, *Lepidoptera* caterpillars can meet their
nutritional demands through host-encoded digestive enzymes and intrinsic mechanisms to
degrade or tolerate plant toxins (Hammer et al., 2017).

Honeypot ants represent a compelling system for investigating animal-associated gut
microbiomes. This convergently-evolved group comprises at least eight independently evolved
genera spanning three Formicidae subfamilies (Sawh, 2023; Nogueira et al., 2026). In this
study, we focus on the North American honeypot ants in the genus *Myrmecocystus*. This genus
began diversifying approximately 14 million years ago, coinciding with the aridification of the
southwestern United States and northern Mexico (Van Elst et al., 2021). The formation of new
desert habitats created ecological niches that likely shaped the evolution of this group (Van Elst
et al., 2021). To overcome irregular resource availability, honeypot ants have evolved
specialized workers, known as repletes, that act as living food storage units to sustain the
colony during dry, nutrient-limited seasons (Nogueira et al., 2026). Honeypot ant foragers collect
extrafloral nectar, honeydew, and insect fluids to provide them to the repletes through
trophallaxis, a regurgitation food-sharing process common to social insects (Meurville &
LeBoeuf, 2021). As repletes accumulate the collected resources, the crops become enlarged
and the intersegmental membranes of the gaster undergo extensive stretching (Fig. 1B, D-E)
(Conway, 1977). While the basic structure of adult ant digestive tracts are conserved across all
ants, honeypot ants are able to distend their crop far beyond the limits observed in non-replete
ants, facilitating long term storage of liquid resources (Eisner & Brown, 1956). A 2023 study by
Dong et al. found that the liquid stored in Australian honeypot ants, *Camponotus inflatus*, has
antimicrobial properties. Based on whole ant sequencing, they concluded that the microbiome
was dominated by the endosymbiotic bacterial genus *Candidatus Blochmannia* (99.75%) and
the fungal genus *Neocelosporium* (92.77%) (Dong et al., 2023). *Blochmannia* is a well-known
endosymbiont of *Camponotus* ants, residing within bacteriocytes associated with the midgut,
and it contributes to host nutrition and development (Feldhaar et al., 2007). Since *Blochmannia*
is restricted to midgut tissues, we suspect that the microbe responsible for the antimicrobial
properties resides within the crop rather than the midgut (Feldhaar et al., 2007; Sauer et al.,
2002). Therefore, we hypothesize that the gut microbiome of *Myrmecocystus* honeypot ants
contributes to antimicrobial activity within their sugar-rich crop, thereby limiting the growth of
pathogenic bacteria and fungi and preventing spoilage. However, before addressing the
potential function, it is necessary to first characterize the gut microbiome of *Myrmecocystus*
species*.*

In this study, we characterize the gut microbiome of six species of *Myrmecocystus* honeypot
ants: *M. arenarius, M.depilis, M. flaviceps, M. mexicanus, M. mimicus* and *M, wheeleri*. We use
community amplicon sequencing to determine whether a consistent resident gut microbiome
exists. We compare the gut microbiome across scales: between different species within the

genus, between colonies of the same species, between castes within a colony, and between
digestive organs within individual ants. If these ants possess a resident gut microbiome, we
expect to observe the same community members in the same organs within a species, and
potentially across the genus. Establishing whether *Myrmecocystus* hosts a resident gut
microbiome is an important first step in understanding the relationship between these ants and
their gut microbiome.

Materials and Methods
Field Sampling
Myrmecocystus ants were collected at five time points from 2022-2025 (Tab. S1), either from
private property with permission or Bureau of Land Management (BLM) land, throughout the
Southwestern United States (Tab. S1). Initial species identification was conducted onsite using
worker morphological characteristics. We later confirmed ant identification through sequencing
and analysis of the COI gene (Van Elst et al., 2021). Overall, we collected six *Myrmecocystus*
species including *M. arenarius, M. depilis, M. flaviceps, M. mexicanus, M. mimicus*, and *M.*
wheeleri. Our goal was to collect 10 repletes per colony, although this was not always
achievable (Tab. S1). Alongside replete and worker samples, we also collected queens, alates,
and brood however these collections were inconsistent and are saved for future examinations.
105

To collect repletes and workers, we excavated colonies until we found chambers containing
repletes. Compared to repletes, regular workers were easy to find and were located both in the
replete chambers and throughout the rest of the colony chambers and tunnels. Typically, replete
chambers would be found at least 1 m underground, but on rare occasions we would find them
closer to the surface. Holes were carefully hand dug with small shovels, and were filled in once
collections were completed. We removed repletes and workers with featherweight forceps or
with fan paintbrushes into containers. Ants were not fed and kept alive until the dissection
process. Colony fragments were maintained together prior to dissection, allowing possible
trophallactic exchange to continue during this period. Ants were first surface sterilized with 70%
ethanol followed by sterile phosphate buffered solution. Dissections were carried out under
sterile conditions next to an open flame, using a dissecting scope, on a sterile petri dish. To
minimize microbial contamination, all tools and equipment were surface sterilized repeatedly
between each dissection using 70% ethanol and by passing them through the flame. The ants
were surface sterilized and dissected by first removing the head, then the external membrane of
the gaster was opened using sterile ultrafine forceps, and the digestive organs (crop and
midgut) were transferred with contents intact and not leaking into individual sterile tubes. If any
organ ruptured during dissection, the sample was discarded and not used for DNA extraction or
subsequent analyses. Dissected midguts and crops were placed into tubes containing 100 µL of
TES buffer (10mM pH 7.5 Tris, 10mM pH 8 EDTA, 0.5% SDS) for DNA extractions. For the
repletes, dissected midguts were also placed into 100 µL of TES buffer. Because the replete
crops contained a relatively large amount of fluid, we reserved some material for future
proteomics, metabolomics and microbial isolation. However, this study is focused on
characterizing the microbiome community composition, and so we describe only the methods
for conducting community amplicon sequencing.
130

DNA Extractions and Sequencing
For individual ant crop and midgut samples stored in 100 µL of TES buffer (10mM pH 7.5 Tris,
10mM pH 8 EDTA, 0.5% SDS), our custom phenol/chloroform extraction protocol was followed.
Samples were incubated with 36 µL of lysozyme (100 mg/mL in 10mM Tris pH 7.5; Thermo
Scientific, Waltham, MA, USA) for 24 hours at 37°C. Then, 220 µL of TES and 5 µL proteinase K
(800 U/ml; New England Biolabs, Ipswich, MA, USA) were added to our samples. Samples were
incubated at 56°C for 20 minutes. Following incubation, 250 µL of Sigma Aldrich 3M sodium
acetate was added, and samples were frozen at -80°C for at least 15 minutes. Once the
samples had thawed, samples were spun at 14,000 rpm for 15 minutes at 4°C. The supernatant
is then transferred to a new tube, and a 1:1 volume of phenol:chloroform is added. Samples
were spun at 14,000 rpm for 15 minutes at 4°C. The supernatant was transferred to a new tube,
and a chloroform wash was performed. Before precipitating, 10% sodium acetate and 90% cold
isopropanol were added. Samples were spun at 14,000 rpm for 15 minutes at 4°C to pellet the
DNA. Then we performed two ethanol washes with 70% ethanol. The DNA was then
resuspended in 50 µL TE (10mM pH 7.5 Tris, 1mM EDTA). DNA concentrations were quantified
with Qubit HS dsDNA (Thermo Fisher Scientific, Waltham, MA, USA). Extracted DNA samples
were then shipped to the University of Connecticut's Microbial Analysis, Resources, and
Services (UCONN MARS) facility for 16S rRNA gene amplicon sequencing and ITS amplicon
sequencing.
150
According to the UCONN MARS facility protocol, the 16S rRNA V4 region was amplified using
515F and 806R with Illumina adapters and dual 8-basepair indices (Kozich et al., 2013). The
ITS2 region was amplified with ITS3 and ITS4 primers (White et al., 1990) using the same dual
indexing design as the V4. PCR reactions were performed with an initial denaturation of 95°C
for 3.5 minutes, followed by 30 cycles of 30 seconds at 95.0°C, 30 seconds at 50.0°C, and 90
seconds at 72.0°C, with a final extension at 72.0°C for 10 minutes. PCR products were pooled
for quantification and visualization using the QIAxcel DNA Fast Analysis (Qiagen, Hilden,
Germany). PCR products were normalized based on the concentration of DNA from 250-400
bp, then pooled using the epMotion 3075 liquid handling robot. The pooled PCR products were
cleaned using Mag-Bind Beads (0.8x ratio; Omega Bio-Tek, Norcross, GA, USA) and
sequenced on an Illumina MiSeq platform (Illumina; San Diego, CA, USA) with a v2 2x250 base
pair kit. We only conducted ITS sequencing for the first set of collected samples. Because we
found almost no fungi in our samples, we chose to focus on bacteria and archaea only in
subsequent sequencing runs.
165
166
Amplicon and Statistical Analysis
We processed, aligned, and categorized sequence reads using DADA2 1.25.2 in R (Callahan et
al., 2016). The DADA2 pipeline was followed exactly, with the addition of the decontam package
to identify and remove potential contaminants (Davis et al., 2018). After filtering, trimming, and
merging reads, amplicon sequence variants (ASV) were taxonomically assigned using the
Greengenes v2 database. To construct the ASV-based phylogeny required for weighted
UniFrac analyses, we first made an alignment using MAFFT (Katoh et al., 2002). An ASV
phylogeny was inferred from these sequences using the GTR+G model in IQtree v2.4.0 (Minh et

al., 2020). The resulting tree was rooted in R using the phangorn package (Schliep, 2011). The
rooted tree was incorporated into a phyloseq object (McMurdie & Holmes, 2013).

Weighted UniFrac distances were calculated using the phyloseq package, and ordinations were
performed using non-metric multidimensional scaling (NMDS) with the vegan package
(Oksanen et al., 2022). Microbiome composition was visualized using NMDS plots generated
with the plot_ordinate function in phyloseq (McMurdie & Holmes, 2013).

Statistical significance between groups was evaluated using global permutational multivariate
analysis of variance (PERMANOVA) and pairwise differences with ANOSIM in the vegan
package (Oksanen et al., 2022). Analysis of compositions of microbiomes with bias correction
(ANCOM-BC2) was used to identify differentially abundant taxa in each organ, in this case the
log fold changes between the crop and midgut (Lin & Peddada, 2020). ANCOM-BC2 was mainly
used for organ comparison because it allowed us to test whether the repletism phenotype is
associated with differential abundance of specific taxa. Alpha diversity metrics, including
observed richness, Shannon diversity, and inverse Simpson diversity, were calculated to
measure the richness and evenness of microbial communities within the samples, using
phyloseq (Shannon, 1948; Simpson, 1949; McMurdie & Holmes, 2013). Statistical differences in
Shannon diversity index were first assessed using Kruskal-Wallis test, followed by pairwise
Wilcoxon rank-sum tests with Benjamini-Hochberg correction for multiple comparisons
(Wilcoxon, 1945; Kruskal & Wallis, 1952; Benjamini & Hochberg, 1995). Bubble plots were
generated in R using the ggplot2 and reshape2 package to visualize the presence and relative
abundance of microbial taxa across species and across organs (Wickham, 2011, 2016).

Results
Species Differences
Microbiome composition differed strongly across *Myrmecocystus* species, with distinct taxa
characterizing each species (Fig. 2, Fig. 3). In particular, *M. mexicanus* replete crops were
dominated by *Fructilactobacillus*, with the top three ASV belonging to this genus (ASV1, ASV5,
and ASV12) (Fig. 3, Fig S2). In contrast, *Fructilactobacillus* was largely absent from the replete
crops of some species (e.g., *M. arenarius* and *M. wheeleri)* (Fig. 3, Fig S2). These differences
were statistically significant across species and explained a large proportion of variation in
community composition (PERMANOVA p = 0.000999; ANOSIM p = 0.001; Tab. S5).
Myrmecocystus species accounted for approximately 61% of the variation in replete crop
microbiome (R^2 = 0.60726) and 45% in the replete midgut (R^2 = 0.45325; Tab. S5). Shannon
diversity differed significantly among species (Kruskal-Wallis chi-squared = 65.873, p = 7.385e-
13). *Myrmecocystus mexicanus* exhibited significantly lower Shannon diversity than all other
species, whereas some species pairs (e.g., *M. depilis* and *M. mimicus*) did not differ significantly
(Tab. S2).
211
212 Overall, species identity is a substantial factor of microbiome variation. Notably, *M. mexicanus*
213 was characterized by a less diverse, *Fructilactobacillus*-dominated crop microbiome, whereas
214 other species exhibited more compositionally diverse crop microbiomes (Fig. S2).
215

Colony Difference Within Species

Microbiome composition differed among some colonies within *Myrmecocystus* species, but this pattern was not universal (Tab. S5). Significant colony-level differences were observed in *M. arenarius, M. depilis,* and *M. mexicanus* (PERMANOVA, p < 0.01; Tab. S5), whereas *M. flaviceps, M. mimicus,* and *M. wheeleri* showed no detectable colony level differences. Within *M. mexicanus*, collection year (e.g., July 2022, August 2023, and July 2025) explained a relatively small proportion of variation in the crop microbiome (PERMANOVA R^2 = 0.09736; Tab. S6). Despite variation among colonies and across years, *Fructilactobacillus* remained a consistent and dominant genus in *M. mexicanus* crop, with ASV1 present in all samples (100% prevalence). However, the relative abundance of co-dominant taxa shifted across time points, including *Acetobacteraceae* G004843345 (ASV6), *Weizmannia* (ASV21), and lactic acid bacteria such as *Pediococcus* (ASV23) and *Lentilactobacillus* (ASV42). Similarly, *M. depilis* exhibited variation in microbiome composition within dominant genera between collection years, 2023 and 2025 (Tab. S6). The top ASV in *Myrmecocystus depilis* samples collected in 2023 included *Fructilactobacillus* (ASV1), *Acinetobacter* (ASV73), and *Vibrio* A 676397 (ASV7), while 2025 samples were characterized by increased abundance of *Saccharibacter* (ASV11) and other genera.

Here, we observe that colony differences were present in some but not all species, and that collection time can contribute to these patterns. Variations among colonies may partly reflect uneven sampling across species and time points.

Caste Differences (Replete vs. Worker)

Non-metric Multidimensional Scaling (NMDS) allows us to visualize overlap between replete and worker organ communities (Fig. 4). Overall, caste did not strongly structure microbiome composition, as no significant differences were detected between replete and worker samples in either the crop and midgut (PERMANOVA, p > 0.05; Tab. S5). However, caste differences were observed for certain species (Fig. 4; Tab. S5). In *M. depilis, M. flaviceps,* and *M. mexicanus,* worker and replete crop microbiomes differed significantly (PERMANOVA, p < 0.05; Tab. S5), whereas no caste differences were detected in other *Myrmecocystus* species (Tab. S5). These differences were driven by shifts in dominant taxa between castes. For example, in *M. depilis* replete crops contained *Fructilactobacillus* (ASV1: 0-99.28%), *Saccharibacter* (ASV11: 0-82.57%), and *Acinetobacter* (ASV73: 0-60.22%). While *M. depilis* worker crops were characterized by *Vibrio* (ASV7: 0-37.23%), *Pseudoalteromonas* (ASV22: 0-17.31%), and *Haloparvum* (ASV17). Similarly, in *M. flaviceps,* replete crops were dominated by *Haloparvum* (ASV17) and Bacili-associated taxa (ASV362 and ASV224), while worker crops showed higher relative abundance of *Weizmannia* (ASV21) and *Natronomonas* (ASV24). In *M. mexicanus*, both castes were dominated by *Fructilactobacillus*, but worker crops also differed in the relative abundance of the secondary taxon *Acetobacteraceae* G004843345 (ASV6 and ASV19). Shannon diversity also differed between castes in some species. Worker samples had a higher Shannon diversity than repletes in *M. flaviceps* (Wilcoxon test, adjusted p = 0.0378) and *M. mexicanus* (Wilcoxon test, adjusted p = 0.0170). However, no significant caste differences were observed for other species (Fig. S4).

260   Overall, caste did not strongly structure gut microbiomes across *Myrmecocystus* species. While
261   crop communities differed between castes in some species (e.g., *M. depilis, M. flaviceps,* and
262   *M. mexicanus),* the midgut communities remained similar between castes across all species in
263   our analysis.
264
Organ Differences (Crop vs. Midgut)
Microbiome composition differed significantly between the crop and midgut across
Myremecocystus species, independent of species identity (PERMANOVA R^2 = 0.36251, p =
0.000999; Tab. S5). Differences between the crop and midgut were detectable in *M. depilis*, *M.*
mexicanus, *M. mimicus*, and *M. wheeleri* (PERMANOVA and ANOSIM, p < 0.05; Tab. S5).
However, *M. arenarius* and *M. flaviceps* showed no detectable crop-midgut difference (Tab. S5).
Shannon diversity was significantly higher in the midgut than the crop in *M. mexicanus* repletes
(Wilcoxon rank-sum test, FDR-adjusted p = $1.82e^{-13}$) (Fig. 5). While no significant differences in
diversity were observed between organs in other species (all FDR-adjusted p > 0.05) (Fig. 5;
Fig S3). ANCOM-BC2 identified several genera that are differentially enriched in either gut
compartments in *M. depilis* and *M. mexicanus* (Fig. 6; Tab. S3-S4). In both species,
Fructilactobacillus was significantly enriched in the crop (Fig. 6). In *M. mexicanus,* several
additional lactic acid bacteria and Acetobacteraceae were also enriched in the crop, whereas
taxa such as *Vibrio* and *Pseudoaltermonas* were enriched in the midgut (Fig. 6; Tab. S3-S4). No
taxa were identified as differentially abundant between the organs in *M. mimicus, M. arenarius,*
M. flaviceps, and *M. wheeleri.*

Overall, gut compartmentalization was observed in some *Myrmecocystus* species. Particularly
in *M. mexicanus,* where the crop and midgut differed in both community composition and
diversity.

284
Discussion
In this study, we characterize gut microbiomes across six *Myrmecocystus* species by analyzing
the communities across species, colony, caste, and organ. Across our analyses, microbiome
composition was primarily driven by ant species identity. Caste had minimal influence on
microbiome structure, with repletes and workers generally retaining similar bacterial
communities despite differences in morphology, varying tasks, and potential environmental
exposures. While at first we were surprised that there were minimal differences between castes,
in retrospect it makes sense for an ant species that engages in frequent trophallaxis, constantly
sharing the contents of their digestive systems, especially the contents of their crops, also
known as their social stomachs (LeBoeuf, 2017). Organ level differences between gut
microbiomes were only statistically significant in the species *M. mexicanus.* While colony-level
differences between gut microbiome composition were statistically significant for *M. mexicanus,*
M. arenarius, and M. depilis they were not consistent across all species. Among all the species
examined, *M. mexicanus* showed a specialized resident microbiome, with replete crops
dominated almost entirely by *Fructilactobacillus.* This monodominance is striking when
compared to other species within the *Myrmecocystus* genus, which generally retained more
diverse crop microbial communities. Gut compartmentalization can play a role in shaping the
microbial community. For example, in *Cephalotes* turtle ants, specific gut compartments support

specialized and conserved symbiont communities (Flynn et al., 2021). Here, we observe
evidence of compartmentalization in the *M. mexicanus,* where the crop was dominated by
Fructilactobacillus, while the midgut exhibited more microbial diversity.
306
Lactic acid and acetic acid bacteria
The complete dominance of *Fructilactobacillus* was only observed in *M. mexicanus* replete
crops but it was also abundant in other species, just to a lesser degree (e.g., *M. depilis*). Where
there was less *Fructilactobacillus,* there were often other genera of lactic acid or acetic acid
bacteria present in high abundance (Fig 3). Lactic acid bacteria, particularly, are often found in
the guts of sugar-feeding social insects. In aphid-tending *Formicine* ants and *Lasius* ants, their
infrabuccal pockets and crops are abundant in lactic acid bacteria such as *Lactobacillus*,
Weissella, and *Fructobacillus* that are capable of catabolizing sugars commonly found in
hemipteran honeydew, including sucrose, trehalose, melezitose, and raffinose (Zheng et al.,
2022). Beyond catabolizing sugar-rich foods, lactic acid bacteria can inhibit microbial
competitors by producing lactic acid and other antimicrobial compounds, lowering pH, and
generating acidic conditions that many bacteria and fungi cannot tolerate, as demonstrated in
Mycocepurus smithii (Kellner et al., 2015). *Myrmecocystus* ants belong to the subfamily
Formicinae, which are known to produce and consume formic acid from modified poison glands.
Some ants in this subfamily have been shown to control their gut microbiomes by creating an
acidic crop environment through formic acid consumption (Tragust et al., 2020).
Fructilactobacillus, other lactic acid bacteria and acetic acid bacteria, which are acid tolerant,
could be thriving due to these conditions in *Myrmecocystus* guts or they could be contributing to
further acidification to control their environment in their favor. *Fructilactobacillus* is widely used
in the food industry for the fermentation of dairy, meat, sourdough, and vegetables (Hammes &
Vogel, 1995; Koga et al., 1998; Kellner et al., 2015; Byun et al., 2025). Acetic acid (vinegar) is
also used in food production and preservation, both with live culture fermentation and with
industrially derived vinegar (Sengun & Karabiyikli, 2011; De Roos & De Vuyst, 2018). Acetic
acid bacteria inhibit the growth of foodborne pathogens (e.g., *Salmonella* spp.), as well as
agents of food spoilage such as yeasts and molds (Levine & Fellers, 1940).
332
The acid-loving bacteria that dominate the digestive systems of *Myrmecocystus* ants could be
commensal, not providing any benefit or harm to their host, but simply growing in an ideal niche.
However, they could also be providing benefits to their hosts. First, they may facilitate the
metabolism of sugar-rich diets by efficiently catabolizing carbohydrates (e.g., hemipteran
honeydew and extrafloral nectar) stored in replete crops into sugars or other molecules that are
more easily metabolized by the ants. Second, they may contribute to host fitness by acidifying
the gut environment to inhibit the growth of pathogenic microbes. And third, their function may
be analogous to lacto fermentation for human food preservation, where *Fructilactobacillus*
prevents spoilage of nutrients in replete crops. In future work, we will examine the growth and
metabolism of *Myrmecocystus* lactic acid and acetic acid bacteria, starting with
Fructilactobacillus in *M. mexicanus* guts.
344

Halophiles

We also detected halophilic archaea and bacteria in multiple species collected in California, including *Haloparvum, Haloarcula, Halorubrum, Halobacterium* and *Natronomonas*. The high osmotic pressure generated by concentrated sugars in replete crops may favor the proliferation of halotolerant taxa, not because they are necessarily adapted to high salt environments, but because they are physiologically adapted to high osmotic pressures. Both high-salt and high-sugar environments reduce water activity and impose similar constraints on microbial growth (Kim et al., 2013). These halophiles could be adapted to the high sugar conditions within the crop. In *Myrmecocystus,* replete crops are concentrated in carbohydrate, composed primarily of fructose and glucose, with smaller amounts of maltose (Burgett & Young, 1974). Strains of *Tetragenococcus halophilus* typically associated with high-salt environments, have also been detected in sugar-rich beet juice (Justé et al., 2008; Lievens et al., 2015). Justé et al. (2008) found that some strains of *T. halophilus* were adapted to high salt, while others were adapted to high sugar environments, but not both. Halophilic archaea, including *Halobacterium* and *Haloarcula,* are capable of growing on sugars such as glucose, fructose, and sucrose (Johnsen et al., 2001). Together, these observations raise the possibility that halophilic bacteria and archaea detected in *Myrmecocystus* guts represent lineages adapted to high sugar conditions within the replete crop.

Where do the microbes come from?

The variation we see in microbiome composition across *Myrmecocystus spp.* is expected in social insects, where species in the same genus may differ in diet, habitat, foraging behavior, and exposure to environmental conditions (Hölldobler & Wilson, 1990). Even within *M. mexicanus*, variation among colonies and sampling periods suggest that beyond species identity, differences in vegetation, resource availability, and soil composition across habitats may influence microbial assembly across colonies. Similar patterns are observed in other social insects, where microbiome composition reflects environmental context. For example, *Oecophylla smaragdina* weaver ants gut microbiome composition differs between forest and urban habitats primarily due to variations in feeding habits and host diet (Chua et al., 2018).

Diet could contribute to the structuring of *Myrmecocystus* microbiome. *Myrmecocystus* foragers collect a wide range of resources, including gall secretions, aphid and coccid secretions, and extrafloral nectar (Burgett & Young, 1974). While all *Myrmecocystus* are considered scavengers, there are differences in dietary preferences and foraging habits among species (Snelling, 1976). Species that are primarily nocturnal, such as *M. mexicanus*, are thought to largely rely on liquid carbohydrate resources such as nectar and plant exudates (Snelling, 1976; Van Elst et al., 2021). Whereas more diurnal species, such as *M. wheeleri,* are described as more carnivorous, supplementing their diet with live or dead arthropods (Snelling, 1976; Van Elst et al., 2021). Comparative studies on Indo-Pacific ants, identified host dietary niche as the primary driver of microbial diversity and outweighing the contributions from the physical environment (Cunningham et al., 2025). In many cases, microorganisms do not just occupy the crop but actively define its physio-chemical state. In *Apis mellifera* honeybees, their diet, consisting of nectar and honey, introduces high levels of sucrose, glucose, and fructose in the crop, acting as a selective pressure that controls the assembly of local bacterial and fungal

communities (Callegari et al., 2021). Specialized microorganisms like acetic acid bacteria act as
oxygen scavengers to maintain suboxic conditions, while other fermentative bacteria produce
organic acids that acidify the gut environment (Callegari et al., 2021). We suspect that a similar
process is occurring in *Myrmecocystus,* where their diet favors microbes adapted to
carbohydrate metabolism, low pH tolerance, and osmotically stressful environments in the crop.
We expect that many members of the *Myrmecocystus* gut microbiome originate from their diets.
Many lactic acid bacteria that have been found in the crops of these ants have also been
isolated from nectar (Oliphant et al., 2023). It is also likely that halophilic archaea and bacteria
are also found in sources such as nectar with relatively high osmotic pressure. In the human
gut, it has been found that halophilic archaea have been associated with the dietary intake of
fermented and salted foods (Chang et al., 2008; Lee, 2013). We expect that diet could be a
source for microbiome community members, and that the crop conditions help to further shape
the community.
402
Another possible contributor to the origin of *Myrmecocystus* gut microbiome is through contact
of their immediate surroundings, such as soil-associated microbes. From our collected samples,
many of the colony entrances were located in close geographic proximity, which could increase
the likelihood of shared environmental exposure. In *Veromessor andrei* harvester ants, colonies
that are geographically closer harbor more similar microbial communities (Gamboa et al., 2025).
However, microbiome composition in this system is also shaped by host-associated factors
such as diet, life stage, and caste (Gamboa et al., 2025). If soil-associated microbes were a
dominant driver of microbiome assembly in *Myrmecocystus,* we would expect colonies that are
spatially close, regardless of species identity, to have similar microbial communities. Yet, our
results showed that colonies, such as *M. mexicanus* (X9 and X10) and *M. mimicus* (M1 and M2)
collected in July 2025, retained distinct species-specific microbiome compositions despite being
in close proximity (Fig. 3). Therefore, we do not expect that the immediate environment, namely
the soil, is the main contributor to the microbiome of these ants, however, more work will be
necessary to confirm this.
417
Moving forward
Future work will aim to disentangle the ecological and evolutionary drivers shaping these
microbial communities. In particular, we will be conducting field work to distinguish the
contributions of diet and physical environment (e.g., surrounding soil and food sources), as well
as to better understand how different transmission modes contribute to the shaping of the gut
community in *Myrmecocystus* honeypot ants. *Myrmecocystus* ants appear to have species-
specific resident gut microbiomes. Building on this finding the next steps are to examine the
relationship between the honeypot ants and their gut microbiome. We will investigate the
localization of these microorganisms throughout the digestive systems of the ants to better
understand if there are specialized structures for supporting these microorganisms. By
leveraging tools such as fluorescence in-situ hybridization and green fluorescent protein-tagged
microbes we will be able to explore the spatial distribution of lactic acid bacteria like
Fructilactobacillus or halophiles like *Haloparvum.* We will explore the nature of symbiosis
between the microbes and the ants (i.e., is the relationship commensal or mutualistic) through
various *in vivo* and *in vitro* physiological assays. We will couple these experiments with

metabolomics, transcriptomics, and proteomics to help identify the metabolic pathways,
metabolites, and functional roles of gut microbiome community members. Beyond work within
Myrmecocystus, comparative studies across convergently-evolved honeypot ant genera such as
Leptomyrmex, *Camponotus*, and *Tapinolepis* could provide insight into the role of host
physiology in structuring microbiome across independent evolutionary lineages.
438

439   **Acknowledgements**
440   The project was funded by NSF BRC-BIO grant 2312984 and seed funding through Rutgers
441   Global to LK and an NSF PRFB Award 2305685 to CF.
442

443   **Data Availability**
444   All raw sequencing data are available in the NCBI Sequence Read Archive (SRA) under the
445   Bioproject PRJNA1449520: *Myrmecocystus* honeypot ants 16s rRNA.
446

Author Contributions
We followed CRediT's 14 Contributor Roles guidelines. **DVN**: Conceptualization; Data Curation;
Formal Analysis; Investigation; Validation; Visualization; Writing (original draft); Writing (review
and editing). **CF**: Conceptualization; Data Curation; Funding Acquisition; Investigation;
Methodology; Visualization; Writing (original draft). **BN**: Investigation. **IS**: Investigation;
Methodology. **ML**: Investigation; Methodology. **LK**: Conceptualization; Funding Acquisition;
Project Administration; Supervision; Validation; Writing (review and editing)

470
471
472 Figure 1. Honeypot ant colonies contain two castes, workers (A) and repletes (B). Compared to
473 the crops of workers (C), the crops of repletes can expand to huge proportions as they are used
474 to store fluids. (E) Depending on the resources collected, repletes within the same colony or
475 chamber can have abdomens of different colors. Photo credits: (A) D. Caroll from buigguide.net,
476 (B) M.P. Meurville, (C) I. Sawh, (D) C. Francoeur, (E) M. Lanan.

477
Figure 2. Non-metric Multidimensional Scaling (NMDS) ordination based on weighted UniFrac
distances visualizes differences in microbial community composition across *Myrmecocystus*
species, caste, and organ. Each point represents individual samples, species indicated by color,
caste by shape, with panels corresponding to crop or midgut samples. Microbial communities
show species associated clustering, with partial overlap among species. Crop and midgut
occupy distinct regions of ordination space across species. Caste has a minimal effect on
community composition, as no clear separation is observed.

491
Figure 3. The bubble plot depicts the relative abundance (as percentage) of bacterial genera in
replete crop samples across *Myrmecocystus* species. Each point represents a genus within a
sample, with the bubble size indicating its relative abundance, and the color signifying genus
identity. Genus names are grouped by lactic acid bacteria (LAB = orange), acetic acid bacteria
(AAB = yellow), and halophilic archaea and bacteria (blue). The microbial communities in
replete crops exhibit species-specific structuring. In particular *M. mexicanus* replete crops are
strongly dominated by *Fructilactobacillus*, whereas other species show more variable
community composition.
500
501

502
Figure 4. NMDS ordination with weighted UniFrac distances comparing the microbial community
composition across caste replete (red) and worker (blue). Caste did not strongly structure gut
microbiomes across *Myrmecocystus* species. However, within certain species, *M. depilis, M.*
flaviceps, and *M. mexicanus,* replete and worker crops differed significantly in microbial
composition. Whereas the midgut communities remained similar between castes across all
species.

509
Figure 5. Shannon diversity index comparing the microbial diversity between replete crop and
midgut samples across *Myrmecocystus* species. Each point represents individual samples, with
boxplots summarizing the distribution of diversity for each organ within species. The midgut
communities generally exhibit higher diversity than crop communities, however, this difference
is only statistically significant in *M. mexicanus* (Wilcoxon rank-sum test, FDR-adjusted p =
1.82e[-13]). No significant differences in Shannon diversity were observed in other *Myrmecocystus*
species (all FDR-adjusted p > 0.05).
517

518
Figure 6. Differentially abundant bacterial genre between replete crop and midgut samples
identified using ANCOM-BC2. Negative Log$_2$ fold changes (red) indicate genera differentially
abundant in the crop, while positive log$_2$ fold change values (blue) indicate enrichment in the
midgut. (A) *Fructilactobacillus* and *Acinetobacter* were found to be differentially abundant in the
crop of *M. depilis*, while no genera was differentially abundant in the midgut. (B) Lactic acid
bacteria such as *Fructilactobacillus*, *Lentilactobacillus*, *Lacticaseibacillus*, and *Pediococcus*, are
differentially more abundant in the crop of *M. mexicanus* than the midgut.

References

1. Auer, L., Lazuka, A., Sillam-Dussès, D., Miambi, E., O'Donohue, M., & Hernandez-Raquet, G. (2017). Uncovering the Potential of Termite Gut Microbiome for Lignocellulose Bioconversion in Anaerobic Batch Bioreactors. *Frontiers in Microbiology*, *8*, 2623. https://doi.org/10.3389/fmicb.2017.02623

2. Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, *57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

3. Brune, A. (2014). Symbiotic digestion of lignocellulose in termite guts. *Nature Reviews Microbiology*, *12*(3), 168–180. https://doi.org/10.1038/nrmicro3182

4. Burgett, D. M., & Young, R. G. (1974). Lipid Storage by Honey Ant1 Repletes. *Annals of the Entomological Society of America*, *67*(5), 743–744. https://doi.org/10.1093/aesa/67.5.743

5. Byun, J. A., Han, N. R., Yun, E. J., Cheon, S., Song, S., Shim, S., Lee, S. H., & Kim, K. H. (2025). Enhancing lactic acid tolerance in Fructilactobacillus sanfranciscensis via adaptive evolution for sourdough fermentation applications. *International Journal of Food Microbiology*, *428*, 110978. https://doi.org/10.1016/j.ijfoodmicro.2024.110978

6. Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

7. Callegari, M., Crotti, E., Fusi, M., Marasco, R., Gonella, E., De Noni, I., Romano, D., Borin, S., Tsiamis, G., Cherif, A., Alma, A., & Daffonchio, D. (2021). Compartmentalization of bacterial and fungal microbiomes in the gut of adult honeybees. *Npj Biofilms and Microbiomes*, *7*(1), 42. https://doi.org/10.1038/s41522-021-00212-9

8. Chang, H.-W., Kim, K.-H., Nam, Y.-D., Roh, S. W., Kim, M.-S., Jeon, C. O., Oh, H.-M., &

Bae, J.-W. (2008). Analysis of yeast and archaeal population dynamics in kimchi using

denaturing gradient gel electrophoresis. *International Journal of Food Microbiology*,

126(1–2), 159–166. https://doi.org/10.1016/j.ijfoodmicro.2008.05.013

9. Chua, K.-O., Song, S.-L., Yong, H.-S., See-Too, W.-S., Yin, W.-F., & Chan, K.-G. (2018).

Microbial Community Composition Reveals Spatial Variation and Distinctive Core

Microbiome of the Weaver Ant Oecophylla smaragdina in Malaysia. *Scientific Reports*,

8(1), 10777. https://doi.org/10.1038/s41598-018-29159-2

10. Conway, J. R. (1977). Analysis of clear and dark amber repletes of the honey ant,

Myrmecocystus mexicanus hortideorum. *Annals of the Entomological Society of*

America, *70*(3), 367–369. https://doi.org/10.1093/aesa/70.3.367

11. Cunningham, P., Vesey, R., Chaudry-Phipps, H., Fayle, T. M., Diara, U., Chomicki, G.,

Lestina, D., Tuma, J., Templer, P. H., Klimes, P., & Henry, L. M. (2025). Dietary niche

shapes bacterial community in Indo-Pacific ants. *Microbiology Spectrum*, *13*(10),

e01965-25. https://doi.org/10.1128/spectrum.01965-25

12. Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A., & Callahan, B. J. (2018).

Simple statistical identification and removal of contaminant sequences in marker-gene

and metagenomics data. *Microbiome*, *6*(1), 226. https://doi.org/10.1186/s40168-018-

0605-2

13. De Roos, J., & De Vuyst, L. (2018). Acetic acid bacteria in fermented foods and

beverages. *Current Opinion in Biotechnology*, *49*, 115–119.

https://doi.org/10.1016/j.copbio.2017.08.007

14. Dong, A. Z., Cokcetin, N., Carter, D. A., & Fernandes, K. E. (2023). Unique antimicrobial

activity in honey from the Australian honeypot ant (*Camponotus inflatus*). *PeerJ*, *11*,

e15645. https://doi.org/10.7717/peerj.15645

15. Eisner, T., & Brown, W. L. J. (1956). *The evolution and social significance of the ant*
proventriculus. 2, 503–508.

16. Engel, P., & Moran, N. A. (2013). The gut microbiota of insects – diversity in structure
and function. *FEMS Microbiology Reviews*, *37*(5), 699–735.
https://doi.org/10.1111/1574-6976.12025

17. Feldhaar, H., Straka, J., Krischke, M., Berthold, K., Stoll, S., Mueller, M. J., & Gross, R.
(2007). Nutritional upgrading for omnivorous carpenter ants by the endosymbiont
Blochmannia. *BMC Biology*, *5*(1), 48. https://doi.org/10.1186/1741-7007-5-48

18. Flynn, P. J., D'Amelio, C. L., Sanders, J. G., Russell, J. A., & Moreau, C. S. (2021).
Localization of Bacterial Communities within Gut Compartments across *Cephalotes*
Turtle Ants. *Applied and Environmental Microbiology*, *87*(8), e02803-20.
https://doi.org/10.1128/AEM.02803-20

19. Gamboa, D. A., Flynn, P. J., Horna-Lowell, E. S., & Pinter-Wollman, N. (2025). Social
organization and physical environment shape the microbiome of harvester ants. *Animal*
Microbiome, *7*(1), 29. https://doi.org/10.1186/s42523-025-00390-3

20. Hammer, T. J., Janzen, D. H., Hallwachs, W., Jaffe, S. P., & Fierer, N. (2017).
Caterpillars lack a resident gut microbiome. *Proceedings of the National Academy of*
Sciences, *114*(36), 9641–9646. https://doi.org/10.1073/pnas.1707186114

21. Hammes, W. P., & Vogel, R. F. (1995). The genus Lactobacillus. In B. J. B. Wood & W.
H. Holzapfel (Eds.), *The Genera of Lactic Acid Bacteria* (pp. 19–54). Springer US.
https://doi.org/10.1007/978-1-4615-5817-0_3

22. Hölldobler, B., & Wilson, E. O. (1990). *The ants*. Harvard University Press.

23. Hu, Y., Sanders, J. G., Łukasik, P., D'Amelio, C. L., Millar, J. S., Vann, D. R., Lan, Y.,
Newton, J. A., Schotanus, M., Kronauer, D. J. C., Pierce, N. E., Moreau, C. S., Wertz, J.
T., Engel, P., & Russell, J. A. (2018). Herbivorous turtle ants obtain essential nutrients

from a conserved nitrogen-recycling gut microbiome. *Nature Communications*, *9*(1), 964.

https://doi.org/10.1038/s41467-018-03357-y

24. Johnsen, U., Selig, M., Xavier, K. B., Santos, H., & Sch nheit, P. (2001). Different

glycolytic pathways for glucose and fructose in the halophilic archaeon Halococcus

saccharolyticus. *Archives of Microbiology*, *175*(1), 52–61.

https://doi.org/10.1007/s002030000237

25. Jones, J. A., Moczek, A. P., & Newton, I. L. G. (2025). The dung beetle microbiome

complements host metabolism and nutrition. *mSystems*, *10*(11), e01172-25.

https://doi.org/10.1128/msystems.01172-25

26. Justé, A., Lievens, B., Frans, I., Marsh, T. L., Klingeberg, M., Michiels, C. W., & Willems,

K. A. (2008). Genetic and physiological diversity of Tetragenococcus halophilus strains

isolated from sugar- and salt-rich environments. *Microbiology*, *154*(9), 2600–2610.

https://doi.org/10.1099/mic.0.2008/018168-0

27. Katoh, K., Misawa, K., Kuma, K., & Miyata, T. (2002). MAFFT: A novel method for rapid

multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research*,

30(14), 3059–3066. https://doi.org/10.1093/nar/gkf436

28. Kellner, K., Ishak, H. D., Linksvayer, T. A., & Mueller, U. G. (2015). Bacterial community

composition and diversity in an ancestral ant fungus symbiosis. *FEMS Microbiology*

Ecology, *91*(7), fiv073. https://doi.org/10.1093/femsec/fiv073

29. Kim, J. (Jeannie), Enache, E., & Hayman, M. (2013). 17. Halophilic and Osmophilic

Microorganisms. In *Compendium of Methods for the Microbiological Examination of*

Foods. American Public Health Association. https://doi.org/10.2105/MBEF.0222.022

30. Koga, T., Mizobel, T., & Takumi, K. (1998). Antibacterial activity of Lactobacillus species

against Vibrio species. *Microbiological Research*, *153*(3), 271–275.

https://doi.org/10.1016/S0944-5013(98)80011-6

31. Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013).

Development of a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing

Amplicon Sequence Data on the MiSeq Illumina Sequencing Platform. *Applied and*

Environmental Microbiology, *79*(17), 5112–5120. https://doi.org/10.1128/AEM.01043-13

32. Kruskal, W. H., & Wallis, W. A. (1952). *Use of Ranks in One-Criterion Variance Analysis*.

33. LeBoeuf, A. C. (2017). Trophallaxis. *Current Biology*, *27*(24), R1299–R1300.

https://doi.org/10.1016/j.cub.2017.10.047

34. Lee, H.-S. (2013). Diversity of Halophilic Archaea in Fermented Foods and Human

Intestines and Their Application. *Journal of Microbiology and Biotechnology*, *23*(12),

1645–1653. https://doi.org/10.4014/jmb.1308.08015

35. Levine, A. S., & Fellers, C. R. (1940). Action of Acetic Acid on Food Spoilage

Microörganisms. *Journal of Bacteriology*, *39*(5), 499–515.

https://doi.org/10.1128/jb.39.5.499-515.1940

36. Lievens, B., Hallsworth, J. E., Pozo, M. I., Belgacem, Z. B., Stevenson, A., Willems, K.

A., & Jacquemyn, H. (2015). Microbiology of sugar-rich environments: Diversity, ecology

and system constraints. *Environmental Microbiology*, *17*(2), 278–298.

https://doi.org/10.1111/1462-2920.12570

37. Lin, H., & Peddada, S. D. (2020). Analysis of compositions of microbiomes with bias

correction. *Nature Communications*, *11*(1), 3514. https://doi.org/10.1038/s41467-020-

17041-7

38. McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible

Interactive Analysis and Graphics of Microbiome Census Data. *PLOS ONE*, *8*(4),

e61217. https://doi.org/10.1371/journal.pone.0061217

39. Meurville, M.-P., & LeBoeuf, A. C. (2021). *Trophallaxis: The functions and evolution of*

social fluid exchange in ant colonies (Hymenoptera: Formicidae).

https://doi.org/10.25849/MYRMECOL.NEWS_031:001

40. Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von

Haeseler, A., & Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era. *Molecular Biology and Evolution*, *37*(5),

1530–1534. https://doi.org/10.1093/molbev/msaa015

41. Motta, E. V., Gage, A., Smith, T. E., Blake, K. J., Kwong, W. K., Riddington, I. M., &

Moran, N. (2022). Host-microbiome metabolism of a plant toxin in bees. *eLife*, *11*,

e82595. https://doi.org/10.7554/eLife.82595

42. Nogueira, B. R., LeBoeuf, A. C., Barden, P., Nguyen, D. Q. V., & Khadempour, L.

(2026). *Honeypots: A review of repletism across the ants*.

https://doi.org/10.25849/MYRMECOL.NEWS_036:039

43. Oksanen, J., Simpson, G., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P., hara, R.,

Solymos, P., STEVENS, H., Szöcs, E., Wagner, H., Barbour, M., Bedward, M., Bolker,

B., Borcard, D., Carvalho, G., Chirico, M., De Cáceres, M., Durand, S., & Weedon, J.

(2022). *Vegan community ecology package version 2.6-2 April 2022*.

44. Oliphant, S. A., Watson-Haigh, N. S., Sumby, K. M., Gardner, J. M., & Jiranek, V.

(2023). Fructilactobacillus cliffordii sp. Nov., Fructilactobacillus hinvesii sp. Nov.,

Fructilactobacillus myrtifloralis sp. Nov., Fructilactobacillus carniphilus sp. Nov. And

Fructobacillus americanaquae sp. Nov., five novel lactic acid bacteria isolated from

insects or flowers of Kangaroo Island, South Australia. *International Journal of*

Systematic and Evolutionary Microbiology, *73*(2). https://doi.org/10.1099/ijsem.0.005730

45. Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of Phylogenetics and

Evolution in R language. *Bioinformatics*, *20*(2), 289–290.

https://doi.org/10.1093/bioinformatics/btg412

46. Sauer, C., Dudaczek, D., Hölldobler, B., & Gross, R. (2002). Tissue Localization of the

Endosymbiotic Bacterium " *Candidatus* Blochmannia floridanus" in Adults and Larvae of

the Carpenter Ant *Camponotus floridanus. Applied and Environmental Microbiology*,

68(9), 4187–4193. https://doi.org/10.1128/AEM.68.9.4187-4193.2002

47. Sawh, I. (2023). *Exploring repletism in honeypot ants through an evolutionary and*

microbial lens [Rutgers University - Graduate School - Newark].

https://doi.org/10.7282/t3-vwe9-6f82

48. Schliep, K. P. (2011). phangorn: Phylogenetic analysis in R. *Bioinformatics*, *27*(4), 592–

593. https://doi.org/10.1093/bioinformatics/btq706

49. Sengun, I. Y., & Karabiyikli, S. (2011). Importance of acetic acid bacteria in food

industry. *Food Control*, *22*(5), 647–656. https://doi.org/10.1016/j.foodcont.2010.11.008

50. Shannon, C. E. (1948). A mathematical theory of communication. *The Bell System*

Technical Journal, *27*(3), 379–423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x

51. Simpson, E. H. (1949). Measurement of Diversity. *Nature*, *163*(4148), 688–688.

https://doi.org/10.1038/163688a0

52. Snelling, R. (1976). *A Revision of the Honey Ants, Genus Myrmecocystus*

(Hymenoptera: Formicidae). https://www.livre-rare-book.com/book/5472473/EH40591

53. Tragust, S., Herrmann, C., Häfner, J., Braasch, R., Tilgen, C., Hoock, M., Milidakis, M.

A., Gross, R., & Feldhaar, H. (2020). Formicine ants swallow their highly acidic poison

for gut microbial selection and control. *eLife*, *9*, e60287.

https://doi.org/10.7554/eLife.60287

54. Van Elst, T., Eriksson, T. H., Gadau, J., Johnson, R. A., Rabeling, C., Taylor, J. E., &

Borowiec, M. L. (2021). Comprehensive phylogeny of Myrmecocystus honey ants

highlights cryptic diversity and infers evolution during aridification of the American

Southwest. *Molecular Phylogenetics and Evolution*, *155*, 107036.

https://doi.org/10.1016/j.ympev.2020.107036

55. White, Bruns, T., Lee, S., & Taylor, J. (1990). *White, T. J., T. D. Bruns, S. B. Lee, and J.*

W. Taylor. Amplification and direct sequencing of fungal ribosomal RNA Genes for

phylogenetics (pp. 315–322).

56. Wickham, H. (2011). The Split-Apply-Combine Strategy for Data Analysis. *Journal of*

Statistical Software, *40*, 1–29. https://doi.org/10.18637/jss.v040.i01

57. Wickham, H. (2016). *Ggplot2*. Springer International Publishing.

https://doi.org/10.1007/978-3-319-24277-4

58. Wilcoxon, F. (1945). *Individual Comparisons by Ranking Methods*.

59. Zheng, Z., Zhao, M., Zhang, Z., Hu, X., Xu, Y., Wei, C., & He, H. (2022). Lactic Acid

Bacteria Are Prevalent in the Infrabuccal Pockets and Crops of Ants That Prefer Aphid

Honeydew. *Frontiers in Microbiology*, *12*, 785016.

https://doi.org/10.3389/fmicb.2021.785016