Glucose-dependent signalling pathways regulate TE differentiation in bovine embryos

Jinlong Qiu[1] and Roger Sturmey[1#]

1. Centre for Biomedicine, Hull York Medical School, University of Hull, Hull, HU6 7RX, UK

Authors for correspondence (Roger.Sturmey@hyms.ac.uk)

Abstract

Glucose uptake and metabolism increase markedly during the transition from morula to blastocyst. However, the functional roles of glucose during this process in bovine embryos remain incompletely understood. Here, we demonstrate that glucose specifically modulates trophectoderm (TE) differentiation without affecting the inner cell mass (ICM), acting through Hippo signalling via the hexosamine biosynthetic pathway (HBP) and the pentose phosphate pathway (PPP) rather than glycolysis. Consistently, scRNA-seq analysis reveals that glycolysis-related genes show only stage-specific changes, whereas HBP- and PPP-related genes exhibit both stage- and lineage-specific differences between TE and ICM. Furthermore, we found that glucose deprivation does not impair pyruvate uptake or intracellular pyruvate levels in blastocysts, yet it significantly depletes the terminal metabolites of the HBP. Together, these findings reveal that glucose-dependent regulation of TE differentiation is conserved in bovine embryos.

Introduction

Glucose is a key metabolic substrate during early embryonic development [1, 2]. Classical models of glucose metabolism presumed that glucose is predominantly catabolized through glycolysis to generate pyruvate, which enters the tricarboxylic acid (TCA) cycle to satisfy the energetic requirements of cell proliferation and development [3, 4].

However, recent studies in mouse embryos have challenged this traditional bioenergetic view of glucose metabolism [5-7]. These studies demonstrate that, during early preimplantation development, glucose contributes only minimally to glycolytic flux and ATP production [5-7]. Instead, glucose-derived carbon is preferentially channeled into HBP and PPP, supporting biosynthetic processes and redox homeostasis [5, 7].

In particular, HBP-derived O-linked N-acetylglucosaminylation (O-GlcNAc) has been implicated in the regulation of the Hippo signaling pathway, thereby influencing TE-associated transcriptional programs [5, 8, 9]. In parallel, the PPP has been shown to function upstream of mechanistic target of rapamycin (mTOR) signaling to regulate TE specification [5, 10, 11]. Notably, whether this glucose-driven signaling paradigm is conserved in bovine embryos, remains unclear.

In this study, we demonstrate that although glucose-mediated regulation of the TE program is conserved in bovine embryos, their metabolic dependency appears substantially more complex than that observed in murine models. Notably, bovine embryos exhibit a greater tolerance to glucose deprivation, as blastocyst formation can still occur in the absence of glucose. Furthermore, the contributions of HBP and PPP appear to be highly context-dependent, with developmental outcomes varying according to the specific stage at which inhibition occurs and the metabolic nodes targeted. Collectively, our findings highlight a critical role for glucose in TE formation in large mammals while revealing a sophisticated and complex metabolic regulatory network underlying this process.

Materials and methods

Unless specified otherwise, all reagents and chemicals were obtained from Merck (Dorset, UK).

In vitro bovine embryo production

In vitro maturation (IVM), fertilization (IVF), and culture (IVC) of bovine embryos were performed according to previously described protocols with minor modifications [12-14]. IVM, IVF, and SemenPrep media were obtained from IVF Bioscience (Falmouth, UK). Bovine ovaries were harvested from animals slaughtered for commercial purposes unrelated to the study. Ovaries were collected and transported to the laboratory within 2 hours of slaughter. Cumulus–oocyte complexes (COCs) were collected from 3–8 mm follicles on the ovarian surface. COCs were cultured in IVM medium at 38.5°C, 5% CO_2 for 22 hours. Following maturation, groups of 25–30 COCs were co-incubated with spermatozoa (Genus Plc) ($1–5 \times 10^6$/mL) purified from frozen–thawed semen using SemenPrep medium. IVF was carried out at 38.5°C, 5% CO_2 for 10–14 hours. After fertilization, cumulus cells were removed. Presumptive zygotes were subsequently cultured in BO-SOF medium supplemented with either 0 mM or 1.5 mM glucose at 39 °C, 5% CO_2 and 5% O_2.

The BO-SOF formulation was prepared according to the protocol described in our previous publication [15, 16]. The concentrations were selected based on previously published studies [5, 17]. The inhibitors used in this study are listed in Supplementary Table 1. Unless otherwise stated, inhibitors were added from E0.

Metabolic assay

Morula-stage embryos were individually cultured for 24 hours in 5-μL microdrops, after which embryos that successfully developed to the blastocyst stage were identified and their spent culture media were collected. All samples were immediately stored at −78°C until subsequent metabolic assays. Pyruvate concentrations were quantified using enzyme-coupled fluorometric methods adapted from a previously established protocol [15]. All enzymes used for metabolic analyses were obtained from Roche (Burgess Hill, UK). Standard curves were generated from serial dilutions of freshly prepared

metabolite standards. Fluorescence values were calculated as the difference between pre- and post-incubation readings and converted to absolute concentrations. All measurements were performed in technical triplicate, and assays with standard curve R^2 values < 0.98 were excluded from further analysis.

Measurement of Basal Oxygen Consumption Rate (OCR)

The OCR detection protocol is based on previously published articles [16]. Briefly, OCR was assessed using a Seahorse XFp Analyzer (Agilent Technologies). Seahorse analysis was performed following the manufacturer's recommendations for OCR. Embryos were allocated in small groups (6-10 bovine blastocysts per well) in 180 μl of HEPES-buffered SOF medium appropriate for the developmental stage.

JC-1 staining

JC-1 (Thermo Fisher) was initially dissolved in DMSO and subsequently diluted to a final concentration of 1 μM in SOF medium. Embryos were incubated in this solution for 30 minutes under standard culture conditions. Following three washes with HEPES-buffered SOF, the embryos were imaged using a Zeiss LSM710 confocal microscope.

IF

Samples were first washed three times with 0.1% polyvinylpyrrolidone/phosphate-buffered saline (PBS), then fixed with 4% paraformaldehyde/PBS for 30 minutes and then permeabilized with 0.5% Triton X-100/PBS for 30 minutes at room temperature. The blocking step was performed in blocking buffer (PBS containing 10% FBS and 0.1% Triton X-100) for 1 hour. Afterwards, samples were incubated overnight with primary antibodies in blocking buffer at 4°C. After three washes, embryos were incubated with secondary antibodies for 2 hours. Finally, samples were stained with Hoechst3342 (Hoechst) (Merck) for 30 minutes and then mounted. Images were acquired using inverted fluorescence microscope (Zeiss AXIO Vert.A1) or a laser scanning confocal microscope (Zeiss LSM 710) with a 40x objective lens.

FIJI (https://imagej.net/software/fiji/) was used for image visualization, cell counting, and signal intensity measurements. Depending on the experiment, signal intensity was

112 measured and background subtracted to analyze absolute intensity. All antibodies

113 (Supplementary Table S2 and S3) were validated to react with bovine cells [12, 18, 19].

Single-cell RNA-seq data acquisition and processing

The single-cell RNA sequencing (scRNA-seq) dataset of bovine preimplantation

embryos analyzed in this study was obtained from the publicly available Gene

Expression Omnibus (GEO) database under accession number GSE239782 [20]. This

dataset contains transcriptomic profiles of bovine embryos at defined developmental

stages, including morula, early blastocyst, mid-blastocyst, and late blastocyst stages.

Raw count matrices were downloaded and processed using the Seurat R package

(version 4.5.2). The calculation of module scores and all analysis codes were adapted

from previously published studies [21].

Statistical Analysis

124 Unless otherwise noted, all experiments were performed with at least three biological

125 replicates. Differences between groups were evaluated using two-tailed unpaired

126 Student's t-tests. Graphs were generated with GraphPad Prism 10.

Results

Glucose is essential for blastocyst formation

Embryos were cultured from embryonic day 0 (E0) in glucose-free medium (G−) or in

medium supplemented with 1.5 mM glucose (CTRL). No significant differences were

observed in developmental rates at the 4-cell stage (E2.0), the 8–16-cell stage (E3.5), or

the morula stage (E5.0) between G− and CTRL embryos (Fig. 1A, B). However, blastocyst

formation in the G− group was markedly reduced at E6.5 (Fig. 1A, B), and the total cell

numbers of blastocysts were significantly decreased, with this decline already apparent

at the morula stage (Fig. S1A–D). Notably, the few G− embryos that reached the

blastocyst stage failed to hatch normally at E8.5 (Fig. 1C, D). Inhibition of glucose

metabolism using the hexokinase inhibitor 2-deoxy-D-glucose (2-DG) similarly reduced

blastocyst rate, confirming the important role of glucose in blastocyst formation (Fig.

S1F, G).

To dissect the contributions of different glucose metabolic pathways, specific inhibitors

were used to block glycolysis, HBP, and PPP (Fig. S1E). Inhibiting glucose flux into

pyruvate with YZ9 had no effect on blastocyst formation (Fig. 1E, F). In contrast,

blocking the PPP with 6-aminonicotinamide (6-AN) or the HBP with 6-Diazo-5-oxo-L-

norleucine (DON) induced morula-stage developmental arrest, which could not be

rescued by supplementation with their downstream metabolites (Fig. S1H–K). However,

inhibition of the downstream HBP enzyme O-GlcNAc transferase (OGT) using

ST045849(TT04) had no effect on blastocyst formation (Fig. 1G, H). In addition, when

PPP inhibition was delayed until E5.5, embryos were able to reach the blastocyst stage,

although the blastocyst formation rate was significantly reduced (Fig. 1I, J). Together,

these results demonstrate that glucose plays a role in blastocyst formation, with

distinct contributions from its metabolic branches.

Glucose metabolism regulates TE through Hippo pathway

Immunofluorescence analysis was performed on E7.0 blastocysts cultured under CTRL

or G− conditions to assess the abundance of key lineage-specific transcription factors

(Fig 2A). Quantitative analysis revealed that glucose deprivation significantly reduced the

fluorescence intensity of the TE marker CDX2 (Fig 2B) and decreased the proportion of

CDX2$^+$ cells within the blastocyst (Fig 2C). Consistently, expression of an additional TE

marker, GATA3, was also reduced under G- conditions (Fig S 2A, B).

In contrast, neither the expression level of the ICM marker SOX2 nor the proportion of

SOX2$^+$ cells was significantly altered (Fig. 2B, C). Consistently, the expression levels and

cell numbers of the epiblast (EPI) marker NANOG and the primitive endoderm (PrE)

marker GATA6 were comparable between groups (Fig. 2D–F), indicating that glucose

predominantly affects TE differentiation rather than ICM lineage specification.

Given the central role of Hippo signalling in TE specification [22, 23], Hippo pathway

activity was next examined. Immunofluorescence analyses revealed a significant

reduction in phosphorylated YAP (p-YAP) levels in G- embryos (Fig 2I, J). Consistently,

nuclear localisation of YAP, as well as the TE-associated transcription factors TFAP2C and

TEAD4, was decreased under G- conditions (Fig 2G, H, K, L), indicating that glucose

regulates TE through Hippo pathway.

Similar phenotypes were observed following inhibition of glucose metabolism 2-DG.

Treatment with 2-DG significantly reduced total cell numbers and altered the TE-to-ICM

cell ratio in E7.0 blastocysts (Fig S 2E), accompanied by selective suppression of CDX2

expression without affecting SOX2 levels (Fig S 2C–F). Moreover, expression of YAP,

TFAP2C, and p-YAP was significantly decreased following 2-DG treatment (Fig S 2G–J),

supporting the conclusion that glucose specifically regulates TE differentiation via Hippo

signalling.

Glucose availability regulates HBP activity and epigenetic landscapes

To investigate the metabolic mechanisms by which glucose regulates TE differentiation

via the Hippo pathway, we re-analyzed publicly available metabolomics datasets from

bovine blastocysts cultured under CTRL and G− conditions [24]. Glucose deprivation

led to a broad reduction in intermediate metabolites across glycolysis, HBP and PPP

(Fig. S3A). However, the end products of glycolysis and PPP remained unchanged, as

pyruvate and ribose levels were unaffected (Fig. S3A). In contrast, HBP metabolites,

including N-acetylglucosamine (GlcNAc) and UDP-GlcNAc, were markedly reduced

(Fig. S3A), highlighting a selective vulnerability of the HBP to glucose deprivation.

To validate our metabolomics findings, we first measured mitochondrial membrane

potential ($\Delta\Psi m$) to assess the impact of glucose availability on cellular energy

homeostasis. Glucose deprivation significantly reduced the orange/green JC-1

fluorescence ratio in bovine blastocysts, indicative of mitochondrial depolarization (Fig

3A, B). However, basal oxygen consumption rates (OCR) measured by Seahorse

analysis were comparable between G− and CTRL embryos (Fig 3C). This suggests that

while glucose is essential for maintaining mitochondrial membrane potential, the

embryos may sustain basal respiratory flux through the utilization of alternative

metabolic substrates [25, 26].

To determine whether embryos compensate for the absence of glucose by increasing

reliance on exogenous pyruvate, pyruvate consumption from the culture medium was

assessed. Measurement of pyruvate concentrations in spent culture media revealed no

significant differences between CTRL and G- embryos (Fig 3D). Consistently, 2-DG or

YZ9 inhibition did not significantly alter pyruvate concentration as well (Fig 3E, F).

Collectively, these findings indicate that embryos do not trigger a compensatory

increase in exogenous pyruvate uptake to offset the energy deficit caused by glucose

deprivation.

To validate the impact of glucose deprivation on the HBP identified in our metabolomics

re-analysis, we assessed global O-GlcNAc levels via immunofluorescence. Consistent

with the marked depletion of HBP metabolites, O-GlcNAc levels were significantly

reduced in blastocysts cultured under G- conditions or treated with 2-DG (Fig 3G, H and

Fig S 3C, D). These results confirm that restricted glucose availability directly

compromises HBP-mediated protein glycosylation in bovine embryos.

In parallel, metabolomics re-analysis revealed significant alterations in metabolites

associated with amino acid methylation, including those linked to lysine trimethylation

(Kme3) and arginine methylation (ADMA and SDMA) (Fig. S3B). Guided by these

observations, we examined the impact of glucose deprivation and metabolic inhibition

on epigenetic modifications. Both glucose deprivation and 2-DG treatment markedly

reduced ADMA and SDMA levels, accompanied by a pronounced decrease in the

repressive histone mark H3K27me3 (Fig. 3I–N and Fig. S3C–J). These findings indicate

that glucose availability is a key determinant of the epigenetic landscape in bovine

blastocysts.

HBP-Dependent O-GlcNAc Regulates CDX2

To determine whether the HBP signaling axis operates through O-GlcNAc during bovine

blastocyst development, we first confirmed that TT04 treatment significantly reduced

global O-GlcNAc levels (Fig. 4G, H). We then assessed the impact of O-GlcNAc

inhibition on lineage specification. TT04 treatment specifically decreased CDX2

expression without affecting SOX2 levels (Fig. 4A, B), while total cell number and the

relative proportions of TE and ICM cells remained unchanged (Fig. 4C, D).

Given the reduction in CDX2 expression, we next examined the Hippo signaling

pathway. TT04 treatment significantly reduced total YAP protein levels without altering

TFAP2C expression (Fig. 4E, F). In addition, TT04 markedly increased phosphorylated

YAP (p-YAP) levels (Fig. 4G, H), indicating enhanced YAP phosphorylation and

cytoplasmic retention upon O-GlcNAc inhibition.

Collectively, these findings demonstrate that O-GlcNAc acts as a key metabolic signal

that maintains TE identity by stabilizing YAP and preventing its phosphorylation.

A PPP–mTOR Pathway Regulates TFAP2C

Glucose-derived nucleotides generated via the PPP act as nutrient signals that activate

mTOR signaling [5, 27]. To assess mTORC1 activity under glucose deprivation or PPP

inhibition, phosphorylated ribosomal protein S6 (pS236-RPS6) was used as a

downstream readout. Immunofluorescence analysis revealed a marked reduction in

pS236-RPS6 levels in both G− and 6-AN–treated blastocysts compared with CTRL

embryos (Fig. 5A–D), indicating that PPP activity is required to sustain mTORC1

signaling during bovine blastocyst development.

Given that mTOR promotes mRNA translation through phosphorylation of 4E-BP1 and

RPS6 [5, 10, 11], we next examined whether PPP regulates TFAP2C expression via

mTOR. Embryos were treated with the mTOR inhibitor INK128 from E5.5, which

effectively suppressed mTORC1 activity, as evidenced by reduced pS236-RPS6 levels

(Fig. 5G, H). Notably, mTOR inhibition did not affect the overall blastocyst formation rate

(Fig. 5E, F), but significantly reduced TFAP2C and CDX2 expression (Fig. 5I, J).

Quantitative analysis further revealed a reduction in total blastocyst cell number

following INK128 treatment, primarily due to a loss of TE (CDX2$^+$) cells, while ICM

(CDX2$^-$) cell numbers remained unchanged (Fig. 5K). Collectively, these findings

identify mTORC1 as a key glucose-responsive signaling node that links PPP-derived

nucleotide metabolism to TFAP2C expression and TE lineage expansion.

Single-cell transcriptomic profiling reveals lineage-specific metabolic signatures in bovine blastocysts

To further delineate the metabolic dynamics during bovine preimplantation development and corroborate our in vitro findings, we analyzed single-cell transcriptomic data spanning from the morula to the late blastocyst stages. UMAP dimensionality reduction successfully reconstructed the developmental trajectory and clearly segregated the distinct early embryonic cell lineages, including the TE, ICM, EPI, and PE (Fig. S4A). Heatmap analysis of metabolic gene modules revealed that oxidative phosphorylation (OXPHOS), glycolysis, the HBP, and the PPP exhibit distinct stage- and lineage-specific expression patterns during early embryogenesis (Fig. S4C and Fig 6 A, D, G). To robustly quantify these pathway-level changes and overcome the sparsity of single-cell data, we calculated metabolic module scores, which reflect the overall transcriptional enrichment of these entire metabolic networks within individual cells.

To establish a macroscopic understanding of the metabolic landscape, we first evaluated the comprehensive module scores across all identified developmental stages and cell lineages (Fig. S4B). This overview revealed that OXPHOS maintains a consistently high and relatively stable transcriptional baseline throughout preimplantation development, likely fulfilling the basal energetic demands of the embryo. In contrast, the glucose-driven pathways displayed lower overall absolute scores but exhibited highly dynamic fluctuations, prompting us to further dissect their stage- and lineage-specific variations (Fig. S4B).

We next quantitatively assessed the dynamic shifts in these metabolic modules as embryos progressed through preimplantation development. Bar graphs comparing grouped stages revealed that during the transition from the morula to the mid-blastocyst (MB) stage, the expression modules for both OXPHOS and glycolysis generally increased (Fig. S4D and Fig 6B). However, at the late blastocyst (LB) stage, OXPHOS scores underwent a significant decline, whereas glycolysis continued to sharply rise (Fig. S4D and Fig 6B). This divergent pattern indicates a fundamental metabolic switch in energy production strategies, transitioning toward a predominantly

glycolytic state, which is highly consistent with previously reported metabolic shifts in

mammalian embryogenesis [21, 28-30].

In contrast to the continuous trajectory of glycolysis, the dynamic changes of the

glucose-driven anabolic branches, HBP and PPP, were more complex. The HBP score

initially exhibited a transient dip at the early blastocyst (EB) stage before steadily

upregulating and peaking at the LB stage (Fig. 6F). Similarly, the PPP module showed a

high baseline at the morula stage, followed by a decline at the EB stage, and a

subsequent recovery and continuous increase through the MB and LB stages (Fig. 6H).

Collectively, these results indicate that glucose-driven metabolic pathways undergo

pronounced fluctuations during the morula-to-blastocyst transition.

Importantly, we investigated whether these glucose-driven pathways exhibited lineage-

specific partitioning corresponding to our in vitro phenotypes. Quantitative comparison

of module scores revealed highly nuanced, pathway-specific metabolic wiring among

the cell lineages at the mid- and late-blastocyst stages (Fig. S4B). Notably, the

glycolysis scores showed no significant differences between the TE and ICM lineages

(Fig. 6C). This finding perfectly aligns with our earlier in vitro observation that inhibiting

high-flux glycolysis with YZ9 does not selectively impair TE specification (Fig. S2K, M),

confirming that basal glycolysis is broadly utilized rather than acting as a lineage-

determining driver.

In stark contrast, the HBP module was robustly and specifically enriched in the TE

compared to the ICM at the critical mid-blastocyst stage (Fig. 6F). This striking

differential expression provides compelling in vivo transcriptomic evidence that HBP

activity is preferentially partitioned to the TE, mechanistically explaining why TE

specification relies heavily on the HBP-O-GlcNAc-YAP signaling axis.

In addition, the PPP module displayed high activity in both embryonic lineages. Notably,

the module score in the MB ICM was significantly higher than that in MB TE, whereas LB

TE scores were significantly higher than LB PE (Fig. 6I). This dynamic lineage-specific

shift suggests that both ICM and TE rely on PPP during blastocyst development, albeit

with different temporal requirements. The elevated PPP activity likely reflects the

increased demand for nucleotide biosynthesis and redox homeostasis required to

sustain rapid cellular proliferation [4, 5, 31, 32]. Consequently, inhibition of PPP is

expected to affect both lineages and may therefore produce more severe

developmental phenotypes than inhibition of other glucose-derived pathways. This

robust and ubiquitous expression pattern perfectly aligns with our in vitro observations:

pharmacological suppression of the PPP using 6-AN starting from the zygote stage

results in a catastrophic developmental arrest at the morula stage (Fig S1H, I).

Discussion

In this study, we demonstrate that bovine embryos exhibit a markedly higher tolerance

to glucose deprivation compared with murine models. Although the absence of glucose

significantly reduced blastocyst formation rates and compromised overall embryo

quality, a subset of embryos retained the capacity to progress beyond the morula stage

and form blastocysts (Fig. 1A, B). This contrasts sharply with mouse embryos, which

undergo irreversible developmental arrest at the morula stage upon glucose deprivation

[2, 5]. In mice, this developmental arrest is generally attributed to impaired glucose-

dependent regulation of CDX2 expression through the Hippo signaling pathway, which

consequently prevents blastocoel formation [2, 5]. However, in bovine embryos, while

the Hippo pathway–dependent regulation of CDX2 is conserved [23], CDX2 disruption

does not preclude blastocyst formation [14, 33]. Accordingly, although glucose

deficiency markedly suppresses the expression of key TE transcription factors, including

CDX2 and GATA3, bovine embryos can still form blastocysts even under glucose-

deprived conditions (Fig. 2A, 2B and Fig S2A, S2B).

In addition, our findings demonstrate that the three main pathways of glucose

metabolism fulfill unique functional roles during TE differentiation. We found that basal

OCR and exogenous pyruvate uptake remained stable in G− embryos (Fig. 3C–F), and

inhibition of glucose flux into pyruvate using YZ9 did not impair blastocyst formation

and TE specification (Fig. 1E, F). This phenotype may be explained by the distinct

metabolic resilience of bovine embryos. Unlike murine embryos, which contain

relatively low lipid reserves, bovine embryos possess abundant lipid droplets that can

serve as alternative energy and biosynthetic substrates under glucose-deprived

conditions [25, 34, 35]. Moreover, the BO-SOF culture medium used in this study

provides sufficient levels of pyruvate, lactate, and amino acids, which likely

compensate for glucose deprivation by providing alternative metabolic fuels [36-38].

Nevertheless, endogenous lipid reserves and exogenous nutrients in the culture

medium are insufficient to fully offset the absence of glucose. We observed that

glucose deprivation, or the pharmacological inhibition of its metabolism, significantly

attenuated HBP-mediated O-GlcNAc (Figure 3G, H and S3 A, C, D). In murine embryos,

O-GlcNAc has been proposed to primarily facilitate the nuclear stabilization of YAP,

independent of its intracytoplasmic phosphorylation status [5, 8, 39]. When YAP is

phosphorylated, it remains localized in the cytoplasm where it undergoes degradation.

This exclusion from the nucleus inhibits the expression of TE-associated genes [22, 23].

However, in contrast to this model, we observed that inhibition of O-GlcNAc using TT04

significantly increased p-YAP levels (Fig. 4G, 4H). Concurrently, the inhibition also

significantly reduced nuclear YAP accumulation and downregulated CDX2 expression

(Figure 4 A-F). The relationship between glycosylation and YAP phosphorylation during

embryogenesis is largely unexplored. However, mutating the Thr241 O-GlcNAc site

significantly compromises YAP stability while promoting its phosphorylation in tumor

cells [40]. We therefore speculate that inhibition of O-GlcNAc promotes YAP retention in

the cytoplasm in bovine embryos, where it becomes increasingly phosphorylated.

Interestingly, unlike HBP-specific inhibition, glucose deprivation in bovine embryos

resulted in a marked reduction in p-YAP (Fig. 1I, 1J). In cancer cells, AMPK-mediated

site-specific phosphorylation serves as a primary mediator of glucose-deficiency

signaling [41, 42]. Thus, we speculate that glucose deprivation reduces overall

phosphorylation levels, leading to decreased p-YAP.

The role of glucose in regulating phosphorylation is further supported by its impact on

the mTOR pathway [42]. We found that glucose deprivation significantly decreased

phosphorylation of RPS6, a downstream target of mTORC1 (Figure 5A, B). In addition,

inhibition of PPP disrupted mTOR signaling, and both PPP inhibition and mTOR

suppression markedly reduced the expression of the TE-associated transcription factor

TFAP2C (Fig. 5G-K). These findings indicate that PPP activity is essential for maintaining

the mTORC1–TFAP2C signaling axis required for proper TE differentiation.

Furthermore, inhibition of PPP from the zygote stage resulted in developmental arrest at

the morula stage, which could not be rescued by supplementation with downstream

metabolites (Fig. S1H–K). Glucose flux through the PPP serves a critical antioxidant

function via NADPH production [31, 32]. When this pathway is disrupted at the G6PD

level, murine embryos experience elevated ROS levels [5, 31]. While uridine can

mitigate some defects, full phenotypic restoration requires a combinatorial approach

addressing both nucleotide depletion and oxidative stress [5]. We thus speculate that

the irreversible arrest observed in bovine embryos is due to elevated reactive oxygen

species (ROS) levels induced by pathway inhibition, ultimately leading to embryonic

lethality.

Beyond direct signaling cascades, glucose availability profoundly shapes the embryonic

epigenetic landscape. We observed that disrupted glucose metabolism led to

significant reductions in arginine methylation (ADMA and SDMA) alongside a

pronounced decrease in the repressive histone modification H3K27me3 (Fig. 3I–N and

Fig. S3E–H). Protein arginine methyltransferase 1 (PRMT1) accounts for approximately

85% of ADMA formation in cells [43, 44]. By inhibiting phosphorylation-dependent

cytoplasmic retention and proteasomal degradation, PRMT1-mediated methylation

promotes YAP stabilization and nuclear accumulation [45]. Therefore, we speculate that

glucose may interfere with the Hippo pathway by affecting arginine methylation, thereby

influencing TE differentiation. Collectively, our findings highlight a potential link

between glucose metabolism and epigenetic control, providing a new avenue for

investigating the mechanisms underlying lineage specification.

Publicly available data support our model: glucose deprivation selectively impacts

glycosylation-related metabolites without reducing pyruvate levels (Figure S3 A, B).

Furthermore, single-cell analysis shows that the HBP and PPP, but not glycolysis, are

differentially regulated between the TE and ICM (Figure 6 A-H). In conclusion, our results

demonstrate that glucose specifically regulates TE differentiation in bovine embryos,

primarily through its flux into the HBP and PPP.

Acknowledgements

We thank Prof. Henry J Leese for his helpful discussion.

Competing interests

The authors declare no competing or financial interests.

Author contributions

J.Q. conceived the project, designed and performed all experiments, analysed and interpreted the data, and drafted and critically revised the manuscript. R.S. secured funding for the project and critically revised the manuscript.

Funding

There were no project specific funds for this work. Reagents used in this study were provided by the R.S. laboratory through general laboratory funds.

References

1. M.L. Sutton-McDowall, J.G.T., *Metabolism in the pre-implantation oocyte and embryo*. Animal Reproduction, 2015.

2. Brown, J.J. and D.G. Whittingham, *The roles of pyruvate, lactate and glucose during preimplantation development of embryos from F1 hybrid mice in vitro*. Development, 1991. **112**(1): p. 99-105.

3. Werner, C., T. Doenst, and M. Schwarzer, *Chapter 4 - Metabolic Pathways and Cycles*, in *The Scientist's Guide to Cardiac Metabolism*, M. Schwarzer and T. Doenst, Editors. 2016, Academic Press: Boston. p. 39-55.

4. Stincone, A., et al., *The return of metabolism: biochemistry and physiology of the pentose phosphate pathway*. Biol Rev Camb Philos Soc, 2015. **90**(3): p. 927-63.

5. Chi, F., et al., *Glycolysis-Independent Glucose Metabolism Distinguishes TE from ICM Fate during Mammalian Embryogenesis*. Dev Cell, 2020. **53**(1): p. 9-26 e4.

6. Biggers, J.D., D.G. Whittingham, and R.P. Donahue, *The pattern of energy metabolism in the mouse oöcyte and zygote*. Proceedings of the National Academy of Sciences, 1967. **58**(2): p. 560-567.

7. O'Fallon, J.V. and R.W. Wright, Jr., *Quantitative determination of the pentose phosphate pathway in preimplantation mouse embryos*. Biol Reprod, 1986. **34**(1): p. 58-64.

8. Peng, C., et al., *Regulation of the Hippo-YAP Pathway by Glucose Sensor O-GlcNAcylation*. Molecular Cell, 2017. **68**(3): p. 591-604.e5.

9. Scrivener, A.L., et al., *Protein O-GlcNAcylation in reproductive biology and the impact of metabolic disease*. Human Reproduction Update, 2025. **31**(5): p. 512-531.

10. Düvel, K., et al., *Activation of a Metabolic Gene Regulatory Network Downstream of mTOR Complex 1*. Molecular Cell, 2010. **39**(2): p. 171-183.

11. Simcox, J. and D.W. Lamming, *The central moTOR of metabolism*. Developmental Cell, 2022. **57**(6): p. 691-706.

12. Li, S., et al., *Linker histone H1FOO is required for bovine preimplantation development by regulating lineage specification and chromatin structuredagger*. Biol Reprod, 2022. **107**(6): p. 1425-1438.

13. Dang, Y., et al., *The lysine deacetylase activity of histone deacetylases 1 and 2 is required to safeguard zygotic genome activation in mice and cattle*. Development, 2022. **149**(11).

14. Shi, Y., et al., *Functional role of GATA3 and CDX2 in lineage specification during bovine early embryonic development*. Reproduction, 2023.

15. Guerif, F., et al., *A simple approach for COnsumption and RElease (CORE) analysis of metabolic activity in single mammalian embryos*. PLoS One, 2013. **8**(8): p. e67834.

16. Muller, B., et al., *Application of extracellular flux analysis for determining mitochondrial function in mammalian oocytes and early embryos*. Scientific Reports, 2019. **9**(1): p. 16778.

17. Cao, D., et al., *Selective utilization of glucose metabolism guides mammalian gastrulation*. Nature, 2024. **634**(8035): p. 919-928.

18. Wu, X., et al., *TEAD4 regulates KRT8 and YAP in preimplantation embryos in mice*
but not in cattle. Reproduction, 2024. **167**(3).

19. Li, S., et al., *NOTCH signaling pathway is required for bovine early embryonic*
developmentdagger. Biol Reprod, 2021. **105**(2): p. 332-344.

20. Hu, B., et al., *Single-cell RNA-Seq reveals the earliest lineage specification and X*
chromosome dosage compensation in bovine preimplantation embryos. FASEB J,
2024. **38**(4): p. e23492.

21. Malkowska, A., et al., *A hexa-species transcriptome atlas of mammalian*
embryogenesis delineates metabolic regulation across three different
implantation modes. Nature Communications, 2022. **13**(1).

22. Wu, Z. and K.-L. Guan, *Hippo Signaling in Embryogenesis and Development.*
Trends in Biochemical Sciences, 2021. **46**(1): p. 51-63.

23. Gerri, C., et al., *A conserved role of the Hippo signalling pathway in initiation of the*
first lineage specification event across mammals. Development, 2023. **150**(8).

24. Uhde, K., et al., *Exposure to elevated glucose concentrations alters the*
metabolomic profile of bovine blastocysts. PLOS ONE, 2018. **13**(6): p. e0199310.

25. Herrick, J.R., et al., *Developmental and molecular response of bovine embryos to*
reduced nutrients in vitro. Reproduction and Fertility, 2020. **1**(1): p. 51-65.

26. Krisher, R.L. and J.R. Herrick, *Bovine embryo production in vitro: evolution of*
culture media and commercial perspectives. Anim Reprod, 2024. **21**(3): p.
e20240051.

27. Ma, C., et al., *mTOR hypoactivity leads to trophectoderm cell failure by enhancing*
lysosomal activation and disrupting the cytoskeleton in preimplantation embryo.
Cell Biosci, 2023. **13**(1): p. 219.

28. Thompson, J.G., et al., *Oxygen uptake and carbohydrate metabolism by in vitro*
derived bovine embryos. J Reprod Fertil, 1996. **106**(2): p. 299-306.

29. Banliat, C., et al., *The proteomic analysis of bovine embryos developed in vivo or*
in vitro reveals the contribution of the maternal environment to early embryo. BMC
Genomics, 2022. **23**(1): p. 839.

30. Traut, M., et al., *Deregulation of oxidative phosphorylation pathways in embryos*
derived in vitro from prepubertal and pubertal heifers based on whole-
transcriptome sequencing. BMC Genomics, 2024. **25**(1): p. 632.

31. Warzych, E., P. Lipinska, and E. Sell-Kubiak, *Pentose phosphate pathway inhibition*
during in vitro maturation substantially affects the metabolism of bovine COCs
and blastocysts. Theriogenology, 2024. **230**: p. 72-80.

32. Braga, J.V., et al., *The role of the pentose phosphate pathway on reproductive*
functions. Anim Reprod, 2025. **22**(4): p. e20240130.

33. Goissis, M.D. and J.B. Cibelli, *Functional characterization of CDX2 during bovine*
preimplantation development in vitro. Molecular Reproduction and Development,
2014. **81**(10): p. 962-970.

34. Barceló-Fimbres, M. and G.E. Seidel, *Cross-validation of techniques for*
measuring lipid content of bovine oocytes and blastocysts. Theriogenology, 2011.
75(3): p. 434-444.

35. Kajdasz, A., et al., *Lipid Stores and Lipid Metabolism Associated Gene Expression*
in Porcine and Bovine Parthenogenetic Embryos Revealed by Fluorescent Staining
and RNA-seq. Int J Mol Sci, 2020. **21**(18).

36.   Pasquariello, R., et al., *Lipid Enriched Reduced Nutrient Culture Medium Improves*
522   *Bovine Blastocyst Formation*. Reprod Fertil, 2023. **4**(4).
523   37.   Sudano, M.J., et al., *Lipid content and apoptosis of in vitro-produced bovine*
524   *embryos as determinants of susceptibility to vitrification*. Theriogenology, 2011.
525   **75**(7): p. 1211-1220.
526   38.   Ordoñez-Leon, E.A., et al., *Lipid droplet analysis using in vitro bovine oocytes and*
527   *embryos*. Reprod Domest Anim, 2014. **49**(2): p. 306-14.
528   39.   Lei, Y., et al., *Protein O-GlcNAcylation coupled to Hippo signaling drives vascular*
529   *dysfunction in diabetic retinopathy*. Nat Commun, 2024. **15**(1): p. 9334.
530   40.   Zhang, X., et al., *The essential role of YAP O-GlcNAcylation in high-glucose-*
531   *stimulated liver tumorigenesis*. Nature Communications, 2017. **8**(1): p. 15280.
532   41.   Mo, J.S., et al., *Cellular energy stress induces AMPK-mediated regulation of YAP*
533   *and the Hippo pathway*. Nat Cell Biol, 2015. **17**(4): p. 500-10.
534   42.   Santo-Domingo, J., et al., *Glucose-dependent phosphorylation signaling*
535   *pathways and crosstalk to mitochondrial respiration in insulin secreting cells*. Cell
536   Communication and Signaling, 2019. **17**(1): p. 14.
537   43.   Bedford, M.T. and S.G. Clarke, *Protein Arginine Methylation in Mammals: Who,*
538   *What, and Why*. Molecular Cell, 2009. **33**(1): p. 1-13.
539   44.   Di Lorenzo, A. and M.T. Bedford, *Histone arginine methylation*. FEBS Letters, 2011.
540   **585**(13): p. 2024-2031.
541   45.   Hong, X.-L., et al., *Positive feedback between arginine methylation of YAP and*
542   *methionine transporter SLC43A2 drives anticancer drug resistance*. Nature
543   Communications, 2025. **16**(1): p. 87.

Fig 1 Glucose deprivation and inhibition of glucose metabolism impair bovine embryo development.

(A–D) Representative brightfield images of bovine embryos cultured in G– or CTRL media. Quantification of the proportion of embryos developing to the 4-cell (E2.0), 8–16-cell (E3.5), morula (E5.0), and blastocyst (E6.5) stages, and the proportion of hatching embryos at E8.5.

(E, F) Representative images and corresponding blastocyst formation rates of E7.0 embryos cultured in the presence of the glycolysis inhibitor YZ9, relative to CTRL.

(G, H) Representative images and blastocyst formation rates of E7.0 embryos cultured with the HBP inhibitor TT04, relative to CTRL.

(I, J) Representative images and blastocyst formation rates of E7.5 embryos cultured with the PPP inhibitor 6-AN for 48 hours, relative to CTRL.

All percentages were calculated relative to the total number of zygotes cultured per treatment. Data

14 are presented as mean ± SEM from at least six independent replicates (≥15 embryos per group).

15 Statistical significance was determined using a two-tailed unpaired Student's t-test: ns, not

16 significant; **P < 0.01; ****P < 0.0001. Scale bars, 50 μm.

Fig 2 Glucose deprivation suppresses Hippo signaling and impairs TE specification in bovine embryos.

(A–C) Immunofluorescence staining and quantitative analysis of the TE marker CDX2 and ICM marker SOX2 in E7.0 bovine blastocysts cultured under CTRL and G− conditions.

(D–F) Immunofluorescence staining and quantification of EPI marker NANOG and PrE marker GATA6 in E8.5 blastocysts cultured under CTRL and G− conditions.

(G–L) Immunofluorescence staining and quantitative analysis of Hippo pathway components,

including TEAD4 (G, H), p-YAP (I, J), YAP (K, L), and TFAP2C (K, L), in E7.0 blastocysts cultured under

CTRL and G− conditions.

For quantification, relative lineage-specific cell ratios were calculated for TE (CDX2$^+$/Hoechst$^+$), ICM

(SOX2$^+$/Hoechst$^+$), EPI (NANOG$^+$/Hoechst$^+$), and PrE (GATA6$^+$/Hoechst$^+$) populations (n ≥ 3

biological replicates, with ≥3 blastocysts per group). Data are presented as mean ± SEM. Statistical

significance was determined using a two-tailed unpaired Student's t-test: ns, not significant; **P <

0.01; ***P < 0.001; ****P < 0.0001. Scale bars, 100 µm.

Fig 3 Glucose deprivation disrupts mitochondrial function and biosynthetic metabolism in bovine blastocysts.

(A, B) Representative fluorescence images of JC-1 staining in CTRL and G− blastocysts. Orange fluorescence indicates J-aggregates, whereas green fluorescence indicates J-monomers. Quantification of the red/green JC-1 fluorescence ratio.

(C) OCR measured by Seahorse XF analysis in CTRL and G− blastocysts.

(D–F) Pyruvate concentrations in culture medium following 24 h embryo culture under G−, 2-DG-treated, and YZ9-treated conditions, relative to CTRL.

(G, H) Representative immunofluorescence images and corresponding quantification of global O-GlcNAc (green) levels in CTRL and G− blastocysts.

(I-N) Representative immunofluorescence images and quantification of ADMA, SDMA, H3K27me3 in CTRL and G− blastocysts.

Data are presented as mean ± SEM. Statistical significance was determined using a two-tailed unpaired Student's t-test (n ≥ 3 biological replicates, with ≥5 blastocysts per group): ns, not significant; **P < 0.01; ***P < 0.001; ****P < 0.0001. Scale bars, 100 μm.

Fig 4 Inhibition of O-GlcNAc disrupts TE specification and Hippo signaling in bovine blastocysts

(A, B) Representative immunofluorescence images and quantitative analysis of CDX2 and SOX2 in E7.0 embryos cultured under CTRL and TT04 treated conditions.

(C, D) Quantitative analysis of total cell number (Hoechst$^+$), and the proportions of TE (CDX2$^+$/Hoechst$^+$) and ICM (SOX2$^+$/Hoechst$^+$) cells in E7.0 embryos cultured under CTRL and TT04-treated conditions.

(E, F) Representative immunofluorescence images and quantitative analysis of YAP and TFAP2C in E7.0 embryos cultured CTRL and TT04 treated conditions.

(G, H) Representative immunofluorescence images and quantitative analysis of p-YAP and global O-GlcNAc in E7.0 embryos cultured CTRL and TT04-treated conditions.

Data are presented as mean ± SEM. n≥3 independent biological replicates, with ≥5 blastocysts per group. Statistical significance was determined using a two-tailed unpaired Student's t-test. ns, not significant; **$P < 0.01$; ****$P < 0.0001$. Scale bars, 100 μm.

Fig 5 Inhibition of the PPP suppresses mTOR activity and disrupts TE differentiation

(A, B) Representative immunofluorescence images and quantitative analysis of pS236-RPS6 in E7.0

embryos cultured under CTRL or G– conditions.

(C, D) Representative immunofluorescence images and quantitative analysis of pS236-RPS6 and

TFAP2C in E7.5 embryos cultured under CTRL or 6-AN-treated conditions.

(E, F) Representative bright-field images and quantification of blastocyst formation rates in E7.0

embryos cultured under CTRL or INK128-treated conditions. Scale bar, 50 μm.

(G, H) Representative immunofluorescence images and quantitative analysis of pS236-RPS6 in E7.5

embryos cultured under CTRL or INK128-treated conditions.

(I–K) Representative immunofluorescence images and quantitative analysis of CDX2 and TFAP2C in

embryos under CTRL or INK128-treated conditions.

Data are presented as mean ± SEM. n ≥3 independent biological replicates, with ≥5 embryos per

groups. Statistical significance was determined using a two-tailed unpaired Student's t-test: ns, not

significant; *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001. Scale bar, 100 μm

Fig 6 Quantitative analysis of metabolic pathway module scores during bovine preimplantation development.

(A, D, G) Heatmap showing the expression of glycolysis,HBP and PPP module genes.

(B, E, H) Bar graphs comparing the dynamic changes in module scores of glycolysis, HBP and PPP across grouped developmental stages.

(C, F, I) Bar graphs illustrating the differences in module scores among TE, ICM, PE and EPI at the mid- and late-blastocyst stages.

89   Data are presented as mean ± SEM. Statistical significance was determined using a two-tailed

90   unpaired Student's t-test: ns, not significant; $*P < 0.05$; $**P < 0.01$; $***P < 0.001$; $****P < 0.0001$.

91