Title: Resolving cell lineages and gene functions in the developing mouse gastrointestinal

tract using *in utero* transduction

Ziwei Liu[1,‡], Krishnanand Padmanabhan[1,‡], Jingyan He[2,†], Katrin Hector[2], Bettina

Semsch[2,3], Jia Sun[3], Viktoria Knoflach[1], Sarantis Giatrellis[2], Khachatur Dallakyan[1], Christian

Göritz[2], Emma Rachel Andersson[2,*], Ulrika Marklund[1,*]

[1] Department of Medical Biochemistry and Biophysics, Karolinska Institutet, Sweden

[2] Department of Cell and Molecular Biology, Karolinska Institutet, Sweden

[3] Comparative Medicine, Karolinska Institutet, Sweden

† Current Adress: Janelia Research Campus, Howard Hughes Medical Institute,

Ashburn, USA

‡These two authors contributed equally

*These two authors share corresponding authorship

Correspondence:

Ulrika.Marklund@ki.se; Address: Tomtebodavägen 16, 171 77 Stockholm, Sweden.

Emma.Andersson@ki.se; Address: Tomtebodavägen 16, 171 77 Stockholm, Sweden.

Keywords: enteric nervous system, gut patterning, neural crest development, lineage tracing,

in utero transduction

ABSTRACT

How diverse cell lineages emerge and are genetically regulated during organogenesis are central questions in understanding the developmental origins of disease. However, the mouse gut, including its intrinsic enteric nervous system (ENS) derived from migratory neural crest, has remained difficult to experimentally target. Here, we introduce an *in utero* lentiviral nano-injection strategy that enables early and efficient access to progenitor cells of all major cell types within the developing gut as well as gut-innervating ganglia. Leveraging this approach in combination with DNA barcoding and single cell transcriptomics, we resolve clonal relationships in all gut lineages, including epithelial, neural, immune, and mesenchymal cell types. Clonal coupling between distinct subsets of fibroblasts and either pericytes, mesothelial cells, or interstitial cells of Cajal, suggested a developmental logic whereby the mesenchymal compartment arises from a set of fate-biased progenitors. Yet, mesenchymal regionalization along the anterior–posterior axis establishes early, whereas the ENS displays broad clonal dispersion across gut regions and acquires subsequent regional identities. We further adapted the platform for temporally controlled cell-type specific gene manipulation and, as a proof-of-principle, show that induced expression of the proneural factor Ascl1 biases ENS progenitor cells toward neuronal differentiation. Together, this work provides insights into refined spatiotemporal lineage relationships within a multigerm-layer organ and establishes a broadly applicable *in vivo* framework for probing gene function during gastrointestinal and neural crest development.

SIGNIFICANCE

The gastrointestinal tract comprises diverse cell types originating from all three germ layers and includes the neural crest-derived enteric nervous system (ENS). Progress in defining these lineages and their gene regulation is challenged by the limited experimental access to the

developing gut. Here, we establish *in utero* lentiviral transduction as an efficient approach to

resolve clonal lineages and address gene functions in defined gut cell types. We show that

mesenchyme assumes positional allocation early and differentiates through fate-restricted

progenitors, linking specialized mesenchymal cell types to different fibroblasts. In contrast, the

ENS differentiates stochastically and acquires late regional identities. Our study reveals

fundamental principles of multi-lineage organogenesis and provides a framework to dissect the

contribution of developmental programs to visceral dysfunction.

INTRODUCTION

The gastrointestinal (GI) tract is essential for life. Congenital gut malformations including

Hirschsprung disease, pyloric stenosis, and various intestinal atresias result in diverse and life-

long complications. In the general population, 30% suffer from chronic gut dysfunctions,

including disorders of gut-brain interaction (DGBIs) and inflammatory bowel disease (IBD)

(1). Conceivably these conditions may arise from subtle alterations in gut cell composition

already established at fetal stages (2). Supporting this notion, premature birth is an identified

risk factor for developing DGBIs (3). Furthermore, congenital chronic intestinal pseudo-

obstruction (CIPO) can be myopathic, mesenchymopathic or neuropathic (4), underscoring the

critical role of coordinated interactions among multiple gut cell types in maintaining GI

motility. Thus, there is a need to better resolve how different cell lineages in the GI-tract

develop and integrate during early life.

Experimental gene perturbation/manipulation - genetic gain or loss of function – are

essential tools to resolve how gene regulatory networks control cellular differentiation and how

developmental deviations can lead to pathology. Owing to the relative ease with which the

central nervous system (CNS) can be genetically manipulated using *in vivo* embryonic

transduction or electroporation (5), the CNS field has gained intricate knowledge of precise

gene regulatory networks controlling the generation of different neuronal subtypes (6, 7). These manipulations are possible due to the ventricular structure of the brain and the lumen of the spinal cord, making them accessible to plasmids or viral vectors. In contrast, the GI-tract is positioned internally and not possible to target using these common approaches.

Normal gut physiology is critically dependent on neuronal control provided by its intrinsic enteric nervous system (ENS), essential for peristalsis, secretion, and blood flow regulation. These functions are modulated by sympathetic innervation, primarily of celiac ganglion neurons, while sensory information is captured from the gut via dorsal root ganglia (DRG) at the trunk regions, and nodose ganglia in cranial regions. The ENS, sympathetic ganglia, and DRG are formed from neural crest cells (NCC), a transient embryonic population originating from the dorsal neural tube (8). In the mouse, delamination of NCC occurs around embryonic day (E)8 (Theiler stages TS12-13) and thereafter, the migrating cells and the ganglia that form are positioned unfavorably for traditional *in utero* manipulations that are typically performed from E11.5 (5). Thus, experimental systems to rapidly investigate the development of the murine gut, the ENS, and other gut-innervating ganglia have been lacking.

Recently, we developed neural plate targeting with *in utero* nano-injection (NEPTUNE) to target the developing CNS at earlier stages (9, 10). With NEPTUNE, ultrasound is used to guide lentiviral delivery of cargo into the mouse amniotic cavity as early as E7.5, allowing access to 80-95% of cells in the CNS. The technique can also be used to introduce high-diversity lentiviral barcode libraries to resolve refined lineages using next generation single cell lineage tracing (11, 12). As distinct barcodes integrate in the DNA of each transduced cell, clonal relationships can be reconstructed after successive cell divisions by single cell RNA-sequencing (scRNA-seq), revealing how individual progenitors contribute to the generation of diverse cell types. Here, we asked whether the method allows for targeting of the GI-tract and of neural

crest/placodes, whose derivatives innervate the gut, for the purpose of revealing clonally linked

gut cell types and performing genetic experimentation of the developing ENS.

We found that ultrasound-guided *in utero* lentiviral nano-injection at E7.5 transduces

progenitors that contribute to the GI tract, including endodermal (i.e. epithelia), mesodermal

(e.g. smooth muscle cells, fibroblasts and interstitial cells of Cajal; ICCs) and ectodermal

progenitors (e.g. ENS). In addition, the method efficiently targets extrinsic gut-innervating

ganglia (e.g DRG). By combining high-throughput clonal labeling and transcriptomic profiling

of cells in the gut, we identified fate-restricted progenitor cells present at the time of

transduction and reconstructed lineage relationships across differentiating gut cell types and

regions. Of note, we uncovered molecularly defined fibroblast populations that are clonally

coupled with either mesothelial cells, pericytes or ICCs, revealing early diversification paths

within the gut mesenchyme. We find that ENS targeting is most efficient at early E7.5, but that

residual epiblast cells are still present at this stage. Beyond lineage tracing, we show that *in*

utero nano-injections can be utilized for conditional gene perturbation in the ENS. Using the

pan-neuronal Baf53-Cre mouse line, we demonstrate that lentivirus carrying Cre-dependent

cargo is selectively expressed in developing enteric neurons. In Sox10-CreERT2 mice,

transgene expression can instead be induced in ENS progenitors/glia in a temporally controlled

manner. As proof-of-principle, we show that timed overexpression of the proneural

transcription factor *Ascl1* promote neuronal differentiation. Taken together, our results show

that ultrasound-guided *in utero* nano-injection is a versatile tool to investigate organogenesis

and ENS development. Our work offers insight into how neural crest and mesoderm lineages

diversify and regionally pattern within the developing gut.

MATERIAL and METHODS

Mice

The following transgenic and reporter mouse strains were used: Baf53b-Cre (JAX, #027826), Sox10-CreERT2 (SER26) (13), R26-EYFP (JAX #006148) and R26-tdTomato (Ai14; JAX #007908). C57/B6J and CD1 (Charles River Laboratories) wild-type (WT) mice were used. Animals were group-housed, with food and water *ad libitum,* under 12-h light-dark cycle conditions, 22°C ambient temperature and 50% humidity. Animal experiments were approved by the local ethics committee (Stockholm Norra djurförsöksetiska nämnd, Jordbruksverket; N5264/18, N6626-2019, N5237-2023, 8188-2017, 2987-2020 and 315-23).

Lentivirus constructs

Plasmid *LV-H2B-GFP* was a gift from Elaine Fuchs (Addgene #25999). *LV-EF1A-H2B-tdTomato-30N* was previously described (11, 12). *Lenti-DIO-EGFP* (Supplementary Figure 1A) was based on the interim vector *pRRLSIN-cPPT-hPGK-EGFP-WPRE* (Gift from Didier Trono, Addgene #12252). *Lenti-DIO-EGFP-Ascl1* (Supplementary Figure 1B) was constructed based on *pRRLSIN-cPPT-shortCAG-dlox-EGFP_2A(PTV)(rev)-dlox-WPRE* and the GeneArt product *mAscl1* (710bp). Both vectors were constructed, amplified, and characterized by the Viral Vector Facility (VVF) of the Neuroscience Center Zurich (ZNZ). Plasmids and virus were produced and characterized as described in (Supplementary Material and Methods). High-titer vesicular stomatitis virus G (VSV-G)–pseudotyped lentiviral particles were produced either in-house or by a commercial facility (GEG TECH, France) and used at a minimum titer of 1×10^9 transducing units (TU)/mL.

Ultrasound-guided *in utero* nano-injections

A detailed description of the procedure has been reported (9, 10) and is described in (Supplementary Material and Methods). We considered Theiler stages TS11A–C to represent the optimal E7.5 window for ultrasound-guided *in utero* injections (Supplementary Figure 4A, B), while injection at E8 or later failed to efficiently target gut and ENS (Supplementary Figure 4C, D).

Tamoxifen administration

Plug-mated females received tamoxifen (0.1 mg/g body weight; Sigma) dissolved in corn oil/ethanol (9:1) by intraperitoneal injection at E16.0-E16.5. Embryos were collected 26 or 55 hours after tamoxifen administration.

Tissue preparation and immunohistochemistry

E13.5 embryos were decapitated and fixed in 4% PFA in PBS for 4-5 hours at 4 °C. Intestines from E14.5-E18.5 were dissected out and fixed in 4% PFA in PBS at 4 °C for 1.5-2 hours. Samples were washed in PBS three times and dehydrated in 30% sucrose in PBS at 4 °C overnight. Intestines were cut into 1 cm long pieces, representing different regions and embedded for transverse analysis in OCT freezing medium and stored at -80 °C. Samples were sectioned at 12 μm and stored at -20 °C.

Frozen tissue sections were air-dried at RT for 30 min, rinsed with PBS and incubated in blocking solution containing purified donkey anti-mouse Fab (Jackson Laboratories, PA, USA #715-007-003) diluted 1:50 in PBS for 2 hours at room temperature (RT). After rinsing with PBS, the sections were blocked with 2% normal donkey serum (NDS, Jackson) and 0.1% Triton X-100 (Sigma) in PBS for 2 hours, and incubated with primary antibodies diluted in the same solution overnight at 4°C. They were then washed three times with PBS (10 minutes each) and

incubated with secondary antibodies at RT for 1 hour. Tissue was washed three times with PBS

(10 minutes each) and mounted in DAKO mounting medium (Agilent) containing DAPI (4',6-

diamidino-2-phenylindole). Primary and secondary antibodies are specified in (Supplementary

Material and Methods.

Preparation of single-cell RNA sequencing datasets

CD1 embryos injected with LV-EF1A-H2B-tdTomato-30N were harvested at E16.5. Stomach

and 1/3 of the distal small intestine were harvested from the E7.5 experiment, while the

stomach, 2/3 of the small intestine, and the entire large intestine were harvested from the

E7.5^{Early} experiment. Dissociation was performed as described in (Supplementary Material and

Methods). Briefly, tissues were cut, enzymatically processed, triturated and TOM+ DRAQ7-

cells isolated through flow cytometry (Supplementary Figure 1C-H). cDNA libraries were

prepared using 10x Chromium Next GEM Single Cell 3' Reagent Kits Version 3.1 Dual Index

(10x Genomics, CA, USA). The datasets were named according to stage of injection (i) and

analysis (a): iE7.5-aE16.5 and iE7.5^{Early}-aE16.5.

Analysis of single-cell RNA sequencing datasets

A detailed description of the scRNA-seq analysis can be found in (Supplementary Material and

Methods). In brief, E16.5 scRNA-seq datasets were aligned to a combined mouse genome

(GRCm38) with a tdTomato-N transgene containing a 30bp barcode region (11, 12) using

CellRanger. Raw count matrices were processed in Seurat, with ambient RNA contamination

assessed and corrected using SoupX. Low-quality cells and doublets were removed based on

gene/UMI thresholds, mitochondrial contents and DoubletFinder predictions. Data were

normalized using SCTransform, followed by dimensionality reduction, batch correction with

Harmony, clustering and UMAP visualisation. CloneIDs were assigned to cells using the TREX

(TRacking and gene EXpression profiling of clonally related cells) pipeline (11, 12) based on

shared barcodes. Clonal composition and relationships between cell types were analyzed using

statistical comparisons to randomized data, generating clonal coupling scores and correlation-

based clustering to infer lineage relationships.

Data Accession:

scRNA-seq data used for analysis in Figure 1 and Supplementary Figure 2 can be found at

ArrayExpress with the accession number E-MTAB-14817 (11). The scRNA-seq data generated

in this study will be made available at GEO upon publication.

RESULTS

Efficient targeting of gut-innervating ganglia by *in utero* transduction at E7.5

Previous studies have demonstrated that high-titer lentivirus injection into the amniotic cavity

of E7.5 embryos results in widespread transduction of the neural plate and otic placodes (9-11).

Because gut-innervating ganglia originate from neural crest or from placodes, we hypothesized

that these structures could also be targeted. To assess whether E7.5 amniotic cavity lentiviral

injection targets neural plate progenitors that generate gut-innervating neuronal populations

(Figure 1A), we re-analyzed a published scRNA-seq dataset of E9.5 and E10.5 embryos

injected at E7.5 with lentiviruses carrying *tdTomato* (11). The cells in this dataset had been

broadly categorized as "neural tube cells", "epithelial cells", and "neural crest cells" (Figure

1B, Supplementary Figure 2A). To investigate whether developing gut-innervating neurons had

been targeted (Figure 1D), we renormalized and subclustered the "neural crest cells" (4,916

cells), which yielded 14 clusters (Figure 1C, E; Supplementary Figure 2B). We identified three

Tubb3+ neuronal clusters (clusters 4, 8, 12), while remaining clusters displayed high *Sox10*

expression, indicative of neural progenitor or other immature cell states (Figure 1E, F;

Supplementary Figure 2C, D). Amongst *Sox10+* clusters, clusters 0 and 9 were defined as

mesenchymal as they expressed *Prrx1* and *Alx1*, while cluster 13 showed a melanocyte identity

with high *Mitf* and *Dct* expression. Cluster 6 exhibited an autonomic nervous system character

(*Phox2b+*, *Hand2+*, *Ascl1+*, *Ret+*) containing presumptive sympathetic (*Isl1+*) and ENS (*Isl1-*

) cells (Figure 1G). The remaining *Sox10+* clusters lacked unique markers and likely

represented progenitor or glial cells; these were annotated as NCC1-NCC7 (Figure 1E).

Amongst neuron-enriched clusters, cluster 4 displayed a nodose ganglion identity (*Phox2a+,*

Phox2b+, Tbx1+, Tfap2b-). Cluster 8 and 12 expressed a DRG profile (*Isl1+, Neurod1+,*

Six1+, Pou4f1+), with cluster 12 representing an immature state (higher *Sox10* and *Foxd3*

expression) (Figure 1E; Supplementary Figure 2C, D).

To further substantiate cluster identities, we addressed the Hox gene expression patterns

(Supplementary Figure 2E, F). Consistent with a cranial origin, the cluster identified as nodose

ganglion exhibited robust anterior Hox gene expression (e.g. *Hoxa1-4*, *Hoxb1-3*). ENS and

parts of the sympathetic ganglia originate from vagal neural crest, and cluster 6 expressed a

characteristic suite of *Hoxd3-4* and *Hoxb6 (14)*. NCC3 lacked Hox gene expression, indicating

a rostral cranial identity, while the Hox gene pattern of NCC5 suggested they are

progenitors/glia in the nodose ganglion. Posterior Hox gene expression became more

pronounced at E10.5, particularly within DRG clusters and several NCC clusters, indicative of

the anterior-to-posterior emergence of neural crest derivatives. Together, the data demonstrate

that lentiviral injection at E7.5 targets primordia of gut-innervating ganglia, including ENS,

sympathetic ganglia, nodose ganglia and DRG (Figure 1D).

To confirm that gut-innervating ganglia are persistently labelled, we performed

immunohistochemical analysis of E13.5 DRG and E18.5 gut, in embryos injected at E7.5 with

lentivirus carrying H2B-GFP. There was robust transduction of the DRG, with efficient

labelling of both NeuN+ neurons and SOX2+ progenitors/glia from thoracic to sacral levels

(65-83%) (Figure 1H-J). The ENS was uniformly transduced across gut regions, with an

average of 51% HuC/D+ neurons and 44% SOX10+ progenitors/glia (Figure 1K-M;

Supplementary Figure 3A-D). In addition to ENS labelling, we observed GFP across all gut

layers, including in α-SMA+ smooth muscle cells and EpCAM+ epithelial cells (Figure 1N;

Supplementary Figure 3E). This indicates that viral particles in amniotic fluid not only

transduce the neural plate (ectoderm) but are also in contact with endo- and mesodermal germ

cells at E7.5 (Supplementary Figure 4A, B). Indeed, injections during an early developmental

window were crucial for ENS targeting, as injections at E8.0-8.5 failed to label both the ENS

and gut epithelium, with only sporadic labeling of mesenchymal cells observed (Supplementary

Figure 4C, D).

Together, these data demonstrate that E7.5 amniotic cavity lentiviral injection enables

widespread and stable transduction of the GI-tract, encompassing extrinsic gut-innervating

neuronal populations, the ENS, and non-ENS gut cell types.

***In utero* transduction targets all major cell types in the developing gut**

As we discovered that injections not only target the ENS but the whole gut wall, we reasoned

that this approach could be leveraged to determine lineage relationships between gut cell types

derived from all three germ layers. Moreover, by capturing different segments of the gut,

conclusions could be drawn about the timing of anterior-posterior regionalization. To

simultaneously profile the transcriptional identity (cell type) and lineage history (clonality) of

single cells, we applied TREX (11, 12) (Figure 2A, B). TREX relies on *in utero* injection of a

high-diversity lentiviral library encoding a 30N (random nucleotides) barcode and tdTomato

resulting in genomic integration into dividing progenitors, thereby allowing stable inheritance

of the construct by all descendant cells. The expression of the barcode under a synthetic

ubiquitous promoter allows its detection by scRNA-seq, enabling reconstruction of clonal

relationships among captured cells based on shared barcodes (CloneIDs).

We injected the tdTomato-30N barcode lentiviral library into the amniotic cavity of

E7.5 embryos (Figure 2A) and analyzed at E16.5 (dataset thus called iE7.5-aE16.5). TOM+

cells were sorted from stomach and small intestine by flow cytometry and sequenced using the

10x Chromium platform (Figure 2B). After removing low-quality cells and potential doublets,

5,356 cells were retained for analysis, comprising 3,974 cells from the stomach and 1,382 cells

from the small intestine (Figure 2C; Supplementary Figure 5A-C). Cells from both regions were

integrated, resulting in eight major clusters defined by canonical lineage markers,

corresponding to fibroblast-like cells (*Vim+, Col1a1+, Lum+*), smooth muscle-like cells

(*Myh11+, Actg2+, Acta2+*), endothelial cells (*Cd36+, Pecam1+*), foregut basal cells (*Cdh1+,*

Epcam+, Krt5+, Krt15+), simple epithelial cells (*Cdh1+, Epcam+, Krt8+, Krt18+*), ENS

(*Sox10+, Phox2b+, Ret+, Elavl4+*), immune cells (*Cd52+, Ptprc+*) and mesothelial cells

(*Upk3b+, Msln+, Wt1+*) (Figure 2D, E; Supplementary Figure 5D). Further analysis of the

simple epithelial cluster revealed predominantly enterocytes (*Vil1+*), along with smaller

populations of enterochromaffin cells (*Chga+, Tph1+*), goblet cells (*Clca1+, Fcgbp+*), and

intestinal stem cells (*Lgr5+*) (Figure 2F). The 195 ENS cells formed four transcriptionally

distinct clusters representing progenitors, neuroblasts, and neurons (Figure 2G). Notably,

previously discovered neuronal branches characterized by *Etv1* (Branch A) and *Bnc2* (Branch

B) were captured (Figure 2H) (15).

Subclustering of the smooth muscle-like cluster revealed a contractile myofibroblast-

like state (*Trpc3+, Tagln+*) and a proliferative cluster (Figure 2I, J; Supplementary Figure 5D,

E). Subclustering of the fibroblast-like cluster uncovered additional mesenchymal

heterogeneity, including Interstitial cells of Cajal (ICC) (*Kit+, Ano1+*), pericytes (*Rgs5+,*

*Reln+, Pdgfrb+, Pdgfr*a-) and seven transcriptionally distinct fibroblast subtypes (Figure 2I,

J). These included proliferative fibroblasts (*Top2a+, Mki67+*), Fibroblast 1 (*Col6a4+,*

Adamdec1+), Fibroblast 2 (*Fndc1+, Dpt+*), Fibroblast 3 (*Adam12+*), Fibroblast 4 (*Kcnn3+*),

Fibroblast 5 (*Mfap5+, Grm3+*) and Fibroblast 6 (*Lepr+, Sp5+*) (Supplementary Figure 5D-H).

The proliferative fibroblast cluster likely comprised a mixture of all fibroblast subtypes.

Collectively, our results demonstrate that *in utero* lentiviral transduction effectively transduces

progenitors of all major gut cell types, encompassing derivatives of the three germ layers

(Figure 2K).

Barcoded *in utero* transduction reveals clonal lineage structures in the developing gut

The comprehensive capture of diverse gut cell types provided us excellent conditions to reveal

detailed lineage relationships. After barcode extraction, 3,959 cells (73.9%) retained CloneIDs

(Figure 3A, B; Supplementary Figure 6A-C). Cells with or without CloneIDs showed

comparable quality metrics and similar CloneID recovery rates were observed across all cell

types, indicating unbiased barcode dropout (Supplementary Figure 6D-G). Most CloneIDs

consisted of a single barcode, indicating single lentiviral particle transductions, while up to 4

barcodes were detected in a few cells (0.42% of clones) (Supplementary Figure 6H).

In total, 1,658 clones were reconstructed and classified into three categories: singletons

(1,026 clones), single-cell-type clones (302 clones) and multi-cell-type clones (330 clones)

(Figure 3C). Multicellular clones contained on average 4.6 cells and reached a maximum of 31

cells (Supplementary Figure 6I). Systematic assessment of multi-cellular clonal compositions

per major cell type showed that pericytes, ICCs, smooth muscle and mesothelial cells almost

exclusively consisted of multi-cell-type clones (Figure 3C), suggesting substantial lineage

diversification. Instead, ENS and immune cells exhibited only single-cell-type clones,

reflecting early lineage restriction of their respective precursors. Clone sharing was

predominantly found between mesenchymal cell types in the gut, consistent with their shared

mesodermal origin (Figure 3D). However, endothelial and mesothelial cells did not share clones

with each other despite sharing clones with other mesenchymal cell types, indicating that their

progenitor allocation is already distinct at E7.5 (Figure 3D). Cells arising from different germ

layers rarely shared clones; only five clones spanned endodermal and mesodermal lineages,

likely reflecting residual pluripotent mesendoderm cells (16) at the time of injection (Figure

3E).

To systematically resolve lineage relationships among refined cell types, we performed

clonal coupling analysis by calculating z-scores for deviations in observed clone sharing

relative to randomized data. Hierarchical clustering of Pearson correlations between z-scores

segregated cell types into broad groups corresponding to germ-layer origin and revealed

intricate clonal coupling relations between mesoderm-derived populations (Figure 3F, G,

Supplementary Figure 6J, K). Notably, ICCs showed strong clonal coupling with smooth

muscle cells and myofibroblasts (Figure 3F-H). This is consistent with previous lineage-tracing

studies showing that ICCs and myofibroblasts arise from common precursors and that ICCs are

not fibroblasts (17). However, ICCs, myofibroblasts and smooth muscle cells also showed

clonal coupling with Fibroblast 4 (*Kcnn3+*). This fibroblast likely corresponds to *Pdfgra+* cells

of the SIP (smooth muscle cells, ICCs, and PDGFRa (18, 19)), recently renamed muscle-

embedded interstitial fibroblasts; MIFs (20)) (Supplementary Figure 5E, F). Fibroblast 3

(*Adam12+*) showed enriched clonal coupling with mesothelial cells (Figure 3F-H), suggesting

an adventitial fibroblast (AF) phenotype (21), further supported by expression of *Wt1* and *Pi16*

(Supplementary Figure 5E, G). Fibroblast 1 (*Col6a4+, Adamdec1+*) displayed clonal coupling

with pericytes, consistent with the reported localization of *Adamdec1+* myofibroblasts in the

subepithelial space (SEMFs) (Figure 3F-H; Supplementary Figure 5E, H). Notably, clonal

coupling analysis thus segregated fibroblast populations into putative contractile, adventitial

and perivascular trajectories which was less apparent from transcriptome-based clustering alone

(Figure 3H; Supplementary Figure 6L).

Clonal coupling patterns among mesoderm-derived populations indicate that

mesenchymal lineages diverge early into fate-biased progenitor pools that underpin coordinated

tissue patterning during gut morphogenesis (Figure 3H). Collectively, we demonstrate that

single-cell lentiviral barcoding at E7.5 recapitulates canonical developmental hierarchies and

uncovers early cell fate diversification within the gut, corroborating known lineage

relationships while revealing previously unrecognized trajectories contributing to gut

organogenesis.

Nano-injection at early E7.5 improves ENS targeting while revealing residual epiblasts

Attempting to enrich for ENS targeting, we performed another nano-injection of the N30

barcode lentiviral library aimed at an earlier developmental stage (E7.5^{Early}, ~TS11b;

Supplementary Figure 4A, B), reasoning that primordia of neural crest could be physically more

accessible and remain for a longer time before delamination. To gain further information about

gut regionalization, we collected stomach, small intestine, and large intestine at E16.5 to

generate the iE7.5^{Early}-aE16.5 dataset (Figure 4A). Data was processed similarly as in the

previous experiment, and we recovered the same major cell types (Figure 4B; Supplementary

Figure 7A-E). Notably, the earlier targeting increased the proportion of ENS cells by 7.7-fold

(Figure 4C), making ENS the most abundant cell population, and corroborating greater

lentiviral accessibility to neural crest primordia at the stage of injection. Higher-resolution

clustering further resolved subpopulations within immune and ENS lineages, revealing myeloid

(*Lyz2+*, *Mrc1+*, *Cd68+*) and lymphoid (*Trbc1/2+*, *Cd4+*) subpopulations, lymphatic signatures

(*Lyve1+*, *Pecam1+*), refined ENS neuron divisions, and Schwann cell precursors (SCPs)

(Figure 4D, Supplementary Figure 7F, G).

CloneIDs could be assigned to 77.8% of the 21,749 cells (Supplementary Figure 8A-E),

yielding 3,421 clones comprising 2,488 singletons, 394 single-cell-type clones, and 539 multi-

cell-type clones (Figure 4E). The average and maximum size of multicellular clones were larger

in the E7.5^{Early}-aE16.5 dataset (Supplementary Figure 8F). Consistent with the first experiment,

mesenchymal populations comprised mainly of multi-cell-type clones, whereas ENS, immune

and epithelial populations were more lineage-restricted (Figure 4E). Although most

multicellular clones were composed of a single germ layer, a notable fraction (~9%) of clones

spanned more than one germ layer derivative (~1.5% spanned all three germ layers; 14 clones)

(Figure 4F, G). These cross-germ-layer clones suggest the presence of pluripotent epiblast cells

at E7.5^{Early}, in agreement with recent high-throughput spatial transcriptome analyses of murine

gastrula (22).

The presence of cross-germ-layer clones in the iE7.5^{Early}-aE16.5 dataset complicated

lineage relationship analysis between refined cell types, but we reasoned that analysis using

germ-layer pure clones would still be relevant. Clonal coupling analysis of pure clones (851

clones) recapitulated patterns observed in the iE7.5-aE16.5 dataset (compare Figures 3F with

4H; Supplementary Figure 8G). We next restricted analysis to pure mesoderm clones (639

clones) (Figure 4I; Supplementary Figure 8H). As previously observed (Figure 3G, H), ICCs,

myofibroblasts and Fibroblast 4 (putative MIFs) featured stronger clonal coupling, as did

mesothelial cells and Fibroblast 3 (putative AFs) (Figure 4I). Although Fibroblast 1 (putative

SEMFs) was not adjacent to pericytes in the correlation heatmap, it exhibited the strongest

clonal pericyte coupling of all fibroblasts (Figure 4I; Supplementary Figure 8H). Within the

immune population, *Mrc1+* myeloid and *Trbc1/2+* lymphoid cells shared only few clones,

indicating early lineage specification prior to gut colonization (Figure 4J). The large ENS

dataset allowed detailed clonal analysis of ENS subclusters, revealing that many clones

consisted of multiple ENS states/types (Figure 4K). Notably, SCPs showed clonal coupling

with all ENS subtypes, however less so with the more mature neuron clusters ENC12 and

Branch B2 which may differentiate before SCP arrival to the gut (Supplementary Figure 8G).

Coupling was overall strong between Branch A and B neurons (Figure 4K; Supplementary

Figure 8G, grey box), suggesting no predetermination of either Branch in agreement with an

earlier lineage-tracing study (23).

Together, our results from the E7.5^{Early} injection highlights the importance of precise

embryonic staging for enriched targeting of specific lineage precursors. The optimal window

for ENS targeting appears to coincide with early E7.5 (~TS11b stage), when some epiblasts

cells are still present. This stage is therefore advantageous for efficient and comprehensive ENS

targeting, while posing challenges for lineage analysis.

Clonal sharing between gut segments reveals lineage-specific timing of regional identity

acquisition

Having resolved lineage relationships between and within gut cell types, we next examined the

regional dispersion of clones across the upper and lower GI tract. In the iE7.5-aE16.5 dataset,

21 clones linking the stomach and small intestine were identified. Of these, 18 clones (85.7%)

consisted exclusively of ENS or immune cells, accounting for 25.9% of ENS and 57.9% of

immune clones (Figure 5A-C). These regional dispersions are consistent with the extrinsic

origins and migratory behavior of these lineages: the ENS arises from neural crest precursor

cells that migrate along the entire gut axis (8), while the first wave of embryonic immune cells

originates from the yolk sac and subsequently colonizes the gut (24, 25). By contrast, clones

corresponding to other cell types were predominantly restricted to a single gut segment,

suggesting that most endoderm- and mesoderm-derived progenitors had acquired their regional

identities by E7.5 (~TS11c).

Considering single-germ-layer clones in the iE7.5^{Early}-aE16.5 dataset, we identified 123

region-spanning clones, again mainly from ENS and immune lineages, accounting for 60.7%

of ENS and 66.7% of total immune clones (Figure 5D-F). Despite their abundance, epithelial

or fibroblast clones did not span regions, indicating that their progenitors are already regionally

patterned also at slightly earlier gastrula stages. We next examined the ENS subtype

composition within cross-regional and local clones and found them to be similar. Notably, both

Branch A and B neurons were present in local and cross-regional clones, supporting the innate

stochastic differentiation program in the ENS (Figure 5G). A distinguishing feature of colonic

ENS is its expression of *Pou3f3* (26), while it has not been shown if expression is induced in

vagally derived enteric *Pou3f3-* precursors as they enter the large intestine. Our barcoding

experiment allowed us to clonally link stomach-resident ENS (derived exclusively from the

vagal neural crest) to *Pou3f3$^+$* ENS cells in the colon. We found that stomach/large intestine

clones distributed randomly amongst ENS differentiation stages and neuron identities (Figure

5H). Cells within large intestine-only clones and stomach/large intestine clones showed similar

expression frequency and level of *Pou3f3* (Figure 5I). Together, these findings support that

colonic ENS cells derived from vagal neural crest acquire their colon-specific identity by

responding to local cues.

Identification of cell type-specific regional gene expression differences

Beyond lineage relationships, our regionally resolved scRNA-seq datasets enabled analysis of

transcriptional similarities and differences across GI regions at E16.5. Integration of the two

datasets showed seamless merging (Supplementary Figure 9A, B), indicating consistent

targeting of comparable progenitor populations. Most cell types were detected across all regions

except for stomach-specific foregut basal epithelia, which expressed distinct marker genes

(Supplementary Figure 9C, D). Small intestine epithelium was instead enriched for *Cps1*,

whereas posterior Hox genes marked large intestine epithelium (Supplementary Figure 9D).

Region-specific expression patterns were also observed in mesoderm-derived populations:

Barx1 and *Nr2f1* were restricted to the stomach, while *Pitx1* was detected in both stomach and

large intestine (Supplementary Figure 9E). *Nkx2.3* and *Mab21l2* were confined to intestinal

mesenchyme. Within ectoderm-derived tissue (ENS), *Hoxb5*/*Hoxa4* were enriched across

intestinal regions, whereas *Hoxc4* was limited to the small intestine (Supplementary Figure 9F).

Focused analysis of fibroblast subtypes revealed that Fibroblasts 5 and 6 were enriched

in the stomach compared with the intestines (Supplementary Figure 9G, H). Within these

populations, *Grm3* and *Sp5* showed stomach-specific expression (Supplementary Figure 9I),

whereas Fibroblast 6 uniquely expressed *Thy1* (Cd90), a marker associated with crypt

fibroblasts (Supplementary Figure 9I). *Wnt6*, *Cpne8* and *Ism1* in Fibroblast 6 further supported

a fibroblast type with high signaling capacity, consistent with a crypt/submucosal-associated

niche (Supplementary Figure 9I). These differences may reflect either a greater abundance of

Fibroblast 5 and 6 in the stomach or their earlier differentiation in anterior gut regions.

In summary, our collective E16.5 scRNA-seq data uncovered regionally distributed cell

types, expression patterns and fibroblast states in the embryonic gut, which may contribute to

establishing the functional specialization of individual GI regions.

Cre-dependent lentivirus nano-injection enables cell type-specific targeting

Our data show that progenitors of essentially all gut cell types are transduced by *in utero*

lentiviral nano-injection at E7.5. For applications requiring cell type-specific targeting, this

approach therefore needs adaptation. Previous studies have demonstrated that mini-promoters

can confer cell type-specificity in the developing CNS (10). However, no ENS-specific

promoters have yet been described. Instead, Cre-driver mouse lines have been used to label all

or subsets of ENS cell populations.

We previously showed that Baf53b-Cre mice specifically label neurons in the juvenile

ENS (15). To investigate their utility during embryonic development, Baf53b-Cre; R26-

tdTomato embryos were analyzed at two stages (Supplementary Figure 10A, B). Reporter

expression was restricted to HuC/D+ neurons and absent from SOX10+ progenitor/glial cells.

We found that ~50% of HuC/D+ neurons expressed TOM at E14.5, increasing to ~90% at

E18.5. Baf53b-Cre activity is thus confined to differentiating neurons, with a slight delay

relative to HuC/D expression.

To achieve conditional targeting with *in utero* nano-injection, we designed a lentiviral

vector expressing an inverted double-floxed EGFP reporter, Lenti-DIO-EGFP (Figure 6A,

Supplementary Figure 1A). While AC injection of non-conditional LV-H2B-GFP resulted in

GFP labelling of HuC/D+ neurons, GFP was additionally observed in non-ENS cells as

previously discussed (Figure 6B; 1N). In contrast, AC injection of Lenti-DIO-EGFP into

Baf53b-Cre embryos resulted in GFP expression restricted to HuC/D+ neurons, labelling ~4-

6% at E18.5 (Figure 6B, C). In conclusion, *in utero* nano-injection of conditional lentiviral

vectors enables precise targeting of specific gut cell types, as demonstrated by Baf53b-Cre-

dependent lentiviral reporter expression in enteric neurons.

Timed overexpression of Ascl1 promotes neuronal differentiation of ENS progenitors

Having validated that lentiviral DIO-EGFP expression can be controlled by Cre, we next sought

to use this approach to drive timed gene expression in ENS progenitor cells, which are more

responsive to developmental cues. Sox10-CreERT2 mice mediate efficient, temporally

controlled recombination in ENS progenitors (13, 14, 23). To assess the efficiency and kinetics

of Cre-dependent recombination, tamoxifen was administered to Sox10-CreERT2 females

mated with R26-EYFP males at E16.5, and embryos were harvested 26 hours later. Out of

SOX10+ cells, ~50% expressed EYFP and ~11% of HuC/D+ neurons, indicating efficient

labeling of progenitors/glia and a subset of newly differentiated neurons within a short time

window (Supplementary Figure 10C-E).

As a proof-of-principle, we sought to overexpress *Ascl1*, a proneural transcription factor

essential for enteric neurogenesis, as demonstrated by us and others (27, 28). We generated a

Cre-inducible *Lenti-DIO-EGFP-Ascl1* construct, in which *Ascl1* was linked to EGFP via a self-

cleaving P2A peptide (Figure 6D, Supplementary Figure 1B). Cre-dependent ASCL1

expression was validated using a HEK293FT-Cre cell line (Figure 6E). We next overexpressed

ASCL1 at late embryonic stages using our system, when both glio- and neurogenesis are

ongoing, to assess its ability to promote neuronal differentiation. Sox10-CreERT2 embryos

were injected at E7.5 with lentivirus carrying either DIO-EGFP as controls or DIO-EGFP-

Ascl1, followed by tamoxifen administration at E16.0 and analysis at E18.5 (Figure 6F). As

compared to controls, ASCL1 overexpression reduced the EGFP+ progenitors/glia population

(SOX10+ HuC/D-) from ~20% to ~5% (Figure 6G, H) and increased EGFP+ neuronal

population (HuC/D+) from ~80% to ~95% (Figure 6G, I). Notably, ~25% of HuC/D+ EGFP+

cells in controls co-expressed SOX10, indicative of a transitional population, which was

reduced to <5% upon ASCL1 overexpression (Figure 6G, J).

Together, these findings demonstrate that ASCL1 overexpression during a gliogenic

developmental window is sufficient to bias enteric progenitors toward neuronal differentiation.

More broadly, this approach establishes ultrasound-guided *in utero* delivery of Cre-inducible

lentiviral constructs as a powerful method for temporally controlled, cell type-specific gene

manipulation in developing embryos.

DISCUSSION

Our understanding of gut development and related diseases has advanced substantially through

recent technologies, including scRNA-seq and spatial transcriptomics (15, 26, 29). However,

methods to define lineage relationships and interrogate gene function *in vivo* have lagged

behind, partly due to the inaccessibility of embryonic gut using traditional targeting methods.

Here, we present ultrasound-guided AC nano-injection as an efficient methodology to access

major gut cell populations in mouse embryos, including epithelial and mesenchymal

populations, the ENS, as well as gut-extrinsic ganglia. Combined with high-diversity lentiviral

barcoding, this approach establishes a versatile lineage-tracing platform, and coupled with Cre-

dependent constructs, the approach supports temporally controlled, cell type-specific gene

manipulation in the developing gut.

Barcode-based lineage tracing combined with scRNA-seq overcomes key limitations of

traditional Cre-based lineage tracing approaches as they depend on targeting stem cells defined

by single marker genes with often equivocal specificities that risk confounding interpretation.

In contrast, the method presented here is unbiased and enables comprehensive reconstruction

of lineage relationships. It allowed us to resolve unprecedented details amongst

transcriptionally related mesenchymal populations refining their cellular identities, lineage

relationships and early development. First, we identified three fibroblast states with distinct

clonal relationships to specialized mesenchymal cell types. *Kcnn3+* fibroblasts – putative

contractile fibroblasts (MIFs) – were clonally related to ICCs and myoblast/smooth muscle

cells, suggesting shared progenitors. *Adamdec1+* fibroblasts (putative SEMFs) were linked to

pericytes, whereas *Adam12+* fibroblasts (putative AFs) were clonally related to mesothelial

cells. Notably, although previously suggested marker genes were used to annotate fibroblast

subtypes, no consensus nomenclature currently exists for gut fibroblasts. Clonal analysis

therefore provided important orthogonal support for subtype assignments and supported a

developmental logic whereby stromal gut compartments are assembled by a set of fate-biased

mesenchymal progenitor populations. Second, by analyzing clonal dispersion across gut

regions, we demonstrate that gut mesoderm is already regionally specified along the anterior-

posterior axis by E7.5, coinciding with persistent epiblast cells. Third, we present regional diversity amongst mesenchymal cell types, likely reflecting early programs that shape region-specific gut wall composition. Overall, these findings highlight how lineage information complements transcriptomic and anatomical analyses to better define gut stroma development. Increasing evidence shows that different fibroblast subpopulations may have distinct roles in cancer, inflammation, and fibrosis (30), underscoring the importance of better understanding their heterogeneity and development.

By integrating multiple gut regions, we found that region-spanning clones were common in migratory lineages such as ENS and immune cells. Within the ENS, clonal sharing across stomach and colon supports a model in which vagal neural crest-derived progenitors acquire region-specific features, such as colonic *Pou3f3* expression, in response to local cues. In contrast, the restriction of epithelial and mesenchymal clones to single regions indicates that regional identity of these compartments is established incredibly early. Together with the region-specific transcriptional programs observed in mesenchymal subtypes, these findings suggest that developmental patterning of the gut wall emerges from a combination of early regional allocation and local differentiation programs.

Beyond lineage reconstruction, *in utero* nano-injection enables conditional gene manipulation in defined ENS populations. Using Baf53b-Cre and Sox10-CreERT2 mice, we show that Cre-dependent lentiviral cargo can be activated in differentiating enteric neurons or temporally controlled progenitor/glial populations. Proof-of-principle experiments with *Ascl1* demonstrated that the system can be used to test gene function *in vivo* during defined developmental windows. This approach complements emerging fetal atlases and disease genetics studies, which increasingly identify candidate regulators in transient embryonic cell states but often lack efficient *vivo* validation strategies. We believe that the ability to combine

clonal analysis with targeted perturbation can help shift descriptive cell atlas data to mechanistic

studies of gut development.

Several limitations should also be considered. Our results highlight a strong temporal

component of lineage accessibility and commitment. Targeting at E7.5^{Early} enhanced labeling

of ENS precursors but also increased cross-germ-layer clones, consistent with presence of

epiblast cells. While residual epiblast labeling complicates clonal interpretation, it also provides

a functional readout of the developmental window during which germ-layer restriction is being

consolidated *in vivo*. Precise embryonic staging is thus essential, and inter-embryo variability

can complicate stage estimation, although ultrasound imaging partially mitigates this issue.

Tissue dissociation, single-cell capture, and barcode recovery also introduce dropouts that will

inevitably reduce clonal reconstruction efficiency. In the gene perturbation setting, conditional

targeting efficiencies in the ENS were relatively modest. However, we believe that this

mosaicism may in fact be advantageous for studying cell-intrinsic mechanisms given that ENS

development depends on coordinated proliferation and long-range migration. Broad

perturbation within the ENS may cause large secondary effects at the population level, whereas

mosaic manipulation may reveal how genetically altered cells behave within a largely normal

developing ENS.

Finally, this work has implications for understanding the developmental origins of GI

disease. Gut organogenesis relies on interdependent differentiation of diverse cell populations

from all three germ layers, and subtle deviations in lineage allocation, maturation, or regional

patterning may have lasting functional consequences (31). This is relevant not only to

congenital disorders such as Hirschsprung disease, achalasia, and CIPO, but also to more

common conditions in which early developmental alterations may predispose to later

dysfunction. Some of these disorders may result from *de novo* mutations, in which early

mutations occur in a mosaic fashion (32). It is conceivable that the impact of such mutations

would be more predictable with a clearer understanding of refined cell lineages arising in post-gastrula stages. We propose that this platform will provide a scalable strategy for mechanistic testing of candidate disease regulators, and modeling early cellular events that may contribute to GI disorders and enteric neuropathologies.

ACKNOWLEDGEMENTS

We thank Jonas Frisén and Michael Ratz for designing and sharing the tdTomato 30N barcoded viral plasmids. We also thank Jonas Frisén for access to flow cytometry at Karolinska Institutet. We thank Fernando Lopez-Redondo for laboratory assistance. The authors acknowledge support from the National Genomics Infrastructure in Stockholm funded by Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish Research Council. The authors acknowledge the computational resources provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS), partially funded by the Swedish Research Council through grant agreement no. 2022-06725 and the Centre for Bioinformatics and Biostatistics (CBB) at Karolinska Institutet funded by CIMED Infrastructure and Karolinska Institutet. The imaging was performed at the Biomedicum Imaging Core with support from the Karolinska Institutet. We thank the Viral Vector Facility (VVF) of the Neuroscience Center Zurich (ZNZ) for excellent service, construction and production of Lenti viral plasmids. We thank animal staff from the Department of Comparative Medicine at Karolinska Institutet for help with animals.

U.M. was supported by the Swedish Research Council (Vetenskapsrådet; 2020-01129 and 2022-01570), the European Research Council (divENSify; 101045026), the Knut and Alice Wallenberg Foundation (KAW; 2020.0109), the Strategic Research Area in Neuroscience, the Strategic Research Area in Stem Cell Research and Regenerative Medicine, Karolinska Institutet (KI) Consolidator Grant, and the Brain Foundation (Hjärnfonden; FO2025-0169;

FO2023-0130). E.R.A was supported by KAW (2020.0109), KI Consolidator Grant (2-

195/2021), the Strategic Research Area in Neuroscience, Hjärnfonden (FO2025-0271),

Vetenskapsrådet 3R grants (2024-03113, 2021-03537, 2017-01054), and the European

Research Council (LIMITLESS; 101171156). K.P was supported by Marie Sklodowska Curie

Award (1011544005; DeSTENy). K.H, J.H and Z.L were supported by a KI Grant for Salary

(2-2110;2019-7;2019-3).

FIGURE LEGENDS

Figure 1. Efficient targeting of neural crest/placode populations destined to innervate the

gut by *in utero* transduction at E7.5. (A) Schematic drawing of experimental setup for

ultrasound-guided *in utero* nano-injection. Left-most panel reproduced from Mangold et al

2021 (10), with permission. The right panel was created with BioRender.com. (B) Uniform

manifold approximation and projection (UMAP) plot of cells recovered from E9.5 and E10.5

embryos following E7.5 amniotic cavity injection (11). (C) UMAP plot of reprocessed neural

crest/placode cells (green populations in B), revealing 14 clusters. (D) Schematic drawing of

neural crest/placode-derived ganglia innervating the gut. (E) Dot plot showing expression of

canonical marker genes for neural crest/placode populations across the clusters. Dot size

represents the fraction of cells expressing the gene and color intensity represents mean

expression. (F) Feature plots showing expression of neuronal (*Tubb3+*) and migratory neural

crest/placode (*Sox10+*) markers. (G) Feature plots showing gene expression associated with

nodose ganglia, sympathetic ganglia, DRG and the ENS. (J) Representative

immunofluorescence staining of an E13.5 DRG showing H2B-GFP expression in both SOX2+

neural progenitors (arrowheads) and NeuN+ postmitotic neurons (asterisks). (I, J) Bar graphs

showing proportions of GFP positive NeuN+ neurons and SOX2+ neural progenitors along the

thoracic-sacral axis (n=3 mice). Bars represent the mean ± standard deviation (SD). (K)

Representative immunofluorescence image of E18.5 small intestine showing GFP expression

in HuC/D+ neurons (asterisks) and SOX10+ ENS progenitors and glia (arrowheads). (L, M)

Bar graphs showing proportion of GFP positive HuC/D+ neurons and SOX10+ progenitors/glia

across gut regions (n=4 mice). Bars represent mean ± SD. Each dot represents one animal. (N)

Representative immunofluorescence image of E18.5 gut showing GFP expression in alpha-

SMA+ smooth muscle (arrowheads) and EpCAM+ epithelium (asterisks). Scale bars: 50μm.

Figure 2. *In utero* transduction at E7. 5 targets diverse cell types in the developing gut. (A)

Schematic of injection of lentiviral barcode library into the E7.5 amniotic cavity. (B)

Experimental workflow to generate the iE7.5-aE16.5 dataset consisting of TOM+ cells, flow

sorted from two gut regions. (C) UMAP plot of high-quality gut cells colored by tissue of origin.

Pie chart indicates the proportion of cells from each region. (D) UMAP plot showing 8 major

clusters, annotated based on established marker gene expression. (E) Dot plot displaying

expression of canonical marker genes used to define major clusters. (F) Feature plots displaying

marker gene expression of epithelial cell types. (G) Reprocessed ENS cells revealing four

clusters. (H) Feature plots showing expression of selected markers for ENS states and branches.

(I) UMAP plot showing subclusters within fibroblast-like and smooth muscle-like populations.

(J) Dot plot showing marker gene expression distinguishing fibroblast-like and smooth muscle-

like subclusters. (K) Schematic drawing indicating the diverse gut cell types targeted by *in utero*

nano-injection, showing that all three germ layers are efficiently transduced. Part of 2A, B and

K were created with BioRender.com. (E, J) Dot size represents the percentage of cells

expressing the gene, and color indicates relative mean expression.

Figure 3. Lentiviral barcoding at E7.5 resolves clonal lineage relations in the developing

gut. (A) UMAP plot highlighting proportions of cells with CloneIDs in the iE7.5-aE16.5

dataset. (B) UMAP plot showing 11 major gut cell types after merging subclusters. (C) Stacked

bar plot showing distribution of clone categories across major cell types. Schematic drawing

describing clonal types (right). (D) UpSet plot showing frequencies of clones with different

composition of major cell types. (E) UpSet plot showing minimal clone sharing across germ

layers. Inset chord diagrams in (D, E) illustrate the extent and direction of clonal sharing, with

ribbon width proportional to the number of shared clones. (F, G) Heatmaps showing the

correlation matrix of clonal coupling z-scores, computed as deviations of observed clone

sharing relative to randomized data, for refined gut cell types (F) and restricted to mesoderm-

derived cell types (G). Colored boxes denote germ-layer origin (pink: endoderm; purple:

ectoderm; yellow: mesoderm), or lineage-coupled mesenchymal populations (green: Fibroblast

4 with myofibroblasts, ICC and smooth muscle cells; black: Fibroblast 3 with mesothelial cells;

brown: Fibroblast 1 with pericytes). (G) Schematic summary of lineage relationships amongst

mesenchymal cell types inferred from the clonal coupling analysis. H and left panel of 3C were

created with Biorender.com.

Figure 4. *In utero* transduction at early E7.5 improves ENS targeting but uncovers

presence of epiblast cells. (A) Schematic diagram showing the generation of the iE7.5^{Early}-

aE16.5 dataset consisting of three gut regions. Created with BioRender.com. (B) UMAP plot

showing major cell types. (C) Alluvial plot showing proportions of major cell types in the iE7.5-

aE16.5 and iE7.5^{Early}-aE16.5 datasets. (D) UMAP plot showing refined gut cell type annotations

in the iE7.5^{Early}-aE16.5 dataset. (E) Stacked bar plot showing distribution of clone categories

across major cell types identified in the iE7.5^{Early}-aE16.5 dataset. (F) UpSet plot showing

frequencies of clones with different composition of major cell types. (G) UpSet plot showing

notable clone sharing across germ layers. Inset chord diagrams in (F, G) illustrate the extent

and direction of clonal sharing, with ribbon width proportional to the number of shared clones.

(H, I) Heatmaps showing the correlation matrix of clonal coupling z-scores, computed as

deviations of observed clone sharing relative to randomized data, for refined gut cell types (H)

and restricted to mesoderm-derived cell types (I). Colored boxes denote germ-layer origin

(pink: endoderm; purple: ectoderm; yellow: mesoderm), or lineage-coupled mesenchymal

populations (green: Fibroblast 4 with myofibroblasts, ICC and smooth muscle cells; black:

Fibroblast 3 with mesothelial cells; brown: Fibroblast 1 with pericytes). (J) Chord diagram

showing self- and cross-linking between Mrc1+ myeloid and Trbc1/2+ lymphoid cells. (K)

UpSet plot showing co-occurrence of ENS cell states/types in the same clone. Note that Branch

B1/2 are not exact equivalents of Branch B1/2 defined in Morarach et al, 2021 (15).

Figure 5. Regional clone sharing reveals long-range ENS migration and early regional

patterning of mesenchymal populations along the GI tract. (A) UpSet plot showing co-

occurrence of major cell types in cross-region clones in the iE7.5-aE16.5 dataset. (B) UMAP

plot highlighting cells in cross-region and local clones. (C) Alluvial plot illustrating relationship

between tissue region, cell type, and CloneID of the cross-region clones in (A). (D) UpSet plot

showing co-occurrence of major cell types in the cross-region clones in the iE7.5^{Early}-aE16.5

dataset (restricted to single-germ-layer clones). (E) UMAP plot highlighting cells in cross-

region and local clones. (F) Heatmap showing the distribution of clones across gut regions in

the iE7.5-aE16.5 dataset, with binary indication of presence or absence. Colored bars denote

cell type identities of clones. (G) Heatmap showing clonal composition of ENS cell states/types

for cross-region or local clones, revealing no apparent bias. (H) UMAP plot highlighting the

differentiation state of clones with different dispersion. (I) Violin plot showing comparable

levels of *Pou3f3* expression in colonic ENS cells that are local or share clones with small

intestine and/or stomach. Parts of 5A and D were created with Biorender.com.

Figure 6. Temporal overexpression of Ascl1 in Sox10+ cells promotes neuronal

differentiation (A) Schematic overview of the Lenti-DIO-EGFP vector, before and after Cre-

mediated gene inversion. (B) Representative immunohistochemistry images of intestine from

wild-type mice injected with LV-H2B-GFP showing GFP expression in both HuC/D- cells

(asterisks) and HuC/D+ (arrowheads), and Baf53b-Cre mice injected with Lenti-DIO-EGFP

showing GFP expression restricted to HuC/D+ cells (arrowheads). (C) Bar graph showing the

percentage of HuC/D+ cells expressing GFP across gut regions in Baf53b-Cre embryos injected

with Lenti-DIO-EGFP. (n=5 mice). Bars represent mean ± SD. (D) Schematic drawing of the

Lenti-DIO-EGFP-Ascl1 construct with and without Cre-mediated gene inversion. (E)

Representative immunohistochemistry images showing validation of transgene expression in

Cre-expressing HEK293FT cells. (F) Schematic representation of experimental procedure for

in vivo overexpression of EGFP-Ascl1 or EGFP in ENS progenitors. Part was created with

BioRender.com. (G) Representative confocal images of E18.5 small intestine transduced with

DIO-EGFP or DIO-Ascl1-EGFP vectors showing SOX10 and HuC/D expression patterns in

GFP+ cells. asterisks denote EGFP+ cells and texts indicate their status of expression, with +

or – signs denoting positive or negative expression. (H-J) Bar graphs showing the proportions

of GFP+ cells with SOX10 expression without HuC/D (H), GFP+ cells with HuC/D expression

(I), and GFP+ HuC/D+ cells expressing SOX10 (J), in Baf53b-Cre embryos injected with either

DIO-Ascl1-EGFP or Lenti-DIO-EGFP. (H)*p=0.0136; (I)**p= 0.0086, and (J)* p= 0.011. n=6-

7 mice per condition. SI: small intestine; AC: amniotic cavity; WT: wild-type. Scale bars:

20μm.

REFERENCES:

1.   C. H. Knowles, G. Lindberg, E. Panza, R. De Giorgio, New perspectives in the diagnosis and management of enteric neuropathies. *Nat Rev Gastroenterol Hepatol* **10**, 206–218 (2013).

2.   B. Niesler, S. Kuerten, I. E. Demir, K. H. Schafer, Disorders of the enteric nervous system - a holistic view. *Nat Rev Gastroenterol Hepatol* **18**, 393–410 (2021).

3.   M. P. Sherman, H. Zaghouani, V. Niklas, Gut microbiota, the immune system, and diet influence the neonatal gut-brain axis. *Pediatr Res* **77**, 127–135 (2015).

4.   S. K. Hashmi, R. H. Ceron, R. O. Heuckeroth, Visceral myopathy: clinical syndromes, genetics, pathophysiology, and fall of the cytoskeleton. *Am J Physiol Gastrointest Liver Physiol* **320**, G919–G935 (2021).

5.   R. Dixit *et al.*, Efficient gene delivery into multiple CNS territories using in utero electroporation. *J Vis Exp* 10.3791/2957 (2011).

6.   E. Andersson *et al.*, Identification of intrinsic determinants of midbrain dopamine neurons. *Cell* **124**, 393–405 (2006).

7.   E. Dvoretskova *et al.*, Spatial enhancer activation influences inhibitory neuron identity during mouse embryonic development. *Nat Neurosci* **27**, 862–872 (2024).

8.   T. Uesaka, H. M. Young, V. Pachnis, H. Enomoto, Development of the intrinsic and extrinsic innervation of the gut. *Dev Biol* **417**, 158–167 (2016).

9.   K. Mangold, J. He, S. Stokman, E. R. Andersson, Murine Neural Plate Targeting by In Utero Nano-Injection (NEPTUNE) at Embryonic Day 7.5. *J Vis Exp* 10.3791/63148 (2022).

10.   K. Mangold *et al.*, Highly efficient manipulation of nervous system gene expression with NEPTUNE. *Cell Rep Methods* **1** (2021).

11.   S. de Haan *et al.*, Ectoderm barcoding reveals neural and cochlear compartmentalization. *Science* **388**, 60–68 (2025).

12.   M. Ratz *et al.*, Clonal relations in the mouse brain revealed by single-cell and spatial transcriptomics. *Nat Neurosci* **25**, 285–294 (2022).

13.   C. Laranjeira *et al.*, Glial cells in the mouse enteric nervous system can undergo neurogenesis in response to injury. *J Clin Invest* **121**, 3412–3424 (2011).

14.   F. Memic *et al.*, Transcription and Signaling Regulators in Developing Neuronal Subtypes of Mouse and Human Enteric Nervous System. *Gastroenterology* **154**, 624–636 (2018).

15.   K. Morarach *et al.*, Diversification of molecularly defined myenteric neuron classes revealed by single-cell RNA sequencing. *Nat Neurosci* **24**, 34–46 (2021).

16.   V. P. Masamsetti *et al.*, Lineage contribution of the mesendoderm progenitors in the gastrulating mouse embryo. *Dev Cell* **60**, 1991–2006 e1999 (2025).

17.   S. Torihashi *et al.*, Blockade of kit signaling induces transdifferentiation of interstitial cells of cajal to a smooth muscle phenotype. *Gastroenterology* **117**, 140–148 (1999).

18.   S. A. Baker *et al.*, Distribution and Ca(2+) signalling of fibroblast-like (PDGFR(+)) cells in the murine gastric fundus. *J Physiol* **591**, 6193–6208 (2013).

19.   S. Schneider *et al.*, Single Nucleus Sequencing of Human Colon Myenteric Plexus-Associated Visceral Smooth Muscle Cells, Platelet Derived Growth Factor Receptor Alpha Cells, and Interstitial Cells of Cajal. *Gastro Hep Adv* **2**, 380–394 (2023).

20.   S. E. Viitala E.W, Miihkinen M, Chalkidi N, Stavropoulou A, Lemmetyinen TT, Päivinen P, Kaptrio T, Hagström J, Haglund C, Mäkelä TP, Aittokallio T, Katajisto P, Koliaraki V, Ollila S. (2025) High-resolution integrative analysis allows characterization and spatial annotation of normal and cancer-associated colon fibroblasts. (bioRxiv).

21. Y. Rinkevich *et al.*, Identification and prospective isolation of a mesothelial precursor
lineage giving rise to smooth muscle cells and fibroblasts for mammalian internal
organs, and their vasculature. *Nat Cell Biol* **14**, 1251–1260 (2012).

22. L. T. G. Harland *et al.*, A spatiotemporal atlas of mouse gastrulation and early
organogenesis to explore axial patterning and project in vitro models onto in vivo space.
Cell Rep **44**, 116047 (2025).

23. R. Lasrado *et al.*, Lineage-dependent spatial and functional organization of the
mammalian enteric nervous system. *Science* **356**, 722–726 (2017).

24. C. C. Bain, A. Schridde, Origin, Differentiation, and Function of Intestinal
Macrophages. *Front Immunol* **9**, 2733 (2018).

25. W. H. Neo, A. L. M. Lie, M. Z. H. Fadlullah, G. Lacaud, Contributions of Embryonic
HSC-Independent Hematopoiesis to Organogenesis and the Adult Hematopoietic
System. *Front Cell Dev Biol* **9**, 631699 (2021).

26. C. M. Wright *et al.*, scRNA-Seq Reveals New Enteric Nervous System Roles for
GDNF, NRTN, and TBX3. *Cell Mol Gastroenterol Hepatol* **11**, 1548–1592 e1541
(2021).

27. E. Blaugrund *et al.*, Distinct subpopulations of enteric neuronal progenitors defined by
time of development, sympathoadrenal lineage markers and Mash-1-dependence.
Development **122**, 309–320 (1996).

28. F. Memic *et al.*, Ascl1 Is Required for the Development of Specific Neuronal Subtypes
in the Enteric Nervous System. *J Neurosci* **36**, 4339–4350 (2016).

29. M. Zilbauer *et al.*, A Roadmap for the Human Gut Cell Atlas. *Nat Rev Gastroenterol*
Hepatol **20**, 597–614 (2023).

30. N. Chalkidi, C. Paraskeva, V. Koliaraki, Fibroblasts in intestinal homeostasis, damage,
and repair. *Front Immunol* **13**, 924866 (2022).

31.   Z. Zhang *et al.*, Impairment of stromal-epithelial regenerative cross-talk in

820   Hirschsprung disease primes for the progression to enterocolitis. *Sci Transl Med* **17**,

821   eadp4679 (2025).

822   32.   M. Mohiuddin, R. F. Kooy, C. E. Pearson, De novo mutations, genetic mosaicism and

823   human disease. *Front Genet* **13**, 983668 (2022).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6