Title: *In utero* transduction resolves gut cell lineages and enables conditional gene

perturbation in the developing enteric nervous system

Short Title: Resolving and targeting gut cell lineages

Ziwei Liu[1,‡], Krishnanand Padmanabhan[1,‡], Jingyan He[2,†], Katrin Hector[2], Bettina

Semsch[2,3], Jia Sun[3], Viktoria Knoflach[1], Sarantis Giatrellis[2], Khachatur Dallakyan[1], Christian

Göritz[2], Emma Rachel Andersson[2,*], Ulrika Marklund[1,*]

[1] Department of Medical Biochemistry and Biophysics, Karolinska Institutet, Sweden

[2] Department of Cell and Molecular Biology, Karolinska Institutet, Sweden

[3] Comparative Medicine, Karolinska Institutet, Sweden

† Current Adress: Janelia Research Campus, Howard Hughes Medical Institute,

Ashburn, USA

‡These two authors contributed equally

*These authors share corresponding authorship

Correspondence: Ulrika.Marklund@ki.se; Address: Tomtebodavagen 16, 171 77 Stockholm,

Sweden. Emma.Andersson@ki.se; Address: Tomtebodavagen 16, 171 77 Stockholm,

Sweden.

Disclosures: The authors declare no conflict of interest.

ABSTRACT

Background and Aims: Understanding the etiology of gastrointestinal malfunction requires insights into how heterogeneous gut cell lineages arise during development and the identification of genes controlling these processes. While *in utero* genetic manipulation has advanced knowledge in other organs, the gut and its intrinsic enteric nervous system (ENS) remain difficult to target. We aimed to establish an approach to resolve precise lineage relationships and achieve conditional gene perturbation of developing gut cell types *in vivo*.

Methods: We performed ultrasound-guided *in utero* nano-injection of lentiviral vectors into the mouse amniotic cavity at embryonic day 7.5. By combining lentiviral DNA barcoding with single cell RNA-sequencing, we reconstructed clonal relationships between distinct gut cell types. We used Cre-dependent lentiviral vectors in transgenic mice to achieve cell type-specific expression and tested whether gene overexpression can direct developmental processes.

Results: Lentiviral delivery efficiently targeted gut-innervating ganglia and all major gut cell types, including neural, epithelial, immune, and mesenchymal populations. Barcode-based lineage tracing revealed clonal relationships within and across gut cell types. Notably, specialized mesenchymal populations, including pericytes, mesothelial cells, and interstitial cells of Cajal (ICCs) were each linked to different fibroblast subpopulations. Regional analyses demonstrated early establishment of anterior–posterior identity of mesenchyme, whereas ENS and immune cells exhibited long-range clonal dispersion. Cre-dependent targeting enabled selective gene expression in ENS progenitors or neurons, and temporally controlled overexpression of Ascl1 promoted neuronal differentiation.

Conclusions: *In utero* lentiviral transduction expands experimental access to the developing gastrointestinal tract, enabling lineage-resolved analysis and conditional gene manipulation of

gut cell types, including the ENS. This approach enables mechanistic investigation of gut

organogenesis and offers a framework to study developmental origins of disease.

Keywords: Enteric Nervous System, *in utero* lineage tracing, mesenchymal cell types, single

cell RNA-sequencing, gastrointestinal development

INTRODUCTION

The gastrointestinal (GI) tract is essential for life. Congenital gut malformations including

Hirschsprung disease, pyloric stenosis, and various intestinal atresias result in diverse and

life-long complications. In the general population, 30% suffer from chronic gut dysfunctions,

including disorders of gut-brain interaction (DGBIs) and inflammatory bowel disease (IBD). [1]

It is conceivable that many of those conditions arise from subtle alterations in gut cell

composition already established at fetal stages.[2] Indeed, premature birth is a risk factor for

developing DGBIs.[3] Chronic intestinal pseudo-obstruction (CIPO) is often congenital and can

be myopathic, mesenchymopathic or neuropathic, underscoring that intestinal motility

depends on functional interactions among various gut cell types. Increasing evidence shows

that various fibroblast types play important regulatory roles in cancer, inflammation, and

fibrosis.[4] Gut dysfunctions are also common co-morbidities in central nervous system (CNS)

disorders, including autism spectrum disorder, likely resulting from neurodevelopmental

issues in the enteric nervous system (ENS).[5] Thus, there is a need to better resolve how

different cell lineages in the GI-tract develop.

Experimental gene perturbation/manipulation - genetic gain or loss of function – are

essential tools to resolve how gene regulatory networks control cellular differentiation and

how developmental deviations can lead to pathology. Owing to the relative ease with which

the CNS can be genetically manipulated using *in vivo* embryonic transduction or

electroporation[6], the CNS field has gained intricate knowledge of precise gene regulatory

networks controlling the generation of different neuronal subtypes.[7, 8] These manipulations are

possible due to the ventricular structure of the brain and the lumen of the spinal cord, making

them accessible to plasmids or viral vectors. In contrast, the GI-tract is positioned internally

and not possible to target using these common approaches.

Gut physiology is critically dependent on neuronal control, with the ENS embedded

within the gut wall, essential for peristalsis, secretion, and blood flow regulation. These

functions are modulated by sympathetic innervation, primarily of celiac ganglion neurons,

while sensory information is captured from the gut via dorsal root ganglia (DRG) at the trunk

regions, and nodose ganglia in cranial regions. The ENS, sympathetic ganglia, and DRG are

formed from neural crest cells (NCC), a transient embryonic population originating from the

dorsal neural tube.[9] In the mouse, delamination of NCC occurs around embryonic day (E)8

(Theiler stages TS12-13) and thereafter, the migrating cells and the ganglia that form are

positioned unfavorably for traditional *in utero* manipulations that are typically performed

from E11.5.[6] Thus, experimental systems to rapidly investigate the development of the murine

gut, the ENS, and other gut-innervating ganglia have been lacking.

Recently, we developed neural plate targeting with *in utero* nano-injection

(NEPTUNE) to target the developing CNS at earlier stages.[10, 11] With NEPTUNE, ultrasound

is used to guide lentiviral delivery of cargo into the mouse amniotic cavity as early as E7.5,

allowing access to 80-95% of cells in the CNS. The technique can also be used to introduce

high-diversity lentiviral barcode libraries to resolve refined lineages using next generation

single cell lineage tracing.[12, 13] As distinct barcodes integrate in the DNA of each transduced

cell, clonal relationships can be reconstructed after successive cell divisions by single cell

RNA-sequencing (scRNA-seq), revealing how individual progenitors contribute to the

generation of diverse cell types. Here, we asked whether the method allows for targeting of

the GI-tract and of neural crest/placodes, whose derivatives innervate the gut, for the purpose

of revealing clonally linked gut cell types and performing genetic experimentation of the

developing ENS.

We found that ultrasound-guided *in utero* lentiviral nano-injection at E7.5 transduces

progenitors that contribute to the GI tract, including endodermal (i.e. epithelia), mesodermal

(e.g. smooth muscle cells, fibroblasts and interstitial cells of Cajal; ICCs) and ectodermal

progenitors (e.g. ENS). In addition, the method efficiently targets extrinsic gut-innervating

ganglia (e.g DRG). By combining high-throughput clonal labeling and transcriptomic

profiling of cells in the gut, we identified fate-restricted progenitor cells present at the time of

transduction and reconstructed lineage relationships across differentiating gut cell types. Of

note, we uncovered molecularly defined fibroblast populations that are clonally coupled with

either mesothelial cells, pericytes or ICCs, revealing early diversification paths within the gut

mesenchyme. We find that ENS targeting is most efficient at early E7.5, but that residual

epiblast cells are still present at this stage. Beyond lineage tracing, we show that *in utero*

nano-injections can be utilized for conditional gene perturbation in the ENS. Using the pan-

neuronal Baf53-Cre mouse line, we demonstrate that lentivirus carrying Cre-dependent cargo

is selectively expressed in developing enteric neurons. In Sox10-CreERT2 mice, transgene

expression can instead be induced in ENS progenitors/glia in a temporally controlled manner.

As proof-of-principle, we show that timed overexpression of the proneural transcription factor

Ascl1 biases progenitors toward neuronal over glial differentiation. Taken together, our results

show that ultrasound-guided *in utero* nano-injection is a versatile tool to investigate gut

organogenesis and ENS development.

MATERIAL and METHODS

Mice

The following transgenic and reporter mouse strains were used: Baf53b-Cre (JAX, #027826),

Sox10-CreERT2 (SER26)[14], R26-EYFP (JAX #006148) and R26-tdTomato (Ai14; JAX

#007908). C57/B6J and CD1 (Charles River Laboratories) wild-type (WT) mice were used.

Animals were group-housed, with food and water *ad libitum,* under 12-h light-dark cycle

conditions, 22°C ambient temperature and 50% humidity. Animal experiments were approved

by the local ethics committee (Stockholm Norra djurfo☐rso☐ksetiska na☐mnd,

Jordbruksverket; N5264/18, N6626-2019, N5237-2023, 8188-2017, 2987-2020 and 315-23).

Lentivirus constructs

Plasmid *LV-H2B-GFP* was a gift from Elaine Fuchs (Addgene #25999). *LV-EF1A-H2B-*

tdTomato-30N was previously described.[12, 13] *Lenti-DIO-EGFP* (Supplementary Figure 1A)

was based on the interim vector *pRRLSIN-cPPT-hPGK-EGFP-WPRE* (Gift from Didier

Trono, Addgene #12252). *Lenti-DIO-EGFP-Ascl1* (Supplementary Figure 1B) was

constructed based on *pRRLSIN-cPPT-shortCAG-dlox-EGFP_2A(PTV)(rev)-dlox-WPRE* and

the GeneArt product *mAscl1* (710bp). Both vectors were constructed, amplified, and

characterized by the Viral Vector Facility (VVF) of the Neuroscience Center Zurich (ZNZ).

Plasmids and virus were produced and characterized as described in (Supplementary Material

and Methods). High-titer vesicular stomatitis virus G (VSV-G)–pseudotyped lentiviral

particles were produced either in-house or by a commercial facility (GEG TECH, France) and

used at a minimum titer of $1 \times 10\square$ transducing units (TU)/mL.

Ultrasound-guided *in utero* nano-injections

A detailed description of the procedure has been reported[10, 11] and is described in

(Supplementary Material and Methods). We considered Theiler stages TS11A–C to represent

the optimal E7.5 window for ultrasound-guided *in utero* injections (Supplementary Figure 4A,

B), while injection at E8 or later failed to efficiently target gut and ENS (Supplementary

Figure 4C, D).

Tamoxifen administration

Plug-mated females received tamoxifen (0.1 mg/g body weight; Sigma) dissolved in corn

oil/ethanol (9:1) by intraperitoneal injection at E16.0-E16.5. Embryos were collected 26 or 55

hours after tamoxifen administration.

Tissue preparation and immunohistochemistry

E13.5 embryos were decapitated and fixed in 4% PFA in PBS for 4-5 hours at 4 °C. Intestines

from E14.5-E18.5 were dissected out and fixed in 4% PFA in PBS at 4 °C for 1.5-2 hours.

Samples were washed in PBS three times and dehydrated in 30% sucrose in PBS at 4 °C

overnight. Intestines were cut into 1 cm long pieces, representing different regions and

embedded for transverse analysis in OCT freezing medium and stored at -80 °C. Samples

were sectioned at 12 μm and stored at -20 °C.

Frozen tissue sections were air-dried at RT for 30 min, rinsed with PBS and incubated

in blocking solution containing purified donkey anti-mouse Fab (Jackson Laboratories, PA,

USA #715-007-003) diluted 1:50 in PBS for 2 hours at room temperature (RT). After rinsing

with PBS, the sections were blocked with 2% normal donkey serum (NDS, Jackson) and 0.1%

Triton X-100 (Sigma) in PBS for 2 hours, and incubated with primary antibodies diluted in

the same solution overnight at 4°C. They were then washed three times with PBS (10 minutes

each) and incubated with secondary antibodies at RT for 1 hour. Tissue was washed three

times with PBS (10 minutes each) and mounted in DAKO mounting medium (Agilent)

containing DAPI (4',6-diamidino-2-phenylindole). Primary and secondary antibodies are

specified in (Supplementary Material and Methods.

Preparation of single-cell RNA sequencing datasets

CD1 embryos injected with LV-EF1A-H2B-tdTomato-30N were harvested at E16.5. Stomach

and 1/3 of the distal small intestine were harvested from the E7.5 experiment, while the

stomach, 2/3 of the small intestine, and the entire large intestine were harvested from the

E7.5^{Early} experiment. Dissociation was performed as described in (Supplementary Material

and Methods). Briefly, tissues were cut, enzymatically processed, triturated and TOM+

DRAQ7-cells isolated through flow cytometry (Supplementary Figure 1C-H). cDNA libraries

were prepared using 10x Chromium Next GEM Single Cell 3' Reagent Kits Version 3.1 Dual

Index (10x Genomics, CA, USA). The datasets were named according to stage of injection (i)

and analysis (a): iE7.5-aE16.5 and iE7.5^{Early}-aE16.5.

Analysis of single-cell RNA sequencing datasets

A detailed description of the scRNA-seq analysis can be found in (Supplementary Material

and Methods). In brief, E16.5 scRNA-seq datasets were aligned to a combined mouse genome

(GRCm38) with a tdTomato-N transgene containing a 30bp barcode region[12, 13] using

CellRanger. Raw count matrices were processed in Seurat, with ambient RNA contamination

assessed and corrected using SoupX. Low-quality cells and doublets were removed based on

gene/UMI thresholds, mitochondrial contents and DoubletFinder predictions. Data were

normalized using SCTransform, followed by dimensionality reduction, batch correction with

Harmony, clustering and UMAP visualisation. CloneIDs were assigned to cells using the

TREX (TRacking and gene EXpression profiling of clonally related cells) pipeline[12, 13] based

on shared barcodes. Clonal composition and relationships between cell types were analyzed

using statistical comparisons to randomized data, generating clonal coupling scores and

correlation-based clustering to infer lineage relationships.

Data Accession:

scRNA-seq data used for analysis in Figure 1 and Supplementary Figure 2 can be found at

ArrayExpress with the accession number E-MTAB-14817. The scRNA-seq data generated in

this study will be made available at GEO upon publication.

RESULTS

Efficient targeting of gut-innervating ganglia by *in utero* transduction at E7.5

Previous studies have demonstrated that high-titer lentivirus injection into the amniotic cavity

of E7.5 embryos results in widespread transduction of the neural plate and otic placodes.[10-12]

Because gut-innervating ganglia originate from neural crest or from placodes, we

hypothesized that these structures could also be targeted. To assess whether E7.5 amniotic

cavity lentiviral injection targets neural plate progenitors that generate gut-innervating

neuronal populations (Figure 1A), we re-analyzed a published scRNA-seq dataset of E9.5 and

E10.5 embryos injected at E7.5 with lentiviruses carrying *tdTomato*.[12] The cells in this dataset

had been broadly categorized as "neural tube cells", "epithelial cells", and "neural crest cells"

(Figure 1B, Supplementary Figure 2A). To investigate whether developing gut-innervating

neurons had been targeted (Figure 1D), we renormalized and subclustered the "neural crest

cells" (4,916 cells), which yielded 14 clusters (Figure 1C, E; Supplementary Figure 2B). We

identified three *Tubb3+* neuronal clusters (clusters 4, 8, 12), while remaining clusters

displayed high *Sox10* expression, indicative of neural progenitor or other immature cell states

(Figure 1E, F; Supplementary Figure 2C, D). Amongst *Sox10+* clusters, clusters 0 and 9 were

defined as mesenchymal as they expressed *Prrx1* and *Alx1*, while cluster 13 showed a

melanocyte identity with high *Mitf* and *Dct* expression. Cluster 6 exhibited an autonomic

nervous system character (*Phox2b+*, *Ascl1+*, *Ret+*) containing presumptive sympathetic

(*Hand2-, Isl1+*) and ENS cells (*Hand2+, Isl1-*) (Figure 1G). The remaining *Sox10+* clusters

lacked unique markers and likely represented progenitor or glial cells; these were annotated as

NCC1-NCC7 (Figure 1E). Amongst neuron-enriched clusters, cluster 4 displayed a nodose

ganglion identity (*Phox2a+, Phox2b+, Tbx1+, Tfap2b-*). Cluster 8 and 12 expressed a DRG

profile (*Isl1+, Neurod1+, Six1+, Pou4f1+*), with cluster 12 representing an immature state

(higher *Sox10* and *Foxd3* expression) (Figure 1E; Supplementary Figure 2D).

To further substantiate cluster identities, we addressed the Hox gene expression

patterns (Supplementary Figure 2E-F). Consistent with a cranial origin, the cluster identified

as nodose ganglion exhibited robust anterior Hox gene expression (e.g. *Hoxa1-4*, *Hoxb1-3*).

ENS and parts of the sympathetic ganglia originate from vagal neural crest, and cluster 6

expressed a characteristic suite of *Hoxd3-4* and *Hoxb6*[15]. NCC3 lacked Hox gene expression,

indicating a rostral cranial identity, while the Hox gene pattern of NCC5 suggested they are

progenitors/glia in the nodose ganglion. Posterior Hox gene expression was more pronounced

at E10.5, particularly within DRG clusters and several NCC clusters, indicative of the

anterior-to-posterior emergence of neural crest derivatives. Together, the data demonstrate

that lentiviral injection at E7.5 targets primordia of gut-innervating ganglia, including ENS,

sympathetic ganglia, nodose ganglia and DRG (Figure 1D).

To confirm that gut-innervating ganglia are persistently labelled, we performed

immunohistochemical analysis of E13.5 DRG and E18.5 gut, in embryos injected at E7.5 with

lentivirus carrying H2B-GFP. There was robust transduction of the DRG, with efficient

labelling of both NeuN+ neurons and SOX2+ progenitors/glia from thoracic to sacral levels

(65-83%) (Figure 1H-J). The ENS was uniformly transduced across gut regions, with an

average of 51% HuC/D+ neurons and 44% SOX10+ progenitors/glia (Figure 1K-M;

Supplementary Figure 3A-D). In addition to ENS labelling, we observed GFP across all gut

layers, including in α-SMA+ smooth muscle cells and EpCAM+ epithelial cells (Figure 1N;

Supplementary Figure 3E). This indicates that viral particles in amniotic fluid not only

transduce the neural plate (ectoderm) but are also in contact with endo- and mesodermal germ

cells at E7.5 (Supplementary Figure 4A, B). Indeed, injections during an early developmental

window were crucial for ENS targeting, as injections at E8.0-8.5 failed to label both the ENS

and gut epithelium, with only sporadic labeling of mesenchymal cells observed

(Supplementary Figure 4C, D).

Together, these data demonstrate that E7.5 amniotic cavity lentiviral injection enables

widespread and stable transduction of the GI tract, encompassing extrinsic gut-innervating

neuronal populations, the ENS, and non-ENS gut cell types.

***In utero* transduction targets all major cell types in the developing gut**

As we discovered that injections not only target the ENS but the whole gut wall, we reasoned

that this approach could be leveraged to determine lineage relationships between gut cell

types derived from all three germ layers. Moreover, by capturing different segments of the

gut, conclusions could be drawn about the timing of anterior-posterior regionalization. To

simultaneously profile the transcriptional identity (cell type) and lineage history (clonality) of

single cells, we applied TREX[12, 13] (Figure 2A, B). TREX relies on *in utero* injection of a

high-diversity lentiviral library encoding a 30N (random nucleotides) barcode and tdTomato

resulting in genomic integration into dividing progenitors, thereby allowing stable inheritance

of the construct by all descendant cells. The expression of the barcode under a synthetic

ubiquitous promoter allows its detection by scRNA-seq, enabling reconstruction of clonal

relationships among captured cells based on shared barcodes (CloneIDs).

We injected the tdTomato-30N barcode lentiviral library into the amniotic cavity of

E7.5 embryos (Figure 2A) and analyzed the embryos at E16.5 (dataset thus called iE7.5-

aE16.5). TOM+ cells were sorted from stomach and small intestine by flow cytometry and

sequenced using the 10x Chromium platform (Figure 2B). After removing low-quality cells

and potential doublets, 5,356 cells were retained for analysis, comprising 3,974 cells from the

stomach and 1,382 cells from the small intestine (Figure 2C). Cells from both regions were

integrated, resulting in eight major clusters defined by canonical lineage markers,

corresponding to fibroblast-like cells (*Vim+, Col1a1+, Lum+*), smooth muscle-like cells

(*Myh11+, Actg2+, Acta2+*), endothelial cells (*Cd36+, Pecam1+*), foregut basal cells (*Cdh1+,*

Epcam+, Krt5+, Krt15+), simple epithelial cells (*Cdh1+, Epcam+, Krt8+, Krt18+*), ENS

(*Sox10+, Phox2b+, Ret+, Elavl4+*), immune cells (*Cd52+, Ptprc+*) and mesothelial cells

(*Upk3b+, Msln+, Wt1+*) (Figure 2D, E; Supplementary Figure 5A-C). Further analysis of the

simple epithelial cluster revealed predominantly enterocytes (*Vil1+*), along with smaller

populations of enterochromaffin cells (*Chga+, Tph1+*), goblet cells (*Clca1+, Fcgbp+*), and

intestinal stem cells (*Lgr5+*) (Figure 2F). The 195 ENS cells formed four transcriptionally

distinct clusters representing progenitors, neuroblasts, and neurons (Figure 2G). Notably,

previously discovered neuronal branches characterized by *Etv1* (Branch A) and *Bnc2* (Branch

B) were captured (Figure 2H).[16]

Subclustering of the smooth muscle-like cluster revealed a contractile myofibroblast-

like state (*Trpc3+, Tagln+*) and a proliferative cluster (Figure 2I, J; Supplementary Figure

5D). Subclustering of the fibroblast-like cluster uncovered additional mesenchymal

heterogeneity, including Interstitial cells of Cajal (ICC) (*Kit+, Ano1+*), pericytes (*Rgs5+,*

*Reln+, Pdgfrb+, Pdgfr*a-) and seven transcriptionally distinct fibroblast subtypes (Figure 2I,

J). These comprised proliferative fibroblasts (*Top2a+, Mki67+*), Fibroblast 1 (*Col6a4+,*

Adamdec1+), Fibroblast 2 (*Fndc1+, Dpt+*), Fibroblast 3 (*Adam12+*), Fibroblast 4 (*Kcnn3+*),

Fibroblast 5 (*Mfap5+, Grm3+*) and Fibroblast 6 (*Lepr+, Sp5+*) (Supplementary Figure 5D-

H). The proliferative fibroblast cluster likely comprised a mixture of all fibroblast subtypes.

Collectively, our results demonstrate that *in utero* lentiviral transduction effectively

transduces progenitors of all major gut cell types, encompassing derivatives of the three germ

layers (Figure 2K).

Barcoded *in utero* transduction reveals clonal lineage structures in the developing gut

The comprehensive capture of diverse gut cell types provided us excellent conditions to reveal

detailed lineage relationships. After barcode extraction, 3,959 cells (73.9%) retained

CloneIDs (Figure 3A, B; Supplementary Figure 6A-C). Cells with or without CloneIDs

showed comparable quality metrics and similar CloneID recovery rates were observed across

all cell types, indicating unbiased barcode dropout (Supplementary Figure 6D-G). Most

CloneIDs consisted of a single barcode, indicating single lentiviral particle transductions,

while up to 4 barcodes were detected in a few cells (0.42% of clones) (Supplementary Figure

6H).

In total, 1,658 clones were reconstructed and classified into three categories:

singletons (1,026 clones), single-cell-type clones (302 clones) and multi-cell-type clones (330

clones) (Figure 3C). Multicellular clones contained on average 4.6 cells and reached a

maximum of 31 cells (Supplementary Figure 6I). Systematic assessment of clonal

compositions per major cell type showed that pericytes, ICCs, smooth muscle and mesothelial

cells almost exclusively consisted of multi-cell-type clones (Figure 3C), suggesting

substantial lineage diversification. Instead, ENS and immune cells exhibited only single-cell-

type clones, reflecting early lineage restriction of their respective precursors. Clone sharing

was predominantly found between mesenchymal cell types in the gut, consistent with their

shared mesodermal origin (Figure 3D). However, endothelial and mesothelial cells did not

share clones with each other despite sharing clones with other mesenchymal cell types,

indicating their progenitor allocation is already distinct at E7.5 (Figure 3D). Cells arising from

different germ layers rarely shared clones; only five clones spanned endodermal and

mesodermal lineages, likely reflecting residual pluripotent mesendoderm cells[17] at the time of

injection (Figure 3E).

To systematically resolve lineage relationships among refined cell types, we

performed clonal coupling analysis by calculating z-scores for deviations in observed clone

sharing relative to randomized data. Hierarchical clustering of Pearson correlations between

z-scores segregated cell types into broad groups corresponding to germ-layer origin and

revealed intricate clonal coupling relations between mesoderm-derived populations (Figure

3F-G, Supplementary Figure 6J-K). Notably, ICCs showed strong clonal coupling with

smooth muscle cells and myofibroblasts (Figure 3GF-H). This is consistent with previous

lineage-tracing studies showing that ICCs and myofibroblasts arise from common precursors

and that ICCs are not fibroblasts.[18] However, ICCs, myofibroblasts and smooth muscle cells

also showed clonal coupling with Fibroblast 4 (*Kcnn3+*). This fibroblast likely corresponds to

Pdfgra+ cells of the SIP (smooth muscle cells, ICCs, and PDGFRa[19, 20]), recently renamed

muscle-embedded interstitial fibroblasts; MIFs[21]) (Supplementary Figure 5E, F). Fibroblast 3

(*Adam12+*) showed enriched clonal coupling with mesothelial cells (Figure 3F,G), suggesting

an adventitial fibroblast (AF) phenotype[22], further supported by expression of *Wt1* and *Pi16*

(Supplementary Figure 5E, G). Fibroblast 1 (*Col6a4+, Adamdec1+*) displayed clonal

coupling with pericytes, consistent with the reported localization of *Adamdec1+*

myofibroblasts in the subepithelial space (SEMFs) (Figure 3F, G; Supplementary Figure 5E,

H). Notably, clonal coupling analysis thus segregated fibroblast populations into putative

contractile, adventitial and perivascular trajectories which was less apparent from

transcriptome-based clustering alone (Figure 3H; Supplementary Figure 6L).

Clonal coupling patterns among mesoderm-derived populations indicate that

mesenchymal lineages diverge early into fate-biased progenitor pools that underpin

coordinated tissue patterning during gut morphogenesis (Figure 3H). Collectively, we

demonstrate that single-cell lentiviral barcoding at E7.5 recapitulates canonical developmental

hierarchies and uncovers early cell fate diversification within the gut, corroborating known

lineage relationships while revealing previously unrecognized trajectories contributing to gut

organogenesis.

353   Nano-injection at early E7.5 improves ENS targeting while revealing residual epiblasts

Attempting to enrich for ENS targeting, we performed another nano-injection of the N30

barcode lentiviral library aimed at an earlier developmental stage (E7.5$^{\text{Early}}$, ~TS11b;

Supplementary Figure 4A, B), reasoning that primordia of neural crest could be physically

more accessible and remain for a longer time before delamination. To gain further

information about gut regionalization, we collected stomach, small intestine, and large

intestine at E16.5 to generate the iE7.5$^{\text{Early}}$-aE16.5 dataset (Figure 4A). Data was processed

similarly as in the previous experiment, and we recovered the same major cell types (Figure

4B; Supplementary Figure 7A-E). Notably, the earlier targeting increased the proportion of

ENS cells by 7.7-fold (Figure 4C), making ENS the most abundant cell population, and

corroborating greater lentiviral accessibility to neural crest primordia at the stage of injection.

Higher-resolution clustering further resolved subpopulations within immune and ENS

lineages, revealing myeloid (*Lyz2+*, *Mrc1+*, *Cd68+*) and lymphoid (*Trbc1/2+*, *Cd4+*)

subpopulations, lymphatic signatures (*Lyve1+*, *Pecam1+*), refined ENS neuron divisions, and

Schwann cell precursors (SCPs) (Figure 4D, Supplementary Figure 7F, G).

CloneIDs could be assigned to 77.8% of the 21,749 cells (Supplementary Figure 8A-

E), yielding 3,421 clones comprising 2,488 singletons, 394 single-cell-type clones, and 539

multi-cell-type clones (Figure 4E). The average and maximum size of multicellular clones

were larger in the E7.5^{Early}-aE16.5 dataset (Supplementary Figure 8F). Consistent with the

first experiment, mesenchymal populations comprised mainly of multi-cell-type clones,

whereas ENS, immune and epithelial populations were more lineage-restricted (Figure 4E).

Although most multicellular clones were composed of a single germ layer, a notable fraction

(~9%) of clones spanned more than one germ layer derivative (~1.5% spanned all three germ

layers; 14 clones) (Figure 4F, G). These cross-germ-layer clones suggest the presence of

pluripotent epiblast cells at E7.5^{Early}, in agreement with recent high-throughput spatial

transcriptome analyses of murine gastrula.[23]

The presence of cross-germ-layer clones in the iE7.5^{Early}-aE16.5 dataset complicated

lineage relationship analysis between refined cell types, but we reasoned that analysis using

germ-layer pure clones would still be relevant. Clonal coupling analysis of pure clones (851

clones) recapitulated patterns observed in the iE7.5-aE16.5 dataset (compare Figures 3F with

4H; Supplementary Figure 8G). We next restricted analysis to pure mesoderm clones (639

clones) (Figure 4I; Supplementary Figure 8H). As previously observed (Figure 3G, H), ICCs,

myofibroblasts and Fibroblast 4 (putative MIFs) featured stronger clonal coupling, as did

mesothelial cells and Fibroblast 3 (putative AFs) (Figure 4I). Although Fibroblast 1 (putative

SEMFs) was not adjacent to pericytes in the correlation heatmap, it exhibited the strongest

clonal pericyte coupling of all fibroblasts (Figure 4I; Supplementary Figure 8H). Within the

immune population, *Mrc1*+ myeloid and *Trbc1/2*+ lymphoid cells shared only few clones,

indicating early lineage specification prior to gut colonization (Figure 4J). The large ENS

dataset allowed detailed clonal analysis of ENS subclusters, revealing that many clones

consisted of multiple ENS states/types (Figure 4K). Notably, SCPs showed clonal coupling

with all ENS subtypes, however less so with the more mature neuron clusters ENC12 and

Branch B2 which may differentiate before SCP arrival to the gut (Supplementary Figure 8G).

Coupling was overall strong between Branch A and B neurons (Figure 4K; Supplementary

Figure 8G), suggesting no predetermination of either Branch in agreement with an earlier

lineage-tracing study[24].

Together, our results from the E7.5^{Early} injection highlights the importance of precise

embryonic staging for enriched targeting of specific lineage precursors. The optimal window

for ENS targeting appears to coincide with early E7.5 (~TS11b stage), when some epiblasts

cells are still present. This stage is therefore advantageous for efficient and comprehensive

ENS targeting, while posing challenges for lineage analysis.

Clonal sharing between gut segments reveals lineage-specific timing of regional identity

acquisition

Having resolved lineage relationships between and within gut cell types, we next examined

the regional dispersion of clones across the upper and lower GI tract. In the iE7.5-aE16.5

dataset, 21 clones linking the stomach and small intestine were identified. Of these, 18 clones

(85.7%) consisted exclusively of ENS or immune cells, accounting for 25.9% of ENS and

57.9% of immune clones (Figure 5A-C). These regional dispersions are consistent with the

extrinsic origins and migratory behavior of these lineages: the ENS arises from neural crest

precursor cells that migrate along the entire gut axis[9], while the first wave of embryonic

immune cells originates from the yolk sac and subsequently colonizes the gut[25, 26]. By contrast,

clones corresponding to other cell types were predominantly restricted to a single gut segment,

suggesting that most endoderm- and mesoderm-derived progenitors had acquired their

regional identities by E7.5 (~TS11c).

Considering single-germ-layer clones in the iE7.5^{Early}-aE16.5 dataset, we identified

123 region-spanning clones, again mainly from ENS and immune lineages, accounting for

60.7% of ENS and 66.7% of total immune clones (Figure 5D-F). Despite their abundance,

epithelial or fibroblast clones did not span regions, indicating that their progenitors are

already regionally patterned also at slightly earlier gastrula stages. We next examined the

ENS subtype composition within cross-regional and local clones and found them to be

similar. Notably, both Branch A and B neurons were present in local and cross-regional

clones, supporting the innate stochastic differentiation program in the ENS (Figure 5G). A

distinguishing feature of colonic ENS is its expression of *Pou3f3*[27], while it has not been

shown if expression is induced in vagally derived enteric *Pou3f3-* precursors as they enter the

large intestine. Our barcoding experiment allowed us to clonally link stomach☐resident ENS

(derived exclusively from the vagal neural crest) to *Pou3f3*☐ ENS cells in the colon. We

found that stomach/large intestine clones distributed randomly amongst ENS differentiation

stages and neuron identities (Figure 5H). Cells within large intestine-only clones and

stomach/large intestine clones showed similar expression frequency and level of *Pou3f3*

(Figure 5I). Together, these findings support that colonic ENS cells derived from vagal neural

crest acquires their colon-specific identity by responding to local cues.

Identification of cell type-specific regional gene expression differences

Beyond lineage relationships, our regionally resolved scRNA-seq datasets enabled analysis of

transcriptional similarities and differences across GI regions at E16.5. Integration of the two

datasets showed seamless merging (Supplementary Figure 9A, B), indicating consistent

targeting of comparable progenitor populations. Most cell types were detected across all

regions except for stomach-specific foregut basal epithelia, which expressed distinct marker

genes (Supplementary Figure 9C, D). Small intestine epithelium was instead enriched for

Cps1, whereas posterior Hox genes marked large intestine epithelium (Supplementary Figure

9D). Region-specific expression patterns were also observed in mesoderm-derived

populations: *Barx1* and *Nr2f1* were restricted to the stomach, while *Pitx1* was detected in both

stomach and large intestine (Supplementary Figure 9E). *Nkx2.3* and *Mab21l2* were confined

to intestinal mesenchyme. Within ectoderm-derived tissue (ENS), *Hoxb5*/*Hoxa4* were

enriched across intestinal regions, whereas *Hoxc4* was limited to the small intestine

(Supplementary Figure 9F).

Focused analysis of fibroblast subtypes revealed that Fibroblasts 5 and 6 were

enriched in the stomach compared with the intestines (Supplementary Figure 9G, H). Within

these populations, *Grm3* and *Sp5* showed stomach-specific expression (Supplementary Figure

9I)*,* whereas Fibroblast 6 uniquely expressed *Thy1* (Cd90), a marker associated with crypt

fibroblasts (Supplementary Figure 9I). *Wnt6*, *Cpne8* and *Ism1* in Fibroblast 6 further

supported a fibroblast type with high signaling capacity, consistent with a crypt/submucosal-

associated niche (Supplementary Figure 9I). These differences may reflect either a greater

abundance of Fibroblast 5 and 6 in the stomach or their earlier differentiation in anterior gut

regions.

In summary, our collective E16.5 scRNA-seq data uncovered regionally distributed

cell types, expression patterns and fibroblast states in the embryonic gut, which may

contribute to establishing the functional specialization of individual GI regions.

Cre-dependent lentivirus nano-injection enables cell type-specific targeting

Our data show that progenitors of essentially all gut cell types are transduced by *in utero*

lentiviral nano-injection at E7.5. For applications requiring cell type-specific targeting, this

approach therefore needs adaptation. Previous studies have demonstrated that mini-promoters

can confer cell type-specificity in the developing CNS[11]. However, no ENS-specific

promoters have yet been described. Instead, Cre-driver mouse lines have been used to label all

or subsets of ENS cell populations.

We previously showed that Baf53b-Cre mice specifically label neurons in the juvenile

ENS.[16] To investigate their utility during embryonic development, Baf53b-Cre; R26-

tdTomato embryos were analyzed at two stages (Supplementary Figure 10A, B). Reporter

expression was restricted to HuC/D+ neurons and absent from SOX10+ progenitor/glial cells.

We found that ~50% of HuC/D+ neurons expressed TOM at E14.5, increasing to ~90% at

E18.5. Baf53b-Cre activity is thus confined to differentiating neurons, with a slight delay

relative to HuC/D expression.

To achieve conditional targeting with *in utero* nano-injection, we designed a lentiviral

vector expressing an inverted double-floxed EGFP reporter, Lenti-DIO-EGFP (Figure 6A,

Supplementary Figure 1A). While AC injection of non-conditional LV-H2B-GFP resulted in

GFP labelling of HuC/D+ neurons, GFP was additionally observed in non-ENS cells as

previously discussed (Figure 6B; 1N). In contrast, AC injection of Lenti-DIO-EGFP into

Baf53b-Cre embryos resulted in GFP expression restricted to HuC/D+ neurons, labelling ~4-

6% at E18.5 (Figure 6B, C). In conclusion, *in utero* nano-injection of conditional lentiviral

vectors enables precise targeting of specific gut cell types, as demonstrated by Baf53b-Cre-

dependent lentiviral reporter expression in enteric neurons.

Timed overexpression of Ascl1 promotes neuronal differentiation of ENS progenitors

Having validated that lentiviral DIO-EGFP expression can be controlled by Cre, we next

sought to use this approach to drive timed gene expression in ENS progenitor cells, which are

more responsive to developmental cues. Sox10-CreERT2 mice mediate efficient, temporally

controlled recombination in ENS progenitors.[14, 15, 24] To assess the efficiency and kinetics of

Cre-dependent recombination, tamoxifen was administered to Sox10-CreERT2 females mated

with R26-EYFP males at E16.5, and embryos were harvested 26 hours later. Out of SOX10+

cells, ~50% expressed EYFP and ~11% of HuC/D+ neurons, indicating efficient labeling of

progenitors/glia and a subset of newly differentiated neurons within a short time window

(Supplementary Figure 10D, E).

As a proof-of-principle, we sought to overexpress *Ascl1*, a proneural transcription

factor essential for enteric neurogenesis, as demonstrated by us and others.[28, 29] We generated

a Cre-inducible *Lenti-DIO-EGFP-Ascl1* construct, in which *Ascl1* was linked to EGFP via a

self-cleaving P2A peptide (Figure 6D, Supplementary Figure 1B). Cre-dependent ASCL1

expression was validated using a HEK293FT-Cre cell line (Figure 6E). We next

overexpressed ASCL1 at late embryonic stages using our system, when both glio- and

neurogenesis are ongoing, to assess its ability to promote neuronal differentiation. Sox10-

CreERT2 embryos were injected at E7.5 with lentivirus carrying either DIO-EGFP as controls

or DIO-EGFP-Ascl1, followed by tamoxifen administration at E16.0 and analysis at E18.5

(Figure 6F). As compared to controls, ASCL1 overexpression reduced the EGFP+

progenitors/glia population (SOX10+ HuC/D-) from ~20% to ~5% (Figure 6G, H) and

increased EGFP+ neuronal population (HuC/D+) from ~80% to ~95% (Figure 6G, I).

Notably, ~25% of HuC/D+ EGFP+ cells in controls co-expressed SOX10, indicative of a

transitional population, which was reduced to <5% upon ASCL1 overexpression (Figure 6G,

J).

Together, these findings demonstrate that ASCL1 overexpression during a gliogenic

developmental window is sufficient to bias enteric progenitors toward neuronal

differentiation. More broadly, this approach establishes ultrasound-guided *in utero* delivery of

Cre-inducible lentiviral constructs as a powerful method for temporally controlled, cell type-

specific gene manipulation in developing embryos.

DISCUSSION

Our understanding of gut development and related diseases has advanced substantially

through recent technologies, including scRNA-seq and spatial transcriptomics.[16, 27, 30]

However, methods to define lineage relationships and interrogate gene function *in vivo* have

lagged behind, partly due to the inaccessibility of embryonic gut using traditional targeting

methods. Here, we present ultrasound-guided amniotic cavity nano-injection as an efficient

methodology to access major gut cell populations in mouse embryos, including epithelial and

mesenchymal populations, the ENS, as well as gut-extrinsic ganglia. Combined with high-

diversity lentiviral barcoding, this approach establishes a versatile lineage-tracing platform,

and coupled with Cre-dependent constructs, the approach supports temporally controlled, cell

type-specific gene manipulation in the developing gut.

Barcode-based lineage tracing combined with scRNA-seq overcomes key limitations

of traditional Cre-based lineage tracing approaches as they depend on targeting stem cells

defined by single marker genes with often equivocal specificities that risk confounding

interpretation. In contrast, the method presented here is unbiased and enables comprehensive

reconstruction of lineage relationships. It allowed us to resolve unprecedented details amongst

transcriptionally related mesenchymal populations refining their cellular identities, lineage

relationships and early development. First, we identified three fibroblast states with distinct

clonal relationships to specialized mesenchymal cell types. *Kcnn3+* fibroblasts - putative

contractile fibroblasts (MIFs) – were clonally related to ICCs and myoblast/smooth muscle

cells, suggesting shared progenitors. *Adamdec1+* fibroblasts (putative SEMFs) were linked to

pericytes, whereas *Adam12+* fibroblasts (putative AFs) were clonally related to mesothelial

cells. Notably, although previously suggested marker genes were used to annotate fibroblast

subtypes, no consensus nomenclature currently exists for gut fibroblasts. Clonal analysis

therefore provided important orthogonal support for subtype assignments and supported a

developmental logic whereby stromal gut compartments are assembled by a set of fate-biased

mesenchymal progenitor populations. Second, by analyzing clonal dispersion across gut

regions, we demonstrate that gut mesoderm is already regionally specified along the anterior-

posterior axis by E7.5, coinciding with persistent epiblast cells. Third, we present regional

diversity amongst mesenchymal cell types, likely reflecting early programs that shape region-

specific gut wall composition. Overall, these findings highlight how lineage information

complements transcriptomic and anatomical analyses to better define gut stroma development.

By integrating multiple gut regions, we found that region-spanning clones were

common in migratory lineages such as ENS and immune cells. Within the ENS, clonal

sharing across stomach and colon supports a model in which vagal neural crest-derived

progenitors acquire region-specific features, such as colonic *Pou3f3* expression, in response to

local cues. In contrast, the restriction of epithelial and mesenchymal clones to single regions

indicates that regional identity of these compartments is established incredibly early. Together

with the region-specific transcriptional programs observed in mesenchymal subtypes, these

findings suggest that developmental patterning of the gut wall emerges from a combination of

early regional allocation and local differentiation programs.

Beyond lineage reconstruction, *in utero* nano-injection enables conditional gene

manipulation in defined ENS populations. Using Baf53b-Cre and Sox10-CreERT2 mice, we

show that Cre-dependent lentiviral cargo can be activated in differentiating enteric neurons or

temporally controlled progenitor/glial populations. Proof-of-principle experiments with *Ascl1*

demonstrated that the system can be used to test gene function *in vivo* during defined

developmental windows. This approach complements emerging fetal atlases and disease

genetics studies, which increasingly identify candidate regulators in transient embryonic cell

states but often lack efficient *vivo* validation strategies. We believe that the ability to combine

clonal analysis with targeted perturbation can help shift descriptive cell atlas data to

mechanistic studies of gut development.

Several limitations should also be considered. Our results highlight a strong temporal

component of lineage accessibility and commitment. Targeting at E7.5$^{\text{Early}}$ enhanced labeling

of ENS precursors but also increased cross-germ-layer clones, consistent with presence of

epiblast cells. While residual epiblast labeling complicates clonal interpretation, it also

provides a functional readout of the developmental window during which germ-layer

restriction is being consolidated *in vivo*. Precise embryonic staging is thus essential, and inter-

embryo variability can complicate stage estimation, although ultrasound imaging partially

mitigates this issue. Tissue dissociation, single-cell capture, and barcode recovery also

introduce dropouts that will inevitably reduce clonal reconstruction efficiency. In the gene

perturbation setting, conditional targeting efficiencies in the ENS were relatively modest.

However, we believe that this mosaicism may in fact be advantageous for studying cell-

intrinsic mechanisms given that ENS development depends on coordinated proliferation and

long-range migration. Broad perturbation within the ENS may cause large secondary effects

at the population level, whereas mosaic manipulation may reveal how genetically altered cells

behave within a largely normal developing ENS.

Finally, this work has implications for understanding the developmental origins of GI

disease. Gut organogenesis relies on interdependent differentiation of diverse cell populations

from all three germ layers, and subtle deviations in lineage allocation, maturation, or regional

patterning may have lasting functional consequences.[31] This is relevant not only to congenital

disorders such as Hirschsprung disease, achalasia, and CIPO, but also to more common

conditions in which early developmental alterations may predispose to later dysfunction.

Some of these disorders may result from *de novo* mutations, in which early mutations occur in

a mosaic fashion[32]. It is conceivable that the impact of such mutations would be more

predictable with a clearer understanding of refined cell lineages arising in post-gastrula stages.

We propose that this platform will provide a scalable strategy for mechanistic testing of

candidate disease regulators, and modeling early cellular events that may contribute to GI

disorders and enteric neuropathologies.

ACKNOWLEDGEMENTS

We thank Jonas Frisén and Michael Ratz for designing and sharing the tdTomato 30N

barcoded vector. We also thank Jonas Frisén for access to flow cytometry at Karolinska

Institutet. We thank Fernando Lopez-Redondo for laboratory assistance. The authors

acknowledge support from the National Genomics Infrastructure in Stockholm funded by

Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish

Research Council. The authors acknowledge the computational resources provided by the

National Academic Infrastructure for Supercomputing in Sweden (NAISS), partially funded

by the Swedish Research Council through grant agreement no. 2022-06725 and the Centre for

Bioinformatics and Biostatistics (CBB) at Karolinska Institutet funded by CIMED

Infrastructure and Karolinska Institutet.

U.M. was supported by the Swedish Research Council (Vetenskapsrådet; 2020-01129

and 2022-01570), the European Research Council (divENSify; 101045026), the Knut and

Alice Wallenberg Foundation (KAW; 2020.0109), the Strategic Research Area in

Neuroscience, the Strategic Research Area in Stem Cell Research and Regenerative Medicine,

Karolinska Institutet Consolidator Grant, and the Brain Foundation (Hjärnfonden; FO2023-

0169; FO2023-0130). E.R.A was supported by the Knut and Alice Wallenberg Foundation

(KAW; 2020.0109), Karolinska Institutet Consolidator Grant (2-195/2021), the Strategic

Research Area in Neuroscience, Hjärnfonden (FO2025-0271), Vetenskapsrådet 3R grants

(2024-03113, 2021-03537, 2017-01054), and the European Research Council (ERG-CoG

LIMITLESS; 101171156). K.P was supported by Marie Sklodowska Curie Award

(1011544005; DeSTENy). K.H, J.H and Z.L were supported by a Karolinska Institutet Grant

for Salary (2-2110;2019-7;2019-3). The imaging was performed at the Biomedicum Imaging

Core with support from the Karolinska Institutet. We thank the Viral Vector Facility of the

Neuroscience Center Zurich for excellent service and production of AAV vectors. We thank

animal staff from the Department of Comparative Medicine at Karolinska Institutet for help

with animals.

Abbreviations: AF, adventitial fibroblasts; CIPO, chronic intestinal pseudo-obstruction;

CNS, central nervous system; DGBIs, disorders of gut-brain interaction; DIO, double-floxed

inverted open reading frame; DRG, dorsal root ganglia; E, embryonic; EGFP, enhanced green

fluorescent protein; ENS, enteric nervous system; EYFP, enhanced yellow fluorescent

protein; GI, gastrointestinal; H2B, histone 2B; IBD, inflammatory bowel disease; LV,

lentiviral; MIFs, muscle-embedded interstitial fibroblasts; NCC, neural crest cells;

NEPTUNE, neural plate targeting with in utero nano-injection; scRNA-seq, single cell RNA

sequencing; SCP, Schwann cell precursor; SEMF, subepithelial myofibroblasts; TREX,

tracking and gene expression profiling of clonally related cells; TS, Theiler stage

FIGURE LEGENDS

Figure 1. Efficient targeting of neural crest/placode populations destined to innervate

the gut by *in utero* transduction at E7.5. (A) Schematic drawing of experimental setup for

ultrasound-guided *in utero* nano-injection. Left-most panel reproduced from Mangold et al

2021[11], with permission. The right panel was created with BioRender.com. (B) Uniform

manifold approximation and projection (UMAP) plot of cells recovered from E9.5 and E10.5

embryos following E7.5 amniotic cavity injection.[12] (C) UMAP plot of reprocessed neural

crest/placode cells (green populations in B), revealing 14 clusters. (D) Schematic drawing of

neural crest/placode-derived ganglia innervating the gut. (E) Dot plot showing expression of

canonical marker genes for neural crest/placode populations across the clusters. Dot size

represents the fraction of cells expressing the gene and color intensity represents mean

expression. (F) Feature plots showing expression of neuronal (*Tubb3+*) and migratory neural

crest/placode (*Sox10+*) markers. (G) Feature plots showing gene expression associated with

nodose ganglia, sympathetic ganglia, DRG and the ENS. (J) Representative

immunofluorescence staining of an E13.5 DRG showing H2B-GFP expression in both

SOX2+ neural progenitors (arrowheads) and NeuN+ postmitotic neurons (asterisks). (I, J) Bar

graphs showing proportions of GFP positive NeuN+ neurons and SOX2+ neural progenitors

along the thoracic-sacral axis (n=3 mice). Bars represent the mean ± standard deviation (SD).

(K) Representative immunofluorescence image of E18.5 small intestine showing GFP

expression in HuC/D+ neurons (asterisks) and SOX10+ ENS progenitors and glia

(arrowheads). (L, M) Bar graphs showing proportion of GFP positive HuC/D+ neurons and

SOX10+ progenitors/glia across gut regions (n=4 mice). Bars represent mean ± SD. Each dot

represents one animal. (N) Representative immunofluorescence image of E18.5 gut showing

GFP expression in alpha-SMA+ smooth muscle (arrowheads) and EpCAM+ epithelium

(asterisks). Scale bars: 50µm.

Figure 2. *In utero* transduction at E7. 5 targets diverse cell types in the developing gut.

(A) Schematic of injection of lentiviral barcode library into the E7.5 amniotic cavity. (B)

Experimental workflow to generate the iE7.5-aE16.5 dataset consisting of TOM+ cells, flow

sorted from two gut regions. (C) UMAP plot of high-quality gut cells colored by tissue of

origin. Pie chart indicates the proportion of cells from each region. (D) UMAP plot showing 8

major clusters, annotated based on established marker gene expression. (E) Dot plot

displaying expression of canonical marker genes used to define major clusters. (F) Feature

plots displaying marker gene expression of epithelial cell types. (G) Reprocessed ENS cells

revealing four clusters. (H) Feature plots showing expression of selected markers for ENS

states and branches. (I) UMAP plot showing subclusters within fibroblast-like and smooth

muscle-like populations. (J) Dot plot showing marker gene expression distinguishing

fibroblast-like and smooth muscle-like subclusters. (K) Schematic drawing indicating the

diverse gut cell types targeted by *in utero* nano-injection, showing that all three germ layers

are efficiently transduced. Part of 2A, B and K were created with BioRender.com. (E, J) Dot

size represents the percentage of cells expressing the gene, and color indicates relative mean

expression.

Figure 3. Lentiviral barcoding at E7.5 resolves clonal lineage relations in the developing

gut. (A) UMAP plot highlighting proportions of cells with CloneIDs in the iE7.5-aE16.5

dataset. (B) UMAP plot showing 11 major gut cell types after merging subclusters. (C)

Stacked bar plot showing distribution of clone categories across major cell types. Schematic

drawing describing clonal types (right). (D) UpSet plot showing frequencies of clones with

different composition of major cell types. (E) UpSet plot showing minimal clone sharing

across germ layers. Inset chord diagrams in (D, E) illustrate the extent and direction of clonal

sharing, with ribbon width proportional to the number of shared clones. (F, G) Heatmaps

showing the correlation matrix of clonal coupling z-scores, computed as deviations of

observed clone sharing relative to randomized data, for refined gut cell types (F) and

restricted to mesoderm-derived cell types (G). Colored boxes denote germ-layer origin (pink:

endoderm; purple: ectoderm; yellow: mesoderm), or lineage-coupled mesenchymal

populations (green: Fibroblast 4 with myofibroblasts, ICC and smooth muscle cells; black:

Fibroblast 3 with mesothelial cells; brown: Fibroblast 1 with pericytes). (G) Schematic

summary of lineage relationships amongst mesenchymal cell types inferred from the clonal

coupling analysis. H and left panel of 3C were created with Biorender.com.

Figure 4. *In utero* transduction at early E7.5 improves ENS targeting but uncovers presence of epiblast cells. (A) Schematic diagram showing the generation of the iE7.5^{Early}-aE16.5 dataset consisting of three gut regions. Created with BioRender.com. (B) UMAP plot showing major cell types. (C) Alluvial plot showing proportions of major cell types in the iE7.5-aE16.5 and iE7.5^{Early}-aE16.5 datasets. (D) UMAP plot showing refined gut cell type annotations in the iE7.5^{Early}-aE16.5 dataset. (E) Stacked bar plot showing distribution of clone categories across major cell types identified in the iE7.5^{Early}-aE16.5 dataset. (F) UpSet plot showing frequencies of clones with different composition of major cell types. (G) UpSet plot showing notable clone sharing across germ layers. Inset chord diagrams in (F, G) illustrate the extent and direction of clonal sharing, with ribbon width proportional to the number of shared clones. (H, I) Heatmaps showing the correlation matrix of clonal coupling z-scores, computed as deviations of observed clone sharing relative to randomized data, for refined gut cell types (H) and restricted to mesoderm-derived cell types (I). Colored boxes denote germ-layer origin (pink: endoderm; purple: ectoderm; yellow: mesoderm), or lineage-coupled mesenchymal populations (green: Fibroblast 4 with myofibroblasts, ICC and smooth muscle cells; black: Fibroblast 3 with mesothelial cells; brown: Fibroblast 1 with pericytes). (J) Chord diagram showing self- and cross-linking between Mrc1+ myeloid and Trbc1/2+ lymphoid cells. (K) UpSet plot showing co-occurrence of ENS cell states/types in the same clone.

Figure 5. Regional clone sharing reveals long-range ENS migration and early regional patterning of mesenchymal populations along the GI tract. (A) UpSet plot showing co-occurrence of major cell types in cross-region clones in the iE7.5-aE16.5 dataset. (B) UMAP plot highlighting cells in cross-region and local clones. (C) Alluvial plot illustrating relationship between tissue region, cell type, and CloneID of the cross-region clones in (A).

(D) UpSet plot showing co-occurrence of major cell types in the cross-region clones in the

iE7.5^{Early}-aE16.5 dataset (restricted to single-germ-layer clones). (E) UMAP plot highlighting

cells in cross-region and local clones. (F) Heatmap showing the distribution of clones across

gut regions in the iE7.5-aE16.5 dataset, with binary indication of presence or absence.

Colored bars denote cell type identities of clones. (G) Heatmap showing clonal composition

of ENS cell states/types for cross-region or local clones, revealing no apparent bias. (H)

UMAP plot highlighting the differentiation state of clones with different dispersion. (I) Violin

plot showing comparable levels of *Pou3f3* expression in colonic ENS cells that are local or

share clones with small intestine and/or stomach. Parts of 5A and D were created with

Biorender.com.

Figure 6. Temporal overexpression of Ascl1 in Sox10+ cells promotes neuronal

differentiation (A) Schematic overview of the Lenti-DIO-EGFP vector, before and after Cre-

mediated gene inversion. (B) Representative immunohistochemistry images of intestine from

wild-type mice injected with LV-H2B-GFP showing GFP expression in both HuC/D- cells

(asterisks) and HuC/D+ (arrowheads), and Baf53b-Cre mice injected with Lenti-DIO-EGFP

showing GFP expression restricted to HuC/D+ cells (arrowheads). (C) Bar graph showing the

percentage of HuC/D+ cells expressing GFP across gut regions in Baf53b-Cre embryos

injected with Lenti-DIO-EGFP. (n=5 mice). Bars represent mean ± SD. (D) Schematic

drawing of the Lenti-DIO-EGFP-Ascl1 construct with and without Cre-mediated gene

inversion. (E) Representative immuohistochemistry images showing validation of transgene

expression in Cre-expressing HEK293FT cells. (F) Schematic representation of experimental

procedure for *in vivo* overexpression of EGFP-Ascl1 or EGFP in ENS progenitors. Part was

created with BioRender.com. (G) Representative confocal images of E18.5 small intestine

transduced with DIO-EGFP or DIO-Ascl1-EGFP vectors showing SOX10 and HuC/D

expression patterns in GFP+ cells. asterisks denote EGFP+ cells and texts indicate their status

of expression, with + or – signs denoting positive or negative expression. (H-J) Bar graphs

showing the proportions of GFP+ cells with SOX10 expression without HuC/D (H), GFP+

cells with HuC/D expression (I), and GFP+ HuC/D+ cells expressing SOX10 (J), in Baf53b-

Cre embryos injected with either DIO-Ascl1-EGFP or Lenti-DIO-EGFP. (H)*p=0.0136;

(I)**p= 0.0086, and (J)* p= 0.011. n=6-7 mice per condition. SI: small intestine; AC:

amniotic cavity; WT: wild-type. Scale bars: 20µm.

REFERENCES:

1. Knowles CH, Lindberg G, Panza E, et al. New perspectives in the diagnosis and
management of enteric neuropathies. Nat Rev Gastroenterol Hepatol 2013;10:206–18.

2. Niesler B, Kuerten S, Demir IE, et al. Disorders of the enteric nervous system - a
holistic view. Nat Rev Gastroenterol Hepatol 2021;18:393–410.

3. Sherman MP, Zaghouani H, Niklas V. Gut microbiota, the immune system, and diet
influence the neonatal gut-brain axis. Pediatr Res 2015;77:127–35.

4. Chalkidi N, Paraskeva C, Koliaraki V. Fibroblasts in intestinal homeostasis, damage,
and repair. Front Immunol 2022;13:924866.

5. Hung LY, Margolis KG. Autism spectrum disorders and the gastrointestinal tract:
insights into mechanisms and clinical relevance. Nat Rev Gastroenterol Hepatol
2024;21:142–163.

6.   Dixit R, Lu F, Cantrup R, et al. Efficient gene delivery into multiple CNS territories
using in utero electroporation. J Vis Exp 2011.

7.   Andersson E, Tryggvason U, Deng Q, et al. Identification of intrinsic determinants of
midbrain dopamine neurons. Cell 2006;124:393–405.

8.   Dvoretskova E, Ho MC, Kittke V, et al. Spatial enhancer activation influences
inhibitory neuron identity during mouse embryonic development. Nat Neurosci
2024;27:862–872.

9.   Uesaka T, Young HM, Pachnis V, et al. Development of the intrinsic and extrinsic
innervation of the gut. Dev Biol 2016;417:158–67.

10.   Mangold K, He J, Stokman S, et al. Murine Neural Plate Targeting by In Utero Nano-
Injection (NEPTUNE) at Embryonic Day 7.5. J Vis Exp 2022.

11.   Mangold K, Masek J, He J, et al. Highly efficient manipulation of nervous system
gene expression with NEPTUNE. Cell Rep Methods 2021;1.

12.   de Haan S, He J, Corbat AA, et al. Ectoderm barcoding reveals neural and cochlear
compartmentalization. Science 2025;388:60–68.

13.   Ratz M, von Berlin L, Larsson L, et al. Clonal relations in the mouse brain revealed by
single-cell and spatial transcriptomics. Nat Neurosci 2022;25:285–294.

14.   Laranjeira C, Sandgren K, Kessaris N, et al. Glial cells in the mouse enteric nervous
system can undergo neurogenesis in response to injury. J Clin Invest 2011;121:3412–
24.

15.   Memic F, Knoflach V, Morarach K, et al. Transcription and Signaling Regulators in
Developing Neuronal Subtypes of Mouse and Human Enteric Nervous System.
Gastroenterology 2018;154:624–636.

16.   Morarach K, Mikhailova A, Knoflach V, et al. Diversification of molecularly defined myenteric neuron classes revealed by single-cell RNA sequencing. Nat Neurosci 2021;24:34–46.

17.   Masamsetti VP, Salehin N, Kim HJ, et al. Lineage contribution of the mesendoderm progenitors in the gastrulating mouse embryo. Dev Cell 2025;60:1991–2006 e9.

18.   Torihashi S, Nishi K, Tokutomi Y, et al. Blockade of kit signaling induces transdifferentiation of interstitial cells of cajal to a smooth muscle phenotype. Gastroenterology 1999;117:140–8.

19.   Baker SA, Hennig GW, Salter AK, et al. Distribution and Ca(2+) signalling of fibroblast-like (PDGFR(+)) cells in the murine gastric fundus. J Physiol 2013;591:6193–208.

20.   Schneider S, Hashmi SK, Thrasher AJ, et al. Single Nucleus Sequencing of Human Colon Myenteric Plexus-Associated Visceral Smooth Muscle Cells, Platelet Derived Growth Factor Receptor Alpha Cells, and Interstitial Cells of Cajal. Gastro Hep Adv 2023;2:380–394.

21.   Viitala E.W SE, Miihkinen M, Chalkidi N, Stavropoulou A, Lemmetyinen TT, Päivinen P, Kaptrio T, Hagström J, Haglund C, Mäkelä TP, Aittokallio T, Katajisto P, Koliaraki V, Ollila S. High-resolution integrative analysis allows characterization and spatial annotation of normal and cancer-associated colon fibroblasts. bioRxiv, 2025.

22.   Rinkevich Y, Mori T, Sahoo D, et al. Identification and prospective isolation of a mesothelial precursor lineage giving rise to smooth muscle cells and fibroblasts for mammalian internal organs, and their vasculature. Nat Cell Biol 2012;14:1251–60.

23.   Harland LTG, Lohoff T, Koulena N, et al. A spatiotemporal atlas of mouse gastrulation and early organogenesis to explore axial patterning and project in vitro models onto in vivo space. Cell Rep 2025;44:116047.

24.   Lasrado R, Boesmans W, Kleinjung J, et al. Lineage-dependent spatial and functional
organization of the mammalian enteric nervous system. Science 2017;356:722–726.

25.   Bain CC, Schridde A. Origin, Differentiation, and Function of Intestinal Macrophages.
Front Immunol 2018;9:2733.

26.   Neo WH, Lie ALM, Fadlullah MZH, et al. Contributions of Embryonic HSC-
Independent Hematopoiesis to Organogenesis and the Adult Hematopoietic System.
Front Cell Dev Biol 2021;9:631699.

27.   Wright CM, Schneider S, Smith-Edwards KM, et al. scRNA-Seq Reveals New Enteric
Nervous System Roles for GDNF, NRTN, and TBX3. Cell Mol Gastroenterol Hepatol
2021;11:1548–1592 e1.

28.   Blaugrund E, Pham TD, Tennyson VM, et al. Distinct subpopulations of enteric
neuronal progenitors defined by time of development, sympathoadrenal lineage
markers and Mash-1-dependence. Development 1996;122:309–20.

29.   Memic F, Knoflach V, Sadler R, et al. Ascl1 Is Required for the Development of
Specific Neuronal Subtypes in the Enteric Nervous System. J Neurosci 2016;36:4339–
50.

30.   Zilbauer M, James KR, Kaur M, et al. A Roadmap for the Human Gut Cell Atlas. Nat
Rev Gastroenterol Hepatol 2023;20:597–614.

31.   Zhang Z, Lee D, Liu L, et al. Impairment of stromal-epithelial regenerative cross-talk
in Hirschsprung disease primes for the progression to enterocolitis. Sci Transl Med
2025;17:eadp4679.

32.   Mohiuddin M, Kooy RF, Pearson CE. De novo mutations, genetic mosaicism and
human disease. Front Genet 2022;13:983668.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6