Tissue-specific adaptation of human T cells is preserved during tissue

inflammation

Nicole B. Potchen[1], Hugh R. MacMillan[1], Eva Domenjo-Vila[1], Andrew J. Konecny[1,2], Alexis K.

Taber[1], Caitlin S. DeJong[1], Geervani Daggupati[1,2], Raj Shree[3], Stephen A. McCartney[3], Shelton

W. Wright[4], Evan W. Newell[1,6], Douglas R. Dixon[5], Martin Prlic[1,6*]

[1] Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center, Seattle, WA, USA.

[2] Department of Immunology, University of Washington, Seattle, WA, USA.

[3] Division of Maternal Fetal Medicine, Department of Obstetrics and Gynecology, University of Washington, Seattle,

WA, USA.

[4] Division of Pediatric Critical Care Medicine, Department of Pediatrics, University of Washington, Seattle, WA, USA.

[5] Department of Bioscience Research and Periodontics, University of Tennessee Health Science Center, College of

Dentistry, Memphis, TN, USA.

[6] Department of Global Health, University of Washington, Seattle, WA, USA.

* Corresponding author

Fred Hutchinson Cancer Center, Vaccine and Infectious Disease Division, Seattle, WA, USA.

mprlic@fredhutch.org

Summary

T cells play an essential role in protecting tissues against pathogens and regulating tissue homeostasis. Previous studies highlight that T cells display tissue-specific phenotypic and functional properties, suggesting that T cells adapt to their local environment. However, whether this holds true in inflamed tissues or whether inflammation disrupts any tissue-specific T cell adaptations remains poorly understood. To address this open question, we examined the T cell compartment – including conventional CD4 and CD8 T cells, regulatory T cells, $\gamma\delta$ T cells, and MAIT cells – from healthy and inflamed human mucosal tissues. Using high-parameter spectral flow cytometry, we examined phenotype *ex vivo* and the functional capacity following stimulation, utilizing conventional gating and unsupervised clustering analysis approaches. Overall, we analyzed 65 tissue samples including mild, moderate, and severely inflamed oral gingiva, healthy and inflamed lung, along with healthy and inflamed tissue from the decidual-placental interface. Across these mucosal barrier tissues, we find that tissue location plays a dominant role in shaping the composition, phenotype, and functional capacity of the T cell compartment. Importantly, these tissue-specific adaptations were largely maintained during states of tissue inflammation. This included the ability to exert tissue repair functions, which was preserved across T cell subsets, even in severely inflamed tissues.

Introduction

Human T cells have distinct phenotypic and functional properties across different tissues and organs. This was initially demonstrated by Farber and colleagues who examined the phenotype and functional properties of conventional CD4 and CD8 T cells isolated from secondary lymphoid tissues, lung, jejunum, ileum and colon of trauma-induced, deceased organ donors[1]. This study furthermore established CD69 expression as biomarkers for tissue-residence for human CD4 T cells, and CD69 and CD103 co-expression to indicate tissue-resident CD8 T cells[1]. Follow-up studies highlighted that expression of CD69 by human CD4 and CD8 T cells is associated with a core transcriptional signature akin to bona fide murine tissue-resident T cells[2]. Human tissue-resident memory CD8 T cells (T_{RM}) show tissue-specific adaptation such as distinct cytotoxic properties across skin, lung, and jejunum, and are also clonally distinct across these tissue sites[3-7]. A recent study indicated that within the intestine, distinct tissue niches drive unique phenotypic and functional properties in T cells, thus providing further compelling evidence that T_{RM} tissue location and functional state are fundamentally linked[8].

Similarly, conventional CD4 T cells (CD4 Tconv) and regulatory T cells (CD4 Treg) also have distinct phenotypic and functional properties across tissues[9-12]. Further evidence points to innate-like T cells also adapting in a tissue-specific manner. Human $\gamma\delta$ T cells differ by tissue location with cells from adult lung, spleen, lymph nodes (LNs), and gut tissue clustering differentially by tissue based on their expression of transcripts associated with effector and repair functions[13]. Human mucosal-associated invariant T (MAIT) cells in tissues have distinct T cell receptor (TCR) repertoires, cytokine expression profiles, and display a more activated phenotype compared to counterparts in the blood[14,15].

Whether tissue-specific adaptations are maintained during tissue inflammation is unclear. Inflammation involves coordinated immune cell recruitment and activation and thus has the potential to profoundly alter the T cell compartment within tissue[16]. Memory phenotype CD8 T

cells can be activated by pro-inflammatory signals alone; this event has been coined bystander-

activation or inflammation-driven activation of T cells[17,18]. Inflammation-driven activation of

memory CD8 T cells leads to increased expression of CD69, PD-1, and TOX, along with

expression of IFNγ and Granzyme B (GzmB)[19-23] which can have protective properties[24,25] but can

also lead to tissue pathologies[26]. Such bystander-activated CD8 T cell-mediated pathology has

been demonstrated in a human cohort following infection with Hepatitis A virus (HAV)[27]. For CD4

T cells, inflammation and associated tissue pathology is often associated with expression of IL-

17, but it is noteworthy that IL-17 should not be simply viewed as a pro-inflammatory cytokine[28]

or indicator of excessive inflammation[29]. Importantly, in addition to mediating tissue protection and

pathology, T cells also play a role in restoring tissue homeostasis via secretion of IL-22 and the

epidermal growth factor receptor (EGFR)-ligand amphiregulin (AREG)[30-32]. Tregs express AREG

in response to IL-18 and IL-33[33], while a recent study demonstrates that human memory CD8 T

cells express AREG in response to TCR stimulation[34].

While there has been substantial progress in our understanding of the T cell compartment

and its tissue-specific functional properties in healthy human tissues, the extent to which

inflammation alters these functional properties remains an open question. We sought to address

if inflammation overrides any tissue-specific T cell adaption in human tissues. Specifically, we

aimed to determine if inflammation alters the composition of the T cell compartment and

associated functional properties, and if it affects the ability of T cells to maintain their tissue-

regenerative functions. To accomplish this, we collected healthy and inflamed tissues from 3

distinct mucosal barrier sites: oral mucosa (gingiva), lung, and the decidual-placental interface

(DPI). We developed a high-parameter flow cytometry panel containing 37 markers to define the

composition, phenotype, and functional capacity of the T cell compartment: CD4 Tconv, CD4

Treg, CD8 CD103+ T_{RM} and CD8 CD103- circulating T cells, along with the innate-like MAIT and

γδ T cell subsets. Overall, we found that the composition, phenotype and function of the T cell

compartment was driven more by tissue location than inflammation, indicating that T cells largely

maintain their tissue-specific adaptations during inflammation. Our data emphasize the need to

assess and consider inflammation in a tissue-specific manner. Furthermore, our data highlight

that T cells maintain their tissue-regenerative capacity even in severely inflamed tissues, thus

pinpointing T cells as a therapeutic target to treat inflammatory diseases.

Results

The inflammation state in the oral mucosa has a moderate effect on composition and phenotype of the T cell compartment

To determine the impact of inflammation on the mucosal T cell compartment, we first examined phenotypic differences across mildly, moderately, and severely inflamed gingival tissues. These tissues are discarded from routine surgeries and represent a wide range of inflammatory states. We assigned clinically healthy gingival tissue and tissue with mild gingivitis as "mild inflammation" since even clinically healthy tissue contains a baseline level of inflammation, which may allow for constant immune surveillance[35]. Next, we assigned tissues with pronounced gingivitis transitioning to periodontitis as "moderate inflammation" and those with periodontitis as "severe inflammation". While gingivitis is reversible with proper dental hygiene, periodontitis is an irreversible, chronic disease state without clinical intervention[36]. A full sample list with clinical information can be found in Supplemental Table 1. A 37-color spectral flow cytometry panel was utilized to examine phenotype (no stimulation) and functional capacity (following stimulation) of the T cells isolated from gingival tissues (**Figure 1A**, *panel and gating; Supplemental Figure 1-2*). As a technical control, a PBMC sample from a single leukapheresis donor was included in every flow cytometry experiment (*Supplemental Figure 3*).

We first examined the frequency of T cells (CD3+ cells) within the CD45+ cell compartment and found no changes between mild, moderate, and severely inflamed gingival tissue (**Figure 1B**). Of note, a small population of naïve T cells (CCR7+ CD45RA+) was present in each tissue sample. These naïve T cells presumably stem from the blood that is inherently present in tissues. We excluded these naïve T cells from our manual gating analysis approach. We next assessed if T cell subset composition is altered with inflammation. We gated on T cells by first defining innate-like T cell subsets within the CD3+ population, followed by subsetting the remaining CD4 and CD8 T cells into conventional CD4 T cells (CD4 Tconv), regulatory CD4 T cells (Treg), CD103+

(tissue-resident phenotype) CD8 T cells and CD103- (circulating) CD8 T cells (*Supplemental*

Figure 1B). We observed stable frequencies of most T cell subsets, including CD4 Tconv, CD4

Treg, and CD103+ CD8 T cells (**Figure 1C**). We noticed a slight reduction in CD103+ CD8 T cells

within the total T cell population (p = 0.1026), but a significant increase in circulating CD103- CD8

T cells in severely inflamed tissue compared to mild inflammation (**Figure 1D**).

We next examined if the expression of activation, co-stimulation, and exhaustion markers

change with inflammation severity to look for evidence of inflammation-driven (bystander)

activation. To visualize similarities and differences in expression of multiple proteins across

different tissues, we chose a heatmap-based approach to displaying expression frequency. For

each protein, the mean expression frequency is shown on all heatmaps, and the median and data

range for each protein are shown in *Supplemental Table 2.* Within conventional CD4 and CD8

subsets, we observed only few changes between mild, moderate, and severely inflamed gingiva

(**Figure 1E**). Pro-inflammatory cytokines can activate memory T cells and increase the expression

of CD69, PD-1, and induce expression of GzmB[20], but we did not observe an inflammation-

associated increase of these proteins (**Figure 1E**). On the contrary, we noted reduced expression

of CD69 in both the CD8 CD103+ and CD103- populations in severely inflamed compared to

mildly inflamed tissues. In the CD4 Tconv population, there were no differences in these

phenotypic markers. In general, we observed relative phenotypic stability across inflammation

status within each T cell subset (*Supplemental Figure 4*).

Next, we examined innate-like T cell subsets including MAIT and $\gamma\delta$ T cells, which can

also be activated by pro-inflammatory stimuli[37,38]. While there were no statistically significant

differences in MAIT cell frequencies and phenotypic markers across inflammation status, we

found a significant reduction in $\gamma\delta$ T cells in severely inflamed compared to moderately inflamed

tissue (**Figure 1F**). Within the moderately inflamed condition, we also observed a pronounced

reduction of PD-1 expression in the $\gamma\delta$ T cell population (**Figure 1G**). The concomitant increase

in the abundance of γδ T cells (as a % of CD3+ T cells) and decreased PD-1 expression could

potentially be due to recruitment of γδ T cells from the blood to moderately inflamed tissue. PD-1

expression frequency was also significantly higher within CD8 CD103- circulating cells during

severe inflammation (**Figure 1E**). While we observed some phenotypic differences in our dataset

as tissue became increasingly inflamed, we noted that the T cell profile within all oral mucosal

tissues remained rather stable even in severely inflamed tissues. Although we did not observe

phenotypic changes indicative of inflammation-driven (bystander) activation of one or more T cell

subset in inflamed tissues, we considered that a smaller proportion of T cells may respond in a

TCR-dependent manner.

Biomarkers of TCR-mediated activation do not significantly change across inflammation

states in the oral mucosa

We next assessed biomarker pairs indicative of TCR-mediated activation[39-41]. We

examined co-expression of CD39+ PD-1+ (for CD8+ cells[41]) as well as ICOS+ PD-1+ (for CD4+

cells[39]) across mild, moderate, and severely inflamed oral mucosal tissues. Across gingiva of

varying inflammation states, there were no differences in frequency of expression of each of these

populations. CD39, PD-1 and ICOS are also expressed on highly activated Tregs, and we

consistently observed the highest frequency of expression in CD4 Tregs (**Figure 2A-B**). This

suggests that as inflammation within the oral mucosa progresses from mild to severe, there is no

evidence for an increase in antigen-mediated responses by T cells. In line with this observation,

no changes in Ki67 expression, an indicator of recent cell proliferation, were observed across

inflammation status (**Figure 2C-D**). Lastly, we examined TIM3 and PD-1 expression, a biomarker

pair indicative of exhaustion, within each T cell subset. We observed unaltered co-expression of

these markers regardless of inflammation state, indicating that the lack of observed changes is

not due to T cell exhaustion (**Figures 2C-D**). Together, these data indicate that biomarkers of T

cell activation and proliferation remained stable across a range of inflammatory disease states

within the gingiva. Overall, we did not detect evidence of inflammation- or TCR-driven activation

of the T cell compartment in the inflamed gingiva. We considered that this could be the result of

a state of T cell dysfunction and thus interrogated T cell functional properties.

Functional capacity for T cell-mediated tissue healing and cytokine production is

maintained in inflamed oral mucosa

To examine how the functional capacity of T cell subsets varies across inflammation, we

stimulated T cells isolated from gingival tissue for 6 hours with PMA/ionomycin (*Figure 1A*). We

define the functional capacity simply as the ability of a T cell to express protein following this brief,

yet broad, activation stimulus. Within CD4 and CD8 subsets, we observed no statistically

significant differences in AREG, IFNγ, TNFα, IL-17A, IL-17F, IL-2, or IL-22 (*Figure 3A,*

Supplemental Figure 5). Overall, there were no changes in the functional capacity of T cell subsets

across mildly, moderately, and severely inflamed gingiva (*Figure 3B*). This was also consistent

within innate like subsets (*Figure 3C*). The mean, median, and data range are reported in

Supplemental Table 2. Within several T cell subsets, we found co-expression of GzmB and AREG

upon stimulation (*Figure 3D*), which may indicate dual functional roles (cytotoxicity and tissue

repair) can occur simultaneously within the same T cell. The highest frequency of GzmB and

AREG co-expression was observed in CD8 CD103+ cells (*Figure 3D*), though these were also

detected to a lesser extent in CD4 Tconv, CD4 Tregs, and innate-like subsets (~1% within CD4

Tconv, Treg and MAIT cells, ~4% in γδ T cells). The frequencies of GzmB+ AREG+ populations

within each T cell subset did not change across inflammation states (*Figure 3D*). Overall, these

data suggest T cells retained functional properties even in severely inflamed gingiva, and that

inflammation did not disrupt tissue-specific adaptation of the T cell compartment in the gingiva.

T cells in both the lung and decidual-placental interference (DPI) display few phenotypic

changes by inflammation status, but high variation by tissue origin

We next wanted to assess whether the phenotypic and functional stability of T cells across

inflammation states is a gingiva-specific phenomenon or could also be extended to other mucosal

tissues in healthy and inflamed states. We decided to examine the lung and the decidual-placental

interface (DPI) as a set of mucosal tissues that are inherently very distinct from the oral mucosa

in function and commensal exposure[42-44]. We assessed lung samples from deceased adult organ

donors (healthy) along with resected lung tissue from patients with interstitial lung disease (ILD)

patients. Briefly, ILD is a group of pulmonary disorders causing inflammation and lung tissue

scarring[45]. Tissue sampling was performed on both upper and lower lung lobes, which yielded

nearly congruent profiles (*Supplemental Figure 6*); the upper lobe data are shown for the

remainder of the figures. Decidual-placental interface (DPI) samples were taken from a cohort of

term (37-41 weeks gestation), uncomplicated pregnancies (healthy) and a cohort of term

pregnancies diagnosed with intra-amniotic inflammation (IAI). IAI, often referred to as

chorioamnionitis, is typically an acute inflammatory response to a bacterial infection[46].

First, we examined the composition of the T cell compartment in lung and DPI (***Figure***

4A*)*. T cells isolated from DPI tissues displayed a higher frequency of naïve cells among all CD3+

CD45+, which are likely derived from the highly vascularized placental tissue. These naïve T cells

were again excluded from manual gating analysis to facilitate a comparison with the memory

phenotype T cells in other tissues. We noted a decrease in MAIT cell (p = 0.020) and $\gamma\delta$ T cell (p

= 0.133) abundance in inflamed over healthy lung (***Figure 4A****)*. We next quantified the number of

markers that are significantly different based on tissue origin vs. inflammation state across each

T cell subset. For this, the source of variation from 2-way ANOVA statistical testing (row factor

and column factor p values) were counted as significant differences if p < 0.05 and totaled for all

subsets. We observed that tissue origin had a greater impact on subset composition and

phenotypic differences than inflammation status. (***Figure 4B****, Supplementary Table 2, markers*

displayed on heatmaps in Supplemental Figure 7). We found an inflammation-associated,

statistically significant increase in CD69+ T cells within several subsets, specifically, CD4 Tregs,

CD8 CD103+, and CD8 CD103- T cells in the placenta, but not the lung (**Figure 4C***, Supplemental*

Figure 7, statistical analysis across tissues excluded from heatmaps). False discovery rate

corrections for multiple comparisons of lung and placenta phenotype were also performed

(*Supplemental Table 3*). The most profound inflammation-associated phenotypic change

occurred within T cells in the lung. We observed a dramatic reduction in CD25 expression on all

T cell subsets in inflamed lungs (**Figure 4C**). Similarly to the oral mucosa, we did not observe an

increase in GzmB in healthy vs. inflamed tissues (lung and DPI) (**Figure 4C**). Furthermore, there

was no evidence of inflammation-associated activation of T cells, TCR-mediated activation,

proliferation, or exhaustion (*Supplemental Figure 8*), but rather, tissue origin predominantly

shaped the general phenotypic profile of T cell subsets.

To further visualize this relationship of phenotype and tissue location vs. inflammation

state across all three tissue sets, we grouped all datapoints from each tissue (irrespective of

inflammation state), color-coded each datapoint according to inflammation state, and compared

the expression of T cell surface markers across gingiva, lung, and placenta within conventional T

cell subsets (**Figures 4D-F**). As highlighted in previous figures, we did not observe inflammation-

associated changes in GzmB expression in CD8 T cell subsets, however there was a significant

difference in GzmB expression in CD8 CD103- T cells in the gingiva compared to lung and DPI

(**Figure 4D**). Among these tissue sets, the gingiva is presumably the tissue that is most

consistently and substantially exposed to microbes and antigen yet still displayed the lowest

baseline GzmB expression in CD8 CD103- T cells. Similarly, the frequency of TIM3+ Tregs was

lowest in the gingiva (**Figure 4E**). TIM3 expression by Tregs is typically observed in highly

activated Tregs in the tumor microenvironment[47]. In our dataset, we observed a nearly 3x increase

in frequency of TIM3 expression in Tregs from the DPI compared to gingiva (**Figure 4E**).

Together, these data suggest that the DPI, although typically considered to be a sterile tissue

environment[42], shows signs of being highly activated and highly suppressive. In line with this

notion, we also observed the highest frequency of CTLA4+ expressing CD8 CD103+ T cells in

the DPI (*Figure 4F*). Together, these data highlight that tissue-specific adaptions in the T cell

compartment exist across these 3 mucosal tissues and were maintained during states of

inflammation. We next wanted to test if the maintenance of tissue-specific adaptation also

extended to the functional capacity of these T cell subsets.

Functional capacity of T cells is primarily shaped by tissue location

As a first step, we examined the functional capacity of T cell subsets across lung and DPI

during health and disease. We found that the expression pattern of cytokines was relatively stable

across inflammatory states. Two statistically significant changes between healthy and inflamed

lung were noted in CD4 Tconv; a decrease in IL-17A and IL-17F expressing cells in inflamed

lungs (*Figure 5A, Supplemental Figure 9, statistical analysis across tissues excluded from*

heatmaps). Full statistical testing (FDR corrections for multiple comparisons is detailed in

Supplementary Table 3. CD4 Tconv from the DPI expressed more IL-2 compared to their lung

counterparts, regardless of inflammation status (*Figure 5A*). Overall, we observed for several T

cell populations (CD4 Tconv, CD8 CD103+, and CD8 CD103-) from the DPI that they express

more IFNγ and TNFα compared to lung (*Figure 5A*).

We next compared the functional capacity across all three tissue sites and inflammation

status, (*Figure 5B, gingiva – top, circles; lung – middle, triangles; placenta – bottom, diamonds*).

These data highlight that the greatest capacity for T cell IFNγ production (*Figure 5B*, left panel)

and IL-2 (*Figure 5B*, middle panel) was observed in the DPI. TNFα was also produced in the

highest quantities by T cells from the DPI and lowest in the lung (*Figure 5C*). In contrast, IL-17A

expression was essentially absent in DPI T cell subsets and most prominent in the oral mucosa

(*Figure 5B*, right panel). Differences in the functional capacity of CD4 Tregs particularly stood out

(*Figure 5B).* The oral mucosa contained a large fraction of Tregs that expressed IL-17A following

stimulation (*Figure 5B*, right panel), while Tregs from the placenta expressed IFNγ and IL-2

(*Figure 5B*, left and center panels). Of note, we observed co-expression of IFNγ and IL-2 for both

CD4 Tconv and Tregs from the DPI (*Figure 5D*) thus potentially suggesting that FoxP3+ CD127-

T cells may be activated CD4 Tconv (explored further in the discussion). Finally, it is noteworthy

that the bias towards certain functional properties in each tissue set was mirrored across several

T cell subsets. For example, gingival Tregs were the most potent IL-17 producers, and this was

true for CD4 T cell subsets, but also MAIT cells and γδ T cells (*Figure 5B*) regardless of

inflammation status. Overall, these data support the notion that the functional capacity of T cell

subsets is linked to tissue location, and that this can be maintained during inflammatory disease.

Functional potential of T cells groups by tissue and not inflammatory status in

dimensionality reduction analysis

Our analysis approach so far was focused on specifically examining select biomarkers of

activation and functional properties**.** We next wanted to compare tissues in a more unsupervised

analysis approach that takes the expression pattern of all proteins assessed in our 37-color panel

into account. Of note, we included all phenotypic and functional markers in the antibody staining

panel in this PCA analysis (TIM3, CD39, ICOS, CD25, CD28, CD27, PD1, Ki67, IL1R1, CD137,

CD40L, CTLA4, IL-2, IL-22, AREG, IL-17A, IFNγ, GzmB, TNFα, IL-17F). We first used principal

component analysis (PCA) to again compare the effect of tissue origin vs. inflammation status on

T cell function (*Figures 6A-D, Supplemental Figure 10)*. Tissue-based differences were

particularly apparent in CD4 Tconv and CD8 CD103+ T cells (*Figures 6A- B*) and were somewhat

less pronounced in MAIT and γδ T cells (*Figures 6C-D*). In contrast, for all subsets, there was

near congruent clustering based on inflammation status.

When determining which parameters are contributing most to determining the principal

components (loading scores), upon stimulation, changes in IFNγ among all T cells contributed the

greatest. The co-stimulation markers CD27 and CD28 were also significant contributors to the

PCA. Importantly, similar findings were observed when PCA was performed with samples directly

ex vivo (*Supplemental Figure 11)*. Taken together, an unsupervised analysis approach further

underlines the notion that tissue inflammation is not sufficient to disrupt tissue-specific

adaptations.

Unsupervised analysis approaches highlight that tissue-adaptation is the primary driver

of phenotypic and functional T cell changes

We next wanted to use an unsupervised approach to assess the relative abundance of T

cells with functional similarity across tissue and inflammation state. An unsupervised approach

allows for discovery of unique populations missed by manual gating of each marker using single-

or double-positive gates. For this approach, we included naïve T cells in the analysis as a

reference population. Briefly, we defined cell types by manual gating, performed

scaling/normalization, and defined global clusters across all batches. (described further in

methods). We assessed each T cell subset using this approach with tissues *directly ex vivo* and

following stimulation. We used our technical PBMC control sample within each experiment to

ensure that variation over time (antibody staining, flow cytometry set-up, etc.) was not a

confounding factor (*Supplemental Figure 3*).

Given the profound functional differences of Tregs between tissues, we initially focused

our analysis on Tregs upon stimulation. We observed ten distinct population clusters; the

expression of each marker is represented in the bubble plot (***Figure 7A***). Clusters 1 and 4 were

comprised of naïve cells based on the expression of CD45RA+ CCR7+, whereas the remaining

clusters consisted of memory subsets.

When examining cluster frequencies across tissues (gingiva, pink; lung, green; placenta,

purple) and noted distinct clustering based on tissue origin, described further below (**Figure 7B**,

top panels). We also examined cluster frequencies across inflammation status (healthy, blue;

inflamed, red), and observed relative stability (**Figure 7B,** *bottom left panels*).

Cluster 5, characterized by high expression of TNFα, was most abundant within the oral

mucosa (**Figure 7C-D**). Cluster 6 contains cells that express the proliferation marker Ki67 and

was found in high frequencies among placental tissues. Cluster 7, which comprised a large

proportion of all Tregs, is made up of cells that express CD39 and few other cytokines and was

prevalent in most samples, but most abundant in the placenta. Cytokines IFNγ and IL-17A are

represented within clusters 8 and 10, respectively. Tregs from the DPI during both health and

inflammation that group with cluster 8 are increased compared to gingiva and lung Tregs (**Figure**

7D). Cluster 10, characterized by the expression of IL-17A and IL1R1, was highest in the gingiva

(**Figure 7C-D**). Though markers like CTLA-4 increase upon stimulation with PMA/ionomycin, our

normalization methods allow for comparison of positive cells and comparison across groups.

Overall, these data are in line with our supervised analysis (**Figure 5B***, Supplemental Figure 9*).

We also examined how CD4 Tconv cells cluster across tissue and inflammation

(*Supplemental Figure 12*). Here, cluster 8 contains IL-17A+ Tconv cells (*Supplemental Figure*

12A). This cluster was most abundant within Tconv from gingival samples (*Supplemental Figure*

12D, bottom right panel). To summarize all clustering data for other T cell subsets *ex vivo* and

upon stimulation, we used clustering frequencies to generate PCA plots (*Supplemental Figure*

13-14). When cluster frequencies were used to determine principal components, distinct tissues

(gingiva, lung, and DPI) form unique groups, whereas healthy and inflamed tissues overlap.

Overall, our unsupervised analysis approach highlights that clusters show tissue-specific

distribution with minimal perturbation based on the inflammation status.

In summary, across all analysis methods, we found that tissue-specific adaptations of T

cell subsets were maintained during tissue inflammation.

Discussion

Our main goal was to assess the extent to which tissue-specific adaptations of T cell subsets are altered during tissue inflammation. We were surprised that markers indicative of inflammation-driven T cell activation did not significantly change in most T cell populations, including MAIT and $\gamma\delta$ T cells, as inflammation severity of the gingiva increased. We considered that this could be due to the baseline level of inflammation that is present even in healthy gingival tissues[35], which could yield unique regulatory mechanisms in the tissue. Since the oral mucosa is colonized by a dynamic microbiome and continuously exposed to dietary and microbial antigens[35], we also examined the expression of biomarker pairs that indicate TCR-mediated activation. ICOS and PD-1, and CD39 and PD-1 have been shown to be indicative of recent TCR activation for human CD4 and CD8 T cells, respectively[39-41]. Neither of these biomarker pairs indicated an association of a TCR-mediated T cell response in CD4 Tconv or a CD8 T cell subset gingiva with increased severity in inflammation. Similarly, Ki67 expression, an indicator of cell proliferation, and expression of PD-1 and TIM3, indictors of T cell exhaustion, did not change with increased inflammation in the gingiva. Overall, our observations highlight that the overall composition, phenotypic, and functional properties of T cell subsets in the gingiva were retained at various states of inflammation.

Because of the unique physiological demands of the gingiva, such as mastication, we asked if the T cell compartment would be similarly resistant to inflammation-mediated changes in other mucosal tissues, specifically the lung and the DPI. CD4 Tconv and CD8 T cell subsets in the healthy lung did not express these biomarker pairs of TCR activation. However, in the DPI we found that a substantial fraction of CD4 and CD8 T cell subsets co-expressed ICOS and PD-1, and CD39 and PD-1, respectively (*Supplemental Figure 8*). These data suggest that T cells in the DPI encounter their antigen more frequently than T cell in the gingiva and lung. This could potentially be related to the presence of fetal-derived, semi-allogenic tissue (the placenta is

made of fetal-derived trophoblasts) or could indicate that recently activated T cells migrate to

the DPI to provide protection. An in-depth analysis of the TCR repertoire in the DPI compared to

other tissues may provide a more definitive answer in future studies. Finally, ICOS, CD39 and

PD-1 are all expressed by highly activated Tregs, and expression of these markers was found

on a large fraction of Tregs across all tissues and inflammation states.

Given the presence of highly activated Tregs in all tissues, we explored potential

functional differences. We found that in gingival tissues, regardless of inflammation status,

about 10% of Tregs expressed IL-17A. IL-17+ Treg have been previously described and are

thought to maintain their ability to suppress other T cells, while secreting IL-17 in response to

TCR-mediated activation, or IL-6 and IL-1β stimulation[48,49]. In contrast, we did not observe IL-

17+ Tregs in lung or DPI. However, we found that a fraction of Tregs in both healthy and

inflamed DPI could express IFNγ, while none of the Tregs in other tissues expressed IFNγ

following stimulation. These cells could be Th1-like Tregs that have so far been mainly

described in peripheral blood in context of human autoimmune diseases and are thought to be

hypo-suppressive[50]. Thus, our observation could indicate that these Tregs also exist in a healthy

human tissue (albeit from a unique environment that contains a semi-allogeneic tissue), but it

important to consider that these could also be recently activated Th1 cells that temporarily

express FoxP3[51]. Follow up studies will be required to transcriptionally define this DPI-enriched

CD4 T cell population. Overall, tissue-associated phenotypes and functional properties were

evident, while inflammation-associated changes were limited in our cross-tissue comparison.

The most prominent phenotypic changes that we observed between a healthy and

inflamed tissue was the striking decrease in CD25 expression on T cell subsets in the inflamed

lung (Figure 4C, *Supplemental Figure 7*). We used a technical batch control to rule out any

technical artifacts. Previous reports indicated an increase in serum soluble CD25 levels in

individuals with ILD (both granulomatous-lymphocytic and rheumatoid-arthritis associated

ILD)[52,53]. ILD has also been clinically associated with increased matrix metalloproteinase (MMP)

expression in serum[54,55], which cleaves CD25 from the surface of T cells. If there is a local

increase in MMP activity in ILD lungs, then this could provide an explanation of the decreased

CD25 during ILD in all T cell subsets.

We specifically looked for functional changes that have previously been associated with

inflammation status. Studies using gingival tissue indicated a potential association between

inflammation and the abundance of Th17 cells[56,57]. We found that the overall composition of the

T cell compartment, including the frequency of CD4 T cells that make IL-17A or IL-17F following

stimulation, is stable even during inflammation. It is important to keep in mind that a hallmark

feature of inflammation is the recruitment of immune cell populations to a site of inflammation,

and that inflammation of the gingiva is associated with an increase in T cell infiltration[58]. Thus,

while the overall number of T cells with a specific function increases in inflamed tissues, this is

not necessarily associated with a skewing in functional properties within the T cell compartment.

Overall, the composition of the T cell compartment was consistent between mildly to severely

inflamed gingiva, with few subsets showing an increase or decrease with inflammation.

Specifically, we observed a reduction in resident CD8 CD103+ T cells within more severely

inflamed gingival tissue with a concomitant proportional increase in circulating CD8 CD103- T

cells. This stable maintenance of function would presumably change, at least temporarily, in

disease-specific conditions with an associated antigen-specific T cell response, such as the

Th17-driven response against *Candida albicans* and *Staphylococcus aureus*[59].

We conclude from these data that a state of tissue inflammation does not per se override

tissue-specific features of the T cell compartment. Importantly, our observations for the T cell

compartment do not necessarily extend to inflammation-associated alterations of other cell

types, particularly innate immune cells, in these tissues. In the gingiva, it is well documented

that neutrophil infiltration is associated with acute inflammation[56,60,61]. In the lungs, similar innate

subsets of macrophages and neutrophils have been reported to drive inflammatory responses in

mice and humans[62,63], and acute chorioamnionitis has also been defined by neutrophil

infiltration[64]. Furthermore, it is important to keep in mind that specific diseases can profoundly

impact T cell phenotype and function during inflammation. Thus, while our data indicate that

tissue inflammation is not inherently sufficient to override tissue-specific T cell adaptation,

different disease settings may display profound T cell changes such as previously observed in

the context of Hepatitis A infection, which leads to increased IL-15 availability and bystander

activation of CD8 T cells[27].

Finally, we want to highlight that we used a brief PMA/Ionomycin stimulation condition to

examine the functional capacity of T cells. We reasoned that the broadly activating nature of this

stimulus will reveal which proteins T cell subsets are able to express, while keeping in mind that

it does not reveal which proteins are secreted in situ. This approach indicated that T cell subsets

retained their ability to express AREG even in chronically inflamed tissue such as the ILD lung

and severely inflamed gingiva. Whether AREG is actually secreted *in situ* by T cells to promote

epithelial cell recovery in the inflamed gingiva is unclear. CD8 T cells appear to require a TCR

signal to secret AREG[34] and given the lack of evidence for CD8 T cell activation (including TCR-

mediated T cell activation) in these tissues, *in situ* AREG secretion by CD8 T cells seems

unlikely. However, given that the functional capacity is maintained in inflamed tissues, novel

therapies could target AREG to elicit expression or other specific functional responses.

In summary, we report that tissue-specific T cell adaptations can be maintained during

states of tissue inflammation. By gaining a more complete understanding of how inflammation

affects the tissue-specific T cell landscape, therapeutic approaches can be designed to develop

treatments with tissue-specific requirements in mind.

Material and Methods

Approvals & Study Population

Tissue Samples and Protocols: All protocols used in this study were approved by the Institutional

Review Board (IRB #0008335, 00008263, 00004036, 00001636). Signed informed consent was

obtained from all participants. A list of all samples, along with relevant clinical information can be

found in Supplementary Table 1. Each sample is from a unique donor.

Oral Mucosa: Oral mucosa samples were obtained from surgically discarded tissue from routine

surgeries, such as osseous surgery, functional or esthetic crown lengthening procedures, tooth

extractions, or impacted tooth uncover procedures. Mild inflammation (clinical health, mild to

moderate gingivitis), moderate inflammation (severe gingivitis, stage I, and stage II periodontitis),

and severe inflammation (stage III and IV periodontitis).

Lung: Healthy lung samples were redirected transplants from deceased organ donors. Donors

without history of lung disease, infection, and mechanical ventilation for >7 days were eligible for

screening. Inflamed lung samples (interstitial lung disease, ILD) were obtained from patients

undergoing lung transplant. Explant lungs and hilar lymph nodes were obtained from patients with

ILD undergoing lung transplant at the University of Washington.

Decidual-Placenta Interface: Both healthy and acutely inflamed (chorioamnionitis) third trimester

placentas were obtained following vaginal delivery or C-section, following labor or no labor

(detailed in Supplementary Table 1).

Technical Control Sample: A consistent leukopak peripheral blood mononuclear cell (PBMC)

sample was used throughout each flow cytometry batch and assessed alongside experimental

samples to ensure longitudinal control. Leukopak (4510-01 Ful Leuko Pack) obtained through

Bloodworks Bio (Seattle, WA).

Isolation of Lymphocytes from Tissues

Oral Mucosa: Following procedures, mucosal tissue was priority shipped overnight with cold

packs to Fred Hutch Cancer Center (FHCC). Upon arrival, tissue was washed with complete

media (RPMI1640 (11875-093, Gibco) with penicillin, streptomycin (15140-122, Gibco), and 10%

fetal bovine serum (FBS, PS-FB2, Peak Serum)). An optimized protocol was adapted from

reference[65], originally developed for processing solid tumors. Tissue was placed in a petri dish

and minced with a scalpel, then enzymatically digested (Collagenase II (C6885, Sigma-Aldrich),

DNAse (20 units/ml, #D502, Sigm) in complete media with reduced 7.5% FBS (RP7.5) in a 25 mL

volume at 37°C for 40 minutes while shaking. Following digest, the sample was passed through

a needle (16x1 ½ Blunt, 302833, BD) 10 times prior to passing through a 100-micron cell strainer.

Remaining tissue pieces were mashed using a syringe plunger, washed with complete media,

and frozen in 1 mL Recovery Cell Culture Freezing Medium (12648010, Gibco) and stored in

liquid nitrogen until future experiments. Cell viability and abundance were quantified prior to

freezing by staining a small sample aliquot (Guava antibodies 7-AAD, CD19, and CD45) for 15

minutes, resuspension in 1% PFA, and collection on the Guava easyCyte. Oral mucosa samples

in this study contained between 731-52,195 T cells, as analyzed by flow cytometry.

Lung: Lung tissue was processed in a similar fashion to the gingiva using the RP7.5 digestion for

50 minutes at 37°C while shaking at 225 RPM. Subsequently cells were washed with RP10,

strained through a 70µm filter and enumerated. Lung tissue samples contained between 1,537-

62,719 T cells.

Decidual-Placenta Interface: Tissue was processed as previously described[66]. Briefly, intact

placentas were biopsied up to a 0.5 cm depth on the maternal facing surface of the placenta

including both decidua basalis and placenta villi. Tissue was digested in digestion media (above)

and incubated at 37°C for 60 minutes with agitation. Concentration for digest was RP7.5 + 700

U/mL Collagenase Type II (Sigma-Aldrich) and 200 U/mL DNase. Cells were filtered and washed

with media. Red blood cells were removed using ACK lysis buffer, followed by resuspension with

media. Maternal blood was collected in tubes containing the anti-coagulant acid citric dextrose.

DPI tissue samples contained between 10,303-102,650 T cells.

Technical Control Sample: Leukopak was processed according to StemCell Technologies

protocol (https://www.stemcell.com/leukopak-processing-protocol.html). Two blood cell lysis

steps at room temperature and two platelet washes were performed prior to counting and freezing

20 million cells per aliquot.

Stimulation Assay

As briefly outlined in Figure 1A, upon experimentation, cryo-preserved single-cell suspension

isolated from tissues or blood (outlined above) were thawed in a 37°C water bath. 12,000 to 2

million cells per condition were plated in a U- or V-bottom 96 well plate in 200 uL complete media.

For assessing phenotype directly _ex vivo_, cells were left untreated for 2 hours at 37°C. GolgiStop

(Monensin, 554724, BD) and GolgiPlug (Protein Transport Inhibitor (Containing Brefeldin A),

555029, BD) were then added per manufacturer recommendations to compare directly to

stimulated samples and was returned to the 37°C incubator for 4 more hours (total of 6 hours).

During functional stimulation assays, samples were similarly thawed and plated. 200 uL complete

media supplemented with PMA (100 ng/mL, P8139, Sigma) and ionomycin (1000 ng/mL, I9657,

Sigma) for 6 hours, including the addition of Golgi stop/plug 2 hours after beginning the

stimulation. Upon completion, samples were stained for analysis by flow cytometry.

Spectral Flow Cytometry

Staining: For all experiments, upon completion of the assay, samples were stained previously

described[67]. Prior to experimentation, the spectral flow panel was tittered, optimized, and tested

to ensure proper design and maximal information could be obtained from each sample, as

detailed in reference[68]. All staining steps were performed in the dark at room temperature in a 96-

bottom plate. Cells were incubated with FcX block (422302, Biolegend) and Zombie NIR dye

(422302, Biolegend) in PBS (14190250, Gibco) for 15 minutes. Antibody master mixes were spun

down at 10,000xg for 4 minutes prior to staining to remove aggregates. Following live/dead

staining, cells were washed, following by surface staining in 50uL (antibody mix in PBS + 2% FBS

and 10% brilliant stain buffer (563794, BD)) for 20 minutes. Cells were then washed twice prior to

fix/perm (100 uL, 554722, BD) according to manufacturer guidelines for 30 minutes. After fixation,

cells were stained intracellularly for select markers noted in the panel (Supplementary Figure 1)

for 30 minutes, assembled in perm/wash buffe and 10% BSB. Finally, cells were washed twice

with perm/wash and resuspended in PBS + 2% FBS and stored at 4°C until acquisition the

following day on the BD Discover S8.

Controls: Single-stain controls were optimized for individual markers by staining either

unstimulated PBMCs, PBMCs following a 6-hour PMA/ionomycin stimulation, or unstimulated

control tissue samples (generally tonsil tumor infiltrate or regional lymph nodes). For two markers

expressed in low abundance in PBMC and tissue (IL-13 and IL-22), capture beads were used for

single-stain control. Control samples were taken through an identical staining protocol to

experimental samples, described above. A reference sample was also used within each

experiment as a longitudinal control (*Supplemental Figure 3*).

Cytometer: Data from all experiments was collected with the BD Discover S8 and the

FACSChorus software platform (BD Biosciences). The S8 is equipped with 5 lasers and 78

detectors. Compensation controls were prepared and used for each individual experiment, aside

from one run containing ILD 11 and 12. Upon collection, data was exported (FCS 3.1 format) and

analyzed using FlowJo version 10.10.0. Files were analyzed in batches by collection day, with

minor gating changes across different batches.

Unsupervised Clustering

Upon acquisition on the Discover S8 (BD), samples were gated (*Supplemental Figure 1*).

Following gating of time, lymphocytes, single cells, and live cells, T cells were gated (CD3+

CD45+) and exported as a new .fcs file in FlowJo. Files were also stripped of parameters out of

scope for this analysis, focusing on time, size channels, and those containing a fluorochrome

(*listed in Supplementary Figure 1*). Upon creation of files containing only T cells, separate streams

were created for each relevant T cell subset assessed in this study: $\gamma\delta$ T cells (TCR$\gamma\delta$+), MAIT

cells (CD161+ Vα7.2+), all CD4 (CD8-), Tconv (FoxP3-), Tregs (FoxP3+ CD127-), all CD8 (CD8+

CD4-), resident CD8 cells (CD103+), and circulating CD8 cells (CD103-). Naïve cells were

included in clustering analysis for all tissues.

We included additional samples in our clustering analysis as reference tissues for tissue-

enriched T cell subsets (*Supplemental Table 1, gray lines*). Lung and associated lymph nodes,

along with maternal blood from placental donors were included in multiple batches. We also

included tumor infiltrates from head and neck squamous cell carcinoma (HNSCC) and associated

lymph nodes. One experimental batch did not include marker CD127 and was excluded from the

Tconv and Treg clustering analyses (n = 1 pediatric lung and associated LNs, 2 inflamed lungs

and associated LNs, 2 tumor infiltrate samples, and 8 tumor associated LNs).

For each clustering stream, we used up to 20,000 T cells per sample. No minimum cell

threshold was utilized. Actual T cell counts for individual samples and primary clustering streams

were considered for count analysis. The following markers were used to develop clusters: CD161,

CD39, CCR7, ICOS, CD25, IL-17F, CD28, CD27, CD45RA, PD-1, Ki67, CD69, TIM3, IFNγ,

CTLA4, IL-13, IL-22, GzmB, TNFα, IL1R1, CD137, AREG, CD40L, IL-17A, CD45RO, IL-2.

Lineage markers and QC parameters were excluded from clustering analysis: CD3, CD45, CD4,

CD8, FoxP3, CD127 or CXCR5, CD103, TCR$\gamma\delta$, Vα7.2, Live/Dead, Autofluorescence. Spectre[69]

was used to determine appropriate arcsinh transformation values for each parameter.

Repeated clustering at the single cell level – necessary to identify and factor out technical

noise, or perceived batch effect – is computationally intractable at the scale of this study.

Therefore, within each of the streams, to arrive at globally defined clusters across all batches, we

cluster the centers of clusters from batch-specific clusterings[70,71]. Specifically, we 'over'-cluster

within each batch by first clustering, then subclustering the cells within each cluster. Next, we

compute the centroids of all (sub)clusters in each batch and cluster these (sub)cluster centroids

(or "metacells" or "supercells") to arrive at a global (meta)clustering assignment for each metacell

from each batch[72]. At each stage we use the Rphenograph algorithm, with Leiden clustering of

shared nearest-neighbor graphs[73,74]. To address moderate drift across batches in some

parameters, we first align repeated control median transformed intensity values across batches

for select parameters, and next apply the same batch- and parameter-specific shifts to the

samples before metaclustering. We use Uniform Manifold Approximate Projection (UMAP) for

dimension reduction and data visualization within an iterative process of adjusting for perceived

batch effects, optimized by refining the metaclustering at a fixed Leiden resolution, before pushing

resolution higher[75].

Statistical Analyses

Statistical testing was performed using GraphPad Prism Software version 10.5. Testing us noted within figure legends. All bar graphs represent the mean with standard deviation. Heatmaps display the mean for each marker for a given sample group. For oral mucosa samples (*Figures 1-3*), significance was determined by testing using ordinary one-way ANOVA with Tukey's multiple comparisons test. For lung and placenta data (*Figures 4-5*), 2-way ANOVA was utilized with Tukey's multiple comparisons if comparing two tissues across inflammation states, or one-way ANOVA if comparing with three tissues with grouped inflammation. All data using one-way ANOVA was also assessed using the Benjamini-Hochberg false discovery rate (FDR) correction with a rate of 10% and maintained statistical significance. Supplementary Table 2 used 2-way ANOVA to compare lung and placenta across inflammation; the Benjamini-Hochberg FDR correction with a rate of 10% was utilized.

Resource Availability

Raw flow cytometry files (.fcs) will be available on ImmPort in Study SDY3484.

Acknowledgements

We thank the Flow Cytometry Shared Resources Core at Fred Hutch Cancer Center. We thank Quinn Peters who provided and assisted with the R script used for PCA and Veronica Davé who consulted on proper statical testing and corrections to be used for these analyses. We also thank all members of the Prlic Lab for critical feedback and discussion.

Author contributions

Conceptualization: N.B.P., M.P. Methodology: N.B.P., H. R. M., M.P. Conducting experiments: N.B.P., E. D. V., A. K. T., C. S. D., G. D. Formal analysis: N.B.P., H. R. M., M.P. Writing; original draft: N.B.P., M.P. Writing; review and editing: all authors. Visualization: N.B.P., H. R. M., M.P. Funding acquisition: N.B.P, M.P. Supervision: M.P.

Funding

NIH grants R56DE032009, R01AI123323 and R01AI179712 (to M.P) and funding from the Oral Mucosa Immunology Consortium at the University of Nebraska (to N.B.P, Windsweep Farm Scholar) supported this work.

Declarations of interest

The authors declare no competing interests.

Supplemental Information

Document S1. Figures S1-S14.

Tables S1-3 contain tissue information, data range of heatmaps, and statistical testing.

Figure Legends

Figure 1: The inflammation state in the oral mucosa has a moderate effect on composition and phenotype of the T cell compartment.

A. Experimental layout, described in detail in methods. Briefly, oral mucosa tissue of varying inflammation states was sent from the clinic to lab, kept at 4°C. Tissue was digested mechanically and enzymatically prior to storage of cryovials with single cell suspension in liquid nitrogen. Upon experiment, samples were thawed and assessed directly ex vivo (*Figures 1-2*) or following a 6h PMA/I stimulation at 37°C (*Figure 3*). **B.** Frequency of CD3+ T cells of CD45+ cells across inflammation state, pre-gated on lymphocytes, single cells, and live (*Supplementary Figure 1*). **C.** Of CD3+ CD45+ cells, frequency of each T cell subset. **D.** Frequency of subset (left to right, CD8+ CD103+ resident, and CD8+ CD103- circulating) of CD3+ CD45+ T cells across inflammation state. For **B-D.** T cell subsetting, n = 9 mild, 13 moderate, 11 severely inflamed oral mucosa tissues. **E.** For each conventional subset, heatmaps depict frequency of activation markers across inflammation status. Phenotyping of T cell subsets must meet minimum 25 cell threshold; sample size is consistent with above unless noted; n = 8 mild, 11 moderate for Tregs. Tconv, CD103+ and CD103- all met minimum threshold. **F.** Frequency of MAIT cells and $\gamma\delta$ T cells of CD3+ CD45+ T cells. **G.** For innate-like T cells, frequency of activation markers across inflammation. Minimum threshold of 25 cells; n = 7 mild, 9 moderate for MAIT cells; n = 8 mild for $\gamma\delta$ T cells. For all plots, ns (not significant), $*p < 0.05$, $***p < 0.001$, $****p < 0.0001$ by ordinary one-way ANOVA with Tukey's multiple comparison test.

Figure 2: Biomarkers of T cell receptor (TCR)-mediated activation do not significantly change across inflammation states in the oral mucosa.

A. Expression of CD39+ PD-1+ (right) and ICOS+ PD-1+ (left) among all T cell subsets across inflammation (blue, mild; yellow, moderate; red, severely inflamed). No statistical differences

within each subset across inflammation. **B.** Representative flow cytometry plots of CD39

(BUV615) by PD-1 (BV605) expression (top) and ICOS (BUV737) by PD-1 (BV605) expression

(bottom) within CD4 Tconv cells isolated from severely inflamed oral mucosal tissue. **C.** From left

to right, expression of Ki67+ and TIM3+ PD-1+ across T cell subsets across inflammation status.

No statistical differences within each subset. **D.** Representative flow cytometry plots of Ki67

(BV650) expression (top) and TIM3 (BB515) by PD-1 (BV605) expression (bottom) within CD4

Tconv cells isolated from moderately inflamed (top) or severely inflamed (bottom) oral mucosal

tissue. No significant differences by ordinary one-way ANOVA with Tukey's multiple comparison

test.

Figure 3: Functional capacity of T cell subsets is maintained in inflamed oral mucosa.

A. Within T cell subset, expression of each molecule across mild, moderate, or severely inflamed

oral mucosa following 6h PMA/I stimulation. n = 9 mild, 13 moderate, 11 severely inflamed oral

mucosa tissues, unless noted based on minimum threshold for analysis of 25 cells. For Tregs, n

= 8 mild; n = 11 moderate. No statistical differences across inflammation status for each marker.

B. Expression of AREG, IFNγ, and IL-2 within each T cell subset; no statical differences across

inflammation. **C.** Expression of each molecule within MAIT and $\gamma\delta$ T cells. **D.** Representative

gating of AREG+ GzmB+ populations within mildly inflamed oral mucosa (left) and co-expression

of AREG+ GzmB+ among CD103+ CD8+ resident T cells (right). For MAIT cells, n = 7 mild, n =

8 moderate, n = 10 severe; $\gamma\delta$ T cells, n = 7 mild, n = 10 severe. No significant differences by

ordinary one-way ANOVA with Tukey's multiple comparison test.

Figure 4: T cells subsets from the lung and decidual-placental interference (DPI) display

few phenotypic changes by inflammation status, but high variation by tissue origin.

A. Among all CD3+ CD45+ cells, frequency of indicated subset within each tissue and disease state. **B.** Statistical differences within tissue type and across inflammation by 2-way ANOVA (overall row and column factors). **C.** For conventional subsets (right to left; Tconv, Treg, CD103+ CD8+ and CD103- CD8+), frequency of activation markers by tissue and disease status. Unless noted, n = 4 healthy lung, n = 10 inflamed lung, n = 10 healthy placenta, n = 8 inflamed placenta. Some samples did not meet minimum threshold of 25 cells; for Tregs, n = 3 healthy lung; within CD103+, n = 3 healthy lung for all figure panels. Within one experiment, n = 8 inflamed lung samples for Tconv and Tregs for this and remaining panels. Statistical testing between tissues is excluded from heatmaps. **D.** Frequency of GzmB+ cells among CD8 CD103- across tissue and inflammation state (healthy/mild = blue squares, moderate gingiva = yellow circles, severe/inflamed = red diamonds). **E.** Frequency of TIM3+ Tregs. **F.** Frequency of CTLA-4+ cells among resident CD8 CD103+ cells. Pre-gated as indicated in Supplemental Figure 1 prior to subsetting and single-positive gating. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001 by ordinary one-way ANOVA with Tukey's multiple comparison test.

Figure 5: Functional capacity of T cells is primarily shaped by tissue location.

A. Upon 6hr stimulation with PMA/ionomycin, expression of cytokines and tissue healing markers across T cell subsets in the lung and placenta. Unless noted, n = 4 healthy lung, n = 10 inflamed lung, n = 10 healthy placenta, n = 8 inflamed placenta. For Tconv and Tregs, n = 8 inflamed lung samples for this and subsequent figure panels. Statistical testing between tissues excluded from heatmaps. **B.** Within each T cell subset, expression of IFNγ (left), IL-2 (middle), and IL-17A (right). For all panels, circles = gingiva, upright triangle = lung, upside down triangle = placenta; for this and subsequent figure panels, healthy/mild = blue, moderate gingiva = yellow, severe/inflamed = red. For MAIT cells, n = 3 healthy lung and n = 9 inflamed lung samples. **C.** TNFα expression across tissues in CD4 Tconv (left) and CD8 CD103- (right), pre-gated based on Supplemental

Figure 1 scheme. **D.** Frequencies of IFNγ+ IL-2+ across tissues in CD4 Tconv (left) and Tregs

(right). *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001 by ordinary one-way ANOVA with

Tukey's multiple comparison test.

Figure 6: Functional potential of T cells groups by tissue and not inflammatory status in

dimensionality reduction analysis.

Principal components determined by frequency of positive cells within gates from the following

parameters: TIM3, CD39, ICOS, CD25, CD28, CD27, PD1, Ki67, IL1R1, CD137, CD40L, CTLA4,

IL-2, IL-22, AREG, IL-17A, IFNγ, GzmB, TNFα, IL-17F. PCA plots of flow cytometry data following

6-hour PMA/Ionomycin stimulation from gingiva (pink), lung (green), and placenta (purple) during

health (blue) and inflammation (red) organized by tissue type (left) and inflammation status (right)

of **A.** CD4 Tconv, **B.** CD8 CD103+, **C.** MAIT cells, **D.** γδ T cells. Moderately inflamed oral mucosa

is grouped with inflammation category. Ellipses represent 95% confidence interval.

Figure 7: Unsupervised analysis approaches highlight that tissue-adaptation is the primary

driver of phenotypic and functional T cell changes.

The following markers were used to develop clusters: CD161, CD39, CCR7, ICOS, CD25, IL-17F,

CD28, CD27, CD45RA, PD-1, Ki67, CD69, TIM3, IFNγ, CTLA4, IL-13, IL-22, GzmB, TNFα, IL1R1,

CD137, AREG, CD40L, IL-17A, CD45RO, IL-2. **A.** Bubble plots of expression of each marker

within each cluster among Tregs upon stimulation. Resolution = 0.75. **B.** UMAPs of clusters

organized by tissue origin and inflammation status. **C.** Stacked bar graphs representing cluster

breakdown for individual samples. **D.** Box plots of frequency of each cluster relative to CD4 Treg

counts out of total T cells from each sample.

References

1. Sathaliyawala, T., Kubota, M., Yudanin, N., Turner, D., Camp, P., Thome, J.J., Bickham, K.L., Lerner, H., Goldstein, M., Sykes, M., et al. (2013). Distribution and compartmentalization of human circulating and tissue-resident memory T cell subsets. Immunity *38*, 187-197. 10.1016/j.immuni.2012.09.020.

2. Kumar, B.V., Ma, W., Miron, M., Granot, T., Guyer, R.S., Carpenter, D.J., Senda, T., Sun, X., Ho, S.H., Lerner, H., et al. (2017). Human Tissue-Resident Memory T Cells Are Defined by Core Transcriptional and Functional Signatures in Lymphoid and Mucosal Sites. Cell Rep *20*, 2921-2934. 10.1016/j.celrep.2017.08.078.

3. Szabo, P.A., Levitin, H.M., Miron, M., Snyder, M.E., Senda, T., Yuan, J., Cheng, Y.L., Bush, E.C., Dogra, P., Thapa, P., et al. (2019). Single-cell transcriptomics of human T cells reveals tissue and activation signatures in health and disease. Nat Commun *10*, 4706. 10.1038/s41467-019-12464-3.

4. Connors, T.J., Matsumoto, R., Verma, S., Szabo, P.A., Guyer, R., Gray, J., Wang, Z., Thapa, P., Dogra, P., Poon, M.M.L., et al. (2023). Site-specific development and progressive maturation of human tissue-resident memory T cells over infancy and childhood. Immunity *56*, 1894-1909 e1895. 10.1016/j.immuni.2023.06.008.

5. Poon, M.M.L., Caron, D.P., Wang, Z., Wells, S.B., Chen, D., Meng, W., Szabo, P.A., Lam, N., Kubota, M., Matsumoto, R., et al. (2023). Tissue adaptation and clonal segregation of human memory T cells in barrier sites. Nat Immunol *24*, 309-319. 10.1038/s41590-022-01395-9.

6. Wong, M.T., Ong, D.E., Lim, F.S., Teng, K.W., McGovern, N., Narayanan, S., Ho, W.Q., Cerny, D., Tan, H.K., Anicete, R., et al. (2016). A High-Dimensional Atlas of Human T Cell Diversity Reveals Tissue-Specific Trafficking and Cytokine Signatures. Immunity *45*, 442-456. 10.1016/j.immuni.2016.07.007.

7. Niessl, J., Muller, T.R., Constantz, C., Cai, C., Nilsen, V., Rivera Ballesteros, O., Adamo, S., Kammann, T., Mouchtaridi, E., Gao, Y., et al. (2025). Tissue origin and virus specificity shape human CD8(+) T cell cytotoxicity. Sci Immunol *10*, eadq4881. 10.1126/sciimmunol.adq4881.

8. Reina-Campos, M., Monell, A., Ferry, A., Luna, V., Cheung, K.P., Galletti, G., Scharping, N.E., Takehara, K.K., Quon, S., Challita, P.P., et al. (2025). Tissue-resident memory CD8 T cell diversity is spatiotemporally imprinted. Nature *639*, 483-492. 10.1038/s41586-024-08466-x.

9. Traxinger, B.R., Richert-Spuhler, L.E., and Lund, J.M. (2022). Mucosal tissue regulatory T cells are integral in balancing immunity and tolerance at portals of antigen entry. Mucosal Immunol *15*, 398-407. 10.1038/s41385-021-00471-x.

10. Munoz-Rojas, A.R., and Mathis, D. (2021). Tissue regulatory T cells: regulatory chameleons. Nat Rev Immunol *21*, 597-611. 10.1038/s41577-021-00519-w.

11. Miragaia, R.J., Gomes, T., Chomka, A., Jardine, L., Riedel, A., Hegazy, A.N., Whibley, N., Tucci, A., Chen, X., Lindeman, I., et al. (2019). Single-Cell Transcriptomics of Regulatory T Cells Reveals Trajectories of Tissue Adaptation. Immunity *50*, 493-504 e497. 10.1016/j.immuni.2019.01.001.

12. Simon, M., Stuve, P., Schmidleithner, L., Bittner, S., Beumer, N., Strieder, N., Schmidl, C., Pant, A., Gebhard, C., Eigenberger, A., et al. (2024). Single-cell chromatin accessibility and transposable element landscapes reveal shared features of tissue-residing immune cells. Immunity *57*, 1975-1993 e1910. 10.1016/j.immuni.2024.06.015.

13. Gray, J.I., Caron, D.P., Wells, S.B., Guyer, R., Szabo, P., Rainbow, D., Ergen, C., Rybkina, K., Bradley, M.C., Matsumoto, R., et al. (2024). Human gammadelta T cells in diverse

tissues exhibit site-specific maturation dynamics across the life span. Sci Immunol *9*,
eadn3954. 10.1126/sciimmunol.adn3954.

14. Kammann, T., Cai, C., Sekine, T., Mouchtaridi, E., Boulouis, C., Nilsen, V., Ballesteros,
O.R., Muller, T.R., Gao, Y., Raineri, E.J.M., et al. (2024). MAIT cell heterogeneity across
paired human tissues reveals specialization of distinct regulatory and enhanced effector
profiles. Sci Immunol *9*, eadn2362. 10.1126/sciimmunol.adn2362.

15. Garner, L.C., Amini, A., FitzPatrick, M.E.B., Lett, M.J., Hess, G.F., Filipowicz Sinnreich,
M., Provine, N.M., and Klenerman, P. (2023). Single-cell analysis of human MAIT cell
transcriptional, functional and clonal diversity. Nat Immunol *24*, 1565-1578.
10.1038/s41590-023-01575-1.

16. Mueller, S.N., Gebhardt, T., Carbone, F.R., and Heath, W.R. (2013). Memory T cell
subsets, migration patterns, and tissue residence. Annu Rev Immunol *31*, 137-161.
10.1146/annurev-immunol-032712-095954.

17. Maurice, N.J., Taber, A.K., and Prlic, M. (2021). The Ugly Duckling Turned to Swan: A
Change in Perception of Bystander-Activated Memory CD8 T Cells. J Immunol *206*, 455-
462. 10.4049/jimmunol.2000937.

18. Lee, H., Jeong, S., and Shin, E.C. (2022). Significance of bystander T cell activation in
microbial infection. Nat Immunol *23*, 13-22. 10.1038/s41590-021-00985-3.

19. Sekine, T., Perez-Potti, A., Nguyen, S., Gorin, J.B., Wu, V.H., Gostick, E., Llewellyn-
Lacey, S., Hammer, Q., Falck-Jones, S., Vangeti, S., et al. (2020). TOX is expressed by
exhausted and polyfunctional human effector memory CD8(+) T cells. Sci Immunol *5*.
10.1126/sciimmunol.aba7918.

20. Maurice, N.J., Berner, J., Taber, A.K., Zehn, D., and Prlic, M. (2021). Inflammatory signals
are sufficient to elicit TOX expression in mouse and human CD8+ T cells. JCI Insight *6*.
10.1172/jci.insight.150744.

21. Berg, R.E., and Forman, J. (2006). The role of CD8 T cells in innate immunity and in
antigen non-specific protection. Curr Opin Immunol *18*, 338-343.
10.1016/j.coi.2006.03.010.

22. Berg, R.E., Cordes, C.J., and Forman, J. (2002). Contribution of CD8+ T cells to innate
immunity: IFN-gamma secretion induced by IL-12 and IL-18. Eur J Immunol *32*, 2807-
2816. 10.1002/1521-4141(2002010)32:10<2807::AID-IMMU2807>3.0.CO;2-0.

23. Kambayashi, T., Assarsson, E., Lukacher, A.E., Ljunggren, H.G., and Jensen, P.E. (2003).
Memory CD8+ T cells provide an early source of IFN-gamma. J Immunol *170*, 2399-2408.
10.4049/jimmunol.170.5.2399.

24. Chu, T., Tyznik, A.J., Roepke, S., Berkley, A.M., Woodward-Davis, A., Pattacini, L.,
Bevan, M.J., Zehn, D., and Prlic, M. (2013). Bystander-activated memory CD8 T cells
control early pathogen load in an innate-like, NKG2D-dependent manner. Cell Rep *3*, 701-
708. 10.1016/j.celrep.2013.02.020.

25. Yajima, T., Nishimura, H., Ishimitsu, R., Yamamura, K., Watase, T., Busch, D.H., Pamer,
E.G., Kuwano, H., and Yoshikai, Y. (2001). Memory phenotype CD8(+) T cells in IL-15
transgenic mice are involved in early protection against a primary infection with Listeria
monocytogenes. Eur J Immunol *31*, 757-766. 10.1002/1521-4141(200103)31:3<757::aid-
immu757>3.0.co;2-q.

26. Whiteside, S.K., Snook, J.P., Williams, M.A., and Weis, J.J. (2018). Bystander T Cells: A
Balancing Act of Friends and Foes. Trends Immunol *39*, 1021-1035.
10.1016/j.it.2018.10.003.

27. Kim, J., Chang, D.Y., Lee, H.W., Lee, H., Kim, J.H., Sung, P.S., Kim, K.H., Hong, S.H.,
Kang, W., Lee, J., et al. (2018). Innate-like Cytotoxic Function of Bystander-Activated
CD8(+) T Cells Is Associated with Liver Injury in Acute Hepatitis A. Immunity *48*, 161-173
e165. 10.1016/j.immuni.2017.11.025.

28. McGeachy, M.J., Cua, D.J., and Gaffen, S.L. (2019). The IL-17 Family of Cytokines in
Health and Disease. Immunity *50*, 892-906. 10.1016/j.immuni.2019.03.021.
29. Li, J., Casanova, J.L., and Puel, A. (2018). Mucocutaneous IL-17 immunity in mice and
humans: host defense vs. excessive inflammation. Mucosal Immunol *11*, 581-589.
10.1038/mi.2017.97.
30. Sun, L., Su, Y., Jiao, A., Wang, X., and Zhang, B. (2023). T cells in health and disease.
Signal Transduct Target Ther *8*, 235. 10.1038/s41392-023-01471-y.
31. Singh, N., Baby, D., Rajguru, J.P., Patil, P.B., Thakkannavar, S.S., and Pujari, V.B. (2019).
Inflammation and cancer. Ann Afr Med *18*, 121-126. 10.4103/aam.aam_56_18.
32. Turner, D.L., and Farber, D.L. (2014). Mucosal resident memory CD4 T cells in protection
and immunopathology. Front Immunol *5*, 331. 10.3389/fimmu.2014.00331.
33. Arpaia, N., Green, J.A., Moltedo, B., Arvey, A., Hemmers, S., Yuan, S., Treuting, P.M.,
and Rudensky, A.Y. (2015). A Distinct Function of Regulatory T Cells in Tissue Protection.
Cell *162*, 1078-1089. 10.1016/j.cell.2015.08.021.
34. Delacher, M., Schmidleithner, L., Simon, M., Stuve, P., Sanderink, L., Hotz-Wagenblatt,
A., Wuttke, M., Schambeck, K., Ruhland, B., Hofmann, V., et al. (2024). The effector
program of human CD8 T cells supports tissue remodeling. J Exp Med *221*.
10.1084/jem.20230488.
35. Konkel, J.E., Cox, J.R., and Wemyss, K. (2024). Bite-sized immunology; damage and
microbes educating immunity at the gingiva. Mucosal Immunol *17*, 1141-1150.
10.1016/j.mucimm.2024.07.004.
36. Kinane, D.F., Stathopoulou, P.G., and Papapanou, P.N. (2017). Periodontal diseases. Nat
Rev Dis Primers *3*, 17038. 10.1038/nrdp.2017.38.
37. Paprckova, D., Salyova, E., Michalik, J., and Stepanek, O. (2023). Bystander activation in
memory and antigen-inexperienced memory-like CD8 T cells. Curr Opin Immunol *82*,
102299. 10.1016/j.coi.2023.102299.
38. Konecny, A.J., Huang, Y., Setty, M., and Prlic, M. (2024). Signals that control MAIT cell
function in healthy and inflamed human tissues. Immunol Rev *323*, 138-149.
10.1111/imr.13325.
39. Duhen, R., Fesneau, O., Samson, K.A., Frye, A.K., Beymer, M., Rajamanickam, V., Ross,
D., Tran, E., Bernard, B., Weinberg, A.D., and Duhen, T. (2022). PD-1 and ICOS
coexpression identifies tumor-reactive CD4+ T cells in human solid tumors. J Clin Invest
132. 10.1172/JCI156821.
40. Duhen, T., Duhen, R., Montler, R., Moses, J., Moudgil, T., de Miranda, N.F., Goodall, C.P.,
Blair, T.C., Fox, B.A., McDermott, J.E., et al. (2018). Co-expression of CD39 and CD103
identifies tumor-reactive CD8 T cells in human solid tumors. Nat Commun *9*, 2724.
10.1038/s41467-018-05072-0.
41. Simoni, Y., Becht, E., Fehlings, M., Loh, C.Y., Koo, S.L., Teng, K.W.W., Yeong, J.P.S.,
Nahar, R., Zhang, T., Kared, H., et al. (2018). Bystander CD8(+) T cells are abundant and
phenotypically distinct in human tumour infiltrates. Nature *557*, 575-579. 10.1038/s41586-
018-0130-2.
42. de Goffau, M.C., Lager, S., Sovio, U., Gaccioli, F., Cook, E., Peacock, S.J., Parkhill, J.,
Charnock-Jones, D.S., and Smith, G.C.S. (2019). Human placenta has no microbiome but
can contain potential pathogens. Nature *572*, 329-334. 10.1038/s41586-019-1451-5.
43. Briana, D.D., Papaevangelou, V., and Malamitsi-Puchner, A. (2021). The jury is still out
on the existence of a placental microbiome. Acta Paediatr *110*, 2958-2963.
10.1111/apa.16048.
44. Whiteside, S.A., McGinniss, J.E., and Collman, R.G. (2021). The lung microbiome:
progress and promise. J Clin Invest *131*. 10.1172/JCI150473.
45. Maher, T.M. (2024). Interstitial Lung Disease: A Review. JAMA *331*, 1655-1665.
10.1001/jama.2024.3669.

46.   Jung, E., Romero, R., Suksai, M., Gotsch, F., Chaemsaithong, P., Erez, O., Conde-
874   Agudelo, A., Gomez-Lopez, N., Berry, S.M., Meyyazhagan, A., and Yoon, B.H. (2024).
875   Clinical chorioamnionitis at term: definition, pathogenesis, microbiology, diagnosis, and
876   treatment. Am J Obstet Gynecol *230*, S807-S840. 10.1016/j.ajog.2023.02.002.
877   47.   Sakuishi, K., Ngiow, S.F., Sullivan, J.M., Teng, M.W., Kuchroo, V.K., Smyth, M.J., and
878   Anderson, A.C. (2013). TIM3(+)FOXP3(+) regulatory T cells are tissue-specific promoters
879   of T-cell dysfunction in cancer. Oncoimmunology *2*, e23849. 10.4161/onci.23849.
880   48.   Voo, K.S., Wang, Y.H., Santori, F.R., Boggiano, C., Wang, Y.H., Arima, K., Bover, L.,
881   Hanabuchi, S., Khalili, J., Marinova, E., et al. (2009). Identification of IL-17-producing
882   FOXP3+ regulatory T cells in humans. Proc Natl Acad Sci U S A *106*, 4793-4798.
883   10.1073/pnas.0900408106.
884   49.   Beriou, G., Costantino, C.M., Ashley, C.W., Yang, L., Kuchroo, V.K., Baecher-Allan, C.,
885   and Hafler, D.A. (2009). IL-17-producing human peripheral regulatory T cells retain
886   suppressive function. Blood *113*, 4240-4249. 10.1182/blood-2008-10-183251.
887   50.   Dominguez-Villar, M., Baecher-Allan, C.M., and Hafler, D.A. (2011). Identification of T
888   helper type 1-like, Foxp3+ regulatory T cells in human autoimmune disease. Nat Med *17*,
889   673-675. 10.1038/nm.2389.
890   51.   Wang, J., Ioan-Facsinay, A., van der Voort, E.I., Huizinga, T.W., and Toes, R.E. (2007).
891   Transient expression of FOXP3 in human activated nonregulatory CD4+ T cells. Eur J
892   Immunol *37*, 129-138. 10.1002/eji.200636435.
893   52.   Galant-Swafford, J., Catanzaro, J., Achcar, R.D., Cool, C., Koelsch, T., Bang, T.J., Lynch,
894   D.A., Alam, R., Katial, R.K., and Fernandez Perez, E.R. (2024). Approach to diagnosing
895   and managing granulomatous-lymphocytic interstitial lung disease. EClinicalMedicine *75*,
896   102749. 10.1016/j.eclinm.2024.102749.
897   53.   Cao, S., Liu, X., Li, Y., Yang, Y., Cai, X., Cong, S., Li, Z., Li, Y., Hong, Y., Su, Y., et al.
898   (2024). Serum sCD25 is an indicator for rheumatoid arthritis-associated interstitial lung
899   disease. Clin Exp Rheumatol *42*, 633-641. 10.55563/clinexprheumatol/3iqvk3.
900   54.   Chen, H., Tang, J., Liang, J., Huang, D., Pan, C., Liu, S., Du, X., and Tao, L. (2023).
901   Clinical association study on the matrix metalloproteinase expression in the serum of
902   patients with connective tissue disease complicated with interstitial lung disease. Arch
903   Rheumatol *38*, 367-374. 10.46497/ArchRheumatol.2023.9547.
904   55.   Sheu, B.C., Hsu, S.M., Ho, H.N., Lien, H.C., Huang, S.C., and Lin, R.H. (2001). A novel
905   role of metalloproteinase in cancer-mediated immunosuppression. Cancer Res *61*, 237-
906   242.
907   56.   Kim, T.S., Silva, L.M., Theofilou, V.I., Greenwell-Wild, T., Li, L., Williams, D.W., Ikeuchi,
908   T., Brenchley, L., Genomics, N.N., Computational Biology, C., et al. (2023). Neutrophil
909   extracellular traps and extracellular histones potentiate IL-17 inflammation in periodontitis.
910   J Exp Med *220*. 10.1084/jem.20221751.
911   57.   Dutzan, N., Kajikawa, T., Abusleme, L., Greenwell-Wild, T., Zuazo, C.E., Ikeuchi, T.,
912   Brenchley, L., Abe, T., Hurabielle, C., Martin, D., et al. (2018). A dysbiotic microbiome
913   triggers T(H)17 cells to mediate oral mucosal immunopathology in mice and humans. Sci
914   Transl Med *10*. 10.1126/scitranslmed.aat0797.
915   58.   Woodward Davis, A.S., Roozen, H.N., Dufort, M.J., DeBerg, H.A., Delaney, M.A., Mair,
916   F., Erickson, J.R., Slichter, C.K., Berkson, J.D., Klock, A.M., et al. (2019). The human
917   tissue-resident CCR5(+) T cell compartment maintains protective and functional
918   properties during inflammation. Sci Transl Med *11*. 10.1126/scitranslmed.aaw8718.
919   59.   Zielinski, C.E., Mele, F., Aschenbrenner, D., Jarrossay, D., Ronchi, F., Gattorno, M.,
920   Monticelli, S., Lanzavecchia, A., and Sallusto, F. (2012). Pathogen-induced human TH17
921   cells produce IFN-gamma or IL-10 and are regulated by IL-1beta. Nature *484*, 514-518.
922   10.1038/nature10957.

60.   Ginesin, O., Coyac, B.R., Doppelt-Flikshtain, O., Mayer, Y., Gabay, E., Berg, T., Bar-On,
Y., and Zigdon-Giladi, H. (2025). Macrophage Depletion Reduces Bone Loss and Alters
Inflammatory Responses: A Mouse Peri-Implantitis Model. J Clin Periodontol *52*, 898-906.
10.1111/jcpe.14120.

61.   Moutsopoulos, N.M., Konkel, J., Sarmadi, M., Eskan, M.A., Wild, T., Dutzan, N.,
Abusleme, L., Zenobia, C., Hosur, K.B., Abe, T., et al. (2014). Defective neutrophil
recruitment in leukocyte adhesion deficiency type I disease causes local IL-17-driven
inflammatory bone loss. Sci Transl Med *6*, 229ra240. 10.1126/scitranslmed.3007696.

62.   Misharin, A.V., Morales-Nebreda, L., Reyfman, P.A., Cuda, C.M., Walter, J.M., McQuattie-
Pimentel, A.C., Chen, C.I., Anekalla, K.R., Joshi, N., Williams, K.J.N., et al. (2017).
Monocyte-derived alveolar macrophages drive lung fibrosis and persist in the lung over
the life span. J Exp Med *214*, 2387-2404. 10.1084/jem.20162152.

63.   Prasse, A., Pechkovsky, D.V., Toews, G.B., Jungraithmayr, W., Kollert, F., Goldmann, T.,
Vollmer, E., Muller-Quernheim, J., and Zissel, G. (2006). A vicious circle of alveolar
macrophages and fibroblasts perpetuates pulmonary fibrosis via CCL18. Am J Respir Crit
Care Med *173*, 781-792. 10.1164/rccm.200509-1518OC.

64.   Zhang, C., Cao, J., Xu, M., Wu, D., Li, W., and Chang, Y. (2023). The role of neutrophils
in chorioamnionitis. Front Immunol *14*, 1198831. 10.3389/fimmu.2023.1198831.

65.   Leelatian, N., Doxie, D.B., Greenplate, A.R., Mobley, B.C., Lehman, J.M., Sinnaeve, J.,
Kauffmann, R.M., Werkhaven, J.A., Mistry, A.M., Weaver, K.D., et al. (2017). Single cell
analysis of human tissues and solid tumors with mass cytometry. Cytometry B Clin Cytom
92, 68-78. 10.1002/cyto.b.21481.

66.   Maurice, N.J., Erickson, J.R., DeJong, C.S., Mair, F., Taber, A.K., Frutoso, M., Islas, L.V.,
Vigil, A., Lawler, R.L., McElrath, J., et al. (2025). Cytokine and metabolite networks shape
T cell residency and functionality at the term human maternal-fetal interface. J Immunol
214, 2221-2237. 10.1093/jimmun/vkaf093.

67.   Mair, F., Erickson, J.R., Frutoso, M., Konecny, A.J., Greene, E., Voillet, V., Maurice, N.J.,
Rongvaux, A., Dixon, D., Barber, B., et al. (2022). Extricating human tumour immune
alterations from tissue inflammation. Nature *605*, 728-735. 10.1038/s41586-022-04718-
w.

68.   Mair, F., and Tyznik, A.J. (2019). High-Dimensional Immunophenotyping with
Fluorescence-Based Cytometry: A Practical Guidebook. Methods Mol Biol *2032*, 1-29.
10.1007/978-1-4939-9650-6_1.

69.   Ashhurst, T.M., Marsh-Wakefield, F., Putri, G.H., Spiteri, A.G., Shinko, D., Read, M.N.,
Smith, A.L., and King, N.J.C. (2022). Integration, exploration, and analysis of high-
dimensional single-cell cytometry data using Spectre. Cytometry A *101*, 237-253.
10.1002/cyto.a.24350.

70.   Putri, G.H., Howitt, G., Marsh-Wakefield, F., Ashhurst, T.M., and Phipson, B. (2024).
SuperCellCyto: enabling efficient analysis of large scale cytometry datasets. Genome Biol
25, 89. 10.1186/s13059-024-03229-3.

71.   Bilous, M., Herault, L., Gabriel, A.A., Teleman, M., and Gfeller, D. (2024). Building and
analyzing metacells in single-cell genomics data. Mol Syst Biol *20*, 744-766.
10.1038/s44320-024-00045-6.

72.   Liu, X., Song, W., Wong, B.Y., Zhang, T., Yu, S., Lin, G.N., and Ding, X. (2019). A
comparison framework and guideline of clustering methods for mass cytometry data.
Genome Biol *20*, 297. 10.1186/s13059-019-1917-7.

73.   Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden: guaranteeing
well-connected communities. Sci Rep *9*, 5233. 10.1038/s41598-019-41695-z.

74.   Levine, J.H., Simonds, E.F., Bendall, S.C., Davis, K.L., Amir el, A.D., Tadmor, M.D., Litvin,
O., Fienberg, H.G., Jager, A., Zunder, E.R., et al. (2015). Data-Driven Phenotypic

Dissection of AML Reveals Progenitor-like Cells that Correlate with Prognosis. Cell *162*,
184-197. 10.1016/j.cell.2015.05.047.
75. Becht, E., McInnes, L., Healy, J., Dutertre, C.A., Kwok, I.W.H., Ng, L.G., Ginhoux, F., and
Newell, E.W. (2018). Dimensionality reduction for visualizing single-cell data using UMAP.
Nat Biotechnol. 10.1038/nbt.4314.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7