Tissue-specific adaptation of human T cells is preserved during tissue

inflammation

Nicole B. Potchen[1], Hugh R. MacMillan[1], Eva Domenjo-Vila[1], Andrew J. Konecny[1,2], Alexis K.

Taber[1], Caitlin S. DeJong[1], Geervani Daggupati[1,2], Swati Shree[3], Stephen A. McCartney[3], Shelton

W. Wright[4], Evan W. Newell[1,6], Douglas R. Dixon[5], Martin Prlic[1,6*]

[1] Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center, Seattle, WA, USA.

[2] Department of Immunology, University of Washington, Seattle, WA, USA.

[3] Division of Maternal Fetal Medicine, Department of Obstetrics and Gynecology, University of Washington, Seattle,

WA, USA.

[4] Division of Pediatric Critical Care Medicine, Department of Pediatrics, University of Washington, Seattle, WA, USA.

[5] Department of Bioscience Research and Periodontics, University of Tennessee Health Science Center, College of

Dentistry, Memphis, TN, USA.

[6] Department of Global Health, University of Washington, Seattle, WA, USA.

* Corresponding author

Fred Hutchinson Cancer Center, Vaccine and Infectious Disease Division, Seattle, WA, USA.

mprlic@fredhutch.org

Summary

T cells play an essential role in protecting tissues against pathogens and regulating tissue homeostasis. Previous studies highlight that T cells display tissue-specific phenotypic and functional properties, suggesting that T cells adapt to their local environment. Whether inflammation disrupts tissue-specific T cell adaptation remains poorly understood. To address this open question, we examined the T cell compartment – including conventional CD4 and CD8 T cells, regulatory T cells, $\gamma\delta$ T cells, and MAIT cells – from healthy and inflamed human mucosal tissues. Using high-parameter spectral flow cytometry, we examined phenotype *ex vivo* and the functional capacity following stimulation, utilizing conventional gating and unsupervised clustering analysis approaches. Overall, we analyzed 65 tissue samples including mild, moderate, and severely inflamed oral gingiva, healthy and inflamed lung, along with healthy and inflamed tissue from the decidual-placental interface. Across these mucosal barrier tissues, we find that tissue location plays a dominant role in shaping the composition, phenotype, and functional capacity of the T cell compartment. Importantly, these tissue-specific adaptations were largely maintained during states of tissue inflammation. This included the ability to exert tissue repair functions, which was preserved across T cell subsets, even in severely inflamed tissues.

Introduction

Human T cells have distinct phenotypic and functional properties across different tissues and organs. This was initially demonstrated by Farber and colleagues who examined the phenotype and functional properties of conventional CD4 and CD8 T cells isolated from secondary lymphoid tissues, lung, jejunum, ileum and colon of trauma-induced, deceased organ donors[1]. This study furthermore established CD69 expression as a biomarker for tissue-residence for human CD4 T cells, and CD69 and CD103 co-expression to indicate tissue-residence for CD8 T cells[1]. Follow-up studies highlighted that expression of CD69 by human CD4 and CD8 T cells is associated with a core transcriptional signature akin to bona fide murine tissue-resident T cells[2]. Human tissue-resident memory CD8 T cells (T_{RM}) show tissue-specific adaptation such as distinct cytotoxic properties across skin, lung, and jejunum, and are also clonally distinct across these tissue sites[3-7]. A recent study indicated that within the intestine, distinct tissue niches drive unique phenotypic and functional properties in CD8 T cells, thus providing further compelling evidence that T_{RM} tissue location and functional state are fundamentally linked[8]. Similarly, conventional CD4 T cells (CD4 Tconv) and regulatory T cells (CD4 Treg) also have distinct phenotypic and functional properties across tissues[9-12]. Further evidence points to innate-like T cells also adapting in a tissue-specific manner. Human $\gamma\delta$ T cells differ by tissue location with cells from adult lung, spleen, lymph nodes (LNs), and gut tissue clustering differentially by tissue based on their expression of transcripts associated with effector and repair functions[13]. Finally, human mucosal-associated invariant T (MAIT) cells across tissues have distinct T cell receptor (TCR) repertoires and cytokine expression profiles[14,15].

Whether tissue-specific adaptations are maintained during tissue inflammation is unclear. Inflammation involves coordinated immune cell recruitment and activation and thus has the potential to profoundly alter the T cell compartment within tissue[16]. Memory phenotype CD8 T cells can be activated by pro-inflammatory signals alone; this event has been coined bystander-

activation or inflammation-driven activation of T cells[17,18]. Inflammation-driven activation of

memory CD8 T cells leads to increased expression of CD69, PD-1, and TOX, along with

expression of IFNγ and Granzyme B (GzmB)[19-23] which can have protective properties[24,25] but can

also lead to tissue pathologies[26]. Such bystander-activated CD8 T cell-mediated pathology has

been demonstrated in a human cohort following infection with Hepatitis A virus (HAV)[27]. For CD4

T cells, inflammation and associated tissue pathology is often associated with expression of IL-

17, but it is noteworthy that IL-17 should not be simply viewed as a pro-inflammatory cytokine[28]

or indicator of excessive inflammation[29]. Importantly, in addition to mediating tissue protection and

pathology, T cells also play a role in restoring tissue homeostasis via secretion of IL-22 and the

epidermal growth factor receptor (EGFR)-ligand amphiregulin (AREG)[30-32]. Tregs express AREG

in response to IL-18 and IL-33[33], while a recent study demonstrates that human memory CD8 T

cells express AREG in response to TCR stimulation[34].

While there has been substantial progress in our understanding of the T cell compartment

and its tissue-specific functional properties in healthy human tissues, the extent to which

inflammation alters these functional properties remains an open question. We sought to address

if inflammation overrides any tissue-specific T cell adaption in human tissues. Specifically, we

aimed to determine if inflammation alters the composition of the T cell compartment and

associated functional properties, and if it affects the ability of T cells to maintain their tissue-

regenerative functions. To accomplish this, we collected healthy and inflamed tissues from 3

distinct mucosal barrier sites: oral mucosa (gingiva), lung, and the decidual-placental interface

(DPI). We developed a high-parameter flow cytometry panel containing 37 markers to define the

composition, phenotype, and functional capacity of the T cell compartment: CD4 Tconv, CD4

Treg, CD8 CD103+ T_{RM} and CD8 CD103- circulating T cells, along with the innate-like MAIT and

γδ T cell subsets. Overall, we found that the composition, phenotype and function of the T cell

compartment was driven more by tissue location than inflammation, indicating that T cells largely

maintain their tissue-specific adaptations during inflammation. Our data emphasize the need to

assess and consider inflammation in a tissue-specific manner. Furthermore, our data highlight

that T cells maintain their tissue-regenerative capacity even in severely inflamed tissues, thus

pinpointing T cells as a therapeutic target to treat inflammatory diseases.

Results

The inflammation state in the oral mucosa has a modest effect on composition and phenotype of the T cell compartment

To determine the impact of inflammation on the mucosal T cell compartment, we first examined phenotypic differences across mildly, moderately, and severely inflamed gingival tissues. These tissues are discarded from routine surgeries and represent a wide range of inflammatory states. We assigned clinically healthy gingival tissue and tissue with mild gingivitis as "mild inflammation" since even clinically healthy tissue contains a baseline level of inflammation, which may allow for constant immune surveillance[35]. Next, we assigned tissues with pronounced gingivitis transitioning to periodontitis as "moderate inflammation" and those with periodontitis as "severe inflammation". While gingivitis is reversible with proper dental hygiene, periodontitis is an irreversible, chronic disease state without clinical intervention[36]. A full sample list with clinical information can be found in Supplemental Table 1. A 37-color spectral flow cytometry panel was utilized to examine phenotype (no stimulation) and functional capacity (following stimulation) of the T cells isolated from gingival tissues (**Figure 1A***, panel and gating; Supplemental Figure 1-2*). As a technical control, a PBMC sample from a single leukapheresis donor was included in every flow cytometry experiment (*Supplemental Figure 3*).

We first examined the frequency of T cells (CD3+ cells) within the CD45+ cell compartment and found no changes between mild, moderate, and severely inflamed gingival tissue (**Figure 1B**). Of note, a small population of naïve T cells (CCR7+ CD45RA+) was present in each tissue sample. These naïve T cells presumably stem from the blood that is inherently present in the tissue vasculature. We excluded these naïve T cells from our manual gating analysis approach. We further analyzed the T cell compartment by first defining innate-like T cell subsets within the CD3+ population, followed by subsetting the remaining CD4 and CD8 T cells into conventional CD4 T cells (CD4 Tconv), regulatory CD4 T cells (Treg), CD103+ (tissue-resident phenotype)

CD8 T cells and CD103- (circulating) CD8 T cells (*Supplemental Figure 1B*). We then assessed if T cell subset composition is altered with inflammation. We observed stable frequencies of most T cell subsets, including CD4 Tconv, CD4 Treg, and CD103+ CD8 T cells (**Figure 1C**). We noticed a slight reduction in CD103+ CD8 T cells within the total T cell population (p = 0.1026), but a significant increase in circulating CD103- CD8 T cells in severely inflamed tissue compared to mild inflammation (**Figure 1D**).

We next examined if the expression of activation, co-stimulation, and exhaustion markers change with inflammation severity. To visualize similarities and differences in expression of multiple proteins across different tissues, we chose a heatmap-based approach to displaying expression frequency. For each protein, the mean expression frequency is shown on all heatmaps, and the median and data range for each protein are shown in *Supplemental Table 2.* Within conventional CD4 and CD8 T cell subsets, we observed only few changes between mild, moderate, and severely inflamed gingiva (**Figure 1E**). Pro-inflammatory cytokines can increase the expression of CD69, PD-1, and GzmB by memory CD8 T cells [20], but we did not observe an inflammation-associated increase of these proteins (**Figure 1E**). On the contrary, we noted reduced expression of CD69 in both the CD8 CD103+ and CD103- populations in severely inflamed compared to mildly inflamed tissues. In the CD4 Tconv population, there were no differences in these phenotypic markers. In general, we observed relative phenotypic stability across inflammation status within each T cell subset (*Supplemental Figure 4*).

Next, we examined innate-like T cell subsets including MAIT and γδ T cells, which can also be activated by pro-inflammatory stimuli[37,38]. While there were no statistically significant differences in MAIT cell frequencies and phenotypic markers across inflammation status, we found a significant reduction in γδ T cells in severely inflamed compared to moderately inflamed tissue (**Figure 1F**). Within the moderately inflamed condition, we also observed a pronounced reduction of PD-1 expression in the γδ T cell population (**Figure 1G**). The concomitant increase

in the abundance of $\gamma\delta$ T cells (as a % of CD3+ T cells) and decreased PD-1 expression could

potentially be due to recruitment of $\gamma\delta$ T cells from the blood to moderately inflamed tissue. While

we observed some phenotypic differences in our dataset as tissue became increasingly inflamed,

we noted that the T cell profile within all oral mucosal tissues remained overall rather stable even

in severely inflamed tissues. We next considered that a smaller proportion of T cells may respond

in a TCR-dependent manner.

Biomarkers of TCR-mediated activation do not significantly change across inflammation

states in the oral mucosa

We next assessed biomarker pairs indicative of TCR-mediated activation[39-41]. We

examined co-expression of CD39+ PD-1+ (for CD8+ cells[40,41]) as well as ICOS+ PD-1+ (for CD4+

cells[39]) across mild, moderate, and severely inflamed oral mucosal tissues. Across gingiva of

varying inflammation states, there were no differences in frequency of expression of each of these

populations. CD39, PD-1 and ICOS are also expressed on highly activated Tregs, and we

consistently observed the highest frequency of expression in CD4 Tregs (***Figure 2A-B***). In line

with this observation that TCR-mediated activation did not change across inflammation states, we

also observed no changes in Ki67 expression, an indicator of recent cell proliferation, across

inflammation status (***Figure 2C-D***). Lastly, we examined TIM3 and PD-1 expression, a biomarker

pair indicative of T cell exhaustion. We observed unaltered co-expression of these markers

regardless of inflammation state (***Figures 2C-D***). Together, these data indicate that biomarkers

of T cell activation and proliferation remained stable across a range of inflammatory disease

states within the gingiva. Overall, we did not detect evidence of inflammation- or TCR-driven

activation of the T cell compartment in the inflamed gingiva. To assess the ability of the T cells to

respond to activation we next interrogated T cell functional properties.

Functional capacity for T cell-mediated tissue healing and cytokine production is maintained in inflamed oral mucosa

To examine if the functional capacity of T cell subsets varies across inflammation, we stimulated T cells isolated from gingival tissue for 6 hours with PMA/ionomycin (**Figure 1A**). We define the functional capacity simply as the ability of a T cell to express protein following this brief, yet broad, activation stimulus. Within CD4 and CD8 subsets, we observed no statistically significant differences in AREG, IFNγ, TNFα, IL-17A, IL-17F, IL-2, or IL-22 expression across inflammation states (**Figure 3A**, *Supplemental Figure 5*). Even the distribution of datapoints for each T cell subset was similar across mildly, moderately, and severely inflamed gingiva (**Figure 3B**).

Similarly, the functional capacity of MAIT and $\gamma\delta$ T cells was stable across inflammation states (**Figure 3C**). The mean, median, and data range are reported in Supplemental Table 2. Within several T cell subsets, we found co-expression of GzmB and AREG upon stimulation, which may indicate dual functional roles (cytotoxicity and tissue repair) can occur simultaneously within the same T cell. The highest frequency of GzmB and AREG co-expression was observed in CD8 CD103+ cells (**Figure 3D**), though these were also detected to a lesser extent in CD4 Tconv, CD4 Tregs, and innate-like subsets (~1% within CD4 Tconv, Treg and MAIT cells, ~4% in $\gamma\delta$ T cells). Overall, these data suggest T cell subsets retained their functional capacity even in severely inflamed gingiva.

T cells from lung and decidual-placental interference (DPI) differ based on tissue origin regardless of inflammation status

We next wanted to assess whether the phenotypic and functional stability of T cells across inflammation states is a gingiva-specific phenomenon or could also be extended to other mucosal tissues in healthy and inflamed states. We decided to examine the lung and the decidual-placental

interface (DPI) as a set of mucosal tissues that are inherently very distinct from the oral mucosa

in function and commensal exposure[42-44]. We assessed lung samples from deceased adult organ

donors (healthy) along with resected lung tissue from patients with interstitial lung disease (ILD).

Briefly, ILD is a group of pulmonary disorders causing inflammation and lung tissue scarring[45].

Tissue sampling was performed on both upper and lower lung lobes, which yielded nearly

congruent profiles (*Supplemental Figure 6*); the upper lobe data are shown for the remainder of

the figures. Decidual-placental interface (DPI) samples were taken from a cohort of term (37-41

weeks gestation), uncomplicated pregnancies (healthy) and a cohort of term pregnancies

diagnosed with intra-amniotic inflammation (IAI). IAI, often referred to as chorioamnionitis, is

typically an acute inflammatory response to a bacterial infection[46].

First, we examined the composition of the T cell compartment in lung and DPI (***Figure***

4A). T cells isolated from DPI tissues displayed a higher frequency of naïve cells among all CD3+

CD45+, which are likely derived from the highly vascularized placental tissue. These naïve T cells

were again excluded from manual gating analysis to facilitate a comparison with the memory

phenotype T cells in other tissues. We noted a decrease in MAIT cell (p = 0.020) and $\gamma\delta$ T cell (p

= 0.133) abundance in inflamed over healthy lung, but did not observe significant inflammation-

associated changes in the DPI (***Figure 4A***).

Next, for each protein, the mean expression frequency is shown on all heatmaps in **Figure**

4B, Supplemental Figure 7, and the median and data range for each protein are shown in

Supplemental Table 2. We noted an inflammation-associated, statistically significant increase in

CD69+ T cells within CD4 Tregs, CD8 CD103+, and CD8 CD103- T cells in the placenta, but not

the lung (***Figure 4B***, *Supplemental Figure 7, statistical analysis across tissues excluded from*

heatmaps). False discovery rate corrections for multiple comparisons of lung and placenta

phenotype were also performed (*Supplemental Table 3*). A profound inflammation-associated

phenotypic change occurred within T cells in the lung. We observed a dramatic reduction in CD25

expression on all T cell subsets in inflamed lungs (***Figure 4B***). Similarly to the oral mucosa, we

did not observe an increase in GzmB in healthy vs. inflamed tissues (lung and DPI) (***Figure 4B***),

and did not observe evidence for TCR-mediated activation, proliferation, or exhaustion

(*Supplemental Figure 8*). We next wanted to further quantify how tissue origin (lung and DPI) vs.

inflammation state affect protein expression of T cells. To accomplish this, we took the source of

variation from 2-way ANOVA statistical testing (row factor and column factor p values with p <

0.05). We observed that tissue origin had a greater impact on subset composition and phenotypic

differences than inflammation status. (*Supplementary Table 2, markers displayed on heatmaps*

in Supplemental Figure 7).

So far, the data indicate that tissue origin predominantly shaped the general phenotypic

profile of T cell subsets. We next combined all 3 tissue types for our subsequent analysis steps.

To further visualize this relationship of phenotype and tissue location vs. inflammation state

across all three tissue sets, we grouped all datapoints from each tissue (irrespective of

inflammation state), color-coded each datapoint according to inflammation state, and compared

the expression of T cell surface markers across gingiva, lung, and placenta within conventional T

cell subsets (***Figures 4C-E***). As highlighted in previous figures, we did not observe inflammation-

associated changes in GzmB expression in CD8 T cell subsets, however there was a significant

difference in GzmB expression in CD8 CD103- T cells in the gingiva compared to lung and DPI

(***Figure 4C***). Among these tissue sets, the gingiva is presumably the tissue that is most

consistently and substantially exposed to microbes and antigen yet still displayed the lowest

baseline GzmB expression in CD8 CD103- T cells. Similarly, the frequency of TIM3+ Tregs was

lowest in the gingiva (***Figure 4D***). TIM3 expression by Tregs is typically observed in highly

activated Tregs in the tumor microenvironment[47]. In our dataset, we observed a nearly 3x increase

in frequency of TIM3 expression in Tregs from the DPI compared to gingiva (***Figure 4D***).

Together, these data suggest that the DPI, although typically considered to be a sterile tissue

environment[42], shows signs of being concomitantly activated and suppressive. In line with this

notion, we also observed the highest frequency of CTLA4+ expressing CD8 CD103+ T cells in

the DPI (**Figure 4E**). Together, these data highlight that tissue-specific phenotypic adaptations

in the T cell compartment across these mucosal tissues were maintained during states of

inflammation. We next wanted to test if the functional capacity of these T cell subsets was similarly

maintained in all tissues during inflammation.

Functional capacity of T cells is primarily shaped by tissue location

As a first step, we examined the functional capacity of T cell subsets across lung and DPI

during health and disease. We found that the expression pattern of cytokines was relatively stable

across inflammatory states. Two statistically significant changes between healthy and inflamed

lung were noted in CD4 Tconv; a decrease in IL-17A and IL-17F expressing cells in inflamed

lungs (**Figure 5A**, *Supplemental Figure 9, statistical analysis across tissues excluded from*

heatmaps). Full statistical testing (FDR corrections for multiple comparisons) is detailed in

Supplementary Table 3.

We next compared the functional capacity across all three tissue sites and inflammation

status, (**Figure 5B**, *gingiva – top, circles; lung – middle, triangles; placenta – bottom, diamonds*).

These data highlight that the greatest capacity for T cell IFNγ production (**Figure 5B**, left panel)

and IL-2 (**Figure 5B**, middle panel) was observed in the DPI. TNFα was also produced in the

highest quantities by T cells from the DPI and lowest in the lung (**Figure 5C**). In contrast, IL-17A

expression was essentially absent in DPI T cell subsets and most prominent in the oral mucosa

(**Figure 5B**, right panel). Differences in the functional capacity of CD4 Tregs particularly stood out

(**Figure 5B**). The oral mucosa contained a large fraction of Tregs that expressed IL-17A following

stimulation (**Figure 5B**, right panel), while Tregs from the placenta expressed IFNγ (**Figure 5B**,

left panel). Of note, we observed co-expression of IFNγ and IL-2 for both CD4 Tconv and Tregs

from the DPI (**Figure 5D**) thus potentially suggesting that some FoxP3+ CD127- T cells may be

activated CD4 Tconv (explored further in the discussion). Finally, it is noteworthy that the bias

towards certain functional properties in each tissue set was mirrored across several T cell

subsets. For example, gingival Tregs were the most potent IL-17 producers, and this was true for

CD4 T cell subsets, but also MAIT cells and γδ T cells (**Figure 5B**) regardless of inflammation

status. Overall, these data support the notion that the functional capacity of T cell subsets is linked

to tissue location, and that this can be maintained during inflammatory disease.

Functional capacity of T cells across tissues and inflammation state by dimensionality

reduction analysis

Our analysis approach so far was focused on specifically examining select biomarkers of

activation and functional properties**.** We next wanted to compare tissues in a more unsupervised

analysis approach. Of note, we performed this analysis separately for stimulated and unstimulated

samples. Stimulation – as expected – altered the expression of some cell surface biomarkers. We

reasoned that the presence vs. absence of stimulation-induced changes could also be informative

and included all phenotypic and functional markers in the antibody staining panel in this PCA

analysis (TIM3, CD39, ICOS, CD25, CD28, CD27, PD1, Ki67, IL1R1, CD137, CD40L, CTLA4, IL-

2, IL-22, AREG, IL-17A, IFNγ, GzmB, TNFα, IL-17F). We first used principal component analysis

(PCA) to compare the effect of tissue origin vs. inflammation status on T cell functional capacity

(**Figures 6A-D**, *Supplemental Figure 10)*. Tissue-based differences in functional capacity were

particularly apparent in CD4 Tconv and CD8 CD103+ T cells (**Figures 6A-B**) and were somewhat

less pronounced in MAIT and γδ T cells (**Figures 6C-D**). In contrast, for all subsets, there was

near congruent clustering based on inflammation status.

When determining which parameters are contributing most to determining the principal

components (loading scores), upon stimulation, changes in IFNγ expression contributed the

greatest for all T cells. Differences in expression of the co-stimulatory molecules CD27 and CD28

were also significant contributors to the PCA. Importantly, similar findings were observed when

PCA was performed with unstimulated samples directly *ex vivo* (*Supplemental Figure 11)*. Taken

together, an unsupervised analysis approach further underlines the notion that tissue

inflammation is not sufficient to disrupt tissue-specific adaptation of the T cell compartment.

An unsupervised analysis approach to compare the T cell compartment across tissues

and inflammation

We next wanted to use an unsupervised approach to assess the relative abundance of T

cells with functional similarity across tissue and inflammation state. An unsupervised approach

also allows for discovery of phenotypically/functionally unique populations potentially missed by

manual gating of each marker using single- or double-positive gates. Of note, for this approach,

we included naïve T cells in the analysis as a reference population. Briefly, T cell subsets were

first defined by manual gating. We then performed scaling/normalization and defined global

clusters across all samples (described further in methods). We assessed each T cell subset using

this approach with tissues analyzed directly *ex vivo* and following stimulation. Given the profound

differences of Treg functional capacity between tissues, we first focused our analysis on

stimulated Tregs. We examined cluster frequencies across tissues (gingiva, pink; lung, green;

placenta, purple) (***Figure 7A***, *top row panels*) and across inflammation status (healthy, blue;

inflamed, red) (***Figure 7A,*** *bottom left panels*).

We next assessed the characteristics of the observed ten distinct population clusters using

a bubble plot (***Figure 7B***). The bubble plot highlighted that clusters 1 and 4 were comprised of

naïve cells based on the expression of CD45RA+ CCR7+, whereas the remaining clusters

consisted of memory subsets. The relative abundance of each cluster combined for all donors is

displayed in **Figure 7C**, and then further broken down by single donor in **Figure 7D**. Tregs that

expressed the cytokines IFNγ and IL-17A were in clusters 8 and 10, and – in line with our

observations from manual gating – most abundant in placenta and oral mucosa, respectively.

Overall, these data indicate that our manual gating analysis captured differences in Treg

functional capacity across tissues (**Figure 5B**, *Supplemental Figure 9*). We next performed this

analysis for other T cell subsets. Clustering for CD4 Tconv cells across tissue and inflammation

is shown in *Supplemental Figure 12*; the results are also in line with our manual gating analysis.

To summarize all clustering data for other T cell subsets *ex vivo* and upon stimulation, we used

clustering frequencies to generate PCA plots (*Supplemental Figure 13-14*). When cluster

frequencies were used to determine principal components, distinct tissues (gingiva, lung, and

DPI) form unique groups, whereas healthy and inflamed tissues overlap. Overall, our

unsupervised analysis approach further highlights that clusters show tissue-specific

characteristics with minimal alterations due to inflammation status.

In summary, across all analysis methods, we found that tissue-specific adaptations of T

cell subsets were maintained during tissue inflammation.

Discussion

Our main goal was to assess the extent to which tissue-specific adaptation of the T cell

compartment is altered during tissue inflammation. We initially sought to address this question

by examining oral mucosal (gingiva) tissue, but subsequently extended our study to include

other mucosal tissues to ensure that our conclusions would not be limited to a single tissue. In

the gingiva, we observed that markers indicative of inflammation-driven T cell activation did not

significantly change in most T cell populations, including MAIT and $\gamma\delta$ T cells, as inflammation

severity of the gingiva increased. We considered that this could be due to the baseline level of

inflammation that is present even in healthy gingival tissues[35], which could yield unique

regulatory mechanisms in the tissue. Since the oral mucosa is colonized by a dynamic

microbiome and continuously exposed to dietary and microbial antigens[35], we also examined

the expression of biomarker pairs that indicate TCR-mediated activation. ICOS and PD-1, and

CD39 and PD-1 have been shown to be indicative of recent TCR activation for human CD4 and

CD8 T cells, respectively[39-41]. Neither of these biomarker pairs indicated an association of a

TCR-mediated T cell response in CD4 Tconv or a CD8 T cell subset gingiva with increased

severity in inflammation. Similarly, Ki67 expression, an indicator of cell proliferation, and

expression of PD-1 and TIM3, indictors of T cell exhaustion, did not change with increased

inflammation in the gingiva. Overall, our observations highlight that the overall composition and

the phenotypic properties of T cell subsets in the gingiva were largely maintained at various

states of inflammation.

Because of the unique physiological demands of the gingiva, including but not limited to

mastication, we asked if the T cell compartment would be similarly resistant to inflammation-

mediated changes in other mucosal tissues, specifically the lung and the decidual-placental

interface (DPI). CD4 Tconv and CD8 T cell subsets in the healthy lung did not express

biomarker pairs of TCR activation. However, in the DPI we found that a substantial fraction of

CD4 and CD8 T cell subsets co-expressed ICOS and PD-1, and CD39 and PD-1, respectively

(*Supplemental Figure 8*). These data could suggest that T cells in the DPI encounter their

antigen more frequently than T cell in the gingiva and lung. This could potentially be related to

the presence of fetal-derived, semi-allogenic tissue (the placenta is made of fetal-derived

trophoblasts) or could indicate that recently activated T cells migrate to the DPI to provide

protection. An in-depth analysis of the TCR repertoire in the DPI compared to other tissues may

provide a more definitive answer in future studies. Finally, ICOS, CD39 and PD-1 are all

expressed by highly activated Tregs, and expression of these markers was found on a large

fraction of Tregs across all tissues and inflammation states.

Given the presence of highly activated Tregs in all tissues, we explored whether Tregs

displayed functional differences. We found that in gingival tissues, regardless of inflammation

status, about 10% of Tregs expressed IL-17A. IL-17+ Treg have been previously described and

are thought to maintain their ability to suppress other T cells, while secreting IL-17 in response

to TCR-mediated activation, or IL-6 and IL-1β stimulation[48,49]. Importantly, both IL-6 and IL-1β

are expressed in healthy human gingiva[50,51]. In contrast, we did not observe IL-17+ Tregs in

lung or DPI. However, we found that a fraction of Tregs in both healthy and inflamed DPI could

express IFNγ, while none of the Tregs in other tissues expressed IFNγ following stimulation.

These cells could be Th1-like Tregs that have so far been mainly described in peripheral blood

in context of human autoimmune diseases and are thought to be hypo-suppressive[52]. Thus, our

observation could indicate that these Tregs also exist in a healthy human tissue (albeit from a

unique environment that contains a semi-allogeneic tissue), but it important to consider that

these could potentially be recently activated Th1 cells that temporarily express FoxP3[53-55].

Alternatively, these IFN γ^+ Tregs could be specialized to suppress Th1 responses as previously

shown in a mouse model system[56]. Follow up studies will be required to distinguish between

these possibilities and transcriptionally define this DPI-enriched Treg population. Overall, tissue-

associated phenotypes and functional properties were evident for Tregs, while inflammation-

associated changes were limited in our cross-tissue and inflammation-state comparison.

One of the most prominent phenotypic changes that we observed between a healthy and

inflamed tissue was the striking decrease in CD25 expression on T cell subsets in the inflamed

lung (Figure 4C, *Supplemental Figure 7*). Our PBMC batch control (*Supplemental Figure 3*)

allowed us to rule out technical reasons for these phenotype (wrong titer of the antibody, etc.).

Previous reports indicated an increase in serum soluble CD25 levels in individuals with ILD

(both granulomatous-lymphocytic and rheumatoid-arthritis associated ILD)[57,58]. ILD has also

been clinically associated with increased matrix metalloproteinase (MMP) expression in

serum[59,60], which cleaves CD25 from the surface of T cells. If there is a local increase in MMP

activity in ILD lungs, then this could provide an explanation of the decreased CD25 during ILD in

all T cell subsets.

We specifically looked for functional changes that have previously been associated with

inflammation status. Studies using gingival tissue indicated a potential association between

inflammation and the abundance of Th17 cells[61,62]. We found that the overall composition of the

T cell compartment, including the frequency of CD4 T cells that make IL-17A or IL-17F following

stimulation, is relatively stable even during inflammation. It is important to keep in mind that a

hallmark feature of inflammation is the recruitment of immune cell populations to a site of

inflammation, and that inflammation of the gingiva is associated with an increase in T cell

infiltration[63]. Thus, while the overall number of T cells with a specific function increases in

inflamed tissues, this is not necessarily associated with a skewing in functional properties within

the T cell compartment. Overall, the composition of the T cell compartment was consistent

between mildly to severely inflamed gingiva, with few subsets showing an increase or decrease

with inflammation. Specifically, we observed a reduction in resident CD8 CD103+ T cells within

more severely inflamed gingival tissue with a concomitant proportional increase in circulating

CD8 CD103- T cells. Of note, our study does not suggest that stability is always maintained.

This stable maintenance of function would presumably change, at least temporarily, in disease-

specific conditions with an associated antigen-specific T cell response, such as the Th17-driven

response against *Candida albicans* and *Staphylococcus aureus*[64].

Overall, a state of tissue inflammation does not per se override tissue-specific features

of the T cell compartment. Importantly, our observations for the T cell compartment do not

necessarily extend to inflammation-associated alterations of other cell types, particularly innate

immune cells, in these tissues. In the gingiva, it is well documented that neutrophil infiltration is

associated with acute inflammation[61,65,66]. In the lungs, similar innate subsets of macrophages

and neutrophils have been reported to drive inflammatory responses in mice and humans[67,68],

and acute chorioamnionitis has also been defined by neutrophil infiltration[69]. Again, it is

important to keep in mind that specific diseases can profoundly impact T cell phenotype and

function during inflammation. Thus, while our data indicate that tissue inflammation is not

inherently sufficient to override tissue-specific T cell adaptation, different disease settings may

display profound T cell changes such as previously observed in the context of Hepatitis A

infection, which leads to increased IL-15 availability and bystander activation of CD8 T cells[27].

Finally, we want to highlight that we used a brief PMA/Ionomycin stimulation condition to

examine the functional capacity of T cells. We reasoned that the broadly activating nature of this

stimulus will reveal which proteins T cell subsets are able to express, while keeping in mind that

it does not reveal which proteins are secreted in situ. This approach indicated that T cell subsets

retained their ability to express AREG even in chronically inflamed tissue such as the ILD lung

and severely inflamed gingiva. Whether AREG is actually secreted *in situ* by T cells to promote

epithelial cell recovery in the inflamed gingiva is unclear. CD8 T cells appear to require a TCR

signal to secret AREG[34] and given the lack of evidence for CD8 T cell activation (including TCR-

mediated T cell activation) in these tissues, *in situ* AREG secretion by CD8 T cells seems

unlikely. However, given that the functional capacity is maintained in inflamed tissues, novel

therapies could target AREG to elicit expression or other specific functional responses.

In summary, we report that T cells are able to maintain their tissue-specific core

identities during states of tissue inflammation. By gaining a more complete understanding of

how inflammation affects the tissue-specific T cell landscape, therapeutic approaches can be

designed to develop treatments to restore homeostasis with tissue-specific requirements in

mind.

Material and Methods

Approvals & Study Population

Tissue Samples and Protocols: All protocols used in this study were approved by the Institutional Review Board (IRB #0008335, 00008263, 00004036, 00001636). Signed informed consent was obtained from all participants. A list of all samples, along with relevant clinical information can be found in Supplementary Table 1. Each sample is from a unique donor.

Oral Mucosa: Oral mucosa samples were obtained from surgically discarded tissue from routine surgeries, such as osseous surgery, functional or esthetic crown lengthening procedures, tooth extractions, or impacted tooth uncover procedures. Mild inflammation (clinical health, mild to moderate gingivitis), moderate inflammation (severe gingivitis, stage I, and stage II periodontitis), and severe inflammation (stage III and IV periodontitis).

Lung: Healthy lung samples were redirected transplants from deceased organ donors. Donors without history of lung disease, infection, and mechanical ventilation for >7 days were eligible for screening. Inflamed lung samples (interstitial lung disease, ILD) were obtained from patients undergoing lung transplant. Explant lungs and hilar lymph nodes were obtained from patients with ILD undergoing lung transplant at the University of Washington.

Decidual-Placenta Interface: Both healthy and acutely inflamed (chorioamnionitis) third trimester placentas were obtained following vaginal delivery or C-section, following labor or no labor (detailed in Supplementary Table 1).

Technical Control Sample: A consistent leukopak peripheral blood mononuclear cell (PBMC) sample was used throughout each flow cytometry batch and assessed alongside experimental

samples to ensure longitudinal control. Leukopak (4510-01 Ful Leuko Pack) obtained through

Bloodworks Bio (Seattle, WA).

Isolation of Lymphocytes from Tissues

Oral Mucosa: Following procedures, mucosal tissue was priority shipped overnight with cold

packs to Fred Hutch Cancer Center (FHCC). Upon arrival, tissue was washed with complete

media (RPMI1640 (11875-093, Gibco) with penicillin, streptomycin (15140-122, Gibco), and 10%

fetal bovine serum (FBS, PS-FB2, Peak Serum)). An optimized protocol was adapted from

reference[70], originally developed for processing solid tumors. Tissue was placed in a petri dish

and minced with a scalpel, then enzymatically digested (Collagenase II (C6885, Sigma-Aldrich),

DNAse (20 units/ml, #D502, Sigm) in complete media with reduced 7.5% FBS (RP7.5) in a 25 mL

volume at 37°C for 40 minutes while shaking. Following digest, the sample was passed through

a needle (16x1 ½ Blunt, 302833, BD) 10 times prior to passing through a 100-micron cell strainer.

Remaining tissue pieces were mashed using a syringe plunger, washed with complete media,

and frozen in 1 mL Recovery Cell Culture Freezing Medium (12648010, Gibco) and stored in

liquid nitrogen until future experiments. Cell viability and abundance were quantified prior to

freezing by staining a small sample aliquot (Guava antibodies 7-AAD, CD19, and CD45) for 15

minutes, resuspension in 1% PFA, and collection on the Guava easyCyte. Oral mucosa samples

in this study contained between 731-52,195 T cells, as analyzed by flow cytometry.

Lung: Lung tissue was processed in a similar fashion to the gingiva using the RP7.5 digestion for

50 minutes at 37°C while shaking at 225 RPM. Subsequently cells were washed with RP10,

strained through a 70µm filter and enumerated. Lung tissue samples contained between 1,537-

62,719 T cells.

Decidual-Placenta Interface: Tissue was processed as previously described[71]. Briefly, intact

placentas were biopsied up to a 0.5 cm depth on the maternal facing surface of the placenta

including both decidua basalis and placenta villi. Tissue was digested in digestion media (above)

and incubated at 37°C for 60 minutes with agitation. Concentration for digest was RP7.5 + 700

U/mL Collagenase Type II (Sigma-Aldrich) and 200 U/mL DNase. Cells were filtered and washed

with media. Red blood cells were removed using ACK lysis buffer, followed by resuspension with

media. Maternal blood was collected in tubes containing the anti-coagulant acid citric dextrose.

DPI tissue samples contained between 10,303-102,650 T cells.

Technical Control Sample: Leukopak was processed according to StemCell Technologies

protocol (https://www.stemcell.com/leukopak-processing-protocol.html). Two blood cell lysis

steps at room temperature and two platelet washes were performed prior to counting and freezing

20 million cells per aliquot.

Stimulation Assay

As briefly outlined in Figure 1A, upon experimentation, cryo-preserved single-cell suspension

isolated from tissues or blood (outlined above) were thawed in a 37°C water bath. 12,000 to 2

million cells per condition were plated in a U- or V-bottom 96 well plate in 200 uL complete media.

For assessing phenotype directly _ex vivo_, cells were left untreated for 2 hours at 37°C. GolgiStop

(Monensin, 554724, BD) and GolgiPlug (Protein Transport Inhibitor (Containing Brefeldin A),

555029, BD) were then added per manufacturer recommendations to compare directly to

stimulated samples and was returned to the 37°C incubator for 4 more hours (total of 6 hours).

During functional stimulation assays, samples were similarly thawed and plated. 200 uL complete

media supplemented with PMA (100 ng/mL, P8139, Sigma) and ionomycin (1000 ng/mL, I9657,

Sigma) for 6 hours, including the addition of Golgi stop/plug 2 hours after beginning the

stimulation. Upon completion, samples were stained for analysis by flow cytometry.

Spectral Flow Cytometry

Staining: For all experiments, upon completion of the assay, samples were stained previously described[72]. Prior to experimentation, the spectral flow panel was tittered, optimized, and tested to ensure proper design and maximal information could be obtained from each sample, as detailed in reference[73]. All staining steps were performed in the dark at room temperature in a 96-bottom plate. Cells were incubated with FcX block (422302, Biolegend) and Zombie NIR dye (422302, Biolegend) in PBS (14190250, Gibco) for 15 minutes. Antibody master mixes were spun down at 10,000xg for 4 minutes prior to staining to remove aggregates. Following live/dead staining, cells were washed, following by surface staining in 50uL (antibody mix in PBS + 2% FBS and 10% brilliant stain buffer (563794, BD)) for 20 minutes. Cells were then washed twice prior to fix/perm (100 uL, 554722, BD) according to manufacturer guidelines for 30 minutes. After fixation, cells were stained intracellularly for select markers noted in the panel (Supplementary Figure 1) for 30 minutes, assembled in perm/wash buffe and 10% BSB. Finally, cells were washed twice with perm/wash and resuspended in PBS + 2% FBS and stored at 4°C until acquisition the following day on the BD Discover S8.

Controls: Single-stain controls were optimized for individual markers by staining either unstimulated PBMCs, PBMCs following a 6-hour PMA/ionomycin stimulation, or unstimulated control tissue samples (generally tonsil tumor infiltrate or regional lymph nodes). For two markers expressed in low abundance in PBMC and tissue (IL-13 and IL-22), capture beads were used for single-stain control. Control samples were taken through an identical staining protocol to experimental samples, described above. A reference sample was also used within each experiment as a longitudinal control (*Supplemental Figure 3*).

Cytometer: Data from all experiments was collected with the BD Discover S8 and the

FACSChorus software platform (BD Biosciences). The S8 is equipped with 5 lasers and 78

detectors. Compensation controls were prepared and used for each individual experiment, aside

from one run containing ILD 11 and 12. Upon collection, data was exported (FCS 3.1 format) and

analyzed using FlowJo version 10.10.0. Files were analyzed in batches by collection day, with

minor gating changes across different batches.

Unsupervised Clustering

Upon acquisition on the Discover S8 (BD), samples were gated (*Supplemental Figure 1*).

Following gating of time, lymphocytes, single cells, and live cells, T cells were gated (CD3+

CD45+) and exported as a new .fcs file in FlowJo. Files were also stripped of parameters out of

scope for this analysis, focusing on time, size channels, and those containing a fluorochrome

(*listed in Supplementary Figure 1*). Upon creation of files containing only T cells, separate streams

were created for each relevant T cell subset assessed in this study: $\gamma\delta$ T cells (TCR$\gamma\delta$+), MAIT

cells (CD161+ Vα7.2+), all CD4 (CD8-), Tconv (FoxP3-), Tregs (FoxP3+ CD127-), all CD8 (CD8+

CD4-), resident CD8 cells (CD103+), and circulating CD8 cells (CD103-). Naïve cells were

included in clustering analysis for all tissues.

We included additional samples in our clustering analysis as reference tissues for tissue-

enriched T cell subsets (*Supplemental Table 1, gray lines*). Lung and associated lymph nodes,

along with maternal blood from placental donors were included in multiple batches. We also

included tumor infiltrates from head and neck squamous cell carcinoma (HNSCC) and associated

lymph nodes. One experimental batch did not include marker CD127 and was excluded from the

Tconv and Treg clustering analyses (n = 1 pediatric lung and associated LNs, 2 inflamed lungs

and associated LNs, 2 tumor infiltrate samples, and 8 tumor associated LNs).

For each clustering stream, we used up to 20,000 T cells per sample. No minimum cell

threshold was utilized. Actual T cell counts for individual samples and primary clustering streams

were considered for count analysis. The following markers were used to develop clusters: CD161,

CD39, CCR7, ICOS, CD25, IL-17F, CD28, CD27, CD45RA, PD-1, Ki67, CD69, TIM3, IFNγ,

CTLA4, IL-13, IL-22, GzmB, TNFα, IL1R1, CD137, AREG, CD40L, IL-17A, CD45RO, IL-2.

Lineage markers and QC parameters were excluded from clustering analysis: CD3, CD45, CD4,

CD8, FoxP3, CD127 or CXCR5, CD103, TCR$\gamma\delta$, Vα7.2, Live/Dead, Autofluorescence. Spectre[74]

was used to determine appropriate arcsinh transformation values for each parameter.

Repeated clustering at the single cell level – necessary to identify and factor out technical

noise, or perceived batch effect – is computationally intractable at the scale of this study.

Therefore, within each of the streams, to arrive at globally defined clusters across all batches, we

cluster the centers of clusters from batch-specific clusterings[75,76]. Specifically, we 'over'-cluster

within each batch by first clustering, then subclustering the cells within each cluster. Next, we

compute the centroids of all (sub)clusters in each batch and cluster these (sub)cluster centroids

(or "metacells" or "supercells") to arrive at a global (meta)clustering assignment for each metacell

from each batch[77]. At each stage we use the Rphenograph algorithm, with Leiden clustering of

shared nearest-neighbor graphs[78,79]. To address moderate drift across batches in some

parameters, we first align repeated control median transformed intensity values across batches

for select parameters, and next apply the same batch- and parameter-specific shifts to the

samples before metaclustering. We use Uniform Manifold Approximate Projection (UMAP) for

dimension reduction and data visualization within an iterative process of adjusting for perceived

batch effects, optimized by refining the metaclustering at a fixed Leiden resolution, before pushing

resolution higher[80].

Statistical Analyses

Statistical testing was performed using GraphPad Prism Software version 10.5. Testing us noted within figure legends. All bar graphs represent the mean with standard deviation. Heatmaps display the mean for each marker for a given sample group. For oral mucosa samples (*Figures 1-3*), significance was determined by testing using ordinary one-way ANOVA with Tukey's multiple comparisons test. For lung and placenta data (*Figures 4-5*), 2-way ANOVA was utilized with Tukey's multiple comparisons if comparing two tissues across inflammation states, or one-way ANOVA if comparing with three tissues with grouped inflammation. All data using one-way ANOVA was also assessed using the Benjamini-Hochberg false discovery rate (FDR) correction with a rate of 10% and maintained statistical significance. Supplementary Table 2 used 2-way ANOVA to compare lung and placenta across inflammation; the Benjamini-Hochberg FDR correction with a rate of 10% was utilized.

Resource Availability

Raw flow cytometry files (.fcs) were submitted to ImmPort and assigned as Study SDY3484.

Acknowledgements

We thank the Flow Cytometry Shared Resources Core at Fred Hutch Cancer Center. We thank M. Quinn Peters who provided and assisted with the R script used for PCA and Veronica Davé who consulted on proper statical testing and corrections to be used for these analyses. We also thank all members of the Prlic Lab for critical feedback and discussion.

Author contributions

Conceptualization: N.B.P., M.P. Methodology: N.B.P., H. R. M., M.P. Conducting experiments: N.B.P., E. D. V., A. K. T., C. S. D., G. D. Formal analysis: N.B.P., H. R. M., M.P. Writing; original draft: N.B.P., M.P. Writing; review and editing: all authors. Visualization: N.B.P., H. R. M., M.P. Funding acquisition: N.B.P, M.P. Supervision: M.P.

Funding

NIH grants R56DE032009, R01AI123323 and R01AI179712 (to M.P) and funding from the Oral Mucosa Immunology Consortium at the University of Nebraska (to N.B.P, Windsweep Farm Scholar) supported this work.

Declarations of interest

The authors declare no competing interests.

617   ## Supplemental Information

618   Document S1. Figures S1-S14.

619   Tables S1-3 contain tissue information, data range of heatmaps, and statistical testing.

 Figure Legends

Figure 1: Inflammation in the oral mucosa modestly affects composition and phenotype of

the T cell compartment.

A. Experimental layout, described in detail in methods. Briefly, oral mucosa tissue of varying

inflammation states was sent from the clinic to lab, kept at 4°C. Tissue was digested mechanically

and enzymatically prior to storage of cryovials with single cell suspension in liquid nitrogen. Upon

experiment, samples were thawed and assessed directly ex vivo (*Figures 1-2*) or following a 6h

PMA/I stimulation at 37°C (*Figure 3*). **B.** Frequency of CD3+ T cells of CD45+ cells across

inflammation state, pre-gated on lymphocytes, single cells, and live (*Supplementary Figure 1*). **C.**

Of CD3+ CD45+ cells, frequency of each T cell subset. **D.** Frequency of subset (left to right,

CD8+ CD103+ resident, and CD8+ CD103- circulating) of CD3+ CD45+ T cells across

inflammation state. For **B-D.** T cell subsetting, n = 9 mild, 13 moderate, 11 severely inflamed oral

mucosa tissues. **E.** For each conventional subset, heatmaps depict frequency of activation

markers across inflammation status. Phenotyping of T cell subsets must meet minimum 25 cell

threshold; sample size is consistent with above unless noted; n = 8 mild, 11 moderate for Tregs.

Tconv, CD103+ and CD103- all met minimum threshold. **F.** Frequency of MAIT cells and $\gamma\delta$ T

cells of CD3+ CD45+ T cells. **G.** For innate-like T cells, frequency of activation markers across

inflammation. Minimum threshold of 25 cells; n = 7 mild, 9 moderate for MAIT cells; n = 8 mild for

$\gamma\delta$ T cells. For all plots, ns (not significant), *p < 0.05, ***p < 0.001, ****p < 0.0001 by ordinary

one-way ANOVA with Tukey's multiple comparison test.

Figure 2: Biomarkers of T cell receptor (TCR)-mediated activation do not significantly

change across inflammation states in the oral mucosa.

A. Expression of CD39+ PD-1+ (right) and ICOS+ PD-1+ (left) among all T cell subsets across

inflammation (blue, mild; yellow, moderate; red, severely inflamed). No statistical differences

within each subset across inflammation. **B.** Representative flow cytometry plots of CD39

(BUV615) by PD-1 (BV605) expression (top) and ICOS (BUV737) by PD-1 (BV605) expression

(bottom) within CD4 Tconv cells isolated from severely inflamed oral mucosal tissue. **C.** From left

to right, expression of Ki67+ and TIM3+ PD-1+ across T cell subsets across inflammation status.

No statistical differences within each subset. **D.** Representative flow cytometry plots of Ki67

(BV650) expression (top) and TIM3 (BB515) by PD-1 (BV605) expression (bottom) within CD4

Tconv cells isolated from moderately inflamed (top) or severely inflamed (bottom) oral mucosal

tissue. No significant differences by ordinary one-way ANOVA with Tukey's multiple comparison

test.

Figure 3: Functional capacity of T cell subsets is maintained in inflamed oral mucosa.

A. Within T cell subset, expression of each molecule across mild, moderate, or severely inflamed

oral mucosa following 6h PMA/I stimulation. n = 9 mild, 13 moderate, 11 severely inflamed oral

mucosa tissues, unless noted based on minimum threshold for analysis of 25 cells. For Tregs, n

= 8 mild; n = 11 moderate. No statistical differences across inflammation status for each marker.

B. Expression of AREG, IFNγ, and IL-2 within each T cell subset; no statical differences across

inflammation. **C.** Expression of each molecule within MAIT and $\gamma\delta$ T cells. **D.** Representative

gating of AREG+ GzmB+ populations within mildly inflamed oral mucosa (left) and co-expression

of AREG+ GzmB+ among CD103+ CD8+ resident T cells (right). For MAIT cells, n = 7 mild, n =

8 moderate, n = 10 severe; $\gamma\delta$ T cells, n = 7 mild, n = 10 severe. No significant differences by

ordinary one-way ANOVA with Tukey's multiple comparison test.

Figure 4: T cells from lung and decidual-placental interference (DPI) differ based on tissue

origin regardless of inflammation status

A. Among all CD3+ CD45+ cells, frequency of indicated subset within each tissue and disease state. **B.** For conventional subsets (right to left; Tconv, Treg, CD103+ CD8+ and CD103- CD8+), frequency of activation markers by tissue and disease status. Unless noted, n = 4 healthy lung, n = 10 inflamed lung, n = 10 healthy placenta, n = 8 inflamed placenta. Some samples did not meet minimum threshold of 25 cells; for Tregs, n = 3 healthy lung; within CD103+, n = 3 healthy lung for all figure panels. Within one experiment, n = 8 inflamed lung samples for Tconv and Tregs for this and remaining panels. Statistical testing between tissues is excluded from heatmaps. **C.** Frequency of GzmB+ cells among CD8 CD103- across tissue and inflammation state (healthy/mild = blue squares, moderate gingiva = yellow circles, severe/inflamed = red diamonds). **D.** Frequency of TIM3+ Tregs. **E.** Frequency of CTLA-4+ cells among resident CD8 CD103+ cells. Pre-gated as indicated in Supplemental Figure 1 prior to subsetting and single-positive gating. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001 by ordinary one-way ANOVA with Tukey's multiple comparison test.

Figure 5: Functional capacity of T cells is primarily shaped by tissue location.

A. Upon 6hr stimulation with PMA/ionomycin, expression of cytokines and tissue healing markers across T cell subsets in the lung and placenta. Unless noted, n = 4 healthy lung, n = 10 inflamed lung, n = 10 healthy placenta, n = 8 inflamed placenta. For Tconv and Tregs, n = 8 inflamed lung samples for this and subsequent figure panels. Statistical testing between tissues excluded from heatmaps. **B.** Within each T cell subset, expression of IFNγ (left), IL-2 (middle), and IL-17A (right). For all panels, circles = gingiva, upright triangle = lung, upside down triangle = placenta; for this and subsequent figure panels, healthy/mild = blue, moderate gingiva = yellow, severe/inflamed = red. For MAIT cells, n = 3 healthy lung and n = 9 inflamed lung samples. **C.** TNFα expression across tissues in CD4 Tconv (left) and CD8 CD103- (right), pre-gated based on Supplemental Figure 1 scheme. **D.** Frequencies of IFNγ+ IL-2+ across tissues in CD4 Tconv (left) and Tregs

(right). *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001 by ordinary one-way ANOVA with

Tukey's multiple comparison test.

Figure 6: Functional capacity of T cells across tissues and inflammation state by

dimensionality reduction analysis

Principal components determined by frequency of positive cells within gates from the following

parameters: TIM3, CD39, ICOS, CD25, CD28, CD27, PD1, Ki67, IL1R1, CD137, CD40L, CTLA4,

IL-2, IL-22, AREG, IL-17A, IFNγ, GzmB, TNFα, IL-17F. PCA plots of flow cytometry data following

6-hour PMA/Ionomycin stimulation from gingiva (pink), lung (green), and placenta (purple) during

health (blue) and inflammation (red) organized by tissue type (left) and inflammation status (right)

of **A.** CD4 Tconv, **B.** CD8 CD103+, **C.** MAIT cells, **D.** γδ T cells. Moderately inflamed oral mucosa

is grouped with inflammation category. Ellipses represent 95% confidence interval.

Figure 7: An unsupervised analysis approach to compare the T cell compartment across

tissues and inflammation.

The following markers were used to develop clusters: CD161, CD39, CCR7, ICOS, CD25, IL-17F,

CD28, CD27, CD45RA, PD-1, Ki67, CD69, TIM3, IFNγ, CTLA4, IL-13, IL-22, GzmB, TNFα, IL1R1,

CD137, AREG, CD40L, IL-17A, CD45RO, IL-2. **A.** UMAPs of clusters organized by tissue origin

and inflammation status. **B.** Bubble plots of expression of each marker within each cluster among

Tregs upon stimulation. Resolution = 0.75. **C.** Box plots of frequency of each cluster relative to

CD4 Treg counts out of total T cells from each sample. **D.** Stacked bar graphs representing cluster

breakdown for individual samples.

REFERENCES

1.   Sathaliyawala, T., Kubota, M., Yudanin, N., Turner, D., Camp, P., Thome, J.J., Bickham, K.L., Lerner,
H., Goldstein, M., Sykes, M., et al. (2013). Distribution and compartmentalization of human
circulating and tissue-resident memory T cell subsets. Immunity *38*, 187-197.
10.1016/j.immuni.2012.09.020.

2.   Kumar, B.V., Ma, W., Miron, M., Granot, T., Guyer, R.S., Carpenter, D.J., Senda, T., Sun, X., Ho,
S.H., Lerner, H., et al. (2017). Human Tissue-Resident Memory T Cells Are Defined by Core
Transcriptional and Functional Signatures in Lymphoid and Mucosal Sites. Cell Rep *20*, 2921-2934.
10.1016/j.celrep.2017.08.078.

3.   Szabo, P.A., Levitin, H.M., Miron, M., Snyder, M.E., Senda, T., Yuan, J., Cheng, Y.L., Bush, E.C.,
Dogra, P., Thapa, P., et al. (2019). Single-cell transcriptomics of human T cells reveals tissue and
activation signatures in health and disease. Nat Commun *10*, 4706. 10.1038/s41467-019-12464-
3.

4.   Connors, T.J., Matsumoto, R., Verma, S., Szabo, P.A., Guyer, R., Gray, J., Wang, Z., Thapa, P., Dogra,
P., Poon, M.M.L., et al. (2023). Site-specific development and progressive maturation of human
tissue-resident memory T cells over infancy and childhood. Immunity *56*, 1894-1909 e1895.
10.1016/j.immuni.2023.06.008.

5.   Poon, M.M.L., Caron, D.P., Wang, Z., Wells, S.B., Chen, D., Meng, W., Szabo, P.A., Lam, N., Kubota,
M., Matsumoto, R., et al. (2023). Tissue adaptation and clonal segregation of human memory T
cells in barrier sites. Nat Immunol *24*, 309-319. 10.1038/s41590-022-01395-9.

6.   Wong, M.T., Ong, D.E., Lim, F.S., Teng, K.W., McGovern, N., Narayanan, S., Ho, W.Q., Cerny, D.,
Tan, H.K., Anicete, R., et al. (2016). A High-Dimensional Atlas of Human T Cell Diversity Reveals
Tissue-Specific Trafficking and Cytokine Signatures. Immunity *45*, 442-456.
10.1016/j.immuni.2016.07.007.

7.   Niessl, J., Muller, T.R., Constantz, C., Cai, C., Nilsen, V., Rivera Ballesteros, O., Adamo, S.,
Kammann, T., Mouchtaridi, E., Gao, Y., et al. (2025). Tissue origin and virus specificity shape
human CD8(+) T cell cytotoxicity. Sci Immunol *10*, eadq4881. 10.1126/sciimmunol.adq4881.

8.   Reina-Campos, M., Monell, A., Ferry, A., Luna, V., Cheung, K.P., Galletti, G., Scharping, N.E.,
Takehara, K.K., Quon, S., Challita, P.P., et al. (2025). Tissue-resident memory CD8 T cell diversity
is spatiotemporally imprinted. Nature *639*, 483-492. 10.1038/s41586-024-08466-x.

9.   Traxinger, B.R., Richert-Spuhler, L.E., and Lund, J.M. (2022). Mucosal tissue regulatory T cells are
integral in balancing immunity and tolerance at portals of antigen entry. Mucosal Immunol *15*,
398-407. 10.1038/s41385-021-00471-x.

10.   Munoz-Rojas, A.R., and Mathis, D. (2021). Tissue regulatory T cells: regulatory chameleons. Nat
Rev Immunol *21*, 597-611. 10.1038/s41577-021-00519-w.

11.   Miragaia, R.J., Gomes, T., Chomka, A., Jardine, L., Riedel, A., Hegazy, A.N., Whibley, N., Tucci, A.,
Chen, X., Lindeman, I., et al. (2019). Single-Cell Transcriptomics of Regulatory T Cells Reveals
Trajectories of Tissue Adaptation. Immunity *50*, 493-504 e497. 10.1016/j.immuni.2019.01.001.

12.   Simon, M., Stuve, P., Schmidleithner, L., Bittner, S., Beumer, N., Strieder, N., Schmidl, C., Pant, A.,
Gebhard, C., Eigenberger, A., et al. (2024). Single-cell chromatin accessibility and transposable
element landscapes reveal shared features of tissue-residing immune cells. Immunity *57*, 1975-
1993 e1910. 10.1016/j.immuni.2024.06.015.

13.   Gray, J.I., Caron, D.P., Wells, S.B., Guyer, R., Szabo, P., Rainbow, D., Ergen, C., Rybkina, K., Bradley,
M.C., Matsumoto, R., et al. (2024). Human gammadelta T cells in diverse tissues exhibit site-
specific maturation dynamics across the life span. Sci Immunol *9*, eadn3954.
10.1126/sciimmunol.adn3954.

14. Kammann, T., Cai, C., Sekine, T., Mouchtaridi, E., Boulouis, C., Nilsen, V., Ballesteros, O.R., Muller,
T.R., Gao, Y., Raineri, E.J.M., et al. (2024). MAIT cell heterogeneity across paired human tissues
reveals specialization of distinct regulatory and enhanced effector profiles. Sci Immunol *9*,
eadn2362. 10.1126/sciimmunol.adn2362.

15. Garner, L.C., Amini, A., FitzPatrick, M.E.B., Lett, M.J., Hess, G.F., Filipowicz Sinnreich, M., Provine,
N.M., and Klenerman, P. (2023). Single-cell analysis of human MAIT cell transcriptional, functional
and clonal diversity. Nat Immunol *24*, 1565-1578. 10.1038/s41590-023-01575-1.

16. Mueller, S.N., Gebhardt, T., Carbone, F.R., and Heath, W.R. (2013). Memory T cell subsets,
migration patterns, and tissue residence. Annu Rev Immunol *31*, 137-161. 10.1146/annurev-
immunol-032712-095954.

17. Maurice, N.J., Taber, A.K., and Prlic, M. (2021). The Ugly Duckling Turned to Swan: A Change in
Perception of Bystander-Activated Memory CD8 T Cells. J Immunol *206*, 455-462.
10.4049/jimmunol.2000937.

18. Lee, H., Jeong, S., and Shin, E.C. (2022). Significance of bystander T cell activation in microbial
infection. Nat Immunol *23*, 13-22. 10.1038/s41590-021-00985-3.

19. Sekine, T., Perez-Potti, A., Nguyen, S., Gorin, J.B., Wu, V.H., Gostick, E., Llewellyn-Lacey, S.,
Hammer, Q., Falck-Jones, S., Vangeti, S., et al. (2020). TOX is expressed by exhausted and
polyfunctional human effector memory CD8(+) T cells. Sci Immunol *5*.
10.1126/sciimmunol.aba7918.

20. Maurice, N.J., Berner, J., Taber, A.K., Zehn, D., and Prlic, M. (2021). Inflammatory signals are
sufficient to elicit TOX expression in mouse and human CD8+ T cells. JCI Insight *6*.
10.1172/jci.insight.150744.

21. Berg, R.E., and Forman, J. (2006). The role of CD8 T cells in innate immunity and in antigen non-
specific protection. Curr Opin Immunol *18*, 338-343. 10.1016/j.coi.2006.03.010.

22. Berg, R.E., Cordes, C.J., and Forman, J. (2002). Contribution of CD8+ T cells to innate immunity:
IFN-gamma secretion induced by IL-12 and IL-18. Eur J Immunol *32*, 2807-2816. 10.1002/1521-
4141(2002010)32:10<2807::AID-IMMU2807>3.0.CO;2-0.

23. Kambayashi, T., Assarsson, E., Lukacher, A.E., Ljunggren, H.G., and Jensen, P.E. (2003). Memory
CD8+ T cells provide an early source of IFN-gamma. J Immunol *170*, 2399-2408.
10.4049/jimmunol.170.5.2399.

24. Chu, T., Tyznik, A.J., Roepke, S., Berkley, A.M., Woodward-Davis, A., Pattacini, L., Bevan, M.J.,
Zehn, D., and Prlic, M. (2013). Bystander-activated memory CD8 T cells control early pathogen
load in an innate-like, NKG2D-dependent manner. Cell Rep *3*, 701-708.
10.1016/j.celrep.2013.02.020.

25. Yajima, T., Nishimura, H., Ishimitsu, R., Yamamura, K., Watase, T., Busch, D.H., Pamer, E.G.,
Kuwano, H., and Yoshikai, Y. (2001). Memory phenotype CD8(+) T cells in IL-15 transgenic mice
are involved in early protection against a primary infection with Listeria monocytogenes. Eur J
Immunol *31*, 757-766. 10.1002/1521-4141(200103)31:3<757::aid-immu757>3.0.co;2-q.

26. Whiteside, S.K., Snook, J.P., Williams, M.A., and Weis, J.J. (2018). Bystander T Cells: A Balancing
Act of Friends and Foes. Trends Immunol *39*, 1021-1035. 10.1016/j.it.2018.10.003.

27. Kim, J., Chang, D.Y., Lee, H.W., Lee, H., Kim, J.H., Sung, P.S., Kim, K.H., Hong, S.H., Kang, W., Lee,
J., et al. (2018). Innate-like Cytotoxic Function of Bystander-Activated CD8(+) T Cells Is Associated
with Liver Injury in Acute Hepatitis A. Immunity *48*, 161-173 e165. 10.1016/j.immuni.2017.11.025.

28. McGeachy, M.J., Cua, D.J., and Gaffen, S.L. (2019). The IL-17 Family of Cytokines in Health and
Disease. Immunity *50*, 892-906. 10.1016/j.immuni.2019.03.021.

29. Li, J., Casanova, J.L., and Puel, A. (2018). Mucocutaneous IL-17 immunity in mice and humans: host
defense vs. excessive inflammation. Mucosal Immunol *11*, 581-589. 10.1038/mi.2017.97.

30.   Sun, L., Su, Y., Jiao, A., Wang, X., and Zhang, B. (2023). T cells in health and disease. Signal
Transduct Target Ther *8*, 235. 10.1038/s41392-023-01471-y.

31.   Singh, N., Baby, D., Rajguru, J.P., Patil, P.B., Thakkannavar, S.S., and Pujari, V.B. (2019).
Inflammation and cancer. Ann Afr Med *18*, 121-126. 10.4103/aam.aam_56_18.

32.   Turner, D.L., and Farber, D.L. (2014). Mucosal resident memory CD4 T cells in protection and
immunopathology. Front Immunol *5*, 331. 10.3389/fimmu.2014.00331.

33.   Arpaia, N., Green, J.A., Moltedo, B., Arvey, A., Hemmers, S., Yuan, S., Treuting, P.M., and
Rudensky, A.Y. (2015). A Distinct Function of Regulatory T Cells in Tissue Protection. Cell *162*,
1078-1089. 10.1016/j.cell.2015.08.021.

34.   Delacher, M., Schmidleithner, L., Simon, M., Stuve, P., Sanderink, L., Hotz-Wagenblatt, A., Wuttke,
M., Schambeck, K., Ruhland, B., Hofmann, V., et al. (2024). The effector program of human CD8 T
cells supports tissue remodeling. J Exp Med *221*. 10.1084/jem.20230488.

35.   Konkel, J.E., Cox, J.R., and Wemyss, K. (2024). Bite-sized immunology; damage and microbes
educating immunity at the gingiva. Mucosal Immunol *17*, 1141-1150.
10.1016/j.mucimm.2024.07.004.

36.   Kinane, D.F., Stathopoulou, P.G., and Papapanou, P.N. (2017). Periodontal diseases. Nat Rev Dis
Primers *3*, 17038. 10.1038/nrdp.2017.38.

37.   Paprckova, D., Salyova, E., Michalik, J., and Stepanek, O. (2023). Bystander activation in memory
and antigen-inexperienced memory-like CD8 T cells. Curr Opin Immunol *82*, 102299.
10.1016/j.coi.2023.102299.

38.   Konecny, A.J., Huang, Y., Setty, M., and Prlic, M. (2024). Signals that control MAIT cell function in
healthy and inflamed human tissues. Immunol Rev *323*, 138-149. 10.1111/imr.13325.

39.   Duhen, R., Fesneau, O., Samson, K.A., Frye, A.K., Beymer, M., Rajamanickam, V., Ross, D., Tran, E.,
Bernard, B., Weinberg, A.D., and Duhen, T. (2022). PD-1 and ICOS coexpression identifies tumor-
reactive CD4+ T cells in human solid tumors. J Clin Invest *132*. 10.1172/JCI156821.

40.   Duhen, T., Duhen, R., Montler, R., Moses, J., Moudgil, T., de Miranda, N.F., Goodall, C.P., Blair,
T.C., Fox, B.A., McDermott, J.E., et al. (2018). Co-expression of CD39 and CD103 identifies tumor-
reactive CD8 T cells in human solid tumors. Nat Commun *9*, 2724. 10.1038/s41467-018-05072-0.

41.   Simoni, Y., Becht, E., Fehlings, M., Loh, C.Y., Koo, S.L., Teng, K.W.W., Yeong, J.P.S., Nahar, R., Zhang,
T., Kared, H., et al. (2018). Bystander CD8(+) T cells are abundant and phenotypically distinct in
human tumour infiltrates. Nature *557*, 575-579. 10.1038/s41586-018-0130-2.

42.   de Goffau, M.C., Lager, S., Sovio, U., Gaccioli, F., Cook, E., Peacock, S.J., Parkhill, J., Charnock-Jones,
D.S., and Smith, G.C.S. (2019). Human placenta has no microbiome but can contain potential
pathogens. Nature *572*, 329-334. 10.1038/s41586-019-1451-5.

43.   Briana, D.D., Papaevangelou, V., and Malamitsi-Puchner, A. (2021). The jury is still out on the
existence of a placental microbiome. Acta Paediatr *110*, 2958-2963. 10.1111/apa.16048.

44.   Whiteside, S.A., McGinniss, J.E., and Collman, R.G. (2021). The lung microbiome: progress and
promise. J Clin Invest *131*. 10.1172/JCI150473.

45.   Maher, T.M. (2024). Interstitial Lung Disease: A Review. JAMA *331*, 1655-1665.
10.1001/jama.2024.3669.

46.   Jung, E., Romero, R., Suksai, M., Gotsch, F., Chaemsaithong, P., Erez, O., Conde-Agudelo, A.,
Gomez-Lopez, N., Berry, S.M., Meyyazhagan, A., and Yoon, B.H. (2024). Clinical chorioamnionitis
at term: definition, pathogenesis, microbiology, diagnosis, and treatment. Am J Obstet Gynecol
230, S807-S840. 10.1016/j.ajog.2023.02.002.

47.   Sakuishi, K., Ngiow, S.F., Sullivan, J.M., Teng, M.W., Kuchroo, V.K., Smyth, M.J., and Anderson, A.C.
(2013). TIM3(+)FOXP3(+) regulatory T cells are tissue-specific promoters of T-cell dysfunction in
cancer. Oncoimmunology *2*, e23849. 10.4161/onci.23849.

48.   Voo, K.S., Wang, Y.H., Santori, F.R., Boggiano, C., Wang, Y.H., Arima, K., Bover, L., Hanabuchi, S.,
859   Khalili, J., Marinova, E., et al. (2009). Identification of IL-17-producing FOXP3+ regulatory T cells in
860   humans. Proc Natl Acad Sci U S A *106*, 4793-4798. 10.1073/pnas.0900408106.
861   49.   Beriou, G., Costantino, C.M., Ashley, C.W., Yang, L., Kuchroo, V.K., Baecher-Allan, C., and Hafler,
862   D.A. (2009). IL-17-producing human peripheral regulatory T cells retain suppressive function.
863   Blood *113*, 4240-4249. 10.1182/blood-2008-10-183251.
864   50.   Bamashmous, S., Kotsakis, G.A., Kerns, K.A., Leroux, B.G., Zenobia, C., Chen, D., Trivedi, H.M.,
865   McLean, J.S., and Darveau, R.P. (2021). Human variation in gingival inflammation. Proc Natl Acad
866   Sci U S A *118*. 10.1073/pnas.2012578118.
867   51.   Kerns, K.A., Bamashmous, S., Hendrickson, E.L., Kotsakis, G.A., Leroux, B.G., Daubert, D.D.,
868   Roberts, F.A., Chen, D., Trivedi, H.M., Darveau, R.P., and McLean, J.S. (2023). Localized microbially
869   induced inflammation influences distant healthy tissues in the human oral cavity. Proc Natl Acad
870   Sci U S A *120*, e2306020120. 10.1073/pnas.2306020120.
871   52.   Dominguez-Villar, M., Baecher-Allan, C.M., and Hafler, D.A. (2011). Identification of T helper type
872   1-like, Foxp3+ regulatory T cells in human autoimmune disease. Nat Med *17*, 673-675.
873   10.1038/nm.2389.
874   53.   Wang, J., Ioan-Facsinay, A., van der Voort, E.I., Huizinga, T.W., and Toes, R.E. (2007). Transient
875   expression of FOXP3 in human activated nonregulatory CD4+ T cells. Eur J Immunol *37*, 129-138.
876   10.1002/eji.200636435.
877   54.   Walker, M.R., Kasprowicz, D.J., Gersuk, V.H., Benard, A., Van Landeghen, M., Buckner, J.H., and
878   Ziegler, S.F. (2003). Induction of FoxP3 and acquisition of T regulatory activity by stimulated
879   human CD4+CD25- T cells. J Clin Invest *112*, 1437-1443. 10.1172/JCI19441.
880   55.   Ziegler, S.F. (2007). FOXP3: not just for regulatory T cells anymore. Eur J Immunol *37*, 21-23.
881   10.1002/eji.200636929.
882   56.   Koch, M.A., Tucker-Heard, G., Perdue, N.R., Killebrew, J.R., Urdahl, K.B., and Campbell, D.J. (2009).
883   The transcription factor T-bet controls regulatory T cell homeostasis and function during type 1
884   inflammation. Nat Immunol *10*, 595-602. 10.1038/ni.1731.
885   57.   Galant-Swafford, J., Catanzaro, J., Achcar, R.D., Cool, C., Koelsch, T., Bang, T.J., Lynch, D.A., Alam,
886   R., Katial, R.K., and Fernandez Perez, E.R. (2024). Approach to diagnosing and managing
887   granulomatous-lymphocytic interstitial lung disease. EClinicalMedicine *75*, 102749.
888   10.1016/j.eclinm.2024.102749.
889   58.   Cao, S., Liu, X., Li, Y., Yang, Y., Cai, X., Cong, S., Li, Z., Li, Y., Hong, Y., Su, Y., et al. (2024). Serum
890   sCD25 is an indicator for rheumatoid arthritis-associated interstitial lung disease. Clin Exp
891   Rheumatol *42*, 633-641. 10.55563/clinexprheumatol/3iqvk3.
892   59.   Chen, H., Tang, J., Liang, J., Huang, D., Pan, C., Liu, S., Du, X., and Tao, L. (2023). Clinical association
893   study on the matrix metalloproteinase expression in the serum of patients with connective tissue
894   disease complicated with interstitial lung disease. Arch Rheumatol *38*, 367-374.
895   10.46497/ArchRheumatol.2023.9547.
896   60.   Sheu, B.C., Hsu, S.M., Ho, H.N., Lien, H.C., Huang, S.C., and Lin, R.H. (2001). A novel role of
897   metalloproteinase in cancer-mediated immunosuppression. Cancer Res *61*, 237-242.
898   61.   Kim, T.S., Silva, L.M., Theofilou, V.I., Greenwell-Wild, T., Li, L., Williams, D.W., Ikeuchi, T.,
899   Brenchley, L., Genomics, N.N., Computational Biology, C., et al. (2023). Neutrophil extracellular
900   traps and extracellular histones potentiate IL-17 inflammation in periodontitis. J Exp Med *220*.
901   10.1084/jem.20221751.
902   62.   Dutzan, N., Kajikawa, T., Abusleme, L., Greenwell-Wild, T., Zuazo, C.E., Ikeuchi, T., Brenchley, L.,
903   Abe, T., Hurabielle, C., Martin, D., et al. (2018). A dysbiotic microbiome triggers T(H)17 cells to
904   mediate oral mucosal immunopathology in mice and humans. Sci Transl Med *10*.
905   10.1126/scitranslmed.aat0797.

63. Woodward Davis, A.S., Roozen, H.N., Dufort, M.J., DeBerg, H.A., Delaney, M.A., Mair, F., Erickson,
J.R., Slichter, C.K., Berkson, J.D., Klock, A.M., et al. (2019). The human tissue-resident CCR5(+) T
cell compartment maintains protective and functional properties during inflammation. Sci Transl
Med *11*. 10.1126/scitranslmed.aaw8718.
64. Zielinski, C.E., Mele, F., Aschenbrenner, D., Jarrossay, D., Ronchi, F., Gattorno, M., Monticelli, S.,
Lanzavecchia, A., and Sallusto, F. (2012). Pathogen-induced human TH17 cells produce IFN-
gamma or IL-10 and are regulated by IL-1beta. Nature *484*, 514-518. 10.1038/nature10957.
65. Ginesin, O., Coyac, B.R., Doppelt-Flikshtain, O., Mayer, Y., Gabay, E., Berg, T., Bar-On, Y., and
Zigdon-Giladi, H. (2025). Macrophage Depletion Reduces Bone Loss and Alters Inflammatory
Responses: A Mouse Peri-Implantitis Model. J Clin Periodontol *52*, 898-906. 10.1111/jcpe.14120.
66. Moutsopoulos, N.M., Konkel, J., Sarmadi, M., Eskan, M.A., Wild, T., Dutzan, N., Abusleme, L.,
Zenobia, C., Hosur, K.B., Abe, T., et al. (2014). Defective neutrophil recruitment in leukocyte
adhesion deficiency type I disease causes local IL-17-driven inflammatory bone loss. Sci Transl
Med *6*, 229ra240. 10.1126/scitranslmed.3007696.
67. Misharin, A.V., Morales-Nebreda, L., Reyfman, P.A., Cuda, C.M., Walter, J.M., McQuattie-
Pimentel, A.C., Chen, C.I., Anekalla, K.R., Joshi, N., Williams, K.J.N., et al. (2017). Monocyte-derived
alveolar macrophages drive lung fibrosis and persist in the lung over the life span. J Exp Med *214*,
2387-2404. 10.1084/jem.20162152.
68. Prasse, A., Pechkovsky, D.V., Toews, G.B., Jungraithmayr, W., Kollert, F., Goldmann, T., Vollmer,
E., Muller-Quernheim, J., and Zissel, G. (2006). A vicious circle of alveolar macrophages and
fibroblasts perpetuates pulmonary fibrosis via CCL18. Am J Respir Crit Care Med *173*, 781-792.
10.1164/rccm.200509-1518OC.
69. Zhang, C., Cao, J., Xu, M., Wu, D., Li, W., and Chang, Y. (2023). The role of neutrophils in
chorioamnionitis. Front Immunol *14*, 1198831. 10.3389/fimmu.2023.1198831.
70. Leelatian, N., Doxie, D.B., Greenplate, A.R., Mobley, B.C., Lehman, J.M., Sinnaeve, J., Kauffmann,
R.M., Werkhaven, J.A., Mistry, A.M., Weaver, K.D., et al. (2017). Single cell analysis of human
tissues and solid tumors with mass cytometry. Cytometry B Clin Cytom *92*, 68-78.
10.1002/cyto.b.21481.
71. Maurice, N.J., Erickson, J.R., DeJong, C.S., Mair, F., Taber, A.K., Frutoso, M., Islas, L.V., Vigil, A.,
Lawler, R.L., McElrath, J., et al. (2025). Cytokine and metabolite networks shape T cell residency
and functionality at the term human maternal-fetal interface. J Immunol *214*, 2221-2237.
10.1093/jimmun/vkaf093.
72. Mair, F., Erickson, J.R., Frutoso, M., Konecny, A.J., Greene, E., Voillet, V., Maurice, N.J., Rongvaux,
A., Dixon, D., Barber, B., et al. (2022). Extricating human tumour immune alterations from tissue
inflammation. Nature *605*, 728-735. 10.1038/s41586-022-04718-w.
73. Mair, F., and Tyznik, A.J. (2019). High-Dimensional Immunophenotyping with Fluorescence-Based
Cytometry: A Practical Guidebook. Methods Mol Biol *2032*, 1-29. 10.1007/978-1-4939-9650-6_1.
74. Ashhurst, T.M., Marsh-Wakefield, F., Putri, G.H., Spiteri, A.G., Shinko, D., Read, M.N., Smith, A.L.,
and King, N.J.C. (2022). Integration, exploration, and analysis of high-dimensional single-cell
cytometry data using Spectre. Cytometry A *101*, 237-253. 10.1002/cyto.a.24350.
75. Putri, G.H., Howitt, G., Marsh-Wakefield, F., Ashhurst, T.M., and Phipson, B. (2024). SuperCellCyto:
enabling efficient analysis of large scale cytometry datasets. Genome Biol *25*, 89. 10.1186/s13059-
024-03229-3.
76. Bilous, M., Herault, L., Gabriel, A.A., Teleman, M., and Gfeller, D. (2024). Building and analyzing
metacells in single-cell genomics data. Mol Syst Biol *20*, 744-766. 10.1038/s44320-024-00045-6.
77. Liu, X., Song, W., Wong, B.Y., Zhang, T., Yu, S., Lin, G.N., and Ding, X. (2019). A comparison
framework and guideline of clustering methods for mass cytometry data. Genome Biol *20*, 297.
10.1186/s13059-019-1917-7.

78.   Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden: guaranteeing well-
connected communities. Sci Rep *9*, 5233. 10.1038/s41598-019-41695-z.
79.   Levine, J.H., Simonds, E.F., Bendall, S.C., Davis, K.L., Amir el, A.D., Tadmor, M.D., Litvin, O.,
Fienberg, H.G., Jager, A., Zunder, E.R., et al. (2015). Data-Driven Phenotypic Dissection of AML
Reveals Progenitor-like Cells that Correlate with Prognosis. Cell *162*, 184-197.
10.1016/j.cell.2015.05.047.
80.   Becht, E., McInnes, L., Healy, J., Dutertre, C.A., Kwok, I.W.H., Ng, L.G., Ginhoux, F., and Newell,
E.W. (2018). Dimensionality reduction for visualizing single-cell data using UMAP. Nat Biotechnol.
10.1038/nbt.4314.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7