A leukemia-derived ENL/AF9 chemical probe enhances neuronal stress resilience and ameliorates ALS phenotypes

Luca Lo Piccolo[1,5]*, Chansunee Panto[2], Ranchana Yeewa[3], Ruedeemars Yubolphan[4], Saranyapin Potikanond[4], Salinee Jantrapirom[2,4]*,

[1]Center of Multidisciplinary Technology for Advanced Medicine (CMUTEAM), Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand.

[2]*Drosophila* Centre for Human Diseases and Drug Discovery (DHD), Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand

[3]Department of Biochemistry, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand 50200

[4]Department of Pharmacology, Faculty of Medicine, Chiang Mai University, Chiang Mai, Thailand.

[5]Lead contact

*Correspondence: lopiccolo.l@cmu.ac.th

**Correspondence: salinee.jan@cmu.ac.th

SUMMARY

Chemical perturbation of chromatin reader proteins provides a precise strategy to interrogate epigenetic control of neuronal stress adaptation. ENL and AF9 are YEATS-domain acyl-lysine readers best characterized in leukemia, but their roles in neurons remain unclear. Here, we use the selective YEATS inhibitor SR-0813 to define ENL/AF9 function in neuronal stress responses across *Drosophila* and human systems. SR-0813 phenocopies genetic ENL/AF9 reduction by extending lifespan and enhancing stress tolerance *in vivo*, and improves survival of human neurons under multiple stress conditions, with the strongest effects during endoplasmic reticulum stress. Mechanistically, SR-0813 attenuates PERK-ISR signaling and reduces apoptotic commitment without broadly enhancing proteostasis capacity. Notably, its effects are highly context dependent, conferring protection in stress-signaling-driven models but reduced efficacy or detrimental outcomes under chronic aggregation or mitochondrial stress. These findings identify ENL/AF9 as modulators of stress-response dynamics and highlight YEATS-domain inhibition as a context-dependent strategy to reshape neuronal resilience.

KEYWORDS
ENL/AF9, chromatin reader, chemical probe, integrated stress response, amyotrophic lateral sclerosis.

RUNNING TITLE
ENL/AF9 modulation shapes neuronal stress resilience

1   **GRAPHICAL ABSTRACT**

29   *siRNA* or the YEATS-domain inhibitor SR-0813 suppress the acyl-lysine reader ENL/AF9,
30   extending lifespan and increasing H_2O_2 tolerance in *Drosophila*. In neurons (*Drosophila* and
31   SH-SY5Y), ENL/AF9 inhibition lowers PERK–ISR signaling and apoptosis while maintaining
32   proteostasis. Effects are context dependent: protective under UPR/ISR-dominant stress but
33   potentially detrimental under aggregation or mitochondrial stress. Created with BioRender.
34
35
36

INTRODUCTION

Neuronal identity and function rely on transcriptional programs that must be both stably maintained over long timescales and dynamically responsive to physiological and pathological stimuli. Chromatin-based regulation is now well established as a key determinant of neuronal development and plasticity, and as an important layer controlling stress-responsive transcriptional programs that shape neuronal resilience and vulnerability [1-4]. However, the specific chromatin reader modules that tune these stress programs and their tractability to selective chemical perturbation remain incompletely defined.

One class of candidates with the potential to fulfill this role is the YEATS-domain family of chromatin readers [5]. YEATS-domain proteins have emerged as selective sensors of histone acyl-lysine modifications, linking chromatin state to transcriptional output [6]. ENL (MLLT1) and its paralog AF9 (MLLT3) function within transcriptional elongation complexes, including the super elongation complex (SEC), where YEATS-domain-dependent chromatin engagement promotes RNA polymerase II pause release and amplifies stimulus-responsive gene expression [7-9]. Because histone acylation is tightly coupled to cellular metabolism through acyl-CoA availability, YEATS-domain recognition provides a potential mechanism by which metabolic state can be translated into transcriptional responses [10,11]. While these features have been extensively studied in proliferative systems, particularly in cancer, whether YEATS-domain proteins regulate stress adaptation in post-mitotic neurons remains largely unexplored.

To begin addressing this question *in vivo*, we turned to *Drosophila melanogaster*, which provides a genetically tractable system with strong conservation of chromatin regulatory mechanisms [12]. Flies encode a single ENL/AF9 ortholog, ear (here referred to as dENL/AF9), which contains a conserved YEATS domain and participates in related chromatin-associated complexes [13-15]. We previously found that genetic reduction of *dENL/AF9* extends lifespan and enhances tolerance to oxidative stress, pointing to a role in organismal resilience [16]. Notably, these effects are not shared by other YEATS-domain proteins, as perturbation of the *Drosophila* YEATS2 homolog (also known as D12) produces distinct neuronal phenotypes [17], indicating functional specificity within this protein family. Together, these observations raise the possibility that dENL/AF9 regulates neuronal stress responses through YEATS-dependent transcriptional mechanisms.

These findings prompted two key questions: whether the resilience phenotypes associated with *dENL/AF9* depletion are driven by reduced YEATS-domain activity, and whether this function is conserved in mammalian neuronal systems. To address this, we turned to SR-0813, a selective small-molecule inhibitor of the ENL/AF9 YEATS domains originally developed in the context of leukemia, where ENL-dependent transcription sustains oncogenic gene expression [18]. Acting as a chromatin-competitive ligand, SR-0813 blocks YEATS-domain binding to acyl-lysine marks on histones, thereby disrupting ENL/AF9-dependent transcriptional programs while enabling acute and reversible interrogation of ENL/AF9 chromatin reader activity without broadly targeting other YEATS-domain-containing proteins such as YEATS2 or GAS41 [19]. Although its activity toward dENL/AF9 has not been directly established, this conservation and functional similarity support the use of SR-0813 as a probe in this system when interpreted alongside genetic benchmarks. Here, we combine chemical perturbation with genetic approaches in *Drosophila* and human neuronal models to define how ENL/AF9 YEATS-domain function shapes neuronal stress adaptation across distinct physiological and disease-relevant contexts.

RESULTS

SR-0813 promotes longevity and oxidative stress resistance in *Drosophila* without intrinsic radical scavenging activity

To test whether pharmacological modulation of ENL/AF9 can recapitulate the beneficial effects observed upon genetic depletion, we administered the selective inhibitor SR-0813 to *Drosophila* and assessed phenotypes previously associated with dENL/AF9 reduction. We chose SR-0813 because it is the best-characterized, commercially accessible dual ENL/AF9 YEATS inhibitor, with high biochemical potency, strong selectivity over other YEATS domains, and a well-validated chromatin-competitive mechanism that evicts ENL from target loci without inducing protein degradation [19].

Adult flies were maintained on standard food until day 20 (D20) and subsequently transferred to food containing SR-0813 (1 or 10 µM), with regular media replacement (Figure 1A). Kaplan–Meier survival analysis revealed a significant extension of lifespan at both concentrations compared to untreated controls (Figure 1B), indicating that pharmacological inhibition phenocopies the longevity benefits of dENL/AF9 reduction.

We next examined whether SR-0813 similarly enhances tolerance to oxidative stress. Following 10 days of treatment initiated at D20, flies were exposed to hydrogen peroxide (H_2O_2) (Figure 1C). While short-term exposure (6 h) resulted in minimal mortality with no detectable effect of treatment, prolonged exposure (24 h) led to substantial lethality that was partially rescued by SR-0813, particularly at the lower concentration (Figures 1D and 1E).

Because improved survival under oxidative stress can result from direct antioxidant activity, we evaluated the radical scavenging properties of SR-0813. In both ABTS and DPPH assays, SR-0813 did not exhibit detectable scavenging activity across the tested concentrations (Figures 1F and 1G), in contrast to established antioxidants.

Together, these findings show that SR-0813 recapitulates key phenotypes associated with genetic reduction of dENL/AF9, enhancing lifespan and stress tolerance through mechanisms unlikely to involve direct antioxidant effects. Given that SR-0813 selectively targets the ENL/AF9 YEATS domain, these results suggest that the lifespan extension and increased stress tolerance observed upon dENL/AF9 depletion are at least in part mediated by reduced YEATS-domain activity and consequent transcriptional reprogramming.

SR-0813 enhances stress resilience in differentiated human neurons and attenuates apoptosis under ER and proteotoxic stress

The ability of SR-0813 to phenocopy neuronal ENL/AF9 depletion in *Drosophila* prompted us to examine whether its protective effects extend to human neuronal systems. To this end, we evaluated its impact in retinoic acid (RA)-differentiated SH-SY5Y cells, a widely used model of human neurons.

Differentiation with RA (10 µM, 3 days) established a neuronal-like phenotype suitable for downstream analyses (Figure S1). Under these conditions, treatment did not affect cell viability across the tested dose range (Figure S2), indicating a lack of intrinsic toxicity to neurons. To challenge neuronal resilience, we exposed differentiated cells to multiple stress paradigms relevant to neurodegeneration, including oxidative stress (H_2O_2), proteasomal inhibition (MG132), endoplasmic reticulum (ER) stress (tunicamycin, Tm), and mitochondrial dysfunction (rotenone) (Figure S3). Cells were co-treated with increasing concentrations of the compound during 48 h of stress exposure (Figure 2A).

Treatment enhanced neuronal survival under H_2O_2 (300 µM), MG132 (0.25 µM), and Tm (1 µM), while exacerbating toxicity under rotenone (0.25 µM) (Figures 2B–2E). The most pronounced protective effect was observed under ER stress, where increased survival was detected from low concentrations (0.1 µM) across the entire dose range tested. Under proteasomal stress, a peak protective effect was observed in the range of 0.4–1.6 µM, whereas under oxidative stress, significant benefit was limited to higher concentrations (10 µM) (Figures 2B–2E). Based on these profiles, subsequent analyses focused on Tm- and MG132-induced stress.

To assess whether the survival effects observed in the MTT assay reflect modulation of cell death pathways, we performed Annexin V/propidium iodide (PI) flow cytometry in differentiated SH-SY5Y cells, with single-stain controls used to define gating boundaries (Figures S4C–S4E).

Under ER stress, Tm induced marked apoptotic and necrotic cell death (Figures 2F–2G), accompanied by neurite disruption as observed by light microscopy (Figure S4B). Co-treatment increased the proportion of viable cells while reducing late-stage apoptosis and necrotic populations in a dose-dependent manner, with maximal protection observed at 1.6 µM (Figures 2J–2M and S4B).

We next applied the same approach to proteasomal stress. MG132 induced a more severe phenotype than Tm, with pronounced morphological disruption (Figure S5B). Co-treatment modestly increased the proportion of viable (Annexin V⁻/PI⁻) cells, with a reduction in early apoptotic populations observed primarily at the lower concentration (0.4 µM), while no clear effect was detected on late apoptosis or necrosis (Figure S5A). Thus, consistent with the viability data, the protective effect under proteasomal stress is associated with a limited attenuation of cell death, less pronounced than that observed under ER stress.

Together, these findings indicate that SR-0813 enhances neuronal stress resilience, most prominently under ER stress, in part through attenuation of apoptotic cell death, prompting further investigation into its effects on ER stress signaling pathways.

SR-0813 attenuates ROS accumulation and dampens ER stress, ISR, and apoptotic signaling

Given the pronounced protective effect observed under ER stress conditions, we next sought to define the molecular pathways underlying this phenotype. Because tunicamycin-induced toxicity is closely linked to oxidative stress and activation of maladaptive unfolded protein response (UPR)/integrated stress response (ISR) signaling, we first asked whether SR-0813 modulates these processes.

We began by assessing intracellular reactive oxygen species (ROS), a key contributor to ER stress–associated toxicity. In undifferentiated SH-SY5Y cells, tunicamycin induced a marked increase in ROS levels, as detected by CM-H_2DCFDA confocal imaging, which was progressively reduced by co-treatment in a dose-dependent manner (Figures 3A–3C). To confirm this effect in a neuronal context, ROS levels were quantified in RA-differentiated SH-SY5Y cells by flow cytometry (Figure 3D). Forward- and side-scatter gating was used to define the main cell population and exclude debris (Figure S6B). Tunicamycin-treated cells displayed increased CM-H_2DCFDA fluorescence, whereas co-treatment shifted the fluorescence distribution leftward and reduced the median fluorescence intensity (MFI), consistent with lower ROS levels (Figures 3E–3F and S6A). This effect was reproduced in an independent biological replicate, with a consistent reduction observed at 1.6 µM (Figures S6C–S6D). Together, these data indicate that SR-0813 limits oxidative stress associated with ER dysfunction.

We next investigated whether this reduction in oxidative stress was accompanied by
modulation of ER stress signaling pathways. Immunoblot analysis revealed that expression of
the ER chaperone GRP78/BiP was not altered by co-treatment (Figures 3G and 3M). In
contrast, levels of ERO1L, an oxidoreductase that contributes to oxidative protein folding
within the ER, were reduced at the higher concentration (1.6 µM) (Figures 3G and 3O).
To further dissect UPR signaling, we examined distinct branches of the response. ATF6
levels were not affected by co-treatment (Figure 3H and 3R), indicating that this arm of the
UPR remains largely unchanged. By contrast, markers of the PERK–ISR axis were selectively
attenuated. Co-treatment reduced ATF4 levels and decreased the ratio of phosphorylated
eIF2α to total eIF2α, consistent with suppression of ISR signaling (Figures 3I and 3P–3Q).
This was accompanied by a reduction in CHOP levels at 1.6 µM (Figures 3J and 3S), indicating
dampening of the maladaptive, pro-apoptotic branch of the UPR. In line with this, the Bax/Bcl-
2 ratio was decreased (Figures 3J and 3T), supporting reduced apoptotic commitment,
consistent with the reduced apoptosis observed by Annexin V/propidium iodide staining
(Figures 2F–2M).
Given the link between ER stress and proteostasis impairment, we next examined the
accumulation of ubiquitinated proteins. Co-treatment reduced the levels of polyubiquitinated
(poly-Ub) species under Tm stress (Figure 3K), suggesting improved protein quality control
capacity.
Finally, to determine whether these effects reflect a broader modulation of proteostasis
pathways, we assessed additional proteins involved in ER homeostasis and protein quality
control. The abundance of VCP and UBQLN2 was not altered by co-treatment under
tunicamycin stress (Figures 3G and 3L–3N), indicating that the observed effects occur without
widespread changes in these proteostasis-associated factors.
To determine whether these effects extend beyond ER stress conditions, we performed a
parallel analysis in cells exposed to proteasomal inhibition (MG132), where the protective
effects were more modest (Figure 2). Given that MG132 primarily perturbs proteostasis rather
than directly engaging canonical UPR signaling, we focused this analysis on protein quality
control and apoptotic markers.
Under these conditions, co-treatment did not alter VCP or UBQLN2 levels (Figure S7A–S7C),
consistent with the findings observed under Tm stress. Accumulation of poly-Ub proteins was
only mildly reduced and reached significance at the higher concentration (1.6 µM) (Figure
S7A), indicating a limited impact on proteostasis under these conditions.
Despite this, markers of apoptotic signaling were consistently attenuated, with reduced
CHOP levels and a lower Bax/Bcl-2 ratio observed upon co-treatment (Figure S7D–S7E).
Thus, while the effects under proteasomal stress are less pronounced than those observed
under ER stress, the compound retains the ability to modulate downstream stress signaling
and apoptotic commitment without altering core proteostasis regulators.
Together, these results indicate that SR-0813 reduces ER stress-associated oxidative stress
and PERK-ISR signaling, limiting apoptosis and improving proteostasis. Its effects under
proteostasis impairment are more modest but consistent, without broadly affecting core
proteostasis regulators.
43
44   **ENL/AF9 modulation exerts context-dependent effects across neurodegenerative**
45   **models with selective impact on UBQLN2-associated pathology**
46   Building on the observation that SR-0813 enhances neuronal resilience under ER stress, we
47   next sought to determine whether these effects reflect modulation of ENL/AF9 itself and to
48   what extent this pathway contributes across distinct neurodegenerative contexts. To address

this, we complemented our chemical approach with a genetic strategy in *Drosophila*, enabling direct interrogation of ENL/AF9 function independent of compound-specific effects and across multiple *in vivo* disease models.

To this end, we utilized a panel of transgenic *Drosophila* models expressing human neurodegeneration-associated proteins in the eye under the control of the GMR-GAL4 driver, which directs expression to differentiating photoreceptor neurons. In this system, expression of disease-associated proteins produces a readily quantifiable rough eye phenotype, providing a sensitive *in vivo* readout of neuronal toxicity and genetic modification. The panel encompassed models of major neurodegenerative diseases, including amyotrophic lateral sclerosis (ALS), Alzheimer's disease, Huntington's disease, Parkinson's disease, and spinocerebellar ataxia. Notably, several of these disease-associated proteins have been linked to ER stress, although the underlying mechanisms are heterogeneous and remain incompletely defined [20].

Flies expressing these human transgenes were analyzed in the presence or absence of *dENL/AF9* knockdown, where control animals carried a non-targeting RNAi construct to ensure a matched genetic background (GMR-GAL4 driving two UAS transgenes). Eye degeneration was scored relative to these controls (Figures 4A-E).

Genetic reduction of *dENL/AF9* produced context-dependent effects across neurodegenerative models. A robust suppression of retinal degeneration was observed in flies expressing $UBQLN2^{P497H}$, a mutation associated with familial ALS (Figures 4C-E). More modest improvement was detected in TDP-43^{Q331K} and $(G4C2)_{49}$ repeat models, whereas toxicity was exacerbated in polyglutamine disease models (polyQ), including ATXN1-polyQ$_{82}$ and HTT-polyQ$_{128}$ (Figures 4A-E).

The pronounced rescue of UBQLN2-associated toxicity is consistent with our cellular findings, where ENL/AF9 modulation attenuated ER stress and PERK-ISR signaling, pathways that are strongly implicated in UBQLN2-linked disease. In contrast, as ployQ toxicity is driven by a persistent burden of aggregation-prone proteins that requires sustained chaperone, autophagy, and proteasome activity, these findings suggest that ENL/AF9 inhibition may be beneficial when pathology is dominated by stress-pathway signaling and proteotoxic load, but detrimental when long-term aggregation-buffering capacity is the limiting factor.

Given the pronounced rescue observed in the UBQLN2^{P497H} model, we next sought to evaluate this effect in a more disease-relevant neuronal context. To this end, we generated a recombinant fly line carrying both elav-GAL4 and *UAS-dENL/AF9 RNAi*, enabling pan-neuronal knockdown of *dENL/AF9* (Figure S8A). Knockdown efficiency was modest under standard conditions and was slightly enhanced at 28°C, consistent with the temperature sensitivity of the UAS–GAL4 system [21], although remaining overall partial (Figures S8B-D).

We then assessed locomotor function in flies expressing UBQLN2^{P497H} in neurons in the presence or absence of *dENL/AF9* knockdown (Figures 4F-H). At 25°C, *dENL/AF9* reduction improved climbing performance at early stages of disease progression, with limited benefit at later time points. Increasing the temperature to 28°C, which enhances GAL4-driven knockdown, resulted in a modest but more sustained improvement across disease progression. Notably, UBQLN2^{P497H} flies exhibited a more severe phenotype at 28°C, which may constrain the overall magnitude of the observed rescue.

Together, these findings support a context-dependent role for ENL/AF9 in regulating neuronal stress responses and vulnerability, with selective protective effects in UBQLN2-associated pathology.

SR-0813 ameliorates neuronal dysfunction and stress signaling in a *Drosophila* UBQLN2-ALS model

Having established that genetic reduction of *dENL/AF9* confers protection in UBQLN2-associated pathology, we next tested whether pharmacological inhibition could recapitulate these effects *in vivo*. SR-0813 was administered throughout development to *Drosophila* expressing UBQLN2[P497H], and neuronal function and pathology were assessed.

Pan-neuronal expression of UBQLN2[P497H] in *Drosophila* larvae leads to measurable locomotor (crawling) impairments [20]. We therefore asked whether this phenotype could be modified by SR-0813 treatment. Larvae were therefore exposed to SR-0813 (1 or 10 μM) from early developmental stages, and locomotor behavior was evaluated at the third instar stage (Figure 5A). Treatment resulted in a robust improvement in crawling performance, with increased path length, distance traveled and average speed compared to untreated UBQLN2[P497H] larvae (Figures 5B–5E).

To determine whether this functional improvement was accompanied by structural preservation, we analyzed neuromuscular junction (NMJ) morphology, which is known to be compromised upon pan-neuronal UBQLN2[P497H] expression in flies [22,23]. SR-0813 treatment increased NMJ branch length and bouton number at muscle 4 in abdominal segment A4, indicating partial restoration of synaptic architecture (Figures 5F–5H).

We next examined whether these effects were associated with modulation of disease-relevant molecular pathways *in vivo*. Oxidative stress was assessed in adult brains of UBQLN2[P497H] at 15 days of age (Figure 6A), a stage characterized by progressive neurodegeneration and locomotor decline (Figures 4G–4H and S9). Dihydroethidium (DHE) staining revealed that treatment reduced ROS levels in UBQLN2[P497H] fly brains, as evidenced by decreased fluorescence intensity and ROS index, particularly at 10 μM (Figures 6A–6C). This effect is consistent with the attenuation of oxidative stress observed in neuronal cell models (Figures 3A–3F). Notably, UBQLN2[P497H] protein levels remained unchanged upon treatment, indicating that the observed rescue is not attributable to altered transgene expression or turnover (Figure 6D-E).

We then assessed ER stress and proteostasis-related pathways. Treatment reduced levels of the ISR transcription factor ATF4 (cryptocephalus, Crc in flies) and phosphorylation of eIF2α (Ser51), indicating attenuation of PERK–ISR signaling *in vivo*, consistent with our findings in human neurons (Figure 6D-H). In parallel, levels of the autophagy adaptor Ref(2)P and of polyubiquitinated protein species were reduced in a dose-dependent manner, consistent with decreased accumulation of undegraded autophagic cargo and a lower burden of misfolded proteins, suggestive of an overall enhancement of autophagic/proteostatic clearance (Figures 6D, 6G).

Together with the absence of detectable changes in the proteostasis factors assessed *in vitro*, these findings suggest that ENL/AF9 inhibition reduces proteotoxic burden, coincident with dampened ISR signaling.

Collectively, these findings demonstrate that pharmacological inhibition of ENL/AF9 recapitulates genetic protection *in vivo*, improving neuronal function and attenuating oxidative and ER stress signaling in a UBQLN2-associated ALS model.

SR-0813 differentially modulates locomotor decline across *Drosophila* ALS models

Building on the earlier eye-based genetic modifier analysis, which suggested beneficial interactions in multiple ALS-linked contexts, we next asked whether pharmacological inhibition

could confer protection across diverse ALS models. To address this, we evaluated the effects
of SR-0813 in a panel of *Drosophila* lines expressing ALS-associated proteins.
We first performed a preliminary characterization of locomotor dysfunction to establish a
platform spanning a range of disease severities and genetic etiologies (Figure S8). Pan-
neuronal expression of $(G4C2)_{49}$, TDP-43^{Q331K}, TDP-43^{G298S}, UBQLN2^{P497H}, and SOD1
mutants using elav-GAL4 revealed a spectrum of age-dependent impairments. TDP-43^{Q331K}
induced more pronounced deficits than TDP-43^{G298S}, whereas SOD1^{G94A} and SOD1^{A4V}
exhibited comparatively milder, progressive decline. Based on these profiles and their
translational relevance, we selected $(G4C2)_{49}$, TDP-43^{Q331K}, SOD1^{G94A}, and UBQLN2^{P497H} for
further analysis.
SR-0813 was administered from early developmental stages, and locomotor performance
was assessed longitudinally from day 5 to day 30 post-eclosion (Figure 7). Consistent with
earlier findings, SR-0813 maintained a robust protective effect in UBQLN2^{P497H} flies, improving
climbing performance across disease progression, including at later stages (Figure 7A). In
SOD1^{G94A} flies, treatment conferred a significant benefit at later time points, particularly
beyond day 15, when locomotor impairment becomes more pronounced (Figure 7B and Figure
S8). A modest but significant improvement was also observed in TDP-43^{Q331K} flies,
predominantly at later stages, although treatment did not prevent premature lethality beyond
20 days (Figure 7C).
In contrast, SR-0813 exacerbated locomotor decline in flies expressing $(G4C2)_{49}$ repeats
(Figure 7D), indicating that ENL/AF9 inhibition exerts context-dependent effects across ALS
models. Notably, this effect emerged with age, as treatment did not significantly alter
locomotor performance in young flies but worsened decline as the flies get old.
Overall, this pattern is consistent with SR-0813 reducing proteotoxic stress burden and ISR
signaling, conferring benefit in UBQLN2^{P497H} and SOD1^{G94A} models, but proving less effective
or detrimental when pathology is dominated by sustained aggregation burden requiring long-
term proteostasis support.
28
29
DISCUSSION
ENL/AF9 YEATS-domain proteins are best known as chromatin readers that promote
transcriptional elongation, yet their roles in neuronal stress adaptation have been unclear.
Building on our prior finding that dENL/AF9 depletion extends *Drosophila* lifespan and
improves stress tolerance, we used these phenotypes as benchmarks to test whether the
selective YEATS inhibitor SR-0813 elicits similar organismal outcomes, consistent with
functional engagement of the ENL/AF9 pathway in flies. *In vivo*, SR-0813 phenocopied key
aspects of *dENL/AF9* knockdown: late-onset treatment extended lifespan, enhanced
resistance to oxidative challenge, and ameliorated locomotor decline and NMJs defects in a
UBQLN2-linked ALS model. These convergent genetic and pharmacologic data support a
conserved role for ENL/AF9 in shaping stress resilience at the organismal level and motivated
deeper mechanistic studies in neuronal models.
SR-0813 showed no measurable radical-scavenging activity in cell-free assays, suggesting
its ROS-lowering effects in cells and *in vivo* may arise indirectly from YEATS inhibition
reshaping stress signaling. In cellular models, SR-0813 dampened the
PERK/eIF2α/ATF4/CHOP axis and reduced apoptotic outputs without broadly preventing UPR
induction or altering the proteostasis factors examined. These findings indicate that SR-0813
primarily tunes stress-response signaling rather than globally increasing proteostasis capacity,

consistent with its ability to mitigate degeneration in a UBQLN2-linked ALS model in which toxicity has been associated with amplified ISR engagement [24,25].

The parallel phenotypes produced by YEATS inhibition and genetic knockdown further suggest that ENL/AF9 activity can be modulated through distinct modalities, acutely by disrupting reader function or chronically by reducing expression, highlighting opportunities for future studies using orthogonal YEATS ligands or ENL-directed degradation strategies [26-28] to more comprehensively define ENL/AF9 function in neuronal contexts.

A striking feature of ENL/AF9 inhibition was its context dependence. SR-0813 was protective in paradigms driven by acute ER stress or maladaptive ISR activation, but showed reduced efficacy and in some cases detrimental effects in models associated with sustained mitochondrial dysfunction or aggregation stress, including rotenone exposure and polyglutamine- or repeat-expansion-driven toxicity. These chronic conditions impose a persistent demand for adaptation, requiring continuous engagement of mitochondrial quality control, autophagy, and other proteostasis-support pathways [29-31].

These divergent outcomes are consistent with a trade-off in which ENL/AF9 inhibition is beneficial when it limits excessive ISR output, but becomes unfavorable when long-term survival depends on maintaining transcriptional capacity for sustained adaptation. Such a balance may be particularly relevant for aggregation-prone proteins, where toxicity is strongly shaped by chaperone availability and clearance pathways, and for mitochondrial stress, where ongoing compensatory programs are required to preserve bioenergetic homeostasis [32-34].

This framework aligns with the molecular function of ENL/AF9 in transcriptional elongation complexes, where YEATS-domain-dependent recognition of histone acyl-lysine marks supports RNA polymerase II pause release and amplifies stimulus-responsive transcription [35,36]. In this view, the YEATS domain couples chromatin engagement to elongation control: acyl-lysine recognition promotes locus-specific recruitment and/or stabilization of ENL/AF9, enabling ENL/AF9-containing elongation complexes to increase transcriptional output. Because histone acylation is metabolically sensitive [37,38], ENL/AF9 may thereby link stress- and metabolism-driven chromatin states to the magnitude and persistence of stress-induced transcriptional programs. We therefore propose that ENL/AF9 inhibition modulates the amplitude and persistence of stress-induced transcriptional responses, with outcomes determined by the nature and duration of the insult.

Notably, other YEATS-domain readers appear to support distinct neuronal transcriptional functions; for example, YEATS2, the reader subunit of the ATAC histone acetyltransferase complex, has been implicated in ribosomal and metabolic gene expression [39,40], and its depletion causes neuronal stress and behavioral defects in *Drosophila* [17,41], consistent with a role in maintaining basal biosynthetic capacity. In this light, our data support a model in which YEATS readers are specialized in neurons, with YEATS2 safeguarding core biosynthetic programs and ENL/AF9 tuning stress-responsive gene amplitude.

Overall, these findings extend ENL/AF9 biology beyond its established role in transcriptional control and its current therapeutic focus in cancer, identifying YEATS-domain inhibition as a means to modulate neuronal stress adaptation. The divergent effects observed across models indicate that ENL/AF9 targeting does not confer uniform cytoprotection in neurons; rather, its impact depends on the dominant stress mechanism within a given context-beneficial when pathology is driven by excessive ISR and apoptotic signaling, but less effective or even detrimental when long-term survival relies on sustained proteostasis and metabolic adaptation. Defining the stress conditions and disease settings in which this balance can be favorably shifted will be critical for evaluating the therapeutic potential of ENL/AF9 inhibition in neurodegenerative disorders.

Limitations of the study

This study has several limitations that point to important next steps. First, although SR-0813 is a well-characterized, chromatin-competitive ENL/AF9 YEATS inhibitor, we did not directly quantify its target engagement or pharmacokinetics in Drosophila or mammalian systems, and more detailed exposure and occupancy measurements will be needed to refine dosing windows. Second, our mechanistic work was performed in RA-differentiated SH-SY5Y cells and Drosophila neurons; extending these findings to human iPSC-derived motor neurons and mammalian ALS models will be important to test the generality and translational relevance of the context-dependent effects we observe. Finally, we focused on defined ISR, proteostasis, and apoptotic markers and did not undertake genome-wide chromatin or transcriptomic profiling, so the full spectrum of ENL/AF9-regulated stress-response programs remains to be mapped.

RESOURCE AVAILABILITY

Lead contact
Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Luca Lo Piccolo (lopiccolo.l@cmu.ac.th)

Materials availability
All unique/stable reagents generated in this study are available from the lead contact with a completed materials transfer agreement.

Data and code availability
All data supporting the findings of this study are available from the corresponding author upon reasonable request. No custom code was used for data acquisition or analysis beyond standard software and publicly available packages described in the Methods.

ACKNOWLEDGMENTS

This work was supported by the Fundamental Fund (FF) from the Science, Research and Innovation Promotion Fund (TSRI), Chiang Mai University, FY2568 under Contract No. 207569. This study was partially supported by the FF, FY2568 Contract No. 207573, the Faculty of Medicine, Chiang Mai University grant no FACMED 139/2567, Health Systems Research Institute (HSRI) under grants 66-124 and 68-060, the Mid-Career Research Grant (N42A670768) of the National Research Council of Thailand and PM and other Pollutants Related NCDs from Field to Cell to Bedside (FCB) funding, Chiang Mai University under the grants PM14/2566. We thank Mr. Papon Hitmool, Mr. Sakorn Phongchankhiao, and Mr. Tawan Munwanna (Department of Pharmacology, Faculty of Medicine, Chiang Mai University) for their technical and administrative assistance. We thank Dr. Yoshida Hideki (Kyoto Institute of Technology) and the Bloomington Stock Center for providing fly strains, the Science and Educational Company Limited (SCIED) for training in confocal scanning electron microscopy.

AUTHOR CONTRIBUTIONS

Conceptualization: L.LP. and S.J.; Data curation: L.LP. and S.J.; Formal Analysis: L.LP., C.P., Ra.Y., Ru.Y. and S.J.; Funding acquisition; L.LP.; Investigation: L.LP and S.J.; Methodology: L.LP., Ra.Y., Ru.Y and S.J; Project administration: L.LP.; Resources; L.LP and S.J; Software: L.LP.; Supervision; L.LP; Validation: L.LP. and S.J.; Visualization: L.LP; Writing – original draft: L.LP.; Writing – review & editing: All authors have reviewed and approved the final version of this manuscript.

SUPPLEMENTAL INFORMATION

Document S1. Figures S1–S13, Tables S1-S4

1 **Materials and Methods**
2
***Drosophila* stocks and husbandry**
Drosophila melanogaster stocks were maintained on a standard cornmeal-yeast-glucose
medium at 22°C with 60% humidity under a 12:12 light/dark cycle unless otherwise stated. All
Drosophila stocks and full genotypes used in this study, including GAL4 drivers, UAS-RNAi
lines, and neurodegenerative disease model transgenes, were obtained from the Bloomington
Drosophila Stock Center (BDSC). GMR-Gal4 (X*)* and *GMR-Gal4;+;UAS-GFP* IR are a gift of
Prof. Yoshida (Kyoto Institute of Technology). A full list of fly lines used in this study is provided
in Table S1. The Gal4/UAS binary system was used for targeted, tissue-specific knockdown
or overexpression [42]. Crosses were performed at 25°C unless indicated; for experiments
requiring increased GAL4 activity, flies were maintained at 28°C as described. Adult flies were
collected within 24 h of eclosion and aged under standard density conditions. Pan-neuronal
knockdown or overexpression was achieved using the elav-GAL4 drivers at 25°C. The glass
multiple reporter (GMR)-GAL4 promoter was used for targeting transgene's expression to
developing eyes at 28°C. To minimize genetic background effects, all experimental flies were
backcrossed six times with the *w* strain before use.
18
Compounds and preparation
All compounds were initially prepared as concentrated stock solutions in 100% DMSO and
diluted to working concentrations immediately before use, except where noted. Retinoic acid
and tunicamycin stocks were prepared at 100 mM and 5 mg/mL, respectively. The selective
ENL/AF9 YEATS-domain inhibitor SR-0813 was prepared as 20 mM stock solutions in DMSO.
For cell-based experiments, compounds were applied at final concentrations and durations
indicated in the figure legends. For fly experiments, SR-0813 was administered by
supplementation into food at the indicated concentrations. Unless otherwise specified, drug-
containing food was replaced every 5 days. Details of all compounds and their CAS registry
numbers are provided in Supplementary Table S2.
29
Measurement of radical scavenging activity
Radical scavenging activity of compounds was assessed using complementary DPPH and
ABTS assays, as previously described [43]. Briefly the DPPH assay measured neutralization of
lipophilic free radicals in methanolic solution, while the ABTS assay evaluated both hydrophilic
and lipophilic antioxidant activity in aqueous conditions. For DPPH, a 0.2 mM solution in
methanol was incubated with test compounds in 96-well plates for 30 min in the dark, and
absorbance was measured at 517 nm. For ABTS, radical cations (ABTS•$^+$) were generated
by reaction with potassium persulfate, diluted to an absorbance of 0.70 ± 0.10 at 734 nm, and
incubated with test compounds for 6 min prior to measurement at 734 nm. In both assays,
methanol or water served as negative controls, and Trolox, N-acetylcysteine, and ascorbic
acid were included as positive controls. Radical scavenging activity was calculated as the
percentage reduction in absorbance after correction for intrinsic compound absorbance. All
measurements were performed in technical triplicates across at least three independent
experiments.
44
45 ***Drosophila* lifespan and oxidative stress assays**
46 Lifespan and oxidative stress resistance were assessed in *Drosophila melanogaster* Oregon
47 R males. Adult flies were collected within 24 h of eclosion and maintained on standard food

until day 20 post-eclosion. Flies were then transferred to food containing SR-0813 (1 or 10 µM), while control groups received food supplemented with an equivalent concentration of DMSO. For lifespan analysis, flies were housed in groups of 10 and transferred to fresh vials every 2–3 days, at which time the number of dead individuals was recorded. A total of 100 flies were analyzed per condition across four independent biological replicates. Survival was monitored until all flies had died. Survival curves were generated using the Kaplan–Meier method, and statistical significance was assessed using the log-rank (Mantel–Cox) test. For oxidative stress assays, flies were maintained on SR-0813-containing food for 10 days prior to challenge or left untreated. Flies were then starved for 1 h at 25°C in vials containing water only and subsequently transferred to vials containing 5% H_2O_2. Survival was assessed by counting the number of dead flies at 6 h and 24 h after exposure. Flies were housed in groups of 10, with a total of 100 flies per condition per replicate, and experiments were performed in three independent biological replicates. All experiments were conducted at 25°C under standard husbandry conditions.

DHE staining of fly brains for oxidative stress imaging

Reactive oxygen species (ROS) levels in adult fly brains were assessed using dihydroethidium (DHE) staining, as previously described [44]. Adult flies were collected at the indicated ages and briefly anesthetized on ice. Brains were dissected in cold phosphate-buffered saline (PBS) and immediately incubated with DHE (5 µM) in PBS for 15 min at room temperature in the dark. Following incubation, brains were washed three times in PBS to remove excess dye and mounted in PBS or mounting medium for imaging. Fluorescence was acquired using an Olympus FV3000 confocal microscope with a 10× objective, using identical acquisition settings across all conditions to allow quantitative comparison. Whole brains were analyzed. ROS levels were quantified by measuring fluorescence intensity using ImageJ software. In addition, a ROS index was calculated as total fluorescence intensity × (number of objects / mean number of objects in the control group), using CellSens software (Olympus). Background-subtracted mean fluorescence intensity was calculated for each brain and normalized to control samples. A total of 17 brains were analyzed per condition across three replicates.

***Drosophila* eye degeneration modifier assays**

Genetic modifier screening was performed as previously described [20], with minor modifications. Virgin females of the genotype *GMR-GAL4; +; dENL/AF9 IR* were crossed with males carrying *UAS-linked human neurodegenerative disease genes* (*hNDGs*) listed in Supplementary Table S1. Control crosses were performed using *GMR-GAL4; +; GFP IR* females. Crosses were maintained at 28°C, 55% humidity, under a 12 h light/dark cycle. Male F1 progeny were analyzed for suppression or enhancement of eye degeneration phenotypes relative to controls. A total of 50 males per genotype were examined across three independent parental crosses. Eye degeneration was quantified using a semi-quantitative scoring system. Flies were evaluated for loss of pigmentation, neuronal death (black spots), retinal collapse, and ommatidial fusion. Each phenotype was assigned a score based on severity: 1 point when present, increasing to 2, 3, or 4 points if the affected area exceeded 5%, 30%, or 65%, respectively. Scoring was performed independently by three investigators, and final scores were used to classify genetic interactions as suppressors, enhancers, or neutral relative to controls.

Larval crawling (locomotor) assay

Larval locomotor activity was assessed using a crawling assay as previously described [17], with
minor modifications. Third instar larvae were collected, briefly rinsed in distilled water to
remove food residue, and placed individually on agar plates. After a short acclimation period
(~1 min), larval movement was recorded for 30 s under consistent conditions. Larval
trajectories were analyzed using ImageJ (version 1.53K) with the wrMTrck plugin. Total
distance traveled and path length over the 30 s interval were quantified, and average speed
was calculated accordingly. Data were plotted for each condition. The number of larvae
analyzed per condition is indicated in the corresponding figure legends. Larvae were obtained
from three independent parental crosses. All assays were performed under standardized
conditions at room temperature.
11

Adult negative geotaxis (climbing) assay
Adult locomotor performance was assessed using a negative geotaxis assay as previously
described [45], with minor modifications. Male flies were collected from the indicated genotypes
and maintained under standard conditions at either 25°C or 28°C, as specified in the
corresponding figures. For genetic experiments, flies expressing $UAS\text{-}UBQLN2^{P497H}$ under the
control of elav-GAL4 were combined with either $dENL/AF9\ RNAi$ ($elav{>}dENL/AF9$
$IR{>}UBQLN2^{P497H}$) or control RNAi ($elav{>}GFP\ IR{>}UBQLN2P497H$), and locomotor
performance was assessed longitudinally. For pharmacological experiments, flies expressing
neurodegenerative disease-associated transgenes ($elav{>}UBQLN2^{P497H}$, $elav{>}SOD1^{G94A}$,
$elav{>}TDP\text{-}43^{Q331}K$, or $elav{>}(GGGGCC)_{49}$) were maintained on food supplemented with SR-
0813 (0, 1, or 10 μM), with DMSO as vehicle control. At the indicated ages (5, 10, 15, 20, 25,
and 30 days post-eclosion), flies were gently transferred without anesthesia into transparent
conical tubes (10 flies per tube) and allowed to acclimate for 5 min. Flies were then tapped to
the bottom, and climbing behavior was recorded over a 5 s interval. Each assay was repeated
five times per group at 30 s intervals. Climbing performance was scored based on the height
reached within 5 s using a semi-quantitative scale: 0 (<2.0 cm), 1 (2.0–3.9 cm), 2 (4.0–5.9
cm), 3 (6.0–7.9 cm), 4 (8.0–9.9 cm), and 5 (>10 cm). A climbing index was calculated as the
weighted sum of scores divided by the total number of flies per group. Unless otherwise
specified, 100 flies were analyzed per condition across three independent biological
replicates. Performance was compared to age-matched controls.
32

Neuromuscular junction (NMJ) immunostaining and morphology quantification
Third instar larvae expressing $UBQLN2^{P497H}$ in all neurons (elav-GAL4) were treated with the
indicated compounds or vehicle control, as specified in the corresponding figures. Larvae were
dissected in hemolymph-like saline (HL3) and fixed in 4% paraformaldehyde in PBS for 30
min at room temperature, following established protocols. Samples were then blocked in PBS
containing 2% bovine serum albumin (BSA) and 0.1% Triton X-100. For immunostaining,
neuromuscular junctions were labeled using FITC-conjugated goat anti-horseradish
peroxidase (HRP) antibody (1:1000, Jackson ImmunoResearch), which marks neuronal
membranes, and mouse anti-Discs large (Dlg) antibody (1:300, clone 4F3, DSHB). After
primary incubation, samples were washed and incubated with Alexa Fluor 594-conjugated
anti-mouse secondary antibody to visualize Dlg. Preparations were mounted and imaged
using an Olympus Fluoview FV3000 confocal microscope under identical acquisition settings
across conditions. Quantification was performed at muscle 4 of abdominal segment A4,
focusing on type Ib glutamatergic motor neuron terminals. Bouton number and NMJ length
were measured using ImageJ (version 1.53k). Analyses were performed on 30 larvae derived
from two independent parental crosses, with one NMJ quantified per larva per condition.

Cell culture and neuronal differentiation

Human neuroblastoma SH-SY5Y cells were obtained from American Type Culture Collection (ATCC CRL-2266) and maintained in a 1:1 mixture of EMEM and F-12 medium supplemented with 10% fetal bovine serum and 1% penicillin–streptomycin. Cells were cultured at 37 °C in a humidified incubator with 5% CO_2 and used between passages 6–12. For neuronal differentiation, cells were seeded at 1×10^5 cells per well in 6-well plates and allowed to attach for 24 h. Differentiation was initiated by switching to medium containing 10 μM retinoic acid (RA) and maintained for 3 days, with media refreshed every 48 h. Neuronal differentiation was verified by morphological changes and expression of neuronal markers. All experiments involving cell culture were approved by the Institutional Biosafety Committee (ICB) of the Faculty of Medicine, Chiang Mai University (Thailand), under protocol number CMUIBC0268022.

Chemical treatments and stress paradigms in differentiated SH-SY5Y neurons

RA-differentiated SH-SY5Y neurons were subjected to defined chemical stressors to model distinct cellular insults. Cells were treated for 48 h with hydrogen peroxide (H_2O_2; oxidative stress), MG132 (proteasome inhibition/proteotoxic stress), tunicamycin (Tm) (endoplasmic reticulum stress), or rotenone (mitochondrial dysfunction), at concentrations indicated in figure legends. For co-treatment conditions, cells were exposed to stressors in the presence or absence of SR-0813 at the indicated concentrations. Vehicle (DMSO) controls were included in all experiments. Treatments were performed under standard culture conditions following neuronal differentiation. Cellular responses were quantified after 48 h as described in the corresponding assay sections.

Cell viability assay (MTT)

Cell viability was assessed using the MTT assay (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) in RA–differentiated SH-SY5Y neurons. Cells were seeded in 96-well plates at a density of 3.3×10^3 cells per well and differentiated for 3 days prior to treatment. Neurons were exposed to the indicated stressors (tunicamycin, H_2O_2, MG132, or rotenone) in the presence or absence of SR-0813 at concentrations and durations specified in the figure legends. Following treatment, medium was replaced with MTT solution (0.5 mg/mL) and incubated for 4 h at 37 °C. The solution was then removed, and formazan crystals were solubilized in DMSO. Absorbance was measured at 570 nm using a microplate reader. Cell viability was expressed as a percentage relative to DMSO-treated controls. Unless otherwise specified, each condition included at least six technical replicates across a minimum of two independent biological replicates.

Measurement of intracellular ROS in SH-SY5Y cells

Intracellular reactive oxygen species (ROS) levels were measured in undifferentiated and RA–differentiated SH-SY5Y cells using the fluorescent probe (5-(and-6)-chloromethyl-2′,7′-dichlorodihydrofluorescein diacetate) (CM-H2DCFDA). Cells were treated with Tm (1 μM) in the presence or absence of SR-0813 (0.4 or 1.6 μM), as indicated. For imaging-based analysis, undifferentiated cells were incubated with CM-H2DCFDA, washed, and imaged by confocal microscopy using a 20× objective under identical acquisition settings. Fluorescence intensity was quantified using CellSens (Evident) software and expressed as arbitrary units (a.u.). Experiments were performed in four independent biological replicates. For flow cytometry analysis, differentiated cells were stained with CM-H2DCFDA, harvested, and

analyzed on a DxFLEX flow cytometer (Beckman Coulter, CA, USA) with excitation at 488 nm and fluorescence detected in the FITC channel. At least 10,000 events were acquired per sample. The main cell population was identified based on forward and side scatter (FSC-A and SSC-A) to exclude debris. ROS levels were quantified as the median fluorescence intensity (MFI) of DCF. Unless otherwise specified, experiments were performed with at least three independent biological replicates, and statistical significance was defined as $p < 0.05$.

Apoptosis and necrosis analysis by Annexin V/PI flow cytometry

Annexin V/propidium iodide (PI) staining (FineTest, China) followed by flow cytometry was used to assess apoptosis and necrosis in RA–differentiated SH-SY5Y cells under tunicamycin (1 µM) or MG132-induced stress, in the presence or absence of SR-0813 (0.4 or 1.6 µM). Following treatment, cells were collected, washed with cold PBS, and resuspended in binding buffer. Samples were incubated with fluorophore-conjugated Annexin V and PI for 15 min at room temperature in the dark and analyzed without fixation. Cell populations were gated based on forward and side scatter (FSC-A and SSC-A) to exclude debris. Early apoptotic (Annexin V$^+$/PI$^-$), late apoptotic/necrotic (Annexin V$^+$/PI$^+$), and necrotic (Annexin V$^-$/PI$^+$) populations were defined based on staining profiles. Data are presented as the percentage of cells in each population, with total apoptotic cells calculated as the sum of Annexin V–positive populations. Experiments were performed with at least three independent biological replicates.

Preparation of crude extracts and Western blotting

Crude protein extracts were prepared from RA-differentiated SH-SY5Y cells and adult *Drosophila* melanogaster heads. For SH-SY5Y cells, cultures were washed with PBS, collected, and lysed in RIPA buffer supplemented with protease inhibitors. Lysates were incubated on ice, clarified by centrifugation at 14,000 × g for 20 min at 4 °C, and the supernatant was collected. Protein concentration was determined using a BCA assay. For *Drosophila*, crude extracts were prepared by homogenizing 10 adult male heads per sample in 100 µL of 4× sample buffer. Samples were obtained from three independent biological replicates, each derived from separate parental crosses. Equal amounts of protein (30 µg for SH-SY5Y lysates) or equal volume of crude exctracts (10 µL for *Drosophila* samples) were resolved by SDS–PAGE on 12% polyacrylamide gels and transferred to PVDF membranes using a semi-dry transfer system. Membranes were blocked in 5% skim milk in TBS-T (Tris-buffered saline with 0.1% Tween-20) and incubated overnight with primary antibodies (listed in Supplementary Table S3), followed by HRP-conjugated secondary antibodies (1:3000, 1 h). Membranes were developed using WesternSure Chemiluminescent Reagents (LI-COR) and imaged with the C-DiGit Blot Scanner (LI-COR). Protein levels were normalized to housekeeping controls (e.g., Actin or Lamin C), and phosphorylated proteins were normalized to their corresponding total protein levels. Band intensities were quantified using ImageJ (v1.53k). All experiments were performed with three independent biological replicates, unless otherwise indicated.

Quantitative RT-PCR

Total RNA was extracted from adult *Drosophila melanogaster* heads by pooling 40 heads per sample using the RNeasy Mini Kit (QIAGEN) according to the manufacturer's instructions. Three independent biological replicates were analyzed, each derived from separate parental crosses. RNA concentration and purity were assessed spectrophotometrically. For each sample, 1 µg of total RNA was reverse transcribed into cDNA using SensiFAST cDNA Synthesis Kit (Bioline). Quantitative PCR was performed using SensiFAST SYBR® Lo-ROX

1 Kit on a CFX Opus Real-Time PCR System (Bio-Rad), with specific primers (Supplementary
2 Table 4). Quantitative normalization for each sample was performed by using *ACTIN* (*ACT5C*)
3 or *GAPDH* as an internal control.
4
Statistical analysis
Statistical analyses were performed using GraphPad Prism 11 (GraphPad Software). Data are
presented as mean ± SD, with individual data points shown where appropriate. For
comparisons between two groups, unpaired two-tailed Student's t tests were used, with
Welch's correction applied when variances were unequal. For multiple-group comparisons,
one-way ANOVA was performed followed by the multiple-comparisons test specified in the
figure legend (Dunnett's, Šidák's, or Tukey's). When data did not meet parametric
assumptions, nonparametric tests were applied (Mann–Whitney test for two groups; Kruskal–
Wallis test with Dunn's post hoc test for multiple groups). Survival curves were analyzed using
the log-rank (Mantel–Cox) test. Sample sizes, number of biological replicates, and statistical
tests are indicated in the corresponding figure legends. Statistical significance was defined as
$p < 0.05$.
17

Conflict of Interest: No potential conflicts of interest relevant to this article exist. The funders had no role in the design of the study, the collection/analysis/interpretation of data, or the decision to submit the manuscript.

18
19

References

1. Grabowska, A., Sas-Nowosielska, H., Wojtas, B., Holm-Kaczmarek, D., Januszewicz, E., Yushkevich, Y., Czaban, I., Trzaskoma, P., Krawczyk, K., Gielniewski, B., et al. (2022). Activation-induced chromatin reorganization in neurons depends on HDAC1 activity. Cell Rep *38*, 110352. 10.1016/j.celrep.2022.110352.

2. Yap, E.L., and Greenberg, M.E. (2018). Activity-Regulated Transcription: Bridging the Gap between Neural Activity and Behavior. Neuron *100*, 330-348. 10.1016/j.neuron.2018.10.013.

3. Benito, E., and Barco, A. (2015). The neuronal activity-driven transcriptome. Mol Neurobiol *51*, 1071-1088. 10.1007/s12035-014-8772-z.

4. Gallegos, D.A., Chan, U., Chen, L.F., and West, A.E. (2018). Chromatin Regulation of Neuronal Maturation and Plasticity. Trends Neurosci *41*, 311-324. 10.1016/j.tins.2018.02.009.

5. Li, X., Liu, S., Li, X., and Li, X.D. (2023). YEATS Domains as Novel Epigenetic Readers: Structures, Functions, and Inhibitor Development. ACS Chem Biol *18*, 994-1013. 10.1021/acschembio.1c00945.

6. Li, Y., Zhao, D., Chen, Z., and Li, H. (2017). YEATS domain: Linking histone crotonylation to gene regulation. Transcription *8*, 9-14. 10.1080/21541264.2016.1239602.

7. Kabra, A., and Bushweller, J. (2022). The Intrinsically Disordered Proteins MLLT3 (AF9) and MLLT1 (ENL) - Multimodal Transcriptional Switches With Roles in Normal Hematopoiesis, MLL Fusion Leukemia, and Kidney Cancer. J Mol Biol *434*, 167117. 10.1016/j.jmb.2021.167117.

8. Wan, L., Wen, H., Li, Y., Lyu, J., Xi, Y., Hoshii, T., Joseph, J.K., Wang, X., Loh, Y.E., Erb, M.A., et al. (2017). ENL links histone acetylation to oncogenic gene expression in acute myeloid leukaemia. Nature *543*, 265-269. 10.1038/nature21687.

9. Smith, E., Lin, C., and Shilatifard, A. (2011). The super elongation complex (SEC) and MLL in development and disease. Genes Dev *25*, 661-672. 10.1101/gad.2015411.

10. Dutta, A., Abmayr, S.M., and Workman, J.L. (2016). Diverse Activities of Histone Acylations Connect Metabolism to Chromatin Function. Mol Cell *63*, 547-552. 10.1016/j.molcel.2016.06.038.

11. Sabari, B.R., Zhang, D., Allis, C.D., and Zhao, Y. (2017). Metabolic regulation of gene expression through histone acylations. Nat Rev Mol Cell Biol *18*, 90-101. 10.1038/nrm.2016.140.

12. Llorens-Giralt, P., Camilleri-Robles, C., Corominas, M., and Climent-Canto, P. (2021). Chromatin Organization and Function in Drosophila. Cells *10*. 10.3390/cells10092362.

13. Mohan, M., Herz, H.M., Takahashi, Y.H., Lin, C., Lai, K.C., Zhang, Y., Washburn, M.P., Florens, L., and Shilatifard, A. (2010). Linking H3K79 trimethylation to Wnt signaling through a novel Dot1-containing complex (DotCom). Genes Dev *24*, 574-589. 10.1101/gad.1898410.

14. Smith, E.R., Lin, C., Garrett, A.S., Thornton, J., Mohaghegh, N., Hu, D., Jackson, J., Saraf, A., Swanson, S.K., Seidel, C., et al. (2011). The little elongation complex regulates small nuclear RNA transcription. Mol Cell *44*, 954-965. 10.1016/j.molcel.2011.12.008.

15. Yeewa, R., Chaiya, P., Jantrapirom, S., Shotelersuk, V., and Lo Piccolo, L. (2022). Multifaceted roles of YEATS domain-containing proteins and novel links to neurological diseases. Cell Mol Life Sci *79*, 183. 10.1007/s00018-022-04218-0.

16. Yeewa, R., Pohsa, S., Yamsri, T., Wongkummool, W., Jantaree, P., Potikanond, S., Nimlamool, W., Shotelersuk, V., Lo Piccolo, L., and Jantrapirom, S. (2024). The histone acylation reader ENL/AF9 regulates aging in Drosophila melanogaster. Neurobiol Aging *144*, 153-162. 10.1016/j.neurobiolaging.2024.10.002.

17. Lo Piccolo, L., Yeewa, R., Pohsa, S., Yamsri, T., Calovi, D., Phetcharaburanin, J., Suksawat, M., Kulthawatsiri, T., Shotelersuk, V., and Jantrapirom, S. (2024). FAME4-associating YEATS2 knockdown impairs dopaminergic synaptic integrity and leads to seizure-like behaviours in Drosophila melanogaster. Prog Neurobiol *233*, 102558. 10.1016/j.pneurobio.2023.102558.

18. Garnar-Wortzel, L., Bishop, T.R., Kitamura, S., Milosevich, N., Asiaban, J.N., Zhang, X., Zheng, Q., Chen, E., Ramos, A.R., Ackerman, C.J., et al. (2021). Chemical Inhibition of ENL/AF9 YEATS Domains in Acute Leukemia. ACS Cent Sci *7*, 815-830. 10.1021/acscentsci.0c01550.

19. Erb, M.A. (2023). Small-molecule tools for YEATS domain proteins. Curr Opin Chem Biol *77*, 102404. 10.1016/j.cbpa.2023.102404.

20. Yeewa, R., Sangphukieo, A., Jantaree, P., Wongkummool, W., Yamsri, T., Poompouang, S., Chaiyawat, P., Lo Piccolo, L., and Jantrapirom, S. (2024). ERO1A inhibition mitigates neuronal ER stress and ameliorates UBQLN2(ALS) phenotypes in Drosophila melanogaster. Prog Neurobiol *242*, 102674. 10.1016/j.pneurobio.2024.102674.

21. Duffy, J.B. (2002). GAL4 system in Drosophila: a fly geneticist's Swiss army knife. Genesis *34*, 1-15. 10.1002/gene.10150.

22. Kim, S.H., Nichols, K.D., Anderson, E.N., Liu, Y., Ramesh, N., Jia, W., Kuerbis, C.J., Scalf, M., Smith, L.M., Pandey, U.B., and Tibbetts, R.S. (2023). Axon guidance genes modulate neurotoxicity of ALS-associated UBQLN2. Elife *12*. 10.7554/eLife.84382.

23. Kim, S.H., Stiles, S.G., Feichtmeier, J.M., Ramesh, N., Zhan, L., Scalf, M.A., Smith, L.M., Pandey, U.B., and Tibbetts, R.S. (2018). Mutation-dependent aggregation and toxicity in a Drosophila model for UBQLN2-associated ALS. Hum Mol Genet *27*, 322-337. 10.1093/hmg/ddx403.

24. Halloran, M., Ragagnin, A.M.G., Vidal, M., Parakh, S., Yang, S., Heng, B., Grima, N., Shahheydari, H., Soo, K.Y., Blair, I., et al. (2020). Amyotrophic lateral sclerosis-linked UBQLN2 mutants inhibit endoplasmic reticulum to Golgi transport, leading to Golgi fragmentation and ER stress. Cell Mol Life Sci *77*, 3859-3873. 10.1007/s00018-019-03394-w.

25. Gu, A., Zhang, Y., He, J., Zhao, M., Ding, L., Liu, W., Xiao, J., Huang, J., Liu, M., and Liu, X. (2024). Chronic Oxidative Stress and Stress Granule Formation in UBQLN2 ALS Neurons: Insights into Neuronal Degeneration and Potential Therapeutic Targets. Int J Mol Sci *25*. 10.3390/ijms252413448.

26. Xue, Z., Qin, L., Xuan, H., Luo, K., Huang, M., Xie, L., Su, Y., Xu, L., Harsh, J., Dale, B., et al. (2024). A potent and selective ENL degrader suppresses oncogenic gene expression and leukemia progression. Sci Adv *10*, eado1432. 10.1126/sciadv.ado1432.

27. Asiaban, J.N., Milosevich, N., Chen, E., Bishop, T.R., Wang, J., Zhang, Y., Ackerman, C.J., Hampton, E.N., Young, T.S., Hull, M.V., et al. (2020). Cell-Based Ligand Discovery for the ENL YEATS Domain. ACS Chem Biol *15*, 895-903. 10.1021/acschembio.0c00124.

28. Li, X., Yao, Y., Wu, F., and Song, Y. (2022). A proteolysis-targeting chimera molecule selectively degrades ENL and inhibits malignant gene expression and tumor growth. J Hematol Oncol *15*, 41. 10.1186/s13045-022-01258-8.

29. Mader, B.J., Pivtoraiko, V.N., Flippo, H.M., Klocke, B.J., Roth, K.A., Mangieri, L.R., and Shacka, J.J. (2012). Rotenone inhibits autophagic flux prior to inducing cell death. ACS Chem Neurosci *3*, 1063-1072. 10.1021/cn300145z.

30. Cortes, C.J., and La Spada, A.R. (2015). Autophagy in polyglutamine disease: Imposing order on disorder or contributing to the chaos? Mol Cell Neurosci *66*, 53-61. 10.1016/j.mcn.2015.03.010.

31. Chitiprolu, M., Jagow, C., Tremblay, V., Bondy-Chorney, E., Paris, G., Savard, A., Palidwor, G., Barry, F.A., Zinman, L., Keith, J., et al. (2018). A complex of C9ORF72

and p62 uses arginine methylation to eliminate stress granules by autophagy. Nat Commun *9*, 2794. 10.1038/s41467-018-05273-7.

32. Tenchov, R., Sasso, J.M., and Zhou, Q.A. (2024). Polyglutamine (PolyQ) Diseases: Navigating the Landscape of Neurodegeneration. ACS Chem Neurosci *15*, 2665-2694. 10.1021/acschemneuro.4c00184.

33. Tittelmeier, J., Nachman, E., and Nussbaum-Krammer, C. (2020). Molecular Chaperones: A Double-Edged Sword in Neurodegenerative Diseases. Front Aging Neurosci *12*, 581374. 10.3389/fnagi.2020.581374.

34. Zong, Y., Li, H., Liao, P., Chen, L., Pan, Y., Zheng, Y., Zhang, C., Liu, D., Zheng, M., and Gao, J. (2024). Mitochondrial dysfunction: mechanisms and advances in therapy. Signal Transduct Target Ther *9*, 124. 10.1038/s41392-024-01839-8.

35. Hsu, E., Zemke, N.R., and Berk, A.J. (2021). Promoter-specific changes in initiation, elongation, and homeostasis of histone H3 acetylation during CBP/p300 inhibition. Elife *10*. 10.7554/eLife.63512.

36. Erb, M.A., Scott, T.G., Li, B.E., Xie, H., Paulk, J., Seo, H.S., Souza, A., Roberts, J.M., Dastjerdi, S., Buckley, D.L., et al. (2017). Transcription control by the ENL YEATS domain in acute leukaemia. Nature *543*, 270-274. 10.1038/nature21688.

37. Jo, C., Park, S., Oh, S., Choi, J., Kim, E.K., Youn, H.D., and Cho, E.J. (2020). Histone acylation marks respond to metabolic perturbations and enable cellular adaptation. Exp Mol Med *52*, 2005-2019. 10.1038/s12276-020-00539-x.

38. Nitsch, S., Zorro Shahidian, L., and Schneider, R. (2021). Histone acylations and chromatin dynamics: concepts, challenges, and links to metabolism. EMBO Rep *22*, e52774. 10.15252/embr.202152774.

39. Mi, W., Guan, H., Lyu, J., Zhao, D., Xi, Y., Jiang, S., Andrews, F.H., Wang, X., Gagea, M., Wen, H., et al. (2017). YEATS2 links histone acetylation to tumorigenesis of non-small cell lung cancer. Nat Commun *8*, 1088. 10.1038/s41467-017-01173-4.

40. Zhao, J., Zhao, Z., Zhou, W., Zhang, J., Chen, J., Sun, J., and Li, J. (2025). YEATS2 O-GlcNAcylation promotes chromatin association of the ATAC complex and lung cancer tumorigenesis. J Biol Chem *301*, 110388. 10.1016/j.jbc.2025.110388.

41. Lo Piccolo L., Y.R., Noisagul P., Monteil A., Shotelersuk V, Jantrapirom S. (2025). Dopaminergic neurons are vulnerable to dysregulation of YEATS2-dependent calcium homeostasis. bioRxiv. bioRxiv 2025.09.06.674642.

42. Brand, A.H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. Development *118*, 401-415. 10.1242/dev.118.2.401.

43. Jantrapirom S., S.A., U-on N., Poonsawas P., Wongkumool W, Yeewa R., Lo Piccolo L. (2025). Discovery of a pyrazolopyridine alkaloid inhibitor of ERO1A that mitigates neuronal ER stress and age-related decline. bioRxiv 2025.07.14.664722.

44. Ranchana Yeewa, S.P., Kornravee Photichai, Titaree Yamsri, Natsinee U-on, Wasinee Wongkummool, Phatsara Manussabhorn, Luca Lo Piccolo, Salinee Jantrapirom (2025). Diesel exhaust particles induce lasting and age-dependent damage to the brain in Drosophila melanogaster. biorRxiv. doi.org/10.1101/2025.08.20.671202 .

45. Jantrapirom, S., Lo Piccolo, L., Yoshida, H., and Yamaguchi, M. (2018). Depletion of Ubiquilin induces an augmentation in soluble ubiquitinated Drosophila TDP-43 to drive neurotoxicity in the fly. Biochim Biophys Acta Mol Basis Dis *1864*, 3038-3049. 10.1016/j.bbadis.2018.06.017.

Figure 1   SR-0813 phenocopies genetic ENL/AF9 inhibition to enhance stress resistance in *Drosophila* without direct antioxidant activity. (A) Experimental paradigm for lifespan analysis. Oregon R flies were aged on standard food until day 20 (D20) and then transferred to food containing SR-0813 (1 or 10 µM) or vehicle control, and survival was monitored over time. (B) Kaplan–Meier survival curves of flies treated as described in (A). Statistical significance was determined using the Mantel–Cox test. (C) Experimental paradigm for oxidative stress assays. Flies were aged to D20, treated with SR-0813 (1 or 10 µM) for 10 days, and subsequently exposed to hydrogen peroxide (H_2O_2) 5% for 6 or 24 h before survival assessment. (D) Survival following 6 h (H_2O_2) exposure under the indicated conditions. (E) Survival following 24 h (H_2O_2) exposure under the indicated conditions. Individual values are shown with mean ± SD. (F–G) Radical scavenging activity of SR-0813 measured using (F) 2,2′-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid (ABTS) or (G) 2,2-diphenyl-1-picrylhydrazyl (DPPH) assays across increasing concentrations, with vitamin C (vit C), Trolox, and N-acetylcysteine (NAC) as positive controls. For survival assays, n = 100 flies per condition. Statistical significance in (D) and (E) was determined using one-way ANOVA with appropriate multiple-comparisons testing. Data are presented as mean ± SD.

19
20

1
Figure 2 SR-0813 protects differentiated SH-SY5Y cells from ER and proteotoxic stress–
induced loss of viability. (A) Experimental paradigm. SH-SY5Y cells were differentiated with
retinoic acid (RA; 10 μM) for 3 days and then exposed for 2 days to H_2O_2 (300 μM), MG-132
(0.25 μM), tunicamycin (Tm; 1 μM), or rotenone (0.25 μM), with vehicle (DMSO) or SR-0813
(0–10 μM). (B–E) MTT viability assay of differentiated SH-SY5Y cells exposed to (B) H_2O_2,
(C) MG-132, (D) Tm, or (E) rotenone in the presence of increasing SR-0813 concentrations.
Each dot represents one technical replicate; data are from 3 independent experiments. (F–I)
Representative Annexin V/propidium iodide (PI) flow cytometry plots from cells treated with
(F) DMSO, (G) Tm, (H) Tm + SR-0813 (0.4 μM), or (I) Tm + SR-0813 (1.6 μM). (J–M)
Quantification of Annexin V/PI populations showing (J) live cells (Annexin V^-/PI^-), (K) early
apoptosis (Annexin V^+/PI^-), (L) late apoptosis (Annexin V^+/PI^+), and (M) necrosis (Annexin
V^-/PI^+). Each dot represents one technical replicate; data are from 3 independent

1 experiments. SR- indicates SR-0813. For (B–E) and (J–M), statistical significance was
2 assessed by ordinary one-way ANOVA with Sidak's multiple-comparisons test ($p < 0.05$).
3

Figure 3 SR-0813 reduces tunicamycin-induced ROS and dampens ER stress/ISR and
apoptotic signaling. (A–C) Confocal ROS imaging in undifferentiated SH-SY5Y cells. Cells
were exposed to tunicamycin (Tm; 1 μM) and co-treated with vehicle (DMSO) or SR-0813 (0.4
or 1.6 μM). Intracellular ROS was measured using CM-H$_2$DCFDA. (B) Representative CM-
H$_2$DCFDA fluorescence images (top) and corresponding phase-contrast images (bottom). (C)
Quantification of CM-H$_2$DCFDA fluorescence intensity (a.u.) shown as box-and-whisker plots
with all data points overlaid; whiskers indicate the full range (min to max). (D–F) Flow-

cytometric ROS measurements in differentiated SH-SY5Y cells. (D) Experimental design: differentiated cells were treated with Tm and co-treated with SR-0813 (0, 0.4, or 1.6 µM) or left unstressed (DMSO), then stained with CM-H$_2$DCFDA and analyzed by flow cytometry. (E) Representative histograms (count versus FITC-A). (F) Quantification of median fluorescence intensity (MFI). Points represent biological replicates from 3 independent experiments (n = 3); bars show mean ± SD. Statistical analysis was performed by one-way ANOVA with Brown–Forsythe correction; $p < 0.05$ was considered significant. (G–K) Immunoblot analysis of ER stress/ integrated stress response (ISR), proteostasis, and apoptosis markers in SH-SY5Y cells treated as in (D). β-tubulin was used as a loading control. (L–T) Densitometric quantification of immunoblots in (G–K) from 3 independent experiments (n = 3). Values were normalized to β-tubulin, except p-eIF2α normalized to total eIF2α (EIF2S1); Bax/Bcl-2 is presented as a ratio. Points represent biological replicates; bars show mean ± SD. SR- indicates SR-0813. Statistical significance was assessed by ordinary one-way ANOVA with Sidak's multiple-comparisons test; $p < 0.05$ was considered significant.

1
Figure 4 dENL/AF9 knockdown is a model-dependent modifier of neurodegenerative
phenotypes. (A) Schematic of the genetic modifier screen testing knockdown (KD) of the
Drosophila ENL/AF9 homolog (dENL/AF9) across fly models of neurodegeneration. (B–C)
Representative adult eye images from flies expressing human disease-associated proteins in
the eye under the GMR-GAL4 driver, with or without dENL/AF9 KD. Models included
Alzheimer's disease/amyloid pathology (APP-43), tauopathy (Tau2N4R), Parkinson's disease
(GBA^{N370S}, SNCA^{E46K}, SNCA^{H50Q}), polyglutamine disorders (ATXN1 polyQ82; HTT polyQ128),
and ALS/FTD (G4C2 repeats, TDP-43^{Q331K}, TDP-43^{G298S}, and UBQLN2^{P497H}). Eye
degeneration ("rough eye") served as the primary screening readout. (D–E) Quantification of
rough-eye degeneration scores for each disease model expressed under GMR-GAL4 and co-
expressed with either dENL/AF9 RNAi or a non-targeting control (NTC) inverted repeat (IR).
Bars show mean ± SD. (F) Experimental design for assessing dENL/AF9 KD in a disease-
relevant tissue. Pan-neuronal expression was driven by elav-GAL4, and flies were reared at
25°C or 28°C to modulate GAL4/UAS activity (KD efficiency at 25 and 28°C is shown in Figure
S7). (G) Negative geotaxis (climbing) performance for the indicated genotypes (elav-GAL4)
plotted as climbing score based on distance climbed at 5, 10, 15, 20, 25, and 30 days post-
eclosion. Bars show mean ± SD; a total of 100 flies were analyzed per genotype and time
point. For (D–E) and (G), statistical significance was determined by one-way ANOVA with
Dunnett's multiple-comparisons test; $p < 0.05$ was considered significant. (hNDG: human
neurodegenerative gene; NTC: non-targeting gene)
22
23

Figure 5 SR-0813 improves locomotor function and neuromuscular junction morphology in a *Drosophila* UBQLN2^{P497H} ALS model. (A) Experimental design. UBQLN2^{P497H} was expressed pan-neuronally using elav-GAL4. Parental flies were maintained on standard food supplemented with SR-0813 (1 µM; SR-1 or 10 µM; SR-10) or left untreated (UNT). Progeny were assessed at the third-instar larval stage (L3) by crawling assays, and adults were assayed for climbing performance every 5 days from day 5 to day 30 post - eclosion (D5-D30). (B–E) Larval crawling behavior. (B) Representative 30-s locomotor tracks from L3 foraging larvae (UNT, SR-1, SR-10). Quantification of (C) track length, (D) distance traveled, and (E) average speed over 30 s. Each point represents one larva (UNT n = 60; SR-1 n = 53; SR-10 n = 57). Statistical analysis: ordinary one-way ANOVA with Dunnett's multiple-comparisons test (each SR-0813 condition versus UNT); $p < 0.05$ was considered significant. (F–H) Neuromuscular junction (NMJ) morphology. (F) Representative images of type Ib NMJs on muscle 4 (abdominal segment A4) from L3 elav>UBQLN2^{P497H} larvae reared on UNT, SR-1, or SR-10 food. Presynaptic terminals were labeled with anti-HRP (green) and postsynaptic densities with anti-Discs large (Dlg; red). Quantification of (G) NMJ branch length and (H) bouton number. Each point represents one larva (n = 30 larvae per condition). Statistical analysis: ordinary one-way ANOVA with Dunnett's multiple-comparisons test (each SR-0813 condition versus UNT); $p < 0.05$ was considered significant.

1
Figure 6 SR-0813 reduces oxidative stress and dampens ISR/proteostasis stress
markers in UBQLN2^{P497H} flies. (A) Representative confocal images of adult brains stained
with dihydroethidium (DHE) to assess ROS in UBQLN2^{P497H} flies left untreated (UNT) or
treated with SR-0813 (1 µM, SR-1; 10 µM, SR-10). Flies were reared on standard food and,
after eclosion, maintained on SR-0813–supplemented food (or UNT) for 15 days prior to
imaging. (B–C) Quantification of brain ROS. (B) Mean DHE fluorescence intensity and (C)
ROS index, defined as *total intensity × (N objects / mean N objects in the control group)*,
quantified using CellSense (Olympus FLUOVIEW FV300). Each point represents one brain (n
= 17). Statistical analysis: ordinary one-way ANOVA with Sidak's multiple-comparisons test; p
< 0.05 was considered significant. (D) Immunoblot analysis of head lysates from UBQLN2^{P497H}
flies treated as in (A). HA-UBQLN2 was measured to confirm that phenotypic rescue was not
attributable to reduced UBQLN2 expression. Markers of the integrated stress response (ISR)
included *Drosophila* ATF4 (Cryptocephal, Crc) and phospho-eIF2α (p-eIF2α) with total eIF2α
(eIF2S1). Ref(2)P and ubiquitin (polyubiquitinated species) were assessed as
proteostasis/autophagy-associated readouts. β-tubulin (TUBB) served as a loading control.
(E–H) Densitometric quantification of immunoblots in (D) from 3 independent experiments (n
= 3). (E) HA-UBQLN2, (F) Crc, (G) Ref(2)P, and (H) p-eIF2α normalized to total eIF2α
(eIF2S1). Signals were normalized to β-tubulin (TUBB) unless otherwise indicated. Points
represent biological replicates; bars show mean ± SD. Statistical analysis: ordinary one-way
ANOVA with Dunnett's multiple-comparisons test (each SR-0813 condition versus UNT); p <
0.05 was considered significant.
23

Figure 7 SR-0813 differentially modulates adult locomotor decline across *Drosophila*
ALS models. Adult locomotor function was assessed longitudinally across multiple Drosophila
ALS models to determine whether SR-0813–mediated effects were sustained during
adulthood and across disease progression. Human ALS-associated genes were expressed
pan-neuronally using elav-GAL4, and flies were maintained on standard food supplemented
with SR-0813 (1 µM or 10 µM) or left untreated (UNT). Food was replaced every 5 days, and
negative geotaxis (climbing) performance was measured every 5 days from 5 to 30 days post-
eclosion (D5–D30). "x" marks indicate time points at which flies died. (A–D) Climbing
performance of (A) UBQLN2[P497H], (B) SOD1[G94A], (C) TDP-43[Q331K], and (D) (G4C2)$_{49}$ flies under
UNT, SR-0813 (1 µM), or SR-0813 (10 µM) conditions. Data are shown as bar plots with
individual values from 5 independent replicates (biological replicate vials) at each time point;
a total of 100 flies were tested per genotype/condition. Statistical analysis was performed by
ordinary one-way ANOVA with Dunnett's multiple-comparisons test (each SR-0813 condition
versus UNT); $p < 0.05$ was considered significant. Climbing performance relative to
driver/background controls is shown in Figure S8. Black "x" symbols indicate time points at
which the corresponding genotype had died (no climbing measurement possible).