1   Integrating computational chemistry and machine learning to predict

2   KRAS mutation-induced resistance

Katarzyna Mizgalska[1,2], Konstancja Urbaniak[3], Denis J. Imbody[4], Eric B. Haura[4], Wayne C.

Guida[2], Sergio Branciamore[3], Aleksandra Karolak[1]

[1]Department of Machine Learning, Moffitt Cancer Center, Tampa, FL, United States of America

[2]Department of Chemistry, University of South Florida, Tampa, FL, United States of America

[3]Department of Computational and Quantitative Medicine, City of Hope, Monrovia, CA, United
States of America

[4]Department of Thoracic Oncology, Moffitt Cancer Center, Tampa, FL, United States of
America

*Corresponding author: Aleksandra Karolak (AK)

aleks.karolak@moffitt.org

Abstract

Mutation-induced drug resistance is a major contributor to the failure of targeted cancer therapies, particularly in tumors driven by mutations in the KRAS oncogene. Although covalent inhibitors effectively target KRAS G12C, secondary mutations such as G12C/Y96C, G12C/Y96S, and G12C/Y96D lead to resistance despite leaving the covalent attachment site intact. To predict these resistance outcomes, we developed a computational framework that integrates molecular dynamics-derived structural, energetic, thermodynamic, and contact-based descriptors with machine learning. Features extracted from simulations of treatment-sensitive and treatment-resistant KRAS mutants were used to train logistic regression, random forest, support vector machine, and Bayesian Network classifiers, achieving average accuracies above 90%. Solvent-accessible surface area variability, Lennard-Jones 1,4 energy, mean square displacement, and root mean square fluctuation emerged as the most discriminatory features. Residues G10, E62, and H95 showed the highest predictive value. This approach highlights conformational and solvent-exposure changes as central drivers of KRAS drug resistance and provides a generalizable workflow for other clinically relevant mutant targets.

Author Summary

Mutation-induced resistance is a common challenge across many cancer types and is often associated with aggressive tumor progression and poor therapeutic response. Investigating the dynamic properties of proteins harboring such mutations provides valuable insights into the structural and functional consequences of these alterations, thereby helping to elucidate the mechanisms of drug resistance. Machine learning algorithms are particularly effective at uncovering complex patterns within high-dimensional data, such as molecular dynamics simulation trajectories. Integrating these algorithms with analysis of protein dynamics holds significant potential to aid in drug discovery challenges by reducing both time and resource demands while increasing the likelihood of identifying effective therapeutic candidates. As a proof of concept, we developed a computational framework that integrates molecular dynamics-derived molecular features with machine learning to distinguish treatment-sensitive from treatment-resistant KRAS mutants. KRAS is known for drug resistance arising from secondary mutations that disrupt inhibitor binding despite intact covalent attachment sites. The models achieved over 90% accuracy and identified solvent-exposure and conformational changes at residues G10, E62, and H95 as key predictors of treatment resistance. This workflow offers a generalizable strategy for understanding and forecasting mutation-induced resistance.

Introduction

Mutation-induced drug resistance presents a major challenge in cancer therapy, particularly for targeted treatments. Such resistance can arise through various mechanisms [1], including activation of alternative signaling pathways, eg, BRAF(V600E) inhibition by vemurafenib in colon cancer [2]; impaired apoptosis, as seen in p53(R273W) resistance to 5-fluorouracil [3]; and enhanced DNA repair, eg, BRCA1/2 reversion mutations conferring resistance to polyadenosine diphosphate ribose polymerase inhibitors [4]. This study focuses on a common mechanism of resistance driven by direct mutations in the drug target itself. Such mutations frequently alter the chemical and conformational properties of the target, often within the binding site, disrupting drug-receptor interactions. This mechanism, among others, is especially concerning for one of the most frequently mutated oncogenes, Kirsten rat sarcoma viral oncogene homolog (*KRAS*) [5, 6].

The *KRAS* oncogene encodes a 21 kDa guanosine triphosphate hydrolase (GTPase) K-Ras (KRAS henceforward) that regulates key signaling pathways involved in cell-cycle control [6, 7]. Structurally, KRAS consists of a G-domain (residues 1–166), which binds guanine nucleotides, and a hypervariable region (residues 167–189), which mediates membrane association [7]. Its secondary structure includes 6 β-sheets and 5 α-helices connected by flexible loops. Important regions (shown in **Fig 1**) include the switch I and switch II loops, which interact with effector and regulatory proteins, and the P-loop, which binds nucleotide phosphate groups [7]. KRAS homolog alternates between an inactive guanosine diphosphate– (GDP) bound and an active guanosine triphosphate– (GTP-) bound state. Guanine nucleotide exchange factors (GEFs) promote the release of GDP, allowing GTP, which is more abundant in the cell, to bind and

stabilize the active conformation [8]. Binding of GTP activates KRAS by inducing

conformational changes in the switch I and switch II loops. The activated protein then engages

downstream signaling pathways, including the mitogen-activated protein kinase (MAPK)

pathway, the Ras-like small GTPases pathway, and the phosphoinositide-3-kinase pathway,

which are critical regulators of cell proliferation, survival, growth, and differentiation [9].

Termination of signaling requires GTP hydrolysis, facilitated by GTPase-activating proteins,

returning KRAS to its inactive state [8].

Mutations in signaling pathways such as Ras/Raf/MAPK are frequently implicated in cancer,

with KRAS being one of the most commonly mutated oncogenes [6] accounting for 11.6% of all

carcinomas [5]. In pancreatic ductal adenocarcinoma, KRAS mutations are present in 81.7% of

cases, with G12D as the predominant subtype [5]. In colorectal cancer, KRAS mutations occur in

38% of patients, primarily G12D and G12V[5], while in non–small-cell lung cancer, the

prevalence is 21.2%, with G12C representing 45% of cases [5, 6, 10, 11]. Point mutations at

KRAS residues G12, G13, and Q61 impair GAP-mediated GTP hydrolysis, leading to

constitutive KRAS activation and tumorigenesis [12].

Until recently, KRAS-driven tumors were considered difficult to treat because of KRAS's

picomolar affinity for GTP and the absence of other druggable pockets on its surface. As a result,

therapeutic strategies focused on targeting KRAS indirectly [13, 14]. A breakthrough came in

2013, when Ostrem et al [15] reported the first covalent inhibitor targeting the KRAS G12C

mutant. The compound bound specifically to a cysteine residue within an allosteric pocket

located beneath the switch II region in the inactive (GDP-bound) KRAS conformation. This

breakthrough led to the development of the first KRAS G12C inhibitors approved for clinical

use. In 2021, AMG510 [16] (sotorasib, Lumakras) was approved by the FDA for treating non–

small-cell lung cancer, followed by MRTX849 [17] (adagrasib, Krazati) in 2022. Both drugs

irreversibly bind to the cysteine residue at position 12 and form noncovalent interactions with

other amino acids in the switch II binding site, including tyrosine (Y96), glutamine (Q99), and

histidine (H95). This interaction network stabilizes the inactive, GDP-bound conformation of

KRAS, thereby inhibiting oncogenic signaling [18, 19].

Despite initial clinical success, resistance to sotorasib and adagrasib has emerged through

various mechanisms [20, 21]. One involves RTK-mediated activation of wild-type RAS [22].

Another features secondary mutations in KRAS G12C, such as Y96D [23-25], Y96S [24,

25], and Y96C [24, 26], which confer resistance to both approved drugs. Other mutations, such

as G12C/G13D, R68M, A59S, and A59T, are resistant to sotorasib alone, while Q99L is

associated with resistance to adagrasib [24, 25]. These secondary mutations reactivate ERK

signaling and are likely to promote broader resistance across tumor-cell populations [27].

Current efforts to target KRAS-driven tumors include inhibitors of GDP-bound KRAS G12C and

G12D, as well as proteolysis-targeting chimeras for KRAS G12D and G12V, all in clinical trials

[28]. Alternative strategies focus on the active, GTP-bound form of KRAS and binding regions

outside of switch II to circumvent resistance from secondary mutations [29]. However, the

molecular basis of resistance arising in the inactive-state binding site, particularly in KRAS

secondary mutants, remains poorly understood, underscoring the potential benefit of targeted

approaches against these specific variants.

This study moves beyond prevailing studies focused on conformational analyses [30], single

KRAS mutants [31-33] or drug design [34]. Instead, we investigate the underlying factors of

mutation-induced resistance caused by secondary mutations in the KRAS switch II binding site:

KRAS G12C/Y96D, G12C/Y96C, and G12C/Y96S, which exhibit resistance to both sotorasib

and adagrasib. As both drugs interact with residue Y96 [19], substitution at this site represents a

direct structural change relative to the G12C single mutant, which may influence drug binding.

However, other molecular features may also contribute to resistance. To address this, we

developed a framework combining molecular dynamics– (MD-) derived descriptors with

machine learning (ML) algorithms to identify features that differentiate treatment-resistant

KRAS mutants from treatment-sensitive ones.

Integrating dynamic properties of biological systems with ML offers a novel approach to

tackling complex challenges in drug discovery [35]. While some studies have incorporated MD

descriptors into ML models, they have primarily focused on ligand-based pharmacophore

analyses [36], without accounting for protein or binding site dynamics. Other approaches

describe protein-binding sites statically and apply ML to support ligand design [37]. Although a

few efforts have incorporated protein flexibility and ML, eg, to identify druggable binding sites

[37-39], this approach has not yet been applied to address complex problems such as mutation-

induced drug resistance.

Results

This study investigated mutation-induced drug resistance using the KRAS oncogene as a case

study. Based on experimental treatment responses, KRAS structures were classified either as

treatment-sensitive (G12C) or treatment-resistant (G12C/Y96C, G12C/Y96S, and G12C/Y96D).

Because the secondary mutations occur within the switch II binding site, we first assessed

structural differences in this region using SiteMap (**Fig 1**). One key output, SiteScore, reflects the

likelihood of small-molecule binding, with values near 1 indicating promising binding

sites. KRAS WT exhibited a shallow switch II pocket with a low SiteScore (0.754) and small

volume (123 Å³), suggesting unlikely small-molecule binding, consistent with the absence of

endogenous ligands in native KRAS. In contrast, the G12C single mutant and all secondary

mutants had a SiteScore near 1, but their binding pocket volumes varied: G12C had a smaller

volume (256 Å³), while the secondary mutants ranged from 325 to 353 Å³. This volume increase

is likely attributable to substitution of bulky tyrosine at position 96 with smaller residues

(cysteine, serine, or aspartic acid).

Fig 1. **KRAS mutation points and SiteMap analysis**. The first row shows labeled structural

elements of KRAS forms; the second row shows their respective switch II binding sites. Binding

site properties: binding site surface (cyan), hydrogen bond donors (blue), hydrogen bond

acceptors (red), hydrophobic surface (yellow), and GDP (orange). The binding SiteScore and

volume are reported next to each molecule. For visualization purposes, switch II is approximated

on residues 58–72, switch I on residues 30–40, and P-loop on residues 10–14 as suggested in the

following work [7]. The visualizations were prepared with PyMOL Molecular Graphics System

[40]. Abbreviations: GDP, guanosine diphosphate; KRAS, Kirsten rat sarcoma viral oncogene

homolog. Corresponding PDB structures include 6MBT (KRAS WT); 6OIM (KRAS G12C);

6OIM manually mutated (KRAS G12C/Y96C/D/S).

To capture dynamic and subtle structural changes not evident from static analysis, we performed

MD simulations of each KRAS mutant in the absence of inhibitors. From the MD trajectories,

we extracted a comprehensive set of molecular descriptors reflecting global structural behavior

and localized changes within key regions of the protein. These features were categorized into 4

groups: structural, energetic, thermodynamic, and intra/intermolecular contacts. The descriptors

most strongly associated with resistance were related to the site of secondary mutation (residue

96), and the average solvent-accessible surface area (SASA) of residues G10 and A11. After

removing highly intercorrelated variables (see Methods section), the remaining features were

subjected to statistical and ML analyses.

Univariate statistical analysis

The RMSD relative to the reference structure was calculated to assess system stability. Protein

backbone RMSD plots (**Fig 2**) illustrate structural fluctuations in treatment-sensitive and

treatment-resistant KRAS systems throughout the simulation. The treatment-sensitive systems

reached a maximum RMSD of 0.41 nm (**Fig 2A**), while the treatment-resistant systems exhibited

greater protein fluctuations, with RMSD values rising to 0.50 nm (**Fig 2B**). To better

characterize this variability, we generated violin plots of backbone RMSD distributions for the

most populated conformations (**Fig 2C**). In both groups, medians were broadly distributed

around similar values (0.20 and 0.21 nm), although the treatment-resistant group exhibited

greater variability than the treatment-sensitive group (RMSD range: 0.13–0.39 nm vs 0.13–

0.32 nm). However, the Mann–Whitney U test revealed no statistically significant difference (P

value >.05) in backbone RMSD distributions between groups.

Among all RMSD-based features, only the RMSD of the backbone–GDP complex showed a

statistically significant difference (**Table 1**). While both groups displayed values concentrated

near their medians (0.12 and 0.09 nm, **Fig 2D**), the treatment-sensitive systems were shifted

toward higher RMSD values (mean: 0.13 vs 0.11 nm, mean values not shown) than treatment-

resistant systems, suggesting their greater structural flexibility regarding the protein–GDP

complex.

Fig 2. RMSD over the molecular dynamics simulation. (**A**) KRAS treatment-sensitive

systems. (**B**) KRAS treatment-resistant systems. Data distribution of structural features with

annotated median and *P* values. (**C**) Protein backbone RMSD. (**D**) Protein backbone–GDP

RMSD. (**E**) Protein backbone MSD of switch II binding site residues. (**F**) Protein Rg.

Abbreviations: GDP, guanosine diphosphate; KRAS, Kirsten rat sarcoma viral oncogene

homolog; MSD, mean square deviation; RMSD, root mean square deviation.

Table 1. The results of Mann–Whitney U test. Molecular features with *P* value less than 0.05.

Molecular feature	P-value
SASA G10 SD*	1.50×10^{-17}
SASA H95 AV*	1.75×10^{-12}
LJ 1,4	2.19×10^{-8}
SASA E62 SD	6.00×10^{-8}
RMSF M72	1.20×10^{-7}
SASA I100 AV	1.35×10^{-7}
SASA V103 SD	1.46×10^{-7}
SASA V9 AV	1.83×10^{-7}
SASA Q99 AV	4.72×10^{-6}
SASA K16 AV	1.44×10^{-5}
RMSF Q99	1.64×10^{-5}
RMSF H95	2.83×10^{-5}
RMSF Y64	4.62×10^{-5}
Hb protein-GDP	2.59×10^{-4}
SASA V9 SD	4.20×10^{-4}
SASA M72 SD	4.75×10^{-4}
RMSF I100	5.61×10^{-4}
RMSD backbone-GDP	1.37×10^{-3}
Entropy quasi-harmonic	1.47×10^{-3}
SASA G60 AV	1.99×10^{-3}
SASA D69 SD	2.61×10^{-3}
SASA protein	3.06×10^{-3}
RMSF Q61	3.16×10^{-3}
Coulomb 1,4	3.64×10^{-3}
RMSF D69	3.72×10^{-3}
SASA C12 AV	4.45×10^{-3}
RMSF G10	7.66×10^{-3}
MSD bs*-backbone	1.01×10^{-2}
RMSF E63	1.19×10^{-2}
SASA E63 AV	1.77×10^{-2}

SASA Q61 SD	1.94×10^{-2}
SASA H95 SD	2.22×10^{-2}
Rg protein	2.69×10^{-2}
SASA D69 AV	4.65×10^{-2}
SASA T35 SD	4.84×10^{-2}

*Abbreviations: AV, per average; bs, switch II binding site.

Another statistically significant feature associated with structural displacement was the mean

square displacement (MSD) of the switch II binding site backbone (**Fig 2E**, **Table 1**). Like the

protein–GDP complex, the MSD distribution for the switch II binding site backbone was shifted

toward higher values in the treatment-sensitive systems (median: 4.82 vs 4.47 nm²) and showed

greater variability (mean 4.73 nm² ± 0.86 nm² vs. 4.39 nm² ± 0.77 nm²), compared with the

treatment-resistant group. This indicates that, on average, backbone atoms in the switch II

binding region moved farther from their reference positions in the treatment-sensitive systems. In

contrast, the radius of gyration (Rg), a measure of a protein compactness, showed a slight

increase in the treatment-resistant group values compared to the treatment-sensitive systems

(median: 1.57 with max Rg: 1.61 nm vs 1.56 and 1.59 nm, **Fig 2F**). Although the magnitude of

this difference was small, it yielded a *P* value marginally below .05, suggesting borderline

statistical significance.

Unlike global RMSD, which captures overall structural deviation, root mean square fluctuation

(RMSF) provides insight into residue-level flexibility. **Figs 3A-B** shows RMSF values across the

entire simulation time, with arrows marking peaks above 0.4 nm to highlight the most flexible

residues. In both groups, residues T35, E63, and Y64 consistently exhibited the highest peaks.

Additionally, P34 and E62 showed increased fluctuations in the treatment-sensitive systems,

while E37 and Q61 showed higher flexibility in the treatment-resistant group. Among these, only

Q61, E63, and Y64 exhibited statistically significant RMSF differences between groups (**Table**

1). Analysis of the most populated conformations revealed additional switch II binding site

residues with significant group-wise RMSF differences, including M72, Q99, H95, I100, D69,

and G10 (**Fig 3C**). For 8 out of the 9 reported residues, the treatment-resistant group had higher

median RMSF values than the treatment-sensitive group, indicating greater structural flexibility.

Fig 3. RMSF over the molecular dynamics simulation. Residues with RMSF >0.4 nm are

indicated with arrows. (**A**) KRAS treatment-sensitive systems. (**B**) KRAS treatment-resistant

systems. (**C**) RMSF data distribution for the individual residues in the switch II binding site

ranked by the lowest *P* value, with annotated median values. Abbreviations: KRAS, Kirsten rat

sarcoma viral oncogene homolog; RMSF, root mean square deviation.

The final structural feature analyzed was SASA. Regarding the total SASA of the protein in the

most populated conformations, treatment-resistant systems exhibited slightly higher values

(median: 97.45 nm²) compared with treatment-sensitive systems (95.95 nm²), indicating

increased solvent exposure, statistically significant between groups (**Table 1**). To gain residue-

level insight, SASA was further analyzed within the switch II binding site. Violin plots in

Figs 4A-B highlight the top 5 residues with the lowest *P* values for both average (AV) and SD of

SASA. Across these residues, median SASA values were consistently higher in the treatment-

resistant group, suggesting increased solvent exposure. The most significant difference was

observed in the SASA (SD) for residue 10 ($P = 1.5 \times 10^{-17}$), followed by the SASA (AV) of

residue 95 ($P = 1.8 \times 10^{-12}$) and the SASA (SD) for residue 62 ($P = 6 \times 10^{-8}$), highlighting

substantial variation in solvent accessibility between groups at the individual residue level.

The next group of features included energetic properties: potential energy, kinetic energy, and

both short-range and 1,4 Coulomb and Lennard-Jones (LJ) interaction energies. Among these, LJ

1,4 energy and Coulomb 1,4 energy showed statistically significant differences between

treatment-sensitive and treatment-resistant groups (**Table 1**). Lennard-Jones 1,4 energy had the

third lowest P value among all features, with consistently higher values in the treatment-sensitive

group (median: 2075 mean: 2077 kJ/mol) compared with the treatment-resistant group (2032 and

2031 kJ/mol), highlighting the difference in nonbonded interactions energies between the groups

(**Fig 4C**). Conversely, Coulomb 1,4 energy was significantly elevated in the treatment-resistant

systems (median: 23 369 vs 23 234 kJ/mol, data not shown).

To gain insight into thermodynamic properties of the systems, enthalpy and entropy were

analyzed. Entropy calculated using the quasiharmonic approximation showed a statistically

significant difference between treatment-sensitive and treatment-resistant groups (**Table 1**, **Fig

4D**) with elevated values in the treatment-resistant systems (median: 3485 vs 3434 and mean:

3507 vs 3407 kJ/mol). This finding aligns with the generally increased conformational flexibility

in the treatment-resistant group, as indicated by structural features such as RMSF and SASA.

Intra- and intermolecular contacts were also analyzed. Intramolecular interactions included

contacts between the protein and individual amino acid residues, as well as hydrogen bond

formation within the protein. Intermolecular interactions focused on hydrogen bonds formed

between the protein and GDP. The first group of contacts included residue 12, as cysteine at this

position serves as the covalent binding partner for small-molecule inhibitors targeting the KRAS

G12C mutant. A higher number of contacts between C12 and the rest of the protein were

observed in the treatment-sensitive systems, compared with the treatment-resistant ones (median:

41 vs 37, **Fig 4E**); however, this difference was not statistically significant. Among hydrogen

bond–related features, the number of protein-GDP hydrogen bonds differed significantly

between the 2 groups (**Table 1**). The treatment-resistant systems exhibited a higher median

hydrogen bond count (median:10, range 8-13) than the treatment-sensitive group (9 and range 7-

14), indicating more frequent protein-GDP interactions in the treatment-resistant group on

average, although the treatment-sensitive group exhibited a more disperse range (**Fig 4F**). This

finding is correlated to more stable GDP binding pocket in treatment-resistant group (**Fig 2D**),

which allows for more stable occurrence of hydrogen bonds on average. It was reported in

previous studies [30] that both stronger and weaker GDP binding is possible for KRAS

secondary mutants.

Statistical analysis suggested that the relationships between molecular features are complex, with

no single trend consistently distinguishing the 2 groups. For instance, the treatment-resistant

systems showed increased flexibility in the switch II binding site region, as indicated by RMSF

and SASA, while the treatment-sensitive systems exhibited greater mobility in the same region's

backbone, as measured by MSD. These complex findings highlight the structural variability at

the residue level, making it challenging to draw definitive conclusions about the overall direction

or significance of individual features.

Fig 4. Data distribution of structural features, thermodynamic features and intra- and

intermolecular contacts with annotated median and *P* values. (**A**) SASA (AV) of the switch

II binding site residues ranked by the lowest *P* value. (**B**) SASA (SD) for the switch II binding

site residues ranked by the lowest *P* value. (**C**) LJ 1,4 energy. (**D**) Entropy calculated according

to the quasiharmonic equation. (**E**) Intramolecular contacts count between C12 and the rest of

protein residues. (**F**) Hydrogen bonds count between the protein residues and GDP.

Abbreviations: GDP, guanosine diphosphate; LJ, Lennard-Jones; SASA, solvent-accessible

surface area.

Machine-learning models

Given the complexity of MD simulations data, we applied ML to uncover multivariate patterns that may not be apparent through univariate analysis alone, and to identify the most informative features capable of distinguishing treatment-resistant mutants from treatment-sensitive ones (**Fig 5A**). All predictive ML models (LR, RF, and SVM), achieved high classification performance, with average accuracy and area under the curve exceeding 90% (**Table 2**, **Fig 5B**). Model stability across methods was confirmed through 10-fold cross-validation. Hyperparameter tuning using GridSearchCV yielded parameter values close to the defaults (**Table 2**), further indicating robust and generalizable performance.

Across all ML models, the SASA (SD) for residue G10 (**Fig 5A**) was consistently ranked as the most predictive feature differentiating treatment-resistant from treatment-sensitive KRAS systems. This result aligns with the Mann–Whitney U test, where this feature had the lowest P value among all descriptors, indicating the strongest statistical significance. The SASA (SD) for residue G10, was elevated in treatment-resistant systems (**Fig 4B**), along with its RMSF, which was also statistically significant (**Table 1**).

Fig 5. Machine-learning analyses. (**A**) The top important features selected by the machine learning models: LR, RF, and SVM, ranked in the descending order by mean importance, normalized on a 0 to 1 scale. (**B**) Performance of the ML models. ROC AUC curves for LR, RF, and SVM models. (**C**) BN: a subset of the network highlighting the top features is shown. Node colors correspond to the state of each feature when the *resistance* node is fixed to state 1, with red indicating higher marginal value and blue indicating lower marginal value. (**D**) Comparison of how different feature-state combinations influence the likelihood of treatment resistance

outcome in BN. Each cell shows the marginal probability of treatment resistance when a specific

molecular descriptor is fixed at a given state (0, low; 1, medium; and 2, high). Abbreviations:

AUC, area under the curve; BN, Bayesian network; LR, logistic regression; ML, machine

learning; RF, random forest; ROC, receiver-operating characteristic; SVM, support vector

machine.

Table 2. Performance of the ML Models. Logistic Regression (LR), Random Forest (RF) and
Support Vector Machine (SVM) in terms of standard metrics: accuracy, precision, recall, area
under the curve and specificity. Evaluated using 10-fold cross-validation.

Logistic Regression[a]															
Metric	**Accuracy**			**Precision**			**Recall**			**AUC**			**Specificity**		
Run	**Regularization method**														
	L2	L1	Elastic Net	L2	L1	Elastic Net	L2	L1	Elastic c Net	L2	L1	Elastic Net	L2	L1	Elastic Net
1	0.962	0.923	0.885	0.923	0.857	0.800	1.000	1.000	1.000	1.000	0.863	0.887	0.929	0.857	0.786
2	0.962	1.000	0.962	1.000	1.000	1.000	0.909	1.000	0.909	1.000	1.000	1.000	1.000	1.000	1.000
3	0.923	0.962	0.962	0.846	0.917	0.917	1.000	1.000	1.000	0.982	0.933	0.933	0.867	0.933	0.933
4	1.000	0.962	0.962	1.000	0.917	0.917	1.000	1.000	1.000	1.000	0.945	0.976	1.000	0.933	0.933
5	0.885	0.923	0.923	0.786	0.846	0.846	1.000	1.000	1.000	0.933	0.867	0.867	0.800	0.867	0.867
6	1.000	0.923	0.923	1.000	0.875	0.875	1.000	1.000	1.000	1.000	0.917	0.940	1.000	0.833	0.833
7	0.846	0.962	0.962	0.929	1.000	1.000	0.813	0.938	0.938	0.944	1.000	0.994	0.900	1.000	1.000
8	0.923	0.962	0.962	0.923	0.929	0.929	0.923	1.000	1.000	0.994	0.923	0.929	0.923	0.923	0.923
9	0.962	0.962	1.000	1.000	0.917	1.000	0.909	1.000	1.000	0.958	1.000	1.000	1.000	0.933	1.000
10	1.000	0.923	0.923	1.000	0.882	0.882	1.000	1.000	1.000	1.000	0.824	0.836	1.000	0.818	0.818
Average ±SD	0.946 ± 0.052	0.950 ± 0.026	0.947 ± 0.032	0.941 ± 0.076	0.914 ± 0.053	0.917 ± 0.069	0.955 ± 0.065	0.994 ± 0.020	0.985 ± 0.033	0.981 ± 0.026	0.927 ± 0.062	0.936 ± 0.058	0.942 ± 0.071	0.910 ± 0.064	0.909 ± 0.080
Random Forest[b]															
Run	**Parameters**														
	Default	Optimized	Default	Optimized	Default	Optimized	Default	Optimized	Default	Optimized					
1	0.885	0.885	0.800	0.800	1.000	1.000	0.940	0.935	0.786	0.786					
2	0.962	1.000	1.000	1.000	0.909	1.000	1.000	1.000	1.000	1.000					
3	0.923	0.923	0.846	0.846	1.000	1.000	0.988	0.982	0.867	0.867					
4	0.962	0.962	0.917	0.917	1.000	1.000	1.000	1.000	0.933	0.933					
5	0.808	0.808	0.688	0.688	1.000	1.000	0.948	0.942	0.667	0.667					
6	0.962	0.962	0.933	0.933	1.000	1.000	0.988	0.973	0.917	0.917					
7	0.885	0.923	1.000	1.000	0.813	0.875	0.984	0.975	1.000	1.000					
8	0.962	0.962	0.929	0.929	1.000	1.000	0.964	0.950	0.923	0.923					
9	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000					
10	0.923	0.923	0.882	0.882	1.000	1.000	0.967	0.973	0.818	0.818					
Average ±SD	0.927 ± 0.056	0.935 ± 0.057	0.899 ± 0.100	0.899 ± 0.100	0.972 ± 0.063	0.988 ± 0.040	0.978 ± 0.022	0.973 ± 0.024	0.891 ± 0.109	0.891 ± 0.109					
Support Vector Machine[c]															
Run	**Parameters**														
	Default	Optimized	Default	Optimized	Default	Optimized	Default	Optimized	Default	Optimized					
1	0.885	1.000	0.800	1.000	1.000	1.000	0.982	1.000	0.786	1.000					
2	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000					
3	0.885	0.885	0.786	0.786	1.000	1.000	0.988	0.994	0.800	0.800					
4	0.962	0.962	0.917	0.917	1.000	1.000	1.000	0.994	0.933	0.933					
5	0.808	0.885	0.688	0.786	1.000	1.000	0.921	0.958	0.667	0.800					

6	0.962	0.962	0.933	1.000	1.000	0.929	1.000	1.000	0.917	1.000
7	0.962	0.846	1.000	0.929	0.938	0.813	0.963	0.913	1.000	0.900
8	0.962	0.923	0.929	0.923	1.000	0.923	0.994	0.994	0.923	0.923
9	0.962	0.962	0.917	1.000	1.000	0.909	1.000	0.933	0.933	1.000
10	0.923	1.000	0.882	1.000	1.000	1.000	0.952	1.000	0.818	1.000
Average ±SD	0.931 ± 0.057	0.942 ± 0.055	0.885 ± 0.099	0.934 ± 0.086	0.994 ± 0.020	0.957 ± 0.063	0.980 ± 0.027	0.979 ± 0.032	0.878 ± 0.107	0.936 ± 0.081

[a]**Logistic regression parameters**:
- **L2**: 'C': 1.0, solver: lbfgs (default parameters)
- **L1:** best parameters from GridSearchCV: 'C': 0.1, solver:saga
- **Elastic Net:** best parameters from GridSearchCV: 'C': 0.1, 'l1_ratio': 0.5, solver:saga

[b]**Random forest parameters**:
- **Default**: n_estimators=100, max_depth=None, min_samples_split=2, min_samples_leaf=1,max_features='auto'
- **Optimized:** best parameters from GridSearchCV: n_estimators: 50, max_depth: None, max_features: 'sqrt', min_samples_leaf: 1, min_samples_split: 2

[c]**Support vector machine parameters**:
- **Default:** C: 1.0, gamma: 'scale' kernel: 'rbf'
- **Optimized**: best parameters from GridSearch:C: 100, gamma: 0.001, kernel: 'rbf'

Other top-ranked features included the SASA (AV) for H95 and the SASA (SD) for E62, both of which also had among the lowest *P* values. The SASA (AV) for H95 was notably higher in treatment-resistant systems, similar to the residue G10 (**Fig 4A**). In contrast, the SASA (SD) for E62 was elevated in treatment-sensitive systems (**Fig 4B**). Additionally, its RMSF showed high discriminatory power in the LR model, further highlighting its contribution to conformational dynamics relevant to treatment response.

Additional features prioritized by the models included LJ 1,4 energy, the SASA (AV) of K16, and the MSD of the switch II binding site backbone. All were statistically significant, with LJ 1,4 energy ranked as the third most significant feature overall. LJ 1,4 energy values and MSD of the switch II binding site backbone were both more concentrated at higher levels in the treatment-sensitive systems (**Figs 4C and 2E**, respectively), whereas the SASA (AV) of K16 was elevated in the treatment-resistant systems (**Fig 4A**). For validation, we ran two additional ML models (XGBoost and PLS-DA), which remained in agreement with the results. Models' performance and evaluation can be found in the Supplementary Material (**S1 and S2 Tables**, **S1 and S2 Figs**).

Bayesian Network models helped to understand the underlying structure of relationships between the molecular features. First, BN inference was used to explore how treatment response status

influences the probabilistic behavior of molecular descriptors. When the treatment resistance

node was fixed to either the sensitive or resistant state, the resulting shifts in marginal

probabilities revealed that treatment resistance exerts a structured, nonrandom influence on

specific network regions. The most strongly affected descriptors included the SASA (SD) for

G10 and E62, the SASA (AV) of H95, and LJ 1,4 energy (**Fig 5C**), aligning with the features

previously identified by ML models.

To assess whether these descriptors could, in turn, influence treatment response, each was fixed

to one of its discretized states, and the resulting marginal probabilities of the treatment resistance

node were computed. For comparison, 2 additional descriptors, the MSD of the binding site

backbone and the number of intraprotein hydrogen bonds, were also tested because they showed

little change during resistance-driven inference. This analysis revealed several nonlinear,

descriptor-specific effects. Specifically, reduced values of SASA for G10 (SD) and H95 (AV),

and increased values of SASA for E62 (SD) and LJ 1,4 energy, were associated with higher

marginal probabilities of treatment sensitivity (**Fig 5D**). Conversely, the opposite descriptor

states consistently shifted probabilities toward treatment resistance, reinforcing their importance

in shaping the resistance phenotype (**Fig 5D**). These findings are consistent with the ML results

regarding the molecular features distinguishing between the groups.

To visualize the features identified by ML models, the most populated MD conformations from

both treatment-sensitive and treatment-resistant KRAS systems were examined regarding the

residues G10, E62, and H95 conformations (**Fig 6**). Residues E62 and H95 exhibited distinct

conformational dynamics across the most populated clusters, which may explain the differences

in solvent exposure values and suggest potential alterations in ligand accessibility that warrant

further investigation. Residue G10 was located within a highly flexible region, where

neighboring residues likely contribute to the dynamic variability of its solvent exposure.

Overall, univariate statistical analysis and ML models (LR, RF, SVM, and BN) identified a

consistent set of molecular features that differed significantly between treatment-sensitive and

treatment-resistant systems. There was strong agreement across methods regarding feature

relevance, particularly for structural descriptors. The most influential features were

predominantly related to SASA within the switch II binding site. Across all ML models, the

SASA (SD) for G10 emerged as the top discriminatory feature, followed by the SASA (AV) of

H95, the SASA (SD) for E62, and LJ 1,4 energy. These findings highlight the central role of

residue-level flexibility, solvent exposure and nonbonded interactions in KRAS mutation-

induced drug resistance.

Fig 6. Switch II binding-site dynamics. Aligned most populated conformations from MD

simulations with important residues shown in sticks. (**A**) Side view. Residue A11 was removed

from the ML predictions because of high correlation with target variable and Q61 is known for

its catalytic activity. (**B**) Front view. The most important residues indicated by ML models (G10,

E62, and H95). Treatment-sensitive systems (KRAS G12C) in teal, treatment-resistant systems

(KRAS G12C/Y96C, KRAS G12C/Y96D, and KRAS G12C/Y96S) in salmon. Three examples

from each class are shown for clarity. Abbreviations: KRAS, Kirsten rat sarcoma viral oncogene

homolog; MD; ML, machine learning.

Discussion

In this study, the unit of biological variation was the KRAS variant; we analyzed four variants (G12C; G12C/Y96C, G12C/Y96S, G12C/Y96D) using 126 conformers from multiple PDB starts, replicas, and RMSD clusters, intentionally focusing on, although limited, yet clinically reported, resistance-associated double mutants within the inhibitor-binding site to provide hypothesis-generating results that we will expand with additional sensitive variants in future work. It is important to underline that all MD were performed in the unbound (apo) state; no complexes with inhibitors were simulated. Thus, any conclusions are restricted to conformational/topological shifts and do not establish inhibitor-specific resistance mechanisms. These unbound-state results motivate future testing with representative inhibitors but do not by themselves define resistance.

Our approach identified several residues (G10, K16, E62, and H95) as important contributors to treatment resistance in our models, consistent with their known roles in inhibitor recognition. For example, it has been shown that adagrasib forms a salt bridge with E62 and hydrogen bonds with H95 and K16, while its cyanomethyl group interacts with G10, displacing a conserved water molecule [17]. While sotorasib also forms hydrogen bonds with K16, it exploits a specific H95 orientation enhancing binding affinity to KRAS G12C, forms hydrogen bonds E63, and π-π interactions with Y96 [16]. Additionally, G10 solvation has been extensively discussed in previous structural and MD studies [17, 30, 41]. In our apo-state simulations, resistant systems showed increased solvent exposure of G10 and H95 residues and reduced flexibility of E62, suggesting altered solvation patterns and local rigidity around the binding site. These conformational features, even in the absence of ligand, pre-organize resistant mutants into states less compatible with optimal inhibitor binding.

Interestingly, while our ligand-free simulations of secondary resistant KRAS variants (e.g.,

G12C/Y96C/D/S) revealed mutation-dependent changes within the local contact network around

G10, K16, E62, and H95, independent ligand-bound MD studies have reported parallel shifts in

the contact frequencies of adagrasib and sotorasib with these same residues in mutants such as

G12C/R68S, G12C/K16T, G12C/Y96C, and G12C/Y96D [30], together supporting a model in

which resistance stems from subtle, mutation-induced conformational changes that pre-organize

the inhibitor-binding interface in the apo state and subsequently alter drug contacts upon binding.

In summary, we developed a framework that integrates computational chemistry methods with

machine learning algorithms, using KRAS mutation-induced resistance as a case study. Within

this framework, a dataset of comprehensive molecular descriptors, capturing structural,

energetic, thermodynamic, and contacts-based properties was derived from MD simulations and

analyzed by multiple methods. Predictive ML models, LR, RF, and SVM classified treatment

response based on molecular descriptors, while the BN revealed probabilistic dependencies

between features, enhancing interpretability. Together, these approaches identified subtle yet

meaningful differences in the dynamics of KRAS treatment-sensitive and treatment-resistant

systems, particularly in residue-level flexibility and solvent exposure. The conformational

behavior of key residues within the switch II binding site, as demonstrated in our results, could

be considered in the design of inhibitors for KRAS secondary mutants, resistant to sotorasib and

adagrasib, potentially improving therapeutic outcomes. Our study highlights the potential of ML

to address the complexity of mutation-induced drug resistance.

To date, the integration of protein flexibility or MD-derived features into ML models has not

been applied to problems as complex as mutation-induced drug resistance. Our approach

captures mutation-driven intrinsic conformational changes, not ligand-specific effects. This

reduces mechanism specificity but improves generalizability to other phenotypes and targets. By

incorporating protein dynamics before ligand-based analyses, it becomes possible to uncover

unique, targetable features that support the development of more selective and effective

inhibitors. Given the widespread impact of mutation-induced resistance across cancer types, we

propose extending this framework to other clinically relevant resistance-associated targets in the

future.

Methods

Computational chemistry

To investigate the dynamic behavior of KRAS mutants, we conducted all-atom MD simulations

of the inactive KRAS conformations bound to Mg^{2+} and GDP. The initial protein structures were

obtained from the Protein Data Bank (PDB) [42] under accession codes: 6OIM [17], 6UT0 [16],

and 6MBT [43], and prepared before MD simulations. For the KRAS G12C mutant bound to the

covalent inhibitor sotorasib (PDB ID: 6OIM), crystallographic water molecules were removed

together with the ligand. Missing amino acid residues 105-107 were reconstructed using

MODELLER 10.5 [44]. Three engineered mutations, C51S, C80L, and C118S [15], present in

the structure were reverted back to cysteines using the PyMOL 3.0 mutagenesis panel [40]. The

protein was capped with an acetyl group at the N-terminus and with a methyl amide at the C-

terminus using PyMOL's builder module [40]. This structure served as the basis for building

KRAS double mutant models, which have not yet been experimentally resolved. Subsequently,

KRAS G12C (PDB ID: 6OIM) was manually mutated in PyMOL to construct models of the

double mutants G12C/Y96C, G12C/Y96S, and G12C/Y96D. For the wild-type (WT) KRAS

structure (PDB ID: 6MBT) and the KRAS G12C mutant bound to the covalent inhibitor

adagrasib (PDB ID: 6UT0), the same preparation steps were applied. Additionally, the WT

KRAS structure was mutated at position 12, from glycine to cysteine, to model an additional

KRAS single-mutant system.

All protein models were processed using the CHARMM36 force field [45] in GROMACS [46].

Hydrogen atoms were added to each protein structure, and the protonation states of the residues

were assigned at neutral pH. The GDP molecule was parametrized separately with the

CHARMM General Force Field [47]. To prepare the systems for energy minimization, the

complexes were placed in cubic boxes, extending at least 10 Å from the protein. The solvent was

modeled using the CHARMM-modified TIP3P water model [45]. The protein charges were

neutralized by adding the appropriate amount of Na+ ions, and physiological NaCl salt

concentration 0.1M was ensured. Energy minimization was performed using the steepest descent

algorithm [48].

Next, NVT equilibration was conducted for 100 ps to stabilize the system's temperature at 310 K

with position restraints on protein and ligand heavy atoms to allow the solvent and ions to relax.

Temperature was maintained at 310 K using the V-rescale thermostat [49] with separate coupling

groups for protein and GDP ligand, and solvent/ions. H-bonds were constrained using LINCS

[50], enabling the 2 fs timestep. This was followed by 250 ps of NPT equilibration, continuing

from the NVT step, with the same position restraints and temperature control, while pressure was

maintained at 1 bar using isotropic Berendsen coupling [51]. Production MD were performed for

200 ns with a 2 fs timestep. Temperature was maintained at 310 K using the V-rescale

thermostat, and pressure was maintained at 1 bar using isotropic Parrinello-Rahman coupling.

Coordinates and energies were saved every 10 ps. Each system underwent 3 replica runs, with

different random seeds set before each simulation run. The water shell confirming octahedral

arrangement of water molecules around Mg^{2+} ion is shown in **S3 Fig**.

To extract representative conformations from the trajectories, we performed trajectory clustering

on the 50–200 ns portion of each simulation using the GROMOS [52] method. This method

counts neighboring structures within a specified cutoff and identifies the representative frame as

the one with the smallest average root mean square deviation (RMSD) to all its neighbors.

Clustering was performed for 6 different Cα RMSD cutoffs, ranging from 0.10 nm to 0.15 nm, to

find reasonable distribution of frames for each system. Representative structures from the top 7

clusters (~90% of the trajectory) were extracted from each of the 3 replicas per system for

molecular descriptors calculations. This resulted in 126 conformations across all 6 systems.

Specific elements of each system were treated separately in the calculations, including extended

switch II binding site residues (residues 9, 10, 11, 12, 16, 34, 35, 37, 58, 59, 60, 61, 62, 63, 64,

68, 69, 72, 95, 96, 99, 100, 102, and 103), which were defined as residues within 5Å of any site

point identified in the KRAS G12C switch II binding site by the SiteMap module [53, 54] of

Schrödinger Suite and included motifs from P-loop, switch I region, and adjacent helices. A full

list of molecular descriptors, grouped into categories, is provided in **Table 3**.

Table 3. The complete list of the molecular descriptors extracted from MD simulations.

Molecular dynamics simulation descriptors	
Calculation type	**System elements**
Structural descriptors	
Root Mean Square Deviation (RMSD, nm)*	Calculated as a function of time for: • protein backbone • switch II binding site residues (backbone and sidechains - heavy atoms) • GDP and protein backbone • Mg^{2+} ion and protein backbone
Radius of gyration (Rg, nm)*	Calculated as a function of time for (all-atoms): • protein • protein-GDP • switch II binding site residues
Root Mean Square Fluctuation (RMSF, nm)	Calculated per switch II binding site residue over the trajectory frames extracted for each cluster.
Solvent accessible surface area (SASA, nm^2) using the Double Cubic Lattice Method[55]	Total solvent accessible surface area, estimated solvation free energy, and total volume and density was calculated as a function of time for (all-atoms): • protein • switch II binding site residues

	Additionally, the average solvent accessible surface area and its standard deviation was calculated per each switch II binding site residue over the trajectory frames extracted from each cluster.
Mean Square Displacement (MSD, nm^2)	Calculated as a function of time for: • protein backbone • switch II binding site residues (backbone) • switch II binding site residues (sidechains-heavy atoms).
Energetic features	
Potential energy, total energy, kinetic energy, short-range and 1,4 Coulomb and Lennard-Jones (LJ) energies (kJ/mol)	Extracted for representative frame from the most populated conformational clusters.
Thermodynamic features	
Entropy estimate based on the quasiharmonic approach[56] and Schlitter's formula[57] (kJ/mol)	Estimated as an evolution of the system up to each cluster's representative timeframe.
Enthalpy (kJ/mol)	Extracted for representative frame from the most populated conformational clusters.
Intra- and intermolecular contacts	
Hydrogen bonds count	Calculated as a function of time for (all atoms): • within protein • between protein and GDP • within switch II binding site residues
Number of contacts within 6 Å	Calculated as a function of time for (heavy atoms): • between residue 12 and protein • between residue 12 and switch II binding site residues

490   *First frame of the production run was used as the reference conformation.

Statistical and ML methods

ML approach was implemented to identify the molecular descriptors distinguishing between

treatment-sensitive and treatment-resistant KRAS mutants. The dataset consisted of 3 treatment-

sensitive (resistance = 0) and 3 treatment-resistant (resistance = 1) KRAS systems, each

contributing 63 samples, for a total of 126 samples described by 101 molecular features. To

reduce redundancy, highly target-correlated and intercorrelated features (correlation >75%) were

removed, along with all descriptors directly related to residue 96, the site of the secondary

mutation, to avoid bias toward known changes. The remaining features were standardized using

Z-score scaling to ensure comparable value ranges across inputs.

Before model training, we performed univariate statistical analysis. Feature distributions were

tested for normality using the Shapiro–Wilk test [58], and the Mann–Whitney U test [59] was

used to assess statistical significance between the feature data distribution in the treatment-

sensitive and treatment-resistant groups. Violin plots [60, 61] were used to visualize feature

distributions, with central boxes indicating the interquartile range (IQR, 25th–75th percentile),

median lines, and whiskers extending 1.5× the IQR.

Since the relationships between features and resistance were not assumed to be strictly linear, 3

supervised ML models were applied: logistic regression (LR), random forest (RF), and support

vector machine (SVM), covering both linear and nonlinear classification paradigms.

Hyperparameter optimization was performed using GridSearchCV [62] with 5-fold cross-

validation. For LR, the L1 (Lasso), L2 (Ridge), and Elastic Net [62, 63] regularizations were

tested. The regularization strength C was varied (0.01, 0.1, 1, 10, 100) for all models,

while Elastic Net additionally included tuning of the l1_ratio (0.1, 0.5, 0.7, 1.0). For

the SVM classifier, the grid included C (0.01, 0.1, 1, 10, 100), kernel (linear, rbf, poly), gamma

(0.001, 0.01, 0.1, 1), and degree (2, 3, 4). For the RF model, the parameters tested

were n_estimators (50, 100, 150), max_depth (none, 10, 20), min_samples_split (2, 5,

10), min_samples_leaf (1, 2, 4), and max_features (sqrt, log2, none). Feature importance was

determined by using absolute coefficients for LR, total Gini impurity reduction for RF, and

permutation importance for SVM. Each model was trained on 80% of the data and tested on the

remaining 20%, with performance evaluated using 10-fold cross-validation to ensure

repeatability and prevent overfitting [63]. All analyses were conducted using Python 3.11.11[64].

To analyze probabilistic relationships among molecular descriptors, we used a Bayesian Network

(BN), an interpretable ML model that represents joint probability distributions as a directed

acyclic graph. A universal BN was constructed by combining treatment-sensitive and treatment-

resistant KRAS datasets using BNOmics [65], with 100 random restarts to ensure convergence

and structural robustness. Molecular descriptors were discretized into 3 categories, low (0),

medium (1), and high (2), using a maximum entropy algorithm [65]. The resulting network

captured both direct associations with treatment response and higher-order dependencies among

features. Two types of inference analyses were performed using a belief propagation algorithm.

First, to assess how treatment response affects molecular descriptors, the treatment resistance

node was fixed to either sensitive (0) or resistant (1), and the resulting posterior marginal

probabilities were computed for each feature. These distributions quantify how each node's

probability shifts in response to the fixed phenotype. Second, to evaluate how selected molecular

descriptors influence treatment response, each was fixed to one of its discretized states (0, 1, or

2), and the resulting marginal probability of the treatment resistance node was inferred. These

probabilities were compiled into a heatmap to visualize the effect of each perturbation.

Data and code availability

The GitHub repository containing the code and data can be found at:

https://github.com/Molecular-AI-Group/KatMizgalska/tree/KRAS-MD-descriptors

Author contributions

K.M. performed molecular dynamics simulations, conducted univariate statistical analyses and predictive ML modeling and wrote and edited the manuscript. K.U. and S.B. performed and described all analyses related to the BN. A.K designed the study in consultation with E.H., D.I. and W.G., supervised it, and provided guidance on result interpretation and manuscript editing. All authors reviewed and approved the final manuscript.

Declaration of interests

Authors declare no conflicting interests.

Funding

This research was supported in part by A.K. support funds and the National Institutes of Health under grants R01-LM013138, and California Institute for Regenerative Medicine grant. CIRM EDU4-12772, acknowledged by S.B. and K.U.

554 Acknowledgments

555   Molecular dynamics simulations were performed using HPC computing cluster at H. Lee Moffitt

556   Cancer Center and Research Institute. Editorial assistance was provided by Moffitt Cancer

557   Center's Office of Scientific Publishing by Gerard Hebert.

References

1.   Housman G, Byler S, Heerboth S, Lapinska K, Longacre M, Snyder N, et al. Drug resistance in cancer: an overview. Cancers (Basel). 2014;6(3):1769–92.

2.   Prahallad A, Sun C, Huang S, Di Nicolantonio F, Salazar R, Zecchin D, et al. Unresponsiveness of colon cancer to BRAF(V600E) inhibition through feedback activation of EGFR. Nature. 2012;483(7387):100–3.

3.   Huang Y, Liu N, Liu J, Liu Y, Zhang C, Long S, et al. Mutant p53 drives cancer chemotherapy resistance due to loss of function on activating transcription of PUMA. Cell Cycle. 2019;18(24):3442–55.

4.   Collet L, Hanvic B, Turinetto M, Treilleux I, Chopin N, Le Saux O, et al. BRCA1/2 alterations and reversion mutations in the area of PARP inhibitors in high grade ovarian cancer: state of the art and forthcoming challenges. Front Oncol. 2024;14:1354427.

5.   Yang Y, Zhang H, Huang S, Chu Q. KRAS Mutations in Solid Tumors: Characteristics, Current Therapeutic Strategy, and Potential Treatment Exploration. J Clin Med. 2023;12(2).

6.   Huang L, Guo Z, Wang F, Fu L. KRAS mutation: from undruggable to druggable in cancer. Signal Transduct Target Ther. 2021;6(1):386.

7.   Pantsar T. The current understanding of KRAS protein structure and dynamics. Comput Struct Biotechnol J. 2020;18:189–98.

8.   Bos JL, Rehmann H, Wittinghofer A. GEFs and GAPs: critical elements in the control of small G proteins. Cell. 2007;129(5):865–77.

9.   Li HY, Qi WL, Wang YX, Meng LH. Covalent inhibitor targets KRas(G12C): A new paradigm for drugging the undruggable and challenges ahead. Genes Dis. 2023;10(2):403–14.

10.   Tan AC, Tan DSW. Targeted Therapies for Lung Cancer Patients With Oncogenic Driver Molecular Alterations. J Clin Oncol. 2022;40(6):611–25.

11.   El Osta B, Behera M, Kim S, Berry LD, Sica G, Pillai RN, et al. Characteristics and

outcomes of patients with metastatic KRAS-mutant lung adenocarcinomas: the lung cancer

mutation consortium experience. J Thorac Oncol. 2019;14(5):876–89.

12.   Parker JA, Mattos C. The K-Ras, N-Ras, and H-Ras Isoforms: Unique Conformational

Preferences and Implications for Targeting Oncogenic Mutants. Cold Spring Harb Perspect Med.

2018;8(8).

13.   Blasco MT, Navas C, Martin-Serrano G, Grana-Castro O, Lechuga CG, Martin-Diaz L, et

al. Complete regression of advanced pancreatic ductal adenocarcinomas upon combined

inhibition of EGFR and C-RAF. Cancer Cell. 2019;35(4):573–87 e6.

14.   Gibbs JB, Graham SL, Hartman GD, Koblan KS, Kohl NE, Omer CA, et al.

Farnesyltransferase inhibitors versus Ras inhibitors. Curr Opin Chem Biol. 1997;1(2):197–203.

15.   Ostrem JM, Peters U, Sos ML, Wells JA, Shokat KM. K-Ras(G12C) inhibitors

allosterically control GTP affinity and effector interactions. Nature. 2013;503(7477):548–51.

16.   Canon J, Rex K, Saiki AY, Mohr C, Cooke K, Bagal D, et al. The clinical KRAS(G12C)

inhibitor AMG 510 drives anti-tumour immunity. Nature. 2019;575(7781):217–23.

17.   Fell JB, Fischer JP, Baer BR, Blake JF, Bouhana K, Briere DM, et al. Identification of

the Clinical Development Candidate **MRTX849** , a Covalent KRAS G12C

Inhibitor for the Treatment of Cancer. Journal of Medicinal Chemistry. 2020;63(13):6679–93.

18.   Zhang SS, Nagasaka M. Spotlight on sotorasib (AMG 510) for KRAS (G12C) positive

non-small cell lung cancer. Lung Cancer (Auckl). 2021;12:115–22.

19.   Mahran R, Kapp JN, Valtonen S, Champagne A, Ning J, Gillette W, et al. Beyond

KRAS(G12C): biochemical and computational characterization of sotorasib and adagrasib

binding specificity and the critical role of H95 and Y96. ACS Chem Biol. 2024;19(10):2152–64.

20.   Liu J, Kang R, Tang D. The KRAS-G12C inhibitor: activity and resistance. Cancer Gene

Ther. 2022;29(7):875–8.

21.   Mohanty A, Nam A, Srivastava S, Jones J, Lomenick B, Singhal SS, et al. Acquired

resistance to KRAS G12C small-molecule inhibitors via genetic/nongenetic mechanisms in lung

cancer. Sci Adv. 2023;9(41):eade3816.

22.   Ryan MB, Fece de la Cruz F, Phat S, Myers DT, Wong E, Shahzade HA, et al. Vertical

pathway inhibition overcomes adaptive feedback resistance to KRAS[G12C] inhibition. Clin Cancer

Res. 2020;26(7):1633–43.

23.   Tanaka N, Lin JJ, Li C, Ryan MB, Zhang J, Kiedrowski LA, et al. Clinical acquired

resistance to KRAS[G12C] inhibition through a novel KRAS switch-II pocket mutation and

polyclonal alterations converging on RAS-MAPK reactivation. Cancer Discov.

2021;11(8):1913–22.

24.   Rosen JC, Sacher A, Tsao MS. Direct GDP-KRAS[G12C] inhibitors and mechanisms of

resistance: the tip of the iceberg. Ther Adv Med Oncol. 2023;15:17588359231160141.

25.   Koga T, Suda K, Fujino T, Ohara S, Hamada A, Nishino M, et al. KRAS secondary

mutations that confer acquired resistance to KRAS G12C inhibitors, sotorasib and adagrasib, and

overcoming strategies: insights from in vitro experiments. J Thorac Oncol. 2021;16(8):1321–32.

26.   Awad MM, Liu S, Rybkin, II, Arbour KC, Dilly J, Zhu VW, et al. Acquired resistance to

KRAS[G12C] inhibition in cancer. N Engl J Med. 2021;384(25):2382–93.

27.   Zhao Y, Murciano-Goroff YR, Xue JY, Ang A, Lucas J, Mai TT, et al. Diverse

alterations associated with resistance to KRAS[G12C] inhibition. Nature. 2021;599(7886):679–83.

28.   Isermann T, Sers C, Der CJ, Papke B. KRAS inhibitors: resistance drivers and

combinatorial strategies. Trends Cancer. 2025;11(2):91–116.

29.   Holderfield M, Lee BJ, Jiang J, Tomlinson A, Seamon KJ, Mira A, et al. Concurrent

inhibition of oncogenic and wild-type RAS-GTP for cancer therapy. Nature.

2024;629(8013):919–26.

30.   Tu G, Liu Q, Qiu Y, Leung EL, Yao X. In Silico Study of the Acquired Resistance

Caused by the Secondary Mutations of KRAS G12C Protein Using Long Time Molecular

Dynamics Simulation and Markov State Model Analysis. Int J Mol Sci. 2022;23(22).

31.   Pandey D, Roy KK. Decoding KRAS dynamics: Exploring the impact of mutations and

inhibitor binding. Arch Biochem Biophys. 2025;764:110279.

32.   Mir SA, Nayak B, Aljarba NH, Kumarasamy V, Subramaniyan V, Dhara B. Exploring

KRas protein dynamics: an integrated molecular dynamics analysis of KRas wild and mutant

variants. ACS Omega. 2024;9(28):30665–74.

33.   Mehmood A, Kaushik AC, Wang Q, Li CD, Wei DQ. Bringing structural implications

and deep learning–based drug identification for KRAS mutants. J Chem Inf Model.

2021;61(2):571–86.

34.   Srisongkram T, Weerapreeyakul N. Drug Repurposing against KRAS Mutant G12C: A

Machine Learning, Molecular Docking, and Molecular Dynamics Study. Int J Mol Sci.

2022;24(1).

35.   Stitou M, Koomen J, Imbody D, Liao Y, Monastyrskyi A, Rix U, et al. Does artificial

intelligence need companionship to assist in drug discovery? The Kirsten rat sarcoma virus

study. BJR Artif Intell. 2024;1(1).

36.   Jamal S, Grover A, Grover S. Machine Learning From Molecular Dynamics Trajectories

to Predict Caspase-8 Inhibitors Against Alzheimer's Disease. Front Pharmacol. 2019;10:780.

37.   Xia S, Chen E, Zhang Y. Integrated molecular modeling and machine learning for drug

design. J Chem Theory Comput. 2023;19(21):7478–95.

38.   Cimermancic P, Weinkam P, Rettenmaier TJ, Bichmann L, Keedy DA, Woldeyes RA, et

al. CryptoSite: expanding the druggable proteome by characterization and prediction of cryptic

binding sites. J Mol Biol. 2016;428(4):709–19.

39.   Meller A, Ward M, Borowsky J, Kshirsagar M, Lotthammer JM, Oviedo F, et al.

Predicting locations of cryptic pockets from single protein structures using the PocketMiner

graph neural network. Nat Commun. 2023;14(1):1177.

40.   The PyMOL Molecular Graphics System, Version 3.0 Schrödinger, LLC.

41.   Leini R, Pantsar T. In Silico Evaluation of the Thr58-Associated Conserved Water with

KRAS Switch-II Pocket Binders. J Chem Inf Model. 2023;63(5):1490–505.

42.   Berman HM. The Protein Data Bank. Nucl Acids Res. 2000;28(1):235–42.

43.   Dharmaiah S, Tran TH, Messing S, Agamasu C, Gillette WK, Yan W, et al. Structures of

N-terminally processed KRAS provide insight into the role of N-acetylation. Sci Rep.

2019;9(1):10512–.

44.   Webb B, Sali A. Comparative Protein Structure Modeling Using MODELLER. Curr

Protoc Bioinformatics. 2016;54(1).

45.   Huang J, Rauscher S, Nawrocki G, Ran T, Feig M, de Groot BL, et al. CHARMM36m:

an improved force field for folded and intrinsically disordered proteins. Nat Methods.

2017;14(1):71–3.

46.   Van Der Spoel D, Lindahl E, Hess B, Groenhof G, Mark AE, Berendsen HJC.

GROMACS: fast, flexible, and free. J Comput Chem. 2005;26(16):1701–18.

47.   Vanommeslaeghe K, Hatcher E, Acharya C, Kundu S, Zhong S, Shim J, et al. CHARMM

general force field: a force field for drug-like molecules compatible with the CHARMM all-atom

additive biological force fields. J Comput Chem. 2010;31(4):671–90.

48.   William H. Press SAT, William T. Vetterling, Brian P. Flannery. Numerical Recipes in

FORTRAN: The Art of Scientific Computing: Cambridge University Press; 1986.

49.   Bussi G, Donadio D, Parrinello M. Canonical sampling through velocity rescaling. J

Chem Phys. 2007;126(1).

50.   Hess B. P-LINCS: a parallel linear constraint solver for molecular simulation. J Chem

Theory Comput. 2008;4(1):116–22.

51.   Berendsen HJC, Postma JPM, van Gunsteren WF, DiNola A, Haak JR. Molecular

dynamics with coupling to an external bath. J Chem Phys. 1984;81(8):3684–90.

52.   Daura XG, K. Jaun, B. Seebach, D. van Gunsteren, W. F., Mark AE. Peptide Folding:

When Simulation Meets Experiment. Angew Chem 1999;38:236–40.

53.   Schrödinger Release 2024-3: SiteMap, Schrödinger, LLC, New York, NY, 2024.

54.   Halgren TA. Identifying and characterizing binding sites and assessing druggability. J

Chem Inf Model. 2009;49(2):377–89.

55.   Eisenhaber F, Lijnzaad P, Argos P, Sander C, Scharf M. The double cubic lattice method:

efficient approaches to numerical integration of surface area and volume and to dot surface

contouring of molecular assemblies. J Comput Chem. 1995;16(3):273–84.

56.   Andricioaei I, Karplus M. On the calculation of entropy from covariance matrices of the

atomic fluctuations. J Chem Phys. 2001;115(14):6289–92.

57.   Schlitter J. Estimation of absolute and relative entropies of macromolecules using the

covariance matrix. Chem Phys Lett. 1993;215(6):617–21.

58.   Andrew P. King RJE. Chapter 7 – Inferential Statistics IV: Choosing a Hypothesis Test.

Andrew P. King RJE, editor: Academic Press; 2019.

59.   Nachar N. The Mann-Whitney U: a test for assessing whether two independent samples

come from the same distribution. Tutor Quant Methods Psychol. 2008;4(1):13–20.

60.   Waskom M. Seaborn: statistical data visualization. J Open Source Softw. 2021;6(60).

61.   Hunter JD. Matplotlib: a 2D graphics environment. Comput Sci Eng. 2007;9(3):90–5.

62.   Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-

learn: machine learning in python. J Mach Learn Res. 2011;12(null):2825–30.

63.   Gareth JW, D.; Hastie, T.; Tibshirani, R.;. An Introduction to Statistical Learning:

Springer; 2013.

64.   Van Rossum G, Drake FL. Python 3 Reference Manual: CreateSpace; 2009.

65.   Gogoshin G, Boerwinkle E, Rodin AS. New algorithm and software (BNOmics) for

inferring and visualizing Bayesian Networks from heterogeneous big biological and genetic data.

J Comput Biol. 2017;24(4):340–56.

Supporting Information

S1 Table. XGBoost model evaluation.

S2 Table. PLS-DA model evaluation.

S1 Fig. XGBoost model performance. Left: XGBoost ROC-AUC across 10 iterations. Right: top 15 Feature Importance averaged over 10 iterations calculated using a Permutation technique. This model, even after small-data set adjustment, still relies on one dominant feature SASA (SD) of G10. It agrees with the models studied in the publication.

S2 Fig. PLS-DA model performance. Left: PLS-DA ROC-AUC across 10 iterations. Right: top 15 Feature Importance averaged over 10 iterations calculated using a Variable Importance in Projection (VIP) scores, which estimate each feature contribution to the production. This model does not rely heavily on a few features, but consensus between several equally important. The result agrees with the models studied in the publication.

S3 Fig. Water shell around Mg^{2+}. Left: initial structure (PDB ID: 6OIM, X-ray crystallography). Right: system's starting structure after the NPT equilibration ready for the simulation run.

KRAS WT	KRAS G12C	KRAS G12C/Y96C	KRAS G12C/Y96D	KRAS G12C/Y96S
	TREATMENT-SENSITIVE	TREATMENT-RESISTANT	TREATMENT-RESISTANT	TREATMENT-RESISTANT
SiteScore: 0.754 Volume: 123 A³	SiteScore: 1.051 Volume: 256 A³	SiteScore: 0.976 Volume: 325 A³	SiteScore: 1.028 Volume: 339 A³	SiteScore: 1.021 Volume: 353 A³