Learning latent conformational landscapes encoded in cryo-EM

Haizhao Dai[1,3†], Yingjun Shen[1,3,4†], Qihe Chen[1,3†], Lingqi Li[1,2†], Zhenyang Xu[1,3], Minzhang Li[1,3], Yufan Xie[1], Jingjing Zheng[1,2], Yuan Pei[1,2*], Jiakai Zhang[1,3*], Liping Sun[1,2*], Zhijie Liu[1,2*], Jingyi Yu[1*]

[1]ShanghaiTech University, Shanghai, China.
[2]iHuman Institute, Shanghai, China.
[3]Cellverse Co., Ltd., Shanghai, China.
[4]Shanghai Jiao Tong University, Shanghai, China.

*Corresponding author(s). E-mail(s): peiyuan@shanghaitech.edu.cn; zhangjk@shanghaitech.edu.cn; sunlp@shanghaitech.edu.cn; liuzhj@shanghaitech.edu.cn; yujingyi@shanghaitech.edu.cn;
Contributing authors: daihzh2023@shanghaitech.edu.cn; skrry_12138@sjtu.edu.cn; chenqh2024@shanghaitech.edu.cn; lilq2023@shanghaitech.edu.cn; xuzhy12024@shanghaitech.edu.cn; limzh2022@shanghaitech.edu.cn; xieyf2022@shanghaitech.edu.cn; zhengjj1@shanghaitech.edu.cn;
[†]Equal contributions.

Abstract

Proteins exist as a continuum of conformational states, millions of snapshots of which are captured in every cryo-EM experiment. Current analysis reduces these snapshots to static density maps or discrete classes, discarding the dynamics encoded in the data. Continuous latent-space methods offer a promising alternative, yet whether their learned representations are physically grounded remains unresolved. Here, we show that cryo-EM data can be organized into a probabilistic conformational landscape in which latent density reflects the distribution of molecular states. We demonstrate that this landscape is both physically grounded and experimentally actionable. For integrin $\alpha v\beta 8$, the learned landscape shows strong agreement with independently derived molecular dynamics simulations, supporting its physical plausibility. Additionally, the landscape reveals biologically meaningful organization of molecular states, including a spectrum of states in LIS1-mediated dynein activation, from dominant conformations to low-population intermediates defined by distinct binding stoichiometries, and continuous conformational pathways in KCTD5/CUL3^{NTD}/G$\beta\gamma$ complex that are consistent with prior simulations. Probability-guided particle selection further improves reconstruction quality, closing the loop from landscape to structure determination. These results establish that cryo-EM data encode a latent conformational landscape that can be directly exploited for structural reconstruction, discovery, and interpretation.

1 Introduction

Since the middle of the last century, protein structure has been represented as a single and explicit model defined by atomic coordinates built from density maps. Although such models have proven invaluable for understanding protein function and guiding drug design, they capture mostly a single state of molecules that exist in multiple conformations under physiological conditions. Single-particle cryo-EM inherently captures this structural variability as millions of molecular snapshots. Yet, standard reconstruction approaches assume a single or a few underlying states, averaging over the captured variation into one density map that best fits the data. Discrete classification methods [1, 2] partially resolve this by partitioning single or multiple clusters of particles into a small number of states, but intermediate states and continuous dynamics remain unresolved. Recent advances in continuous heterogeneity analysis [3–8] encode structural variability into learned latent spaces for volume reconstruction. However, the intrinsic structure of these latent spaces,

including the probability density and the transition pathway, lacks established physical grounding. A fundamental question thus arises: can a latent representation faithfully capture the conformational landscape encoded in cryo-EM data?

Here, we show that a latent conformational landscape can be learned directly from cryo-EM images. However, their extremely low signal-to-noise ratios demand latent representations that can reliably separate structural signal from imaging noise [9]. We present CryoUNI, a universal encoder for cryo-EM particle images, trained on diverse particle datasets to map images into a probabilistic latent space in which density reflects the distribution of structural states and defines a relative energy landscape described by Boltzmann statistics. Within this latent space, distinct structural states emerge as density peaks, i.e., energy basins, and transitions between them follow continuous pathways. To systematically analyze this landscape, we introduce WAVE (Watershed Analysis of Variational Embeddings), which automatically identifies conformational states and their transitions from latent density, resolving both dominant conformations and rare intermediates without requiring a predefined number of states.

Across three experimental systems, we demonstrate that the learned landscape is physically grounded and facilitates the discovery of novel structures and their transitions. The latent space of integrin $\alpha v \beta 8$ [10] closely recapitulates the largest structural variations, the leg movement, in the conformational landscape constructed from molecular dynamics simulations. For LIS1-mediated dynein activation [11], the learned landscape reveals not only the known dominant conformations but also previously unresolved low-population intermediates defined by distinct binding stoichiometries. For the KCTD5/CUL3$^{\mathrm{NTD}}$/G$\beta\gamma$ complex [12], continuous conformational dynamics are resolved for the first time from cryo-EM data. Beyond, the learned landscape provides a basis for particle selection, leading to better reconstruction quality for this complex.

Together, these results demonstrate that leveraging CryoUNI, latent conformational landscapes can be learned directly from cryo-EM data. This shift from deterministic and explicit representations to a latent and probabilistic description unlocks a holistic view of cryo-EM data, bringing us closer to understanding the structure-dynamics-function relationship of diverse proteins.

2 CryoUNI learns a latent conformational landscape from cryo-EM

Cryo-EM particle images carry structural information at extremely low signal-to-noise ratios, demanding representations that reliably separate structural signal from imaging noise. We address this by pretraining a universal encoder, CryoUNI, on CryoCRAB-Particle-22M (Fig. 1a), a dataset of 22 million cryo-EM particles spanning five SCOP2 structural classes [13], with molecular weight ranging from below 250 kDa to above 1 MDa. To ensure that the encoder extracts a clean structural signal rather than imaging noise, we adopt a self-supervised training objective [14, 15] customized for cryo-EM particle images. Specifically, CryoUNI is pretrained to denoise particles from one half-dataset using the complementary half as supervision, exploiting the fact that both halves share identical underlying structure but distinct noise (Fig. 1b).

For downstream conformational analysis, the pretrained CryoUNI encoder is adapted to a target dataset *via* a variational autoencoder (VAE) architecture, mapping each particle image into a low-dimensional embedding space through a lightweight linear projector, from which a neural volume decodes the corresponding structural density (Fig. 1c). Notably, since the encoder is trained to suppress noise-driven variation, the density in the latent space is expected to reflect the underlying distribution of structural states, approximating a structural landscape that can be described by Boltzmann statistics.

To recover the structural landscape from the probabilistic latent space, we first apply kernel density estimation to the particle embeddings, followed by the watershed analysis of variational embeddings (WAVE) for automated state identification and transition analysis (Fig. 1d). WAVE identifies local density maxima as discrete conformational states and draws their boundaries, revealing both dominant conformations and rare intermediates. The relative energy between states is then directly inferred from their density ratio *via* Boltzmann statistics, $\Delta G_r = -k_B T \ln(\rho_A / \rho_B)$. Continuous conformational pathways between states are traced along the landscape, revealing the structural transitions and relative population of intermediates (Fig. 1e).

Together, CryoUNI converts cryo-EM particle images into a continuous probabilistic conformational landscape, followed by WAVE analyses to reveal both discrete and continuous structural heterogeneity.

3 The latent space is physically grounded in the molecular conformational landscape

To assess whether the learned latent landscape is physically grounded, we compared it with independent molecular dynamics (MD) simulations on the integrin $\alpha v \beta 8$ in extended-closed conformation [10]. This system exhibits continuous rotation of the flexible leg, which is not successfully resolved in the published

density map. We reproduce such molecular motion through well-converged all-atom MD simulations comprising twenty independent 1-μs trajectories (20 μs aggregate; Fig. 2a). We then characterize it by mapping to a spherical coordinate system anchored at the head-leg hinge center of mass (COM), parameterizing leg orientation *via* polar (θ) and azimuthal (ϕ) angles (Fig. 2b). This coordinate system is applicable to both atomic models and reconstructed cryo-EM volumes, enabling direct comparison across representations. Projection of all sampled 160,000 MD snapshots onto this (θ, ϕ) space reveals that the leg motion is confined to a well-defined thermodynamic landscape (Fig. 2c and Supplementary Fig. 2a).

Applying CryoUNI to the same dataset yields a latent space with a continuous, unimodal distribution (Fig. 2d). The learned latent axes align closely with the physical degrees of freedom captured by MD: PC1 correlates strongly with polar angle θ ($r = 0.982$), and PC2 with azimuthal angle ϕ ($r = 0.963$) (Fig. 2e). This level of correspondence, achieved without imposing biophysical priors, indicates that the latent representation captures the intrinsic conformational coordinates of the system and is therefore physically plausible.

We further evaluate this correspondence by projecting EM-reconstructed conformations into the MD-derived (θ, ϕ) landscape (Fig.2f). Coverage analysis based on highest-density regions (HDR) confirms that the MD conformational landscape encapsulates nearly all CryoUNI-generated samples (Supplementary Fig. 2b). As expected, cryo-EM captures the most populated structural states, while MD additionally further explores surrounding thermal fluctuations.

Finally, we selected three representative conformations (Sample 1–3) spanning the angular range of leg motion from the (θ, ϕ) distribution in Fig. 2f. For each sample point, we extract the corresponding CryoUNI-reconstructed density volume and the MD atomic snapshot with the closest (θ, ϕ) coordinates. The leg orientations observed in the CryoUNI density maps and the MD-derived structures are highly consistent (Fig.2g), as shown for three samples with polar angles θ of 44.68°, 49.69°, and 59.37°, and azimuthal angles ϕ of -26.04°, -24.04°, and -26.02°, respectively. As θ progressively increased, both CryoUNI and MD showed a gradual increase in leg swing amplitude. At each sample point, the rotation direction, amplitude, and local structural details all demonstrate high similarity between the CryoUNI reconstruction and the MD model. Moreover, the leg COM positions (labeled as Leg COMs in Fig.2g) aligned well with the corresponding (θ, ϕ) coordinates. CryoUNI not only recapitulates the global conformational landscape but also reconstructs density maps that faithfully mirror atomic-level dynamics.

Together, these results provide evidence that the latent space of CryoUNI is not a mathematical artifact of dimensionality reduction but rather a deeply physically grounded conformational manifold. It decodes the intrinsic conformational degrees of freedom that govern the structural dynamics of integrin $\alpha v \beta 8$.

4 Structural states and transitions emerge from the latent landscape

We next analyze the intrinsic structure of the latent landscape to resolve molecular heterogeneity (Fig. 1d,e). We introduce WAVE (Watershed Analysis of Variational Embeddings), a landscape-based approach that operates directly on latent density. WAVE automatically identifies conformational states as density peaks and reconstructs transitions by tracing high-density pathways across the landscape, without imposing clustering assumptions or requiring prior knowledge of the number of states. Given the physical correspondence between latent space and molecular conformational space established above, density directly reflects the occupancy, and distances correspond to structural variation. This formulation further defines an effective energy landscape described by Boltzmann statistics, in which high-density regions correspond to low-energy states.

We first evaluated this approach on a simulated Ribosembly benchmark [16], which models the assembly of the 50S ribosomal subunit comprising 16 discrete compositional states. CryoUNI yields a well-organized latent landscape in which individual assemblies occupy compact, high-density regions (Extended Data Fig. 2a). WAVE automatically detects 16 regions as distinct density peaks and recovers all ground-truth states with high accuracy (99.4%). In contrast, other trained-from-scratch baselines achieve lower accuracy despite being provided with the correct number of clusters (Extended Data Fig. 2d), highlighting the importance of the quality of the latent representation itself. These results show that discrete compositional states emerge naturally as topological features of the latent landscape of CryoUNI. Beyond state identification, pairwise latent distances between density peaks strongly correlate with ground-truth structural similarity measured by normalized cross-correlation (Mantel test, $r = -0.887$, $p = 0.001$; Extended Data Fig. 2e). This further confirms that the latent landscape not only separates distinct states but also preserves their structural relationships, with neighboring peaks corresponding to structurally similar assemblies. The complete quantitative comparisons across more baselines are provided in Extended Data Table 1.

To assess scalability to larger compositional mixtures, we evaluated the Tomotwin-100 dataset, which contains 100 distinct structural classes spanning a broad range of molecular weights. CryoUNI produces

a well-organized latent landscape in which WAVE automatically detects all 100 density peaks (Extended Data Fig. 2f). Quantitatively, CryoUNI achieves near-perfect classification accuracy (K-means: 99.98%, GMM: 99.97%, WAVE: 99.96%) regardless of the clustering algorithm used, substantially outperforming all other methods (Extended Data Fig. 2g; Extended Data Table 1). This indicates that the quality of the latent representation is the primary determinant of classification performance. In contrast, applying WAVE to latent spaces learned by other methods leads to substantially degraded performance (Extended Data Table 1), suggesting that their latent density topology does not faithfully reflect the underlying compositional structure.

Finally, we evaluated a dataset dominated by continuous conformational variability (IgG-1D), where discrete state classification is no longer applicable. The CryoUNI latent landscape captures the underlying conformational continuum as a continuous manifold, with WAVE tracing a closed trajectory that recovers the full conformational cycle (Extended Data Fig. 3a,b). To quantitatively assess reconstruction quality, we measured per-particle AUC-FSC, which evaluates how faithfully each particle's structure is recovered. CryoUNI achieves the highest mean AUC-FSC (0.368) with the smallest variance across all methods, significantly outperforming existing approaches (Extended Data Fig. 3c). These results demonstrate that the latent landscape captures not only discrete compositional mixtures but also high-frequency continuous conformational dynamics.

Together, these results demonstrate that structural states and transitions emerge directly from the density and topology of the CryoUNI latent landscape across fundamentally different forms of heterogeneity — from discrete compositional mixtures to continuous conformational dynamics. This establishes a unified representation for analyzing molecular heterogeneity without pre-defined discretization or system-specific assumptions.

5 The latent landscape reveals previously unresolved intermediate states

We next ask whether it can reveal previously unresolved states in experimental data. We analyzed a cryo-EM dataset of LIS1-mediated dynein remodeling under ATP conditions [11], which encompasses multiple stages of the activation pathway. In the global latent landscape, particles organize into well-separated high-density basins corresponding to the three major structural classes along the activation pathway: open-bent (State A), open-straight (State B), and motor-bound (State C) (Fig. 3a). The relative energy profile along the WAVE-traced transition pathway reveals these states as distinct energy minima separated by barriers, outlining the coarse organization of the activation pathway.

Within each major structural class, the landscape further reveals hierarchical organization, with multiple density peaks indicating the presence of sub-states (Extended Data Fig. 4a). We focus here on the open-straight region, where this hierarchical analysis uncovers previously unresolved heterogeneity. The local landscape reveals multiple density peaks with distinct occupancies (Fig. 3b). Particle subsets associated with these peaks yield structurally distinct reconstructions that differ in LIS1 binding stoichiometry: one population (Straight/1x) contains a single LIS1 copy, while a second (Straight/2x) shows additional density consistent with a second LIS1 molecule bound at the dynein stalk-buttress interface (Fig. 3b, inset). This low-population intermediate state forms a distinct density peak rather than merging into the dominant one, and its structural features are largely attenuated in consensus reconstructions. These results demonstrate that the latent landscape can resolve low-population intermediates defined by distinct binding stoichiometries, which conventional methods merge or overlook.

Extending this analysis across all three structural classes reveals a total of 12 sub-states from the latent landscape (Extended Data Fig. 4a,b). Within the motor-bound class, the landscape recovers the autoinhibited ϕ state and early intermediate conformations consistent with previously reported structures [11]. The open-bent class resolves multiple sub-states with distinct LIS1 binding configurations, including variations in stoichiometry that are not successfully separated using conventional approaches. All identified sub-states yield structurally consistent reconstructions at resolutions ranging from 2.75 to 6.44 Å (Extended Data Fig. 4b), confirming that density peaks correspond to genuine conformational populations. Several states achieve improved resolution compared to the original analysis, indicating that landscape-guided particle selection produces more homogeneous subsets. Every previously reported conformation is recovered, together with a new Straight/2x state that has been resolved.

Together, these results demonstrate that the latent landscape unlocks the capability for hierarchical analysis of molecular heterogeneity, from the coarse separation of dominant structural classes to the fine-grained resolution of low-population intermediates defined by distinct binding stoichiometries.

6 The latent landscape enables pathway generation and energy-guided reconstruction

Finally, we ask whether the conformational landscape can improve the reconstruction quality of density maps. In this conformational landscape described by Boltzmann statistics, density ratios directly infer relative energy, making high-density, low-heterogeneity regions a principled basis for energy-guided particle selection. We apply this principle to a cryo-EM dataset of the $KCTD5/CUL3^{NTD}/G\beta\gamma$ complex [12], in which four discrete conformations have been previously resolved but continuous dynamics have only been studied using morphing interpolations. In the latent landscape, particles organize into four well-separated high-density basins corresponding to the reported structural states (Fig. 4a). WAVE automatically identifies these four states with high agreement to the original classification (Fig. 4c), with over 91% of particles in each class correctly assigned.

We first test whether energy-guided particle selection improves reconstruction. For each state, we select particle subsets at increasing relative energy thresholds ($\Delta\Delta G_r$) derived from latent density (Fig. 4a). Across all states, energy-guided selection improves both reconstruction consistency and resolution compared to using the full particle set (Fig. 4b). Subsets at intermediate $\Delta\Delta G_r$ thresholds achieve the best performance, balancing the removal of structurally heterogeneous particles with retention of sufficient signal. These results demonstrate that latent density provides a principled criterion for particle selection without heuristic filtering.

Beyond particle selection, the same landscape captures conformational transitions. The four density basins are connected by continuous pathways, and tracing these pathways identifies two dominant conformational trajectories: A→D→C and A→B→C (Fig. 4d). The relative energy profiles along each trajectory reveal the energy barriers separating intermediate states. These trajectories recover the principal motions previously inferred from morphing interpolation between known states [12], providing direct experimental resolution of continuous conformational dynamics for this complex from cryo-EM data alone.

Together, these results show that latent density simultaneously encodes state occupancy, transition pathways, and reconstruction quality within a single representation, enabling both energy-guided particle selection and direct extraction of conformational dynamics from cryo-EM data.

7 Discussion

We have shown that, leveraging cryoUNI, cryo-EM data can be organized into a probabilistic conformational landscape in which latent density reflects state occupancy and defines a relative energy following Boltzmann statistics. This representation unifies the analysis of discrete compositional states, continuous conformational dynamics, and energy-guided particle selection, providing a physically grounded interpretation of structural heterogeneity directly from experimental data.

CryoUNI is advantageous because it shifts the focus from a computational intermediate for volume generation to a conformational landscape that is directly interpretable in physical terms. That said, previous methods have made substantial progress in modeling structural heterogeneity from cryo-EM data. Discrete classification approaches [1, 2] resolve major conformational states but require specifying the number of classes and cannot capture continuous dynamics. Continuous methods such as cryoDRGN [3], 3DVA [4], 3D Flex [5], and RECOVAR [7] represent variability in learned latent spaces, enabling volume generation along continuous pathways. More recently, foundation-model-based approaches [8, 17] leverage learned priors to improve reconstruction quality. While these methods have advanced heterogeneous reconstruction, the physical interpretability of the latent space itself has been less systematically explored. Our work builds on these advances by showing that, with a sufficiently well-organized latent space, density and distance directly encode conformational state occupancy, structural relationships, and transition pathways, establishing a physically grounded basis for downstream analysis.

The interpretation of latent density as a relative energy connects cryo-EM analysis to statistical mechanics, but several limitations remain. First, the observed state distribution may deviate from true equilibrium occupancy due to preferred orientations, uneven ice thickness, and the particle curation procedure. Second, the latent representation learns a structural signal from noisy 2D projection images, where undesired image features such as membrane and detergent densities may dominate over internal conformational changes. For such systems, incorporating structural priors could help disentangle biologically meaningful variation from environmental contributions. Last, the latent density has not been directly linked to a physical energy function, which is why we report relative rather than absolute energy; incorporating physics-informed constraints could help establish this connection.

More broadly, this work points toward a shift in how cryo-EM data are interpreted: from determining individual structures to characterizing conformational landscapes. Several directions follow naturally. Combining learned conformational landscapes with physics-based simulations may enable quantitative free energy

estimation directly anchored in experimental observation, building on the connection between cryo-EM and molecular dynamics established here. Extending the approach to in situ cryo-electron tomography [18] data could enable characterization of conformational landscapes within their native cellular context. Finally, integrating learned conformational landscapes with complementary biophysical and computational approaches may ultimately connect structural descriptions with the kinetic and thermodynamic principles that govern molecular function.

Methods

Large-scale pretraining

Large-scale particle dataset.

We constructed a large-scale particle image dataset, CryoCRAB-Particle-22M, to train and evaluate CryoUNI. The dataset comprises 22,631,283 particle images extracted from 173,997 micrographs spanning 746 distinct protein species, built upon the CryoCRAB dataset [19]. Particles were extracted using an iteratively optimized scheme: initial picking by particle picking adapted cryoUNI (Extended Data Fig. 1c) was followed by two rounds of 2D classification (200 and 50 classes, respectively), and this picking-curating-retraining cycle was repeated for six iterations to yield a high-purity particle collection. All particles are stored in HDF5 format with float16 precision at their native resolution. To support self-supervised pretraining, each particle is stored as a full-difference pair: the full image is generated by averaging all raw movie frames, while odd and even frame averages provide independent half-images whose subtraction yields a pure noise estimate. During training, geometric augmentation is applied, including random horizontal flipping and random rotation over $\{0°, 90°, 180°, 270°\}$. All images are preprocessed with CTF phase-flipping, contrast normalization, and Z-score standardization following the CryoCRAB protocol [19].

Architecture of CryoUNI.

Our model builds upon the Vision Transformer (ViT) architecture, including modern design choices from recent Transformer models [20] (Extended Data Fig. 1a). Given an input image $x \in \mathbb{R}^{H \times W}$, we first divide it into $H/P \times W/P$ non-overlapping patches with patch size $P = 16$. All patches are linearly projected into D-dimensional embeddings and concatenated with one class token and four register tokens [21] to form the input sequence. This sequence is fed into a stack of L Transformer blocks. Each block consists of a multi-head self-attention mechanism with 2D Rotary Position Embedding (RoPE) [22], followed by a SwiGLU feed-forward network. Residual connections and RMSNorm are applied around each sub-layer for training stability. We use two configurations: small ($D = 384$, $L = 12$) and base ($D = 768$, $L = 12$).

Denoising-reconstruction pretraining.

In the first pretraining stage, we adopt the denoising-reconstruction pretraining strategy from DRACO [15], using odd-even-full triplets generated from the CryoCRAB full-diff pairs as training samples. Take the encoded feature as input, a transformer-based decoder is employed to reconstruct the original image; it projects the D-dimensional encoder features to 576 dimensions and refines them through 8 Transformer blocks. Each patch token is then decoded into a $P \times P$ pixel patch. To accommodate the variable resolution of particle images without information loss from rescaling, we follow the approach of Dehghani et al. [23], dynamically converting images into token sequences at their native resolution. This decoder is discarded after pretraining. The pretraining combines masked image modeling loss on masked regions [24] (against the high-SNR full image) with Noise-to-Noise loss on visible regions (against the paired half-image), as described in [15]. To align with downstream tasks, input particle images are CTF-corrected, while the corresponding CTF modulates the decoder output before computing the loss against the original non-CTF-corrected reference images. The combined loss function is formally defined as:

$$\mathcal{L}_{\text{DRACO}} = \sum_{i \in \text{masked}} \|\text{CTF}_i \cdot \hat{x}_i - x_{\text{full},i}\|_2^2 + \sum_{j \in \text{visible}} \|\text{CTF}_j \cdot \hat{x}_j - x_{\text{half},j}\|_2^2 \tag{1}$$

where $\hat{x}$ represents the decoder output. We use the AdamW optimizer with a base learning rate of 1.5×10^{-4}, weight decay of 0.05, and base EMA rate of 0.995. The image mask ratio is set to 0.75. The model is trained for 800 epochs with a batch size of 4096, including a learning rate warmup for the first 40 epochs.

Denoising pretraining.

In the second pretraining stage, we further enhance the model's denoising capability to capture fine-grained structural details. We leverage odd-even image pairs based on the Noise-to-Noise principle [14] and integrate CryoUNI with the Dense Prediction Transformer (DPT) [25] framework to construct a multi-scale denoising network. Feature maps are extracted from CryoUNI at every Transformer block and transformed into a multi-scale feature pyramid, which is progressively upsampled and fused through cascaded interpolation and residual convolution. To preserve frequency-domain integrity, a two-layer convolutional network preprocesses the input image and adds it to the final fused features before producing the denoised output through another two-layer convolutional network (Extended Data Fig. 1d). The Noise-to-Noise objective is optimized using

the expected loss between the predicted output from one half-image and its paired half-image:

$$\mathcal{L}_{\text{N2N}} = \|f_\theta(x_{\text{odd}}) - x_{\text{even}}\|_2^2 + \|f_\theta(x_{\text{even}}) - x_{\text{odd}}\|_2^2 \tag{2}$$

where f_θ denotes the denoising network. To minimize this loss, we employ the AdamW optimizer with a base learning rate of 1.0e-4, a weight decay of 0.05, and a base exponential moving average (EMA) rate of 0.99. The batch size is 2048, and the model is trained for a total of 50 epochs, with a learning rate warmup applied for the first 5 epochs.

Heterogeneous reconstruction

Model architecture.

To resolve distinct conformational states of proteins from noisy particle images, we developed a heterogeneity reconstruction framework based on a Variational Autoencoder (VAE). This framework adopts the fundamental architecture of cryoDRGN [3] with optimizations in both the encoder and decoder components (Extened Data Fig. 1d). On the encoder side, we integrated a pretrained CryoUNI for end-to-end fine-tuning; it extracts class and patch tokens from images and employs a linear neck to aggregate local features, which are subsequently mapped to a latent space to generate embeddings $\mathbf{z}$ representing conformational heterogeneity. For the decoder, we implemented a volume decoder based on implicit neural representations. Taking 3D coordinates and embeddings $\mathbf{z}$ as inputs, the decoder performs high-frequency encoding of coordinates via random Fourier features with 384 default Fourier embedding channels. The intensities in the Hartley domain are predicted using a 3-layer SwiGLU network with 384 embedding channels and residual connection and LayerNorm.

Adaptation.

The training of the heterogeneous reconstruction model was optimized using the AdamW optimizer. We set the base learning rate to 1×10^{-4}, which was scheduled following a cosine decay strategy. Training was performed on a single NVIDIA A40 GPU with a total batch size of 32 and a default training epochs of 50. To balance reconstruction fidelity with latent space regularization, the VAE objective function was weighted:

$$\mathcal{L}_{\text{VAE}} = \|\hat{x} - x\|_2^2 + \beta \mathcal{D}_{\text{KL}}(q_\phi(z|x) \| p(z)) \tag{3}$$

where $\mathcal{L}_{\text{VAE}}$ measures the data likelihood (reconstruction error in the Hartley domain), $\mathcal{D}_{\text{KL}}$ is the Kullback-Leibler divergence between the approximate posterior $q_\phi(z|x)$ and the prior $p(z)$, and the coefficient β is specifically set to 0.0125.

Watershed analysis of variational embeddings (WAVE)

Density estimation and peak detection.

To characterize the conformational landscape, we project CryoUNI features into a D-dimensional latent space ($D \in \{2, 3, 4\}$) via principal component analysis (PCA), yielding a set of latent coordinates $\{\mathbf{z}_i\}_{i=1}^N$. Similar to RECOVAR [7], a continuous probability density field is then estimated using kernel density estimation (KDE):

$$\hat{p}(\mathbf{z}) = \frac{1}{Nh^D} \sum_{i=1}^N K\left(\frac{\mathbf{z} - \mathbf{z}_i}{h}\right), \tag{4}$$

where $K(\cdot)$ is an isotropic Gaussian kernel and the bandwidth h is determined by Scott's rule. The density field is evaluated on a uniform grid with resolution 256^2 ($D = 2$), 128^3 ($D = 3$), or 64^4 ($D = 4$). Density peaks are identified as local density maxima using a multi-dimensional maximum filter with a 4^D neighborhood. To suppress noise-induced spurious peaks, only maxima with density at least 5% of the global maximum are retained.

Transition pathway analysis.

To trace continuous transition pathways between structural states, we adopt a coarse-to-fine strategy. First, a Traveling Salesman Problem (TSP) solver determines the global ordering of density peaks by minimizing the total path length in latent space, ensuring that the traversal follows physically plausible transitions through high-density regions. Local pathways between adjacent peaks are then refined by solving the Eikonal equation:

$$\|\nabla T(\mathbf{z})\| = c(\mathbf{z}), \quad \text{where} \quad c(\mathbf{z}) = -\log \hat{p}(\mathbf{z}), \tag{5}$$

where $T(\mathbf{z})$ represents the cumulative transition cost. The cost function $c(\mathbf{z})$ is the negative log-density, consistent with the potential energy under Boltzmann statistics, guiding pathways through the highest-density regions of the landscape. We solve this equation on the previously defined grid using the Fast Marching Method (FMM) to extract the most probable transition pathway between states.

Partitioning and particle selection.

To define the boundaries of structural states, we apply a marker-controlled watershed algorithm using the detected density peaks as initial markers. This partitions the latent space into non-overlapping domains, and each particle is assigned to a domain based on its latent coordinate $\mathbf{z}_k$. To enhance conformational purity within each domain, we apply an energy-based filtering criterion. For the i-th state, only particles satisfying the following condition are retained:

$$\hat{p}(\mathbf{z}_k) \geq \hat{p}_i \cdot \exp(-\Delta\Delta G_r), \tag{6}$$

where $\hat{p}_i$ is the peak density of the i-th conformational center and $\Delta\Delta G_r$ is the relative energy threshold. This selects particles within a defined energy range of each density peak.

Metrics

Resolution estimation via Fourier shell correlation (FSC).

The resolution of cryo-EM reconstructions was estimated using the Fourier Shell Correlation (FSC) criterion [26]. The FSC measures the normalized cross-correlation between two independently reconstructed half-maps as a function of spatial frequency. For a given spatial frequency shell k, the FSC is defined as:

$$\mathrm{FSC}(k) = \frac{\sum_{r \in \mathrm{shell}(k)} F_1(r) \cdot F_2^*(r)}{\sqrt{\sum_{r \in \mathrm{shell}(k)} |F_1(r)|^2 \cdot \sum_{r \in \mathrm{shell}(k)} |F_2(r)|^2}} \tag{7}$$

where $F_1(r)$ and $F_2(r)$ are the complex structure factors of the two half-maps at spatial frequency vector r, and $F_2^*(r)$ denotes the complex conjugate of $F_2(r)$. The overall resolution was determined at the gold-standard FSC = 0.143 threshold [27].

To provide a more comprehensive measure of global structural fidelity, we additionally computed the Area Under the Curve of the FSC (AUC-FSC) proposed by Jeon et al. [16]:

$$\mathrm{AUC\text{-}FSC} = \int_0^{k_{\max}} \max(0, \mathrm{FSC}(k)) \, dk \tag{8}$$

where $k_{\max}$ is the Nyquist frequency. By integrating the correlation over all spatial frequency shells, this metric captures the cumulative signal quality across all resolution bands, rather than relying on a single-threshold estimate. To account for the stochastic effects of random particle half-set splitting, we performed multiple independent reconstruction trials for each state and report the mean and variance of the resulting resolution estimates.

Normalized cross-correlation (NCC) and Mantel test.

To assess whether the geometric organization of the CryoUNI latent landscape faithfully preserves the physical relationships between conformational states, we evaluated the correlation between pairwise latent distances and 3D structural similarities. The structural similarity between ground-truth volumes was quantified using the Normalized Cross-Correlation (NCC) score:

$$\mathrm{NCC}(X, Y) = \frac{\sum_i (X_i - \bar{X})(Y_i - \bar{Y})}{\sqrt{\sum_i (X_i - \bar{X})^2 \sum_i (Y_i - \bar{Y})^2}} \tag{9}$$

where X_i and Y_i represent the voxel intensities of two aligned 3D volumes, and $\bar{X}$ and $\bar{Y}$ are their respective mean voxel intensities. The pairwise distance between states in the latent representation was computed as the Euclidean distance between their corresponding detected density peaks:

$$D_{ij} = \|z_i - z_j\|_2 \tag{10}$$

The correspondence between the latent topology and true structural variance was evaluated using the Mantel test [28], which assesses the correlation between the pairwise latent distance matrix and the NCC similarity

matrix while accounting for the non-independence of distance matrix elements. The test yields the Pearson correlation coefficient r and a permutation-based p-value (P_{Mantel}).

Clustering accuracy.

To quantitatively evaluate the ability of the learned latent representations to disentangle distinct conformational states, we measured the clustering accuracy by comparing predicted cluster labels against ground-truth class labels. The clustering accuracy is defined as the maximum proportion of correctly assigned particles over all possible label permutations between the predicted labels $\hat{y}_i$ and the ground-truth labels y_i:

$$\text{Accuracy} = \max_m \frac{1}{N} \sum_{i=1}^{N} \mathbb{I}(y_i = m(\hat{y}_i)) \tag{11}$$

where N is the total number of particles, $\mathbb{I}(\cdot)$ is the indicator function, and m is a permutation mapping that assigns each predicted cluster to a ground-truth class. The optimal mapping m was solved using the Hungarian algorithm.

Adjusted Rand Index (ARI) and Adjusted Mutual Information (AMI).

To evaluate clustering quality on the compositional heterogeneity benchmarks (Ribosembly and Tomotwin-100), we additionally report the Adjusted Rand Index (ARI) and Adjusted Mutual Information (AMI). Both metrics compare predicted cluster assignments against ground-truth labels and are corrected for chance, yielding a value of 0 for random labelling and 1 for perfect agreement. ARI measures the fraction of particle pairs that are consistently co-clustered or separated, adjusted by the expected value under a random permutation model [29]. AMI quantifies the mutual information between the two partitions, normalized by the entropy of the marginals and adjusted for chance agreement [30].

Data

Integrin $\alpha v \beta 8$ complex.

We analyzed the EMPIAR-10345 dataset [10], which captures the integrin $\alpha v \beta 8$ bound to latent transforming growth factor-β (L-TGF-β) and inhibitory antibody fragments. From 1,577 micrographs (after excluding 67 low-quality images), we extracted 559,086 particles using the CryoUNI particle picker (confidence threshold 0.0001) and downsampled to a box size of 128 pixels (3.152 Å/pix). After 2D classification in cryoSPARC, 110,148 clean particles were retained. We performed iterative refinement through three rounds of CryoUNI heterogeneous reconstruction ($z = 8$) with 10-component GMM clustering, progressively removing poorly resolved particles and resolving two compositional states (Fab-bound and unbound). After excluding Fab-lacking particles, a compositionally homogeneous stack of 57,546 Fab-bound particles was re-extracted at the unbinned pixel size (1.345 Å/pix, box size 300) and subjected to final non-uniform refinement (3.97 Å). A third round of heterogeneous reconstruction ($z = 8$) was performed on this stack for the final conformational landscape analysis.

KCTD5/CUL3^{NTD}/G$\beta\gamma$ complex.

We analyzed the EMPIAR-11734 dataset [12] to resolve the conformational landscape of the KCTD5/CUL3$^{\text{NTD}}$/G$\beta\gamma$ complex. We directly imported the curated particle stacks from the original study, yielding 202,484 particles (box size 324, pixel size 1.03 Å/pix) previously assigned to four states: State A (48,977), State B (48,890), State C (57,006), and State D (47,975). CryoUNI heterogeneous reconstruction was trained for 50 epochs with an 8-dimensional latent space ($z = 8$). WAVE partitioned the latent landscape into four conformational basins consistent with the original classification. To resolve continuous transitions between states, we performed PCA on the latent embeddings and extracted transitional volumes along WAVE-traced pathways through the principal component subspace.

LIS1-mediated dynein activation.

We analyzed the EMPIAR-12715 dataset [11] to resolve the conformational landscape of LIS1-mediated dynein remodeling. From 12,753 usable micrographs, the CryoUNI particle picker (confidence threshold 0.0001) identified 4,943,683 particles. After two rounds of 2D classification in cryoSPARC, 965,465 particles were retained. Ab initio reconstruction (6 classes) followed by heterogeneous refinement separated the dataset into three major structural classes: open-bent (230,620 particles), open-straight (346,948 particles), and motor-bound (143,010 particles). Particles were re-extracted at 0.935 Å/pix (box size 352) and refined to 2.64, 3.03, and 2.71 Å, respectively.

To construct the global conformational landscape, poses from the three classes were aligned using cryoSPARC Align 3D, and the combined particle set was used to train a global CryoUNI reconstruction model for 50 epochs. Using WAVE, the global latent landscape recovered the three major classes with refined particle distributions: open-bent (165,014 particles), open-straight (346,896 particles), and motor-bound (208,668 particles). To resolve sub-states within each class, the three subsets were processed independently with CryoUNI for 50 epochs, generating local conformational landscapes. Latent dimensions were optimized per class ($z = 8$ for motor-bound, $z = 4$ for open-bent and open-straight). WAVE partitioned each local landscape into sub-states, and the corresponding particle subsets were re-imported into cryoSPARC for final high-resolution reconstruction.

CryoBench datasets.

We evaluate CryoUNI on three simulated datasets from the CryoBench benchmark [16], each targeting a distinct form of structural heterogeneity. **Ribosembly** comprises 335,240 particle images (128×128, 6 Å/pix, SNR 0.01) modeling the stepwise assembly of the 50S ribosomal subunit across 16 discrete compositional states, providing a benchmark for state identification and latent space organization. **Tomotwin-100** contains 100,000 particle images (128×128, 9 Å/pix, SNR 0.01) spanning 100 distinct structural classes with a broad range of molecular weights, testing scalability to high-complexity compositional mixtures. **IgG-1D** contains 100,000 particle images (128×128, 6 Å/pix, SNR 0.01) simulating continuous conformational variability of an IgG antibody along a single degree of freedom, assessing reconstruction fidelity across the conformational landscape.

For Ribosembly, we train CryoUNI for 25 epochs with latent dimension $D = 4$. For Tomotwin-100, we train CryoUNI for 100 epochs with latent dimension $D = 8$. For IgG-1D, we train CryoUNI for 20 epochs with latent dimension $D = 8$. All models use a batch size of 32 and the AdamW optimizer with a learning rate of 1×10^{-4} and KL divergence weight of 0.0125. All experiments were conducted on a single NVIDIA A40 48G GPU. We compare against cryoDRGN [3], 3DVA [4], RECOVAR [7], OPUS-DSD [6], CryoDECO [8], and 3D Flex [5] as baselines. Detailed configurations for each baseline are provided in the Baselines section below.

Baselines

All methods were run using their official implementations with default hyperparameters unless otherwise noted. Latent embeddings from each method were used for downstream clustering (K-means, GMM) and WAVE analysis.

CryoDRGN.

CryoDRGN [3] models continuous structural heterogeneity using an amortized variational autoencoder. Each particle image is encoded into a low-dimensional latent variable, from which a coordinate-based neural network decodes 3D density volumes in Fourier space. We used the official CryoDRGN implementation (version 3.4.4) with default network architecture. The model was trained for 25 epochs with a latent dimension of 8 on each dataset. Image poses were fixed to the consensus values estimated by cryoSPARC.

3DVA.

3D Variability Analysis (3DVA) [4] fits a linear subspace model to cryo-EM data, representing conformational variability as a set of orthogonal principal modes. Unlike neural-network-based approaches, 3DVA assumes a Gaussian latent distribution and models heterogeneity within a linear framework. We used the 3DVA implementation in cryoSPARC (version 4.5). The analysis was performed with 3 variability components and a filter resolution of 6 Å. The resulting variability coefficients were used as latent coordinates for downstream analysis.

RECOVAR.

RECOVAR [7] estimates a regularized covariance matrix of the heterogeneous 3D density distribution and performs kernel regression in the resulting principal component space to reconstruct conformational variability. We used the official implementation with default parameters. The model was applied with 10 principal components for latent embedding on each dataset.

OPUS-DSD.

OPUS-DSD [6] disentangles dynamics and compositional heterogeneity in cryo-EM data using a deep learning framework with separate latent spaces and decoders for large-scale dynamics and compositional changes. We used the official implementation with default hyperparameters. The model was trained for 25 epochs with a latent dimension of 12 on each dataset.

CryoDECO.

CryoDECO [8] leverages foundation model priors to improve heterogeneous cryo-EM reconstruction, enabling deconstructing of extreme compositional and conformational heterogeneity. We used the official implementation with default hyperparameters. The model was trained for 25 epochs with a latent dimension of 128 on each dataset.

3DFlex.

3DFlex [5] models continuous conformational flexibility by learning a non-rigid deformation field that maps a canonical density to each individual particle conformation. We used the 3D Flex implementation in cryoSPARC (version 4.5) with 2 latent dimensions and default parameters for both mesh preparation and training.

WAVE + CryoSPARC analysis.

Based on the clustering and filtering results obtained via WAVE, the particle subsets corresponding to each cluster were imported into CryoSPARC(version 4.5). For each subset, an initial homogeneous reconstruction was performed to generate a preliminary three-dimensional volume. Subsequently, these volumes were subjected to non-uniform refinement, utilizing cross-validation-based regularization and anisotropic spatial smoothing to achieve high-resolution 3D reconstructions for each identified static state.

Molecular dynamics simulation of integrin $\alpha v \beta 8$

Construction of the integrin $\alpha v \beta 8$ ectodomain model

The cryo-EM structure of integrin $\alpha v \beta 8$ in complex with the Fabs C6D4 and 11D12v2 (PDB ID: 6UJB) was used as the initial experimental model. In the original study, an additional Fab, 11D12v2, was introduced to facilitate high-resolution reconstruction by binding to the αV thigh region. The authors noted that this Fab was used primarily to improve particle alignment and did not affect L-TGF-β binding or the conformation of the integrin headpiece [10]. Thus, 11D12v2 was not considered a functional component relevant to the conformational dynamics investigated here. In addition, the deposited coordinates for 6UJB contain only four chains that could be directly used for subsequent modelling, corresponding to integrin αv, integrin $\beta 8$, the Fab C6D4 heavy chain, and the Fab C6D4 light chain, whereas no usable atomic coordinates were provided for 11D12v2. On the basis of both its auxiliary role in structure determination and the absence of a corresponding deposited atomic model, all subsequent structure reconstruction and molecular dynamics simulations were performed using only these four chains.

In 6UJB, the leg of chain A (integrin αv) and chain B (integrin $\beta 8$) are incomplete. To obtain a complete ectodomain structure, we reconstructed the αv and $\beta 8$ chains using AlphaFold 3, with a previously determined integrin $\alpha v \beta 3$ structure (PDB ID: 8XEN) used as a structural template to guide modelling. The rebuilt αv and $\beta 8$ chains covered both the headpiece and leg regions used in subsequent analyses and were then combined with the experimentally resolved Fab C6D4 chains from 6UJB to generate a complete integrin $\alpha v \beta 8$-Fab ectodomain complex model. Because this study focused on the continuous extracellular conformational heterogeneity of the integrin leg, the transmembrane and intracellular regions were not included in the modelling or simulations.

System preparation

The reconstructed integrin $\alpha v \beta 8$-Fab ectodomain complex was prepared using the Protein Preparation Wizard in Schrödinger Release 2025-4 [31]. During preparation, bond orders were assigned, hydrogen atoms were added, and the hydrogen-bonding network was optimized. The experimentally resolved Ca^{2+} and Mg^{2+} ions were retained throughout to preserve the native metal-coordination environment of the integrin ectodomain. Protonation states of hetero groups were assigned with Epik at pH 7.4 $\pm$ 2.0, followed by restrained minimization of the prepared structure.

The complex was solvated in TIP3P water in a rhombic dodecahedral box with a minimum solute-box distance of 1.2 nm, corresponding to a minimum separation of greater than 2.4 nm between the protein and its nearest periodic image. To mimic physiological conditions, the system was neutralized with counterions and supplemented with 0.15 M NaCl.

All-atom molecular dynamics simulations

All-atom molecular dynamics simulations were performed using GROMACS 2025.1 with the CHARMM36m force field [32, 33]. Following solvation and ion addition, each system was subjected to a two-step energy minimization procedure consisting of steepest-descent and conjugate-gradient minimization. After energy minimization, the system was equilibrated in two successive stages with positional restraints applied to

the protein. First, a 1 ns NVT equilibration was carried out at 310 K with a 2 fs integration time step. Temperature was controlled using the V-rescale thermostat, with the protein and water-and-ions treated as separate coupling groups, each with a coupling constant of 1.0 ps. This was followed by a 1 ns NPT equilibration at 310 K and 1 bar using the same temperature-coupling scheme. Pressure was maintained using the C-rescale barostat with isotropic pressure coupling, a coupling time constant of 5.0 ps, and a compressibility of 4.5×10^{-5} bar^{-1}. In both equilibration stages, non-bonded interactions were treated using the Verlet cutoff scheme; long-range electrostatics were calculated using the particle mesh Ewald (PME) method [34] with a real-space cutoff of 1.2 nm; and van der Waals interactions were treated using a force-switch scheme, beginning at 1.0 nm and reaching zero at 1.2 nm. Bonds involving hydrogen atoms were constrained using the LINCS algorithm[35].

Production simulations were subsequently carried out in the NPT ensemble after release of positional restraints, using the same temperature, pressure, and non-bonded interaction settings as in equilibration. To ensure extensive conformational sampling of the integrin $\alpha v\beta 8$ leg, we initiated 20 independent trajectories from the same starting model using different random initial velocities. Each replicate was simulated for a total of 1 μs, yielding an aggregate sampling time of 20 μs. Together, these trajectories constituted the MD conformational ensemble used for subsequent landscape analysis. To exclude the initial relaxation phase, only the 200-1000 ns portion of each trajectory was used for quantitative analysis, with conformations sampled at 0.1 ns intervals. This yielded a total of 160,000 MD snapshots for subsequent landscape construction and quantitative comparison.

8 Data availability

The raw cryo-EM datasets used in this study are publicly available from EMPIAR under accession codes 10345 (integrin $\alpha v\beta 8$), 11734 (KCTD5/CUL3$^{\text{NTD}}$/G$\beta\gamma$), 12715 (LIS1/dynein), and 13047 (SSU processome). The simulated benchmark datasets (Ribosembly, Tomotwin-100, and IgG-1D) are available through CryoBench [16].

9 Code availability

Source code of CryoUNI, pre-trained model weights, and training, inference scripts are available at https://github.com/Cellverse/cryouni.

References

[1] Scheres SHW. RELION: Implementation of a Bayesian approach to cryo-EM structure determination. Journal of Structural Biology. 2012;180(3):519–530.

[2] Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. Nature Methods. 2017 Mar;14(3):290–296.

[3] Zhong ED, Bepler T, Berger B, Davis JH. CryoDRGN: reconstruction of heterogeneous cryo-EM structures using neural networks. Nature methods. 2021;18(2):176–185.

[4] Punjani A, Fleet DJ. 3D variability analysis: Resolving continuous flexibility and discrete heterogeneity from single particle cryo-EM. Journal of Structural Biology. 2021;213(2):107702.

[5] Punjani A, Fleet DJ. 3DFlex: determining structure and motion of flexible proteins from cryo-EM. Nature Methods. 2023;20(6):860–870.

[6] Luo Z, Ni F, Wang Q, Ma J. OPUS-DSD: deep structural disentanglement for cryo-EM single-particle analysis. Nature Methods. 2023;20(11):1729–1738.

[7] Gilles MA, Singer A. Cryo-EM heterogeneity analysis using regularized covariance estimation and kernel regression. Proceedings of the National Academy of Sciences. 2025;122(9):e2419140122.

[8] Yan Y, Xi Y, Fan S, Wang Y, Tang Z, Yuan F, et al. CryoDECO: Deconstructing Extreme Compositional and Conformational Heterogeneity in Cryo-EM via Foundation Model Priors. LangTaoSha Preprint Server. 2026 Mar;.

[9] Vulović M, Ravelli RBG, van Vliet LJ, Koster AJ, Lazić I, Lücken U, et al. Image formation modeling in cryo-electron microscopy. Journal of Structural Biology. 2013;183(1):19–32.

[10] Campbell MG, Cormier A, Ito S, Seed RI, Bondesson AJ, Lou J, et al. Cryo-EM reveals integrin-mediated TGF-β activation without release from latent TGF-β. Cell. 2020;180(3):490–501.

[11] Nguyen KH, Karasmanis EP, Kendrick AA, Reck-Peterson SL, Leschziner AE. Cryo-EM captures early intermediate steps in dynein activation by LIS1. Nature Communications. 2025;16(1):7054.

[12] Nguyen DM, Rath DH, Devost D, Pétrin D, Rizk R, Ji AX, et al. Structure and dynamics of a pentameric KCTD5/CUL3/G$\beta\gamma$ E3 ubiquitin ligase complex. Proceedings of the National Academy of Sciences. 2024;121(17):e2315018121.

[13] Andreeva A, Kulesha E, Gough J, Murzin AG. The SCOP database in 2020: expanded classification of representative family and superfamily domains of known protein structures. Nucleic Acids Research. 2019 11;48(D1):D376–D382.

[14] Lehtinen J, Munkberg J, Hasselgren J, Laine S, Karras T, Aittala M, et al. Noise2Noise: Learning image restoration without clean data. arXiv preprint arXiv:180304189. 2018;.

[15] Shen Y, Dai H, Chen Q, Zeng Y, Zhang J, Pei Y, et al. DRACO: A Denoising-Reconstruction Autoencoder for Cryo-EM. Advances in Neural Information Processing Systems. 2024;38.

[16] Jeon M, Raghu R, Astore M, Woollard G, Feathers R, Kaz A, et al. CryoBench: Diverse and challenging datasets for the heterogeneity problem in cryo-EM. Advances in Neural Information Processing Systems. 2024;37:89468–89512.

[17] Yan Y, Fan S, Yuan F, Shen H. A comprehensive foundation model for cryo-EM image processing. Nature Methods. 2026;23(1):88–95.

[18] Lučić V, Rigort A, Baumeister W. Cryo-electron tomography: the challenge of doing structural biology in situ. Journal of Cell Biology. 2013;202(3):407–419.

[19] Chen Q, Xu Z, Dai H, Shen Y, Zhang J, Liu Z, et al. A large-scale curated and filterable dataset for cryo-EM foundation model pre-training. Scientific data. 2025;12(1):960.

[20] Grattafiori A, Dubey A, Jauhri A, Pandey A, Kadian A, Al-Dahle A, et al. The Llama 3 herd of models. In: Neural Information Processing Systems. Curran Associates; 2024. .

[21] Darcet T, Oquab M, Mairal J, Bojanowski P. Vision Transformers Need Registers. In: International Conference on Learning Representations (ICLR); 2024. .

[22] Su J, Ahmed M, Lu Y, Pan S, Bo W, Liu Y. Roformer: Enhanced transformer with rotary position embedding. Neurocomputing. 2024;568:127063.

[23] Dehghani M, Mustafa B, Djolonga J, Heek J, Minderer M, Caron M, et al. Patch n'pack: Navit, a vision transformer for any aspect ratio and resolution. Advances in Neural Information Processing Systems. 2023;36:2252–2274.

[24] He K, Chen X, Xie S, Li Y, Dollár P, Girshick R. Masked autoencoders are scalable vision learners. In: Proceedings of the IEEE/CVF conference on computer vision and pattern recognition; 2022. p. 16000–16009.

[25] Ranftl R, Bochkovskiy A, Koltun V. Vision transformers for dense prediction. In: Proceedings of the IEEE/CVF international conference on computer vision; 2021. p. 12179–12188.

[26] Harauz G, van Heel M. Exact filters for general geometry three dimensional reconstruction. Optik. 1986;73(4):146–156.

[27] Rosenthal PB, Henderson R. Optimal determination of particle orientation, absolute hand, and contrast loss in single-particle electron cryomicroscopy. Journal of molecular biology. 2003;333(4):721–745.

[28] Mantel N. The detection of disease clustering and a generalized regression approach. Cancer research. 1967;27(2_Part_1):209–220.

[29] Hubert L, Arabie P. Comparing partitions. Journal of Classification. 1985;2(1):193–218.

[30] Vinh NX, Epps J, Bailey J. Information theoretic measures for clusterings comparison: is a correction for chance necessary? In: Proceedings of the 26th annual international conference on machine learning; 2009. p. 1073–1080.

[31] Madhavi Sastry G, Adzhigirey M, Day T, Annabhimoju R, Sherman W. Protein and ligand preparation: parameters, protocols, and influence on virtual screening enrichments. Journal of computer-aided molecular design. 2013;27(3):221–234.

[32] Huang J, MacKerell Jr AD. CHARMM36 all-atom additive protein force field: Validation based on comparison to NMR data. Journal of computational chemistry. 2013;34(25):2135–2145.

[33] Huang J, Rauscher S, Nawrocki G, Ran T, Feig M, De Groot BL, et al. CHARMM36m: an improved force field for folded and intrinsically disordered proteins. Nature methods. 2017;14(1):71–73.

[34] Darden T, York D, Pedersen L, et al. Particle mesh Ewald: An N log (N) method for Ewald sums in large systems. Journal of chemical physics. 1993;98(12):10089–10092.

[35] Hess B. P-LINCS: A parallel linear constraint solver for molecular simulation. Journal of chemical theory and computation. 2008;4(1):116–122.

[36] Davis JH, Tan YZ, Carragher B, Potter CS, Lyumkis D, Williamson JR. Modular assembly of the bacterial large ribosomal subunit. Cell. 2016;167(6):1610–1622.

[37] Buzovetsky O, Klinge S. Helicase-mediated mechanism of SSU processome maturation and disassembly. Nature. 2025;648(8094):746–754.

[38] Bai Xc, Fernandez IS, McMullan G, Scheres SH. Ribosome structures to near-atomic resolution from thirty thousand cryo-EM particles. eLife. 2013 feb;2:e00461.

[39] Allegretti M, Klusch N, Mills DJ, Vonck J, Kühlbrandt W, Davies KM. Horizontal membrane-intrinsic alpha-helices in the stator a-subunit of an F-type ATP synthase. Nature. 2015 May;521(7551):237–240.

[40] Nguyen THD, Galej WP, Bai Xc, Oubridge C, Newman AJ, Scheres SHW, et al. Cryo-EM structure of the yeast U4/U6.U5 tri-snRNP at 3.7 Å resolution. Nature. 2016;530(7590):298–302.

[41] Lubbe L, Sewell BT, Woodward JD, Sturrock ED. Cryo-EM reveals mechanisms of angiotensin I-converting enzyme allostery and dimerization. The EMBO Journal. 2022 Jul;41(16):EMBJ2021110550.

[42] Oquab M, Darcet T, Moutakanni T, Vo H, Szafraniec M, Khalidov V, et al. DINOv2: Learning Robust Visual Features without Supervision. Transactions on Machine Learning Research Journal. 2024;.

Fig. 1 **CryoUNI learns latent conformational landscapes from single-particle cryo-EM data. a**, Composition of the CryoCRAB-Particle-22M pretraining dataset across SCOP2 structural classes and molecular weight ranges. **b**, Self-supervised pretraining of CryoUNI. Particles from one half-dataset are denoised using the complementary half as supervision, and we demonstrate particle images of 50S ribosomal intermediates [36]. **c**, Variational autoencoder architecture for downstream conformational analysis. A lightweight linear projector maps particle features into a low-dimensional embedding space, from which a neural volume decoder reconstructs the corresponding density volumes. **d**, WAVE (Watershed Analysis of Variational Embeddings) identifies discrete conformational states as local density maxima on the conformational landscape. Relative free energies are inferred from density ratios *via* $\Delta G_r = -k_B T \ln(\rho_A/\rho_B)$. **e**, Continuous conformational trajectories traced between structural states, showing structural evolution and relative population of intermediates of yeast SSU processome assembly intermediates [37].

Fig. 2 Physical grounding of CryoUNI latent conformational landscapes for integrin $\alpha v \beta 8$ dynamics. a, Conformational heterogeneity of the flexible leg of integrin $\alpha v \beta 8$. Overlay of the published consensus cryo-EM map (EMPIAR-10345, grey surface) and representative MD snapshots (green cartoons). The distal leg density remains unresolved due to the continuous rotation captured in the MD ensemble. **b,** Unified spherical coordinate system for leg orientation. The origin is anchored at the head-leg hinge center of mass (COM); leg orientation is parameterized by polar angle θ and azimuthal angle ϕ (illustrated). A representative MD snapshot (green, non-leg regions omitted) and the cryo-EM map (grey surface) are shown. **c,** MD-derived thermodynamic conformational landscape. Two-dimensional probability density of (θ, ϕ) from 160,000 MD snapshots (twenty 1-μs trajectories; 20 μs aggregate). The leg motion is confined to a well-defined continuous region. **d,** Probabilistic latent space distribution of CryoUNI reconstructions. Principal component analysis of 10,000 decoded volumes reveals a continuous, unimodal distribution. **e,** Spontaneous disentanglement of physical coordinates. Pearson correlation between latent axes (PC1, PC2) and spherical coordinates (θ, ϕ). PC1 correlates with θ ($r = 0.982$); PC2 correlates with ϕ ($r = 0.963$). **f,** Projection of CryoUNI conformations onto the MD landscape. CryoUNI-derived conformations (orange contours) overlaid on the MD (θ, ϕ) probability density (green to navy). Three representative conformations are indicated. **g,** 3D structural validation of physical dynamics. CryoUNI density maps at three selected (θ, ϕ) coordinates (Sample 1-3: $\theta = 44.68°$, $49.69°$, $59.37°$; $\phi = -26.04°$, $-24.04°$, $-26.02°$), shown alongside their closest-matching MD atomic models. Leg orientations and local structural details are highly consistent across samples.

Fig. 3 Hierarchical CryoUNI analysis resolves the complete dynein activation pathway and novel stoichiometry from EMPIAR-12715. a, Global latent space representation of the EMPIAR-12715 dataset projected onto the first two principal components (PC 1 and PC 2). **b,** The structural relative energy along the transition pathway in **a**. The color gradient indicates particle density, revealing well-separated regions corresponding to major structural classes such as open-bent (State A), open-straight (State B), and motor-bound (State C) configurations. **c,** Hierarchical analysis of local density structures within open-straight region. Comparisons between the CryoUNI and WAVE + CryoSPARC frameworks are shown. Relative energy boundaries ($\Delta\Delta G_r$ at 1 kT and 2 kT) delineate structurally distinct basins. This progressive refinement resolves finer variability, including distinct LIS1 stoichiometries (1×LIS1 and 2×LIS1) and a newly found straight sub-state featuring a second LIS1-associated density. Structures are colored by subunit: Dynein linker (Warm Terracotta), Dynein motor (Pale Celadon), and LIS1 (Vibrant Ocean).

Fig. 4 Energy-guided reconstruction and conformational trajectory analysis of the KCTD5/CUL3$^{\mathbf{NTD}}$/G$\beta\gamma$ complex. a, CryoUNI latent landscape colored by density, with four detected peaks (A–D) corresponding to the major conformational states. Dashed lines indicate watershed boundaries (black), and relative energy contours at $\Delta\Delta G_{\mathrm{r}} = 1\,kT$ (red) and $2\,kT$ (orange). Trajectories traced by WAVE are shown in purple. **b**, Reconstruction resolution as a function of energy-guided particle selection. For each state, particle subsets are selected at increasing $\Delta\Delta G_r$ thresholds ($1kT$, $2kT$, $3kT$) and at the watershed boundary. Red dashed lines indicate the deposited resolution from the original analysis. Intermediate thresholds yield improved resolution across all states. **c**, Confusion matrix comparing WAVE-identified states with the original classification. Diagonal entries show agreement exceeding 91% for all four states without prior specification of state number. **d**, Continuous conformational pathways connecting the four states. Left: trajectory A→D→C with representative volumes colored by state identity (State A, State D, State C) and the corresponding relative energy profile (ΔG_r) along the pathway. Right: trajectory A→B→C with representative volumes and energy profile. Energy barriers between states are directly resolved from the latent density.

Extended Data Fig. 1 Network architecture of CryoUNI and its downstream applications. a, CryoUNI adopts a modern ViT architecture and undergoes a first-stage denoising-reconstruction pretraining based on the N2N and MAE frameworks. **b**, In the second-stage denoising pretraining, the model incorporates a DPT head while focusing on a specialized frequency-domain protection mechanism to preserve structural integrity. **c**, The particle picking model, built upon the Fast R-CNN object detection framework, is utilized in an iterative "picking-curating-retraining" cycle to generate the large-scale particle dataset. **d**, A CryoDRGN-style heterogeneous reconstruction framework with a specifically designed implicit neural representation volume decoder, which predicts slice intensities in the frequency domain from 3D coordinates and latent variables. Example images are from EMPIAR-10002 [38], EMPIAR-10076 [36], and EMPIAR-10345 [10].

Extended Data Fig. 2 Evaluation of the latent landscape on compositional heterogeneity benchmarks. a, Example particle images from the Ribosembly dataset. **b**, Three-dimensional visualization of the CryoUNI latent landscape for the Ribosembly dataset (16 compositional states). Each high-density region corresponds to a distinct assembly intermediate. Detected peaks are marked by circles and stars. **c**, Representative reconstructed volumes for selected Ribosembly states (8c9c, 8c92, 8c8x). Ground-truth (G.T.) volumes are compared with CryoUNI decoder outputs and WAVE-guided cryoSPARC reconstructions, with achieved resolution indicated below each map. **d**, Clustering accuracy on the Ribosembly dataset across latent representations learned by 3DVA, CryoDRGN, and CryoUNI, using GMM (k=16), K-means (k=16), and WAVE (automatic k). **e**, Correspondence between latent distance and structural similarity. Top: ground-truth pairwise NCC matrix. Bottom: pairwise latent distance matrix between detected peaks. Right: scatter plot showing strong negative correlation between latent distance and structural similarity (Mantel test, $r = -0.887$, $p = 0.001$). **f**, Three-dimensional visualization of the CryoUNI latent landscape for the Tomotwin-100 dataset (100 structural classes), colored by class index. All 100 density peaks are automatically detected by WAVE. **g**, Clustering accuracy (GMM) on the Tomotwin-100 dataset across methods. **h**, Representative reconstructed volumes for selected Tomotwin-100 classes, comparing ground-truth volumes with CryoUNI and WAVE-guided cryoSPARC reconstructions.

Extended Data Fig. 3 Evaluation of the latent landscape on the IgG-1D continuous heterogeneity benchmark. a, Three-dimensional visualization of the CryoUNI latent landscape for the IgG-1D dataset, colored by ground-truth conformational state (0 to 2π). The solid line denotes the trajectory traced by WAVE, recovering the full conformational cycle. **b**, Reconstructed volumes sampled along the WAVE trajectory, colored by conformational state as in **a**. Right: close-up view showing continuous structural variation across the conformational cycle. **c**, Per-particle reconstruction consistency measured by AUC-FSC across methods. CryoUNI achieves the highest mean (0.368) with the smallest variance, significantly outperforming all other methods ($p < 0.0001$).

* Indicates higher resolution than previously reported. ** Denotes a previously uncharacterized structure.

Extended Data Fig. 4 Hierarchical analysis of the conformational landscape for LIS1-mediated dynein activation. a, Local latent landscapes for the three major structural classes: open-straight (peaks 1–3), open-bent (peaks 4–9), and motor-bound (peaks 10–12). Multiple density peaks within each class indicate hierarchical sub-state organization. Dashed lines project peaks onto three PC planes; colored contours indicate relative energy thresholds ($\Delta\Delta G_{\mathrm{r}}$). **b**, Reconstructed volumes for all 12 identified sub-states. Top row: CryoUNI decoder outputs. Bottom row: WAVE-guided cryoSPARC reconstructions, with achieved resolution indicated below each map. Asterisks denote states with improved resolution compared to the original analysis (* moderate improvement; ** novel discovery).

Extended Data Tab. 1 Evaluation on CryoBench's compositional datasets. Performance is measured by Adjusted Rand Index (ARI), Adjusted Mutual Information (AMI), and Accuracy (Acc.) on Ribosembly and Tomotwin-100 datasets. Upward arrows (↑) indicate higher values represent better performance. The best and second-best results across all methods are highlighted in boldface and underlined, respectively.

Method	Clustering Algorithm	# Clusters	Ribosembly ($k = 16$)			Tomotwin-100 ($k = 100$)		
			ARI (↑)	AMI (↑)	Acc. (%,↑)	ARI (↑)	AMI (↑)	Acc. (%,↑)
RECOVAR	GMM	Pre-defined	0.6751	0.8201	69.68	0.3383	0.6828	48.62
	K-means	Manual	0.7317	0.8512	74.89	0.3488	0.6592	48.58
	WAVE	Auto-detected	0.7643	0.8945	77.67	0.0072	0.2884	13.77
OPUS-DSD	GMM	Manual	0.8300	0.9199	78.18	0.9771	0.9923	98.28
	K-means	Manual	0.8656	0.9252	85.22	0.9776	0.9901	98.00
	WAVE	Auto	0.8032	0.9125	82.45	0.5627	0.8727	67.53
CryoDRGN	GMM	Manual	0.9276	0.9301	92.95	0.9684	0.9877	98.33
	K-means	Manual	0.8604	0.9217	83.33	0.9702	0.9890	98.32
	WAVE	Auto	0.9072	0.9406	92.98	0.8506	0.9701	89.40
3DVA	GMM	Manual	0.5664	0.7221	56.64	0.0567	0.3326	13.28
	K-means	Manual	0.5460	0.7150	61.96	0.0575	0.3317	13.47
	WAVE	Auto	0.5652	0.7706	60.00	0.0015	0.0723	2.85
CryoDECO	GMM	Manual	0.7082	0.8408	66.98	0.9751	0.9954	97.06
	K-means	Manual	0.8889	0.9277	86.57	0.9874	0.9977	98.57
	WAVE	Auto	0.7779	0.9081	78.42	0.1523	0.8357	57.90
CryoUNI	GMM	Manual	0.9863	0.9812	97.97	<u>0.9995</u>	<u>0.9996</u>	<u>99.97</u>
	K-means	Manual	**0.9919**	**0.9882**	**99.50**	**0.9995**	**0.9996**	**99.98**
	WAVE	Auto	<u>0.9907</u>	<u>0.9870</u>	<u>99.42</u>	0.9992	0.9993	99.96

Appendix Fig. 1 Visualization of Denoising and Latent Semantics. Qualitative results of CryoUNI across diverse Cryo-EM single-particle datasets (EMPIAR-10023 [39], 10073 [40], 10345 [10], and 10980 [41]). For each example, raw and denoised particles are shown alongside their patch token visualizations. The PCA features are generated by projecting the last layer's patch tokens onto their first three principal components and mapping them to the RGB space (following the DINOv2 [42] protocol). The Multi-Head Self-Attention (MHSA) maps from the last layer illustrate the distribution of the model's focus. Together, these visualizations demonstrate the ability of CryoUNI to distill meaningful structural signals from extremely noisy backgrounds, maintaining a consistent focus on target structures at the center of the image.

Appendix Fig. 2 Convergence of MD simulations and coverage of CryoUNI conformations within the MD-derived landscape. a, Saturation of the MD conformational ensemble. Overlap between landscapes from k random 1-μs trajectories and the full 20-μs ensemble (mean $\pm$ std). Near-unity overlap as $k \to 20$ indicates convergence and sufficient sampling. b, CryoUNI conformations are nested within the MD landscape. Fraction of CryoUNI samples within the MD highest-density region (HDR) versus probability mass threshold ($\log_{10}$ scale). Nearly all CryoUNI conformations lie within the MD-sampled region across thresholds, confirming that cryo-EM captures the most populated states while MD explores surrounding fluctuations.