**Expectation triggers a change-like EEG response without acoustic change**

Haoxuan Xu[1,2,3,4,5#], Wanshun Wen[6#], Yinru Yu[7#], Ishrat Mehmood[1], Zhouyou Dai[1],

Maida Mariam[1], Baorong Zhang[6*] and Xiongjie Yu[1,2,3,4,5*]

[1] Department of Anesthesia, Women's Hospital, Zhejiang University School of

Medicine, Hangzhou, China

[2] Department of Anesthesiology, Shanghai Tenth People's Hospital, Tongji University

School of Medicine, Shanghai, China

[3] Zhejiang Provincial Key Laboratory of Precision Diagnosis and Therapy for Major

Gynecological Diseases, Women's Hospital, Zhejiang University School of Medicine,

Hangzhou, China

[4] College of Biomedical Engineering and Instrument Science, Zhejiang University,

Hangzhou, China

[5] Key Laboratory for Biomedical Engineering of Ministry of Education, Zhejiang

University, Hangzhou, China

[6] Department of Neurology of the Second Affiliated Hospital of Zhejiang University

School of Medicine, Hangzhou, China

[7] Hangzhou Entel Foreign Language School, Hangzhou, Zhejiang, China

[#] Haoxuan Xu, Wanshun Wen, and Yinru Yu contributed equally to this work.

[*] Corresponding Authors: Xiongjie Yu (yuxiongj@gmail.com) and Baorong Zhang

(brzhang@zju.edu.cn)

**Running Title**: Predictive change response

1 **Key Words**: Predictive coding, Temporal uncertainty, Electroencephalography

2 (EEG), Change detection, Click trains

4 **Author Contributions:** Conceptualization: X.Y.; Methodology: H.X., W.W., Y.Y.,

5 X.Y., B.Z.; Software: H.X., W.W.; Formal analysis: H.X., W.W.; Investigation: H.X.,

6 W.W., Y.Y., I.M.; Data curation: H.X., W.W., Y.Y.; Visualization: H.X., W.W.;

7 Writing—original draft: H.X., W.W., Y.Y., X.Y.; Writing—review & editing: H.X.,

8 W.W., Y.Y., I.M., Z.D., M.M., B.Z., X.Y.; Resources: X.Y.; Supervision: X.Y.;

9 Funding acquisition: X.Y.

11 **Competing interests:** The authors declare no competing interests.

13 **Acknowledgments:** This work was supported by Brain Science and Brain-like

14 Intelligence Technology—National Science and Technology Major Project

15 (2022ZD0204600 and 2022ZD0204800) (to X.Y.); National Natural Science

16 Foundation of China 32571216 and 32171044 (to X.Y.);

**Abstract**

Predictive processing theories posit that the auditory system continuously generates expectations and updates internal models when incoming evidence violates those expectations. Whether such updating can be expressed as a temporally precise, change-like neural response at a context-defined moment during continuous stimulation—without any physical change—has remained unclear. Using a transitional click-train paradigm in human EEG, we show that listeners reliably detect subtle midpoint transitions when click timing is fixed, whereas temporal uncertainty markedly weakens this sensory-driven detection. Strikingly, even in a physically unchanged stimulus, variable click timing elicits a robust response time-locked to the nominal midpoint. This endogenous change-like signal is stimulus-statistics dependent—absent in the fixed-timing no-change control—temporally confined to the early post-change window, and strongly state dependent, being amplified during active change detection relative to passive listening. Moreover, within identical no-change trials, the response tracks subjective change reports. Together, these findings show that temporal uncertainty and task engagement jointly gate predictive updating in continuous auditory streams, providing a compact assay to dissociate sensory-driven change responses from context- and state-dependent predictive signals.

**Introduction**

Listening is fundamentally predictive. In natural scenes, the auditory system rarely waits for sounds to "arrive" before interpreting them; instead, it continuously anticipates structure unfolding in time—when a footstep should land, when a syllable should end, when a pattern should turn. Predictive-coding and related inference frameworks formalize this intuition: the brain maintains generative models of sensory causes, evaluates incoming evidence against those models, and updates them when predictions fail [1-3]. One of the most robust population-level signatures of violated expectation is mismatch negativity (MMN), elicited when a deviant breaks an established regularity [4-6]. At the neuronal level, stimulus-specific adaptation (SSA) can likewise be interpreted as a predictive response: repeated inputs become expected and are down-weighted, whereas rare or deviant events elicit relatively preserved responses, consistent with local prediction-error signaling [7-14]. Together, these observations support the modern view that MMN-like signals reflect inference over sequence structure and the consequences of violated expectations, rather than a simple reaction to novelty [4-6,15].

A striking implication of this framework is that neural updating can be aligned to expected moments even when there is little—or no—corresponding acoustic energy. In omission paradigms, the absence of an anticipated sound evokes time-locked cortical activity, demonstrating that prediction-related signals can be anchored to internal timing rather than to stimulus onset or physical change [16,17]. Hierarchical designs further show that prediction errors are sensitive to higher-order structure:

when the brain infers that a deviant should occur at a particular position, omitting that expected deviant can elicit stronger responses than omitting a standard, consistent with multi-level predictions and errors [15,18]. Related work using nonlinguistic structured stimuli has likewise revealed hierarchical temporal processing across the primate thalamocortical system [19]. Moreover, expectations need not be driven solely by external stimulus statistics. Evidence from "phantom" or internally generated percepts suggests that endogenous perceptual states can establish predictions that yield deviance-like responses when violated [20].

Despite these advances, two gaps limit our understanding of predictive updating during continuous stimulation. First, omission-related responses are most often studied in event-based or strongly rhythmic contexts, where entrainment and phase-reset can contribute to time-locked activity, complicating the question of whether prediction-related update signals can be expressed without discrete events and without overt rhythmic scaffolding [16,17]. Second, expectation-induced effects (including "phantom deviant"-like phenomena) are rarely tested when the critical update time is embedded within continuous input and defined primarily by task context—for example, an anticipated midpoint transition—rather than by the occurrence (or omission) of discrete tokens [15,20]. Thus, it remains unclear whether change-related neural machinery can be recruited as a temporally precise, context-defined update signal even when the stimulus contains no physical change, and how such updating depends on stimulus temporal statistics and behavioral engagement.

To address these questions, we used a transitional click-train paradigm in which two continuous segments are concatenated and a nominal change point is fixed at the midpoint. Within the same dataset [21], we identified a midpoint-locked predictive change-like response that emerges without any physical change: in the no-change condition with variable click timing around the same mean (Irreg$_{4-4}$), EEG showed a significant response aligned to the nominal transition, which was context dependent, enhanced by active detection, and absent in the no-change control with fixed click timing (Reg$_{4-4}$). We propose that increased temporal uncertainty reduces the precision of bottom-up evidence and increases reliance on top-down priors about *when* change is likely, enabling an internally timed update signal even when the stimulus remains constant.

1                                  **Results**

We established an experimental context in which participants could form a strong

temporal expectation about when a change might occur. Each trial consisted of a 2-s

click train composed of two consecutive 1-s segments (Train 1 and Train 2), with a

fixed nominal change point at 1 s (Fig. 1A).

**Figure 1. Experimental design and behavioral results. (A) Stimuli and trial**

**structure.** Each trial consisted of a 2-s click train comprising two consecutive 1-s

segments (Train 1 and Train 2), with a fixed nominal change point at 1 s. Eight

stimulus types were presented with equal probability (12.5% each): $Reg_{4-4}$ (no

change), $Reg_{4-4.01/4.02/4.03/4.06}$ (changes in Train 2 click timing), $Irreg_{4-4}$ (no change),

and $Irreg_{4-4.06}$ and $Irreg_{4-8}$ (changes in Train 2 mean timing). In the Irreg conditions,

click timing was drawn from a Gaussian distribution with fixed mean values. Click

duration was 200 μs and the inter-trial interval (ITI) ranged from 2 to 3 s. **(B)**

**Sessions.** Participants first completed a passive listening session without behavioral

report, followed by an active change-detection session in which they reported "change"

or "no change" via key press within 2 s after stimulus offset. **(C) Behavior.**

Change-detection ratio in the active session plotted as a function of Train 2 timing.

Purple bars denote Reg conditions and orange bars denote Irreg conditions. Data are

mean ± s.e.m. across participants (N = 42). **(D) Stimulus proportions and**

**behavioral reports.** Top: programmed proportions of no-change versus change trials.

Bottom: observed proportions of "no change" versus "change" reports pooled across

trials.

Eight stimulus types were randomly interleaved with equal probability (12.5% each):

regular no-change ($Reg_{4-4}$), regular change ($Reg_{4-4.01}$, $Reg_{4-4.02}$, $Reg_{4-4.03}$, $Reg_{4-4.06}$),

irregular no-change ($Irreg_{4-4}$), and irregular change ($Irreg_{4-4.06}$ and $Irreg_{4-8}$) (Fig. 1A).

In the irregular conditions, click timing was sampled from Gaussian distributions with

fixed mean values, and all trials used identical timing of the nominal change point

(Fig. 1A). Because six of the eight trial types contained a change and all

changes—when present—occurred at the fixed midpoint, participants were

predisposed to monitor the 1-s timepoint for a potential transition.

Participants (N = 42) completed two sessions in sequence: a passive listening

session without any behavioral report, followed by an active change-detection session

in which they indicated "change" or "no change" within 2 s after stimulus offset (Fig.

1B). In the active session, change reports depended strongly on Train 2 timing and on

stimulus statistics (Fig. 1C). For fixed-timing trains, the probability of reporting a

change increased monotonically with the magnitude of the Train 2 timing shift

($Reg_{4-4.01/4.02/4.03/4.06}$ relative to $Reg_{4-4}$), indicating high sensitivity to small timing

differences when timing was stable (Fig. 1C). By contrast, when click timing was

variable, change reports were elevated even in the no-change condition ($Irreg_{4-4}$) and

did not reliably separate $Irreg_{4-4.06}$ from $Irreg_{4-4}$ ($p = 0.68$, Wilcoxon signed rank test),

whereas a large timing shift ($Irreg_{4-8}$) was readily detected (Fig. 1C). Together, these

results confirm that temporal variability increases subjective "change-likeness" and

reduces sensitivity to small timing differences, while preserving detection for large

transitions.

The trial composition and behavioral judgments further quantify this

change-expectant context. By design, 75% of trials contained a physical change (six

change types out of eight), and 25% were no-change trials (Fig. 1D, upper).

Behaviorally, participants reported "change" on 64.4% of trials and "no change" on

35.6% of trials (Fig. 1D, lower), indicating a global bias toward change reports that is

consistent with the change-prevalent stimulus context and sets the stage for

expectation-driven midpoint-locked responses in EEG.

We next tested whether a change-like EEG response could emerge without any

physical transition, time-locked to the nominal change point at 1 s. In the active

session, we compared the two no-change stimuli, $Reg_{4-4}$ and $Irreg_{4-4}$ (Fig. 2A–C).

Figure 2. Predictive change-like responses in the no-change stimulus with

variable click timing. (A) Grand-average ERPs (N = 42) from five example channels

(Fz, T7, Cz, T8, and Pz) for the no-change stimuli Reg$_{4-4}$ (purple) and Irreg$_{4-4}$ (orange)

in the active session; shaded bands indicate $\pm$ s.e.m. The vertical dashed line marks

the nominal change point (1 s). The blue bar indicates the RM window (1000–1200

ms post-onset). (B) Scalp distributions of $\Delta$RM for Reg$_{4-4}$ and Irreg$_{4-4}$. The color

scale indicates $\Delta$RM magnitude (warm colors, positive $\Delta$RM; cool colors, negative

$\Delta$RM). Black dots denote electrodes with $\Delta$RM significantly greater than zero

(one-sample $t$-test against zero, FDR-corrected $p < 0.05$). (C) Participant-level mean

$\Delta$RM averaged across the predefined temporal-parietal-occipital ROI. $p$ values above

each condition indicate one-sample $t$-tests against zero; the horizontal comparison

indicates the paired *t*-test between conditions.

Across five example channels, Irreg$_{4-4}$ showed a pronounced change-locked

modulation—most evident at T7, T8, and Pz—whereas Reg$_{4-4}$ exhibited little or no

response at the same timepoint (Fig. 2A). We quantified this effect using ΔRM

computed within the predefined RM window (1000–1200 ms post-onset; blue bar, Fig.

2A). Here, RM denotes the RMS amplitude of the ERP within the RM window

(baseline-corrected using a pre-change interval), and ΔRM denotes the corresponding

baseline-normalized response magnitude used for between-condition comparisons

(see Methods). Scalp topographies revealed significant ΔRM for Irreg$_{4-4}$ (black dots),

primarily over temporal–parietal–occipital regions, whereas Reg$_{4-4}$ showed no

significant change-like effects (Fig. 2B). At the participant level, mean ΔRM

averaged across the predefined temporal–parietal–occipital ROI (23 channels) was

significantly larger for Irreg$_{4-4}$ than for Reg$_{4-4}$ (Fig. 2C), demonstrating a predictive

change-like response expressed in the absence of a physical change and selectively

enhanced when click timing is variable.

To confirm that the paradigm also yields canonical sensory-driven change

responses when a physical change is present, we compared transition and no-change

conditions within each stimulus class. In the active session, Reg$_{4-4.06}$ elicited a robust

change response relative to Reg$_{4-4}$, with clear between-condition differences in

example channels (Supplementary Fig. 1A), widespread significant ΔRM across

electrodes (Supplementary Fig. 1B), and a strong participant-level separation in

average ΔRM (Supplementary Fig. 1C). In contrast, the corresponding contrast under

variable timing (Irreg$_{4–4.06}$ vs Irreg$_{4–4}$) was substantially weaker, though still

detectable at the group level in the active session (Supplementary Fig. 1D–F). These

results establish a positive control: physical transitions produce strong change

responses when timing is stable, but the sensory-driven change response is reduced

when timing is variable.

We then tested whether the predictive change-like response depends on task

engagement by comparing Irreg$_{4–4}$ between passive and active sessions (Fig. 3A–C).

**Figure 3. Task-dependent enhancement of the predictive change-like response in**

**Irreg$_{4–4}$. (A)** Grand-average ERPs (N = 42) from five example channels (Fz, T7, Cz,

T8, and Pz) for Irreg$_{4-4}$ in the passive (light orange) and active (orange) sessions; shaded bands indicate ± s.e.m. The vertical dashed line marks the nominal change point (1 s). The blue bar indicates the RM window (1000–1200 ms post-onset). Vertical gray bar indicates significant difference between conditions ($p < 0.05$, permutation test). **(B)** Scalp distributions of ΔRM for Irreg$_{4-4}$ in the passive and active sessions. The color scale indicates ΔRM magnitude (warm colors, positive ΔRM; cool colors, negative ΔRM). Black dots denote electrodes with ΔRM significantly greater than zero (one-sample *t*-test against zero, FDR-corrected $p < 0.05$). **(C)** Participant-level mean ΔRM averaged across the predefined temporal-parietal-occipital ROI. *p* values above each condition indicate one-sample *t*-tests against zero; the horizontal comparison indicates the paired *t*-test between sessions.

In the same example channels, the change-locked modulation for Irreg$_{4-4}$ was markedly stronger during active detection than during passive listening (Fig. 3A), and permutation testing identified a significant time window in channel T8 (gray shading, Fig. 3A). This task dependence was also evident in ΔRM topographies (Fig. 3B) and in participant-level average ΔRM across channels, which was significantly higher in the active than in the passive session (Fig. 3C). Consistent with this, in the passive session the stable-timing transition remained robust ($p < 0.001$, paired t-test, Reg$_{4-4.06}$ > Reg$_{4-4}$; Supplementary Fig. 2A–C), whereas the variable-timing transition contrast was not significant (Irreg$_{4-4.06}$ vs Irreg$_{4-4}$; Supplementary Fig. 2D–F). Thus, active

engagement selectively enhanced the predictive change-like response in the

variable-timing no-change condition while revealing a broader dissociation between

stable-timing and variable-timing sensory-driven change responses.

Two additional controls strengthened the interpretation that the $Irreg_{4-4}$ effect

reflects a change-like response rather than a generic midpoint artifact. First, the

fixed-timing no-change control ($Reg_{4-4}$) showed no predictive change-like response

and no significant active–passive difference in $\Delta RM$ (Supplementary Fig. 3A–C),

arguing against a general session effect or nonspecific midpoint segmentation. Second,

a time-window comparison aligned to the nominal change point demonstrated

temporal specificity of the $Irreg_{4-4}$ response (Fig. 4).

**Figure 4. Temporal specificity of the predictive change-like response. (A)** ERPs for Irreg4–4 in the active session (N = 42) are shown for all scalp electrodes. For each electrode, three waveform segments aligned to the nominal change point (1 s) are overlaid: pre-change (gray, −200 to 0 ms), early post-change (orange, 0 to 200 ms), and late post-change (black, 300 to 500 ms). **(B)** Scalp distributions of RM for the same three time windows. The color scale indicates RM magnitude. **(C)** Participant-level mean RM averaged across the predefined temporal-parietal-occipital ROI for each time window. Horizontal comparisons indicate paired t-tests between time windows (pre vs early post: $p < 0.001$; early post vs late post: $p = 0.02$; pre vs

late post: $p = 0.06$).

When waveform segments were overlaid across electrodes, a clear deflection was

evident in the early post-change interval (0–200 ms), whereas the pre-change (−200–0

ms) and late post-change (300–500 ms) intervals showed no comparable response

(Fig. 4A). This pattern was mirrored in RM topographies (Fig. 4B) and in

ROI-averaged RM, which was significantly larger in the early post-change window

than in the pre-change and late post-change windows (Fig. 4C). Together, these

controls indicate that the predictive change-like response is specifically time-locked to

the expected change point, rather than reflecting a broad fluctuation across the

change-centered epoch.

Finally, we tested whether the predictive change-like response in $Irreg_{4-4}$ covaries

with participants' subjective reports on physically identical no-change trials. We

therefore separated $Irreg_{4-4}$ trials in the active session according to behavioral report

into trials judged as "change" and trials judged as "no change" (Fig. 5A).

Figure 5. Predictive change-like responses track subjective change reports. (A)

Grand-average ERPs (N = 42) from five example channels (Fz, T7, Cz, T8, and Pz)

for Irreg4–4 in the active session, separated within participant by behavioral report

into trials judged as "change" (red) and trials judged as "no change" (gray); shaded

bands indicate ± s.e.m. The vertical dashed line marks the nominal change point (1 s).

The blue bar indicates the RM window (1000–1200 ms post-onset). (B) Scalp

distributions of ΔRM for "change"-report and "no change"-report trials. The color

scale indicates ΔRM magnitude (warm colors, positive ΔRM; cool colors, negative

ΔRM). Black dots denote electrodes with ΔRM significantly greater than zero

(one-sample t-test against zero, FDR-corrected $p < 0.05$). (C) Participant-level mean

ΔRM averaged across the predefined temporal-parietal-occipital ROI for

"change"-report and "no change"-report trials. *p* values above each condition indicate

one-sample t-tests against zero; the horizontal comparison indicates the paired t-test

between trial types.

Across example channels, "change"-report trials showed a stronger change-locked

modulation around the nominal midpoint than "no change"-report trials (Fig. 5A).

This difference was also apparent in scalp topographies: ΔRM was broadly positive

for "change"-report trials and reached significance relative to zero at multiple

electrodes (black dots), whereas "no change"-report trials showed weaker ΔRM and

did not differ from zero at the ROI level (Fig. 5B–C). At the participant level, mean

ΔRM averaged across the predefined temporal-parietal-occipital ROI was

significantly larger for "change"-report than for "no change"-report trials ($p = 0.03$,

paired t-test; Fig. 5C). Consistent with this, ΔRM for "change"-report trials was

significantly greater than zero ($p < 0.001$, one-sample t-test), whereas ΔRM for "no

change"-report trials was not ($p = 0.48$, one-sample t-test; Fig. 5C). Together, these

results indicate that the predictive change-like response not only depends on stimulus

statistics and task engagement (Figs. 2–3), but also tracks subjective change

judgments even when the stimulus contains no physical transition.

**Discussion**

Our results identify a context-defined, predictive change-like response in human EEG that can emerge in the absence of any physical stimulus transition. Behaviorally, subtle timing changes were reliably detected when click timing was fixed, but sensitivity was markedly reduced when click timing was variable (Fig. 1C). Consistent with the expanded stimulus set and the change-prevalent trial structure, participants were globally biased toward reporting "change" (Fig. 1D). At the neural level, sensory-driven change responses to physical transitions were robust in the fixed-timing condition and attenuated under variable timing (Supplementary Figs. 1–2). Crucially, during active detection the variable-timing no-change stimulus (Irreg$_{4-4}$) elicited a midpoint-locked response that exceeded the fixed-timing no-change control (Reg$_{4-4}$) (Fig. 2A–C), was amplified relative to passive listening (Fig. 3A–C), and remained absent in the fixed-timing no-change control across sessions (Supplementary Fig. 3). Moreover, when aligned to the nominal change point, a clear peak was observed only in the early post-change window, whereas the pre-change and late post-change windows showed no comparable response (Fig. 4). This temporal specificity is consistent with the pre-specified RM window used for ΔRM quantification (1000–1200 ms post-onset). Finally, the predictive response also tracked subjective report: within Irreg$_{4-4}$ trials, ΔRM was larger on trials judged as "change" than on trials judged as "no change" (Fig. 5), linking the neural effect to perceptual decision state even in the absence of a physical transition. Together, these dissociations show that change-like activity can be expressed at an expected timepoint

without a stimulus change, but only under a specific combination of stimulus statistics,

task engagement, and a change-prevalent context.

Predictive coding frames perception as inference: the brain maintains generative

models of the sensory world, compares incoming evidence against those models, and

updates them when expectations fail [2,3]. Mismatch negativity (MMN) provides a

canonical population-level signature of violated expectation, reliably elicited when a

discrete deviant violates an inferred regularity [4-6]. Omission paradigms sharpen the

same logic by demonstrating that time-locked cortical responses can be evoked even

when an expected event is absent, indicating that prediction-related update signals can

be anchored to internal timing rather than stimulus energy [16,17]. However, most

omission effects are demonstrated in rhythmic, event-based sequences, where discrete

token timing and entrainment are integral to the paradigm. Our data extend this

framework to continuous stimulation with a context-defined update timepoint—a

fixed, task-relevant midpoint embedded within a change-prevalent design (Fig. 1A,

Fig. 1D). In this regime, the response we observe is not best interpreted as a reaction

to a missing sound token, but as an internally timed, prediction-error–like update

signal to a missing structural transition at the expected midpoint of a continuous

stream (Fig. 1A; Figs. 2–3). This shift matters because it implies that predictive

machinery can be recruited not only by violations of what occurs, but also by

violations of when an internally defined update is expected during ongoing input [2,3,15].

The result also complements "phantom deviance" evidence showing that endogenous

perceptual states can establish top-down predictions that yield deviance-like responses

when violated [20]. The report-dependent modulation observed in Fig. 5 is consistent

with this view, suggesting that the predictive response is coupled to decision-related

state under uncertainty rather than being a fixed sensory artifact.

A second key feature is gating by temporal uncertainty and task set. The

predictive change-like response was selectively expressed for $Irreg_{4-4}$ and absent for

$Reg_{4-4}$ (Fig. 2A–C; Supplementary Fig. 3), a pattern naturally captured by

precision-weighted accounts in which update signals are scaled by the estimated

reliability (precision) of sensory evidence relative to higher-level priors [2,3,22]. When

click timing is highly consistent ($Reg_{4-4}$), bottom-up evidence is precise and stabilizes

the internal model, suppressing spurious updating at the midpoint despite the broader

task context (Fig. 2A; Supplementary Fig. 3). When timing is variable ($Irreg_{4-4}$),

sensory precision is reduced, increasing the relative influence of context-driven

priors—here, an expectation that something may happen at 1 s given the experimental

setting (Fig. 1A–C). Under these conditions, the absence of a transition at the

expected moment can yield a temporally specific, prediction-error–like update signal

expressed as a midpoint-locked change-like response (Fig. 2A–C) [2,3,5]. The lack of

any corresponding effect in $Reg_{4-4}$, together with the window-specific response

profile quantified in Fig. 4, argues against generic midpoint segmentation, slow drift,

or broad state fluctuations as sufficient explanations (Supplementary Fig. 3; Fig. 4).

The strong active–passive dissociation further supports a task-set contribution: active

detection amplified the response (Fig. 3A–C), consistent with the idea that task

engagement increases the precision of the "midpoint-change" hypothesis and boosts

the gain of mismatch signals at behaviorally relevant timepoints [3,22,23]. In passive

listening, the midpoint carries less behavioral relevance, weakening the

context-defined temporal prior and thereby greatly reducing the predictive response

(Fig. 3A–C). Notably, the report-dependent modulation in Fig. 5 provides convergent

support that the response is coupled to behavioral set rather than reflecting a purely

time-locked sensory transient.

Finally, these results provide a compact empirical separation between (i)

sensory-driven change responses to physical transitions (Supplementary Figs. 1–2)

and (ii) a context- and task-gated predictive change-like response expressed without

physical change when temporal uncertainty is high (Figs. 2–3; Supplementary Fig. 3).

Notably, the effect was temporally confined to the early post-change window and was

modulated by subjective change reports. More broadly, the present paradigm offers a

practical assay for dissociating sensory-driven change detection from context-gated

predictive updating in continuous auditory streams, and it provides a principled

handle for testing how uncertainty and behavioral goals shape prediction-related

neural dynamics in humans [5,6,16,17,23].

**Materials and methods**

**Experimental procedure and participants**

This study was conducted in accordance with the Declaration of Helsinki (2013)

[24]. A total of 42 participants (20 males and 22 females, mean age: 23.36 years old,

standard deviation: 2.55) participated in this experiment. Participants maintained a

stationary head position while listening to auditory stimuli and responding via

keyboard presses. Experiments including healthy participants were approved by the

Institutional Review Board (IRB-20230131-R), and informed consent was obtained

from all participants.

The experiment included two sessions: a passive session followed by an active

session. Eight stimulus types were presented with equal probability in each session,

including regular no-change ($Reg_{4-4}$), regular change ($Reg_{4-4.01}$, $Reg_{4-4.02}$, $Reg_{4-4.03}$,

and $Reg_{4-4.06}$), irregular no-change ($Irreg_{4-4}$), and irregular change ($Irreg_{4-4.06}$ and

$Irreg_{4-8}$). Each stimulus was repeated a minimum of 40 times, in both sessions.

**Auditory Stimuli**

This study employed transitional click trains (Fig. 1A) [21,25,26]. Each click

consisted of a 200-μs monopolar pulse. Each trial comprised a 2-s click train made of

two consecutive 1-s segments, with a fixed nominal change point at 1 s. Eight

stimulus types were used: regular no-change ($Reg_{4-4}$), regular change ($Reg_{4-4.01}$,

$Reg_{4-4.02}$, $Reg_{4-4.03}$, and $Reg_{4-4.06}$), irregular no-change ($Irreg_{4-4}$), and irregular change

($Irreg_{4-4.06}$ and $Irreg_{4-8}$). These stimulus types were randomly interleaved with equal

probability. In regular click trains, inter-click intervals (ICIs) were fixed, whereas in

irregular click trains, ICIs were drawn from a Gaussian distribution with fixed mean

values and satisfied the following constraints:

$$0.3 \times \mu_i \leq I_{i,j} \leq 1.7 \times \mu_i$$

$$1.2 \times \mu_i \leq \sum_{n=j}^{j+1} I_{i,n} \leq 3.1 \times \mu_i$$

$$1.8 \times \mu_i \leq \sum_{n=j}^{j+2} I_{i,n} \leq 4.6 \times \mu_i$$

where $\mu_i$ represents the mean ICI (in milliseconds) of the $i^{\text{th}}$ segment in the

transitional click train and $I_{i,j}$ denotes the $j^{\text{th}}$ ICI in that segment. Transitional click

trains were formed by concatenating two click trains with identical or different ICIs or

mean ICIs. For example, Reg$_{4–4.06}$ consisted of a 1-s regular train with a 4-ms ICI

followed seamlessly by a 1-s regular train with a 4.06-ms ICI. Stimuli were delivered

through a Golden Field M23 speaker, powered by a Creative AE-7 Sound Blaster at a

sampling rate of 384 kHz, and controlled via Psychtoolbox 3 in MATLAB. Sound

intensity was calibrated to a constant 60 dB SPL using a ¼-inch Brüel & Kjær 4954

condenser microphone and PHOTON/RT analyzer.

**Data acquisition**

Electroencephalogram (EEG) data were acquired using a 64-channel NeuroScan

system (Compumedics, Australia). In practice, we only included 60 electrodes in the

analyses. The ground electrode of the 64-channel NeuroScan Quick-Cap was placed

between Fpz and Fz in the frontal area, while the reference electrode was placed

between Cz and CPz. The EEG data were sampled at 1 kHz, and electrode placement

followed the international 10-20 system protocol.

**Data Analysis**

**Preprocessing**. Data analyses were performed using MATLAB R2021b (The

MathWorks, Inc., Natick, MA, USA) and the FieldTrip toolbox [27]. Monopolar

referencing was applied using the default reference electrode positioned between Cz

and Pz. Multichannel EEG data underwent sequential preprocessing: initial bandpass

filtering (two-pass Butterworth, 0.5–40 Hz), followed by epoch extraction from -1 to

3 s relative to trial onset. Independent component analysis (ICA) removed

electrooculogram (EOG) artifacts. Baseline correction subtracted the mean activity

from -200 to 0 ms pre-stimulus onset per trial. Motion artifacts were evaluated via a

relative threshold, flagging samples outside mean $\pm 3 \times$ SD as "bad." Trials exceeding

20% bad samples were discarded as "bad trials," and channels with >10% bad trials

were labeled "bad channels." Bad channels were excluded first (based on all-channel

data), followed by bad trials from remaining channels. Event-related potentials (ERPs)

were computed by averaging epochs per experimental condition, channel, and subject.

Prior to group-level analysis, ERPs were normalized by dividing each channel's data

by the standard deviation of baseline (-200 to 0 ms pre-stimulus onset) per subject to

minimize inter-channel variability.

**Permutation test**. Permutation tests for time-point and channel-wise

comparisons of ERP data between stimuli groups employed a two-tailed cluster-based

approach via the 'ft_timelockstatistics' function in the FieldTrip toolbox for MATLAB.

A two-tailed independent-samples t-test was conducted at each time point and channel

to compute t-values, contrasting Dataset A (e.g., ERP data for $Irreg_{4\text{-}4}$ in the passive

session) and Dataset B (e.g., ERP data for Irreg$_{4\text{-}4}$ in the active session) across 42

participants. A cluster-defining threshold of $p < 0.05$ identified significant time points,

which were grouped into clusters based on temporal adjacency (consecutive

significant time points) and spatial proximity (neighboring electrodes). Condition

labels for Datasets A and B were randomly permuted, generating a new t-value matrix

per iteration; clusters were reformed using the same $p < 0.05$ threshold. The largest

cluster-level statistic (e.g., sum of t-values) from each of 1000 permutations formed

the null distribution. Observed clusters were deemed significant if their statistic

exceeded the 95[th] percentile of this distribution ($p < 0.05$). Tests spanned the time

window of -500 to 2500 ms relative to the transitional click train onset.

**Quantification of EEG response**. To quantify the change response, we defined

the response magnitude (RM) as the root mean square (RMS) of the ERP within the

predefined change-response window. For each channel, RM was calculated in the

specified window (e.g., [1000, 1200] ms relative to stimulus onset, corresponding to

[0, 200] ms relative to the nominal change point at 1 s). The relative response

magnitude (ΔRM) was then obtained by subtracting the RMS of the baseline window

[800, 1000] ms post-onset, corresponding to [-200, 0] ms pre-change. Specifically, for

a channel $i$,

$$\text{RM}_i[t_1, t_2] = \sqrt{\frac{\sum_{t=t_1}^{t_2}[r_i(t)]^2}{n}}$$

where $r_i(t)$ denotes the ERP of the $i^{th}$ channel at time $t$ (ms), $n$ is the number of

samples within the time window $[t_1, t_2]$, and $\text{RM}_i$ is the response magnitude of

1  channel $i$. The relative response magnitude was then defined as

$$\Delta RM_i = RM_i[1000,1200] - RM_i[800,1000].$$

The baseline window immediately preceding the nominal change point represented

the steady-state response to Train 1. To obtain an ROI-based measure of the change

response, $\Delta RM$ was averaged across a predefined set of temporal-parietal-occipital

channels (T7, T8, TP7, TP8, P7, P5, P3, P1, Pz, P2, P4, P6, P8, PO7, PO5, PO3, POz,

PO4, PO6, PO8, O1, Oz, and O2). This ROI was chosen because the canonical change

response in the transitional click-train paradigm is strongest over

temporal-parietal-occipital scalp regions, with pronounced responses reported at

temporal, parietal, and occipital electrodes in an independent EEG study [21], and the

predictive change-like response observed here showed a qualitatively similar posterior

distribution (Fig. 2B). Averaging within this ROI therefore provides a more sensitive

summary measure while avoiding dilution of the effect by electrodes outside the main

response field.

**Statistics**

Behavioral comparisons of change-detection ratio between paired conditions (e.g.,

$Irreg_{4-4}$ and $Irreg_{4-4.06}$) were performed using two-tailed Wilcoxon signed-rank tests.

For EEG analyses, two-tailed one-sample $t$-tests were used to test whether the average

$\Delta RM$ across the predefined temporal-parietal-occipital ROI differed from zero (e.g.,

Fig. 2C, $p$ values above each condition). Unless otherwise noted, pairwise

comparisons of $\Delta RM$ between conditions were performed using two-tailed paired

$t$-tests (e.g., $Irreg_{4-4}$ in the passive versus active session).

To identify channels showing significant change responses, ΔRM was computed

for each channel and subject, as shown in the topographic plots (e.g., Fig. 2B). For

each channel, a two-tailed one-sample $t$-test against zero was performed across

subjects, and the resulting $p$ values were corrected across channels using the

Benjamini-Yekutieli false discovery rate (FDR) procedure [28]. Channels surviving

correction were marked with black dots on the topographic plots.

**Data Availability**

All data used in this study are available on request.

All codes applied in this study are available on a public Github repository (url:

https://github.com:TOMORI233/EEGProcess.git). All source data supporting the

findings of this study are publicly available at Zenodo under the following DOI:

https://doi.org/10.5281/zenodo.15795285 [29].

**Figure legends**

**Figure 1. Experimental design and behavioral results. (A) Stimuli and trial**

**structure.** Each trial consisted of a 2-s click train comprising two consecutive 1-s

segments (Train 1 and Train 2), with a fixed nominal change point at 1 s. Eight

stimulus types were presented with equal probability (12.5% each): $Reg_{4-4}$ (no

change), $Reg_{4-4.01/4.02/4.03/4.06}$ (changes in Train 2 click timing), $Irreg_{4-4}$ (no change),

and $Irreg_{4-4.06}$ and $Irreg_{4-8}$ (changes in Train 2 mean timing). In the Irreg conditions,

click timing was drawn from a Gaussian distribution with fixed mean values. Click

duration was 200 μs and the inter-trial interval (ITI) ranged from 2 to 3 s. **(B)**

**Sessions.** Participants first completed a passive listening session without behavioral

report, followed by an active change-detection session in which they reported "change"

or "no change" via key press within 2 s after stimulus offset. **(C) Behavior.**

Change-detection ratio in the active session plotted as a function of Train 2 timing.

Purple bars denote Reg conditions and orange bars denote Irreg conditions. Data are

mean ± s.e.m. across participants (N = 42). **(D) Stimulus proportions and**

**behavioral reports.** Top: programmed proportions of no-change versus change trials.

Bottom: observed proportions of "no change" versus "change" reports pooled across

trials.

**Figure 2. Predictive change-like responses in the no-change stimulus with**

**variable click timing. (A)** Grand-average ERPs (N = 42) from five example channels

(Fz, T7, Cz, T8, and Pz) for the no-change stimuli $Reg_{4-4}$ (purple) and $Irreg_{4-4}$ (orange)

in the active session; shaded bands indicate ± s.e.m. The vertical dashed line marks

the nominal change point (1 s). The blue bar indicates the RM window (1000–1200

ms post-onset). **(B)** Scalp distributions of ΔRM for $Reg_{4-4}$ and $Irreg_{4-4}$. The color

scale indicates ΔRM magnitude (warm colors, positive ΔRM; cool colors, negative

ΔRM). Black dots denote electrodes with ΔRM significantly greater than zero

(one-sample $t$-test against zero, FDR-corrected $p < 0.05$). **(C)** Participant-level mean

ΔRM averaged across the predefined temporal-parietal-occipital ROI. $p$ values above

each condition indicate one-sample $t$-tests against zero; the horizontal comparison

indicates the paired $t$-test between conditions.

**Figure 3. Task-dependent enhancement of the predictive change-like response in**

**$Irreg_{4-4}$. (A)** Grand-average ERPs (N = 42) from five example channels (Fz, T7, Cz,

T8, and Pz) for $Irreg_{4-4}$ in the passive (light orange) and active (orange) sessions;

shaded bands indicate $\pm$ s.e.m. The vertical dashed line marks the nominal change

point (1 s). The blue bar indicates the RM window (1000–1200 ms post-onset).

Vertical gray bar indicates significant difference between conditions ($p < 0.05$,

permutation test). **(B)** Scalp distributions of $\Delta$RM for $Irreg_{4-4}$ in the passive and active

sessions. The color scale indicates $\Delta$RM magnitude (warm colors, positive $\Delta$RM; cool

colors, negative $\Delta$RM). Black dots denote electrodes with $\Delta$RM significantly greater

than zero (one-sample *t*-test against zero, FDR-corrected $p < 0.05$). **(C)**

Participant-level mean $\Delta$RM averaged across the predefined

temporal-parietal-occipital ROI. *p* values above each condition indicate one-sample

*t*-tests against zero; the horizontal comparison indicates the paired *t*-test between

sessions.

**Figure 4. Temporal specificity of the predictive change-like response.** (**A**) ERPs

for Irreg4–4 in the active session (N = 42) are shown for all scalp electrodes. For each

electrode, three waveform segments aligned to the nominal change point (1 s) are

overlaid: pre-change (gray, −200 to 0 ms), early post-change (orange, 0 to 200 ms),

and late post-change (black, 300 to 500 ms). (**B**) Scalp distributions of RM for the

same three time windows. The color scale indicates RM magnitude. (**C**)

Participant-level mean RM averaged across the predefined temporal-parietal-occipital

ROI for each time window. Horizontal comparisons indicate paired t-tests between

time windows (pre vs early post: $p < 0.001$; early post vs late post: $p = 0.02$; pre vs

late post: $p = 0.06$).

**Figure 5. Predictive change-like responses track subjective change reports. (A)**

Grand-average ERPs (N = 42) from five example channels (Fz, T7, Cz, T8, and Pz)

for Irreg4–4 in the active session, separated within participant by behavioral report

into trials judged as "change" (red) and trials judged as "no change" (gray); shaded

bands indicate ± s.e.m. The vertical dashed line marks the nominal change point (1 s).

The blue bar indicates the RM window (1000–1200 ms post-onset). **(B)** Scalp

distributions of ΔRM for "change"-report and "no change"-report trials. The color

scale indicates ΔRM magnitude (warm colors, positive ΔRM; cool colors, negative

ΔRM). Black dots denote electrodes with ΔRM significantly greater than zero

1    (one-sample t-test against zero, FDR-corrected $p < 0.05$). **(C)** Participant-level mean

2    ΔRM averaged across the predefined temporal-parietal-occipital ROI for

3    "change"-report and "no change"-report trials. *p* values above each condition indicate

4    one-sample t-tests against zero; the horizontal comparison indicates the paired t-test

5    between trial types.

## References

Rao, R. P. & Ballard, D. H. Predictive coding in the visual cortex: a functional interpretation of some extra-classical receptive-field effects. *Nat Neurosci* **2**, 79-87, doi:10.1038/4580 (1999).

Friston, K. A theory of cortical responses. *Philos Trans R Soc Lond B Biol Sci* **360**, 815-836, doi:10.1098/rstb.2005.1622 (2005).

Bastos, A. M. *et al.* Canonical microcircuits for predictive coding. *Neuron* **76**, 695-711, doi:10.1016/j.neuron.2012.10.038 (2012).

Näätänen, R., Gaillard, A. W. & Mäntysalo, S. Early selective-attention effect on evoked potential reinterpreted. *Acta Psychol (Amst)* **42**, 313-329, doi:10.1016/0001-6918(78)90006-9 (1978).

Garrido, M. I., Kilner, J. M., Stephan, K. E. & Friston, K. J. The mismatch negativity: a review of underlying mechanisms. *Clin Neurophysiol* **120**, 453-463, doi:10.1016/j.clinph.2008.11.029 (2009).

Fong, C. Y., Law, W. H. C., Uka, T. & Koike, S. Auditory Mismatch Negativity Under Predictive Coding Framework and Its Role in Psychotic Disorders. *Front Psychiatry* **11**, 557932, doi:10.3389/fpsyt.2020.557932 (2020).

Song, P., Zhai, Y. & Yu, X. Stimulus-specific adaptation (SSA) in the auditory system: Functional relevance and underlying mechanisms. *Neurosci Biobehav Rev* **149**, 105190, doi:10.1016/j.neubiorev.2023.105190 (2023).

Gong, Y. *et al.* Cross-Modal Interaction and Integration Through Stimulus-Specific Adaptation in the Thalamic Reticular Nucleus of Rats. *Neurosci Bull* **38**, 785-795,

1        doi:10.1007/s12264-022-00827-8 (2022).

Song, P. R. *et al.* Adaptation in the Dorsal Belt and Core Regions of the Auditory Cortex in the Awake Rat. *Neuroscience* **455**, 79-88, doi:10.1016/j.neuroscience.2020.11.042 (2021).

Zhai, Y. Y. *et al.* Synaptic Adaptation Contributes to Stimulus-Specific Adaptation in the Thalamic Reticular Nucleus. *Neurosci Bull* **36**, 1538-1541, doi:10.1007/s12264-020-00536-0 (2020).

Xu, X. X., Zhai, Y. Y., Kou, X. K. & Yu, X. Adaptation facilitates spatial discrimination for deviant locations in the thalamic reticular nucleus of the rat. *Neuroscience* **365**, 1-11, doi:10.1016/j.neuroscience.2017.09.022 (2017).

Xu, X., Yu, X., He, J. & Nelken, I. Across-ear stimulus-specific adaptation in the auditory cortex. *Front Neural Circuits* **8**, 89, doi:10.3389/fncir.2014.00089 (2014).

Rui, Y. Y., He, J., Zhai, Y. Y., Sun, Z. H. & Yu, X. Frequency-Dependent Stimulus-Specific Adaptation and Regularity Sensitivity in the Rat Auditory Thalamus. *Neuroscience* **392**, 13-24, doi:10.1016/j.neuroscience.2018.09.015 (2018).

Zhai, Y. Y., Sun, Z. H., Gong, Y. M., Tang, Y. & Yu, X. Integrative stimulus-specific adaptation of the natural sounds in the auditory cortex of the awake rat. *Brain Struct Funct* **224**, 1753-1766, doi:10.1007/s00429-019-01880-2 (2019).

Wacongne, C. *et al.* Evidence for a hierarchy of predictions and prediction errors in human cortex. *Proc Natl Acad Sci U S A* **108**, 20754-20759, doi:10.1073/pnas.1117807108 (2011).

Braga, A. & Schönwiesner, M. Neural Substrates and Models of Omission Responses

and Predictive Processes. *Front Neural Circuits* **16**, 799581, doi:10.3389/fncir.2022.799581 (2022).

Heilbron, M. & Chait, M. Great Expectations: Is there Evidence for Predictive Coding in Auditory Cortex? *Neuroscience* **389**, 54-73, doi:10.1016/j.neuroscience.2017.07.061 (2018).

Strauss, M. *et al.* Disruption of hierarchical predictive coding during sleep. *Proc Natl Acad Sci U S A* **112**, E1353-1362, doi:10.1073/pnas.1501026112 (2015).

Ye, H. *et al.* Hierarchical Temporal Processing in the Primate Thalamocortical System: Insights from Nonlinguistic Structured Stimuli. *Research (Wash D C)* **8**, 0960, doi:10.34133/research.0960 (2025).

Qi, Y. Y. *et al.* Evidence for predictions established by phantom sound. *Neuroimage* **264**, 119766, doi:10.1016/j.neuroimage.2022.119766 (2022).

Xu, H. *et al.* EEG neural indicator of temporal integration in the human auditory brain with clinical implications. *Commun Biol* **8**, 1109, doi:10.1038/s42003-025-08540-8 (2025).

Feldman, H. & Friston, K. J. Attention, uncertainty, and free-energy. *Front Hum Neurosci* **4**, 215, doi:10.3389/fnhum.2010.00215 (2010).

Wacongne, C., Changeux, J. P. & Dehaene, S. A neuronal model of predictive coding accounting for the mismatch negativity. *J Neurosci* **32**, 3665-3678, doi:10.1523/jneurosci.5003-11.2012 (2012).

Duncan, N. Declaration of Helsinki. *World Medical Journal* **59** (2013).

Song, P. *et al.* Temporal merging into pitch with click train in the macaque auditory

cortex. *Natl Sci Rev* **12**, nwaf026, doi:10.1093/nsr/nwaf026 (2025).

26    Song, P. *et al.* A new function of offset response in the primate auditory cortex: marker of temporal integration. *Commun Biol* **7**, 1350, doi:10.1038/s42003-024-07058-9 (2024).

27    Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational intelligence and neuroscience* **2011**, 156869 (2011).

28    Benjamini, Y. & Yekutieli, D. The control of the false discovery rate in multiple testing under dependency. *Annals of statistics*, 1165-1188 (2001).

29    Xu, H.    (Zenodo, 2025).