1

Adolescent social isolation creates a latent vulnerability in maternal care with intergenerational social consequences, rescued by experienced mothers

Jose Francis-Oliveira[a], Rinako Tanaka[a], Matthew Shen[a], Emily Cruvinel[a,b], Shin-ichi Kano[a,c], Minae Niwa[a,c,d,1.]

[a]Department of Psychiatry and Behavioral Neurobiology, University of Alabama at Birmingham School of Medicine, Birmingham, AL, USA

[b]Department of Physiology and Biophysics, Sao Paulo University, Sao Paulo, Brazil

[c]Department of Neurobiology, University of Alabama at Birmingham School of Medicine, Birmingham, AL, USA

[d]Department of Biomedical Engineering, University of Alabama at Birmingham School of Engineering, Birmingham, AL, USA

[1]To whom correspondence may be addressed.

Minae Niwa, Ph.D. Email: mniwa@uabmc.edu.

Author Contributions: M.N. conceived, designed, and supervised the project with input from

S.K. J.F-O., R.T., M.S., E.C., and M.N. performed the experiments and analyzed the data. J.F-

O., and M.N. wrote the initial draft of the manuscript and revised it with input from all co-authors.

Competing Interest Statement: No competing interests to disclose.

Classification: Biological Sciences; Psychological and Cognitive Sciences.

Keywords: Adolescent social isolation; Vulnerability; Maternal behavior; Social behavior;

Cortical circuits.

Abstract

Early-life adversity (ELA) can produce lasting alterations on brain and behavior, yet its effects

on maternal caregiving and intergenerational social outcomes remain poorly understood. Using

a mouse model of mild adolescent social isolation, we demonstrated that females exposed to

isolation in late adolescence exhibited progressive deficits in pup-directed maternal behaviors,

including nursing, licking, nest-building, and pup retrieval, while self-directed behaviors remain

intact. Offspring reared by these dams displayed deficits in adult social behavior, including

reduced sociability, impaired social novelty recognition, and diminished social odor

discrimination, without affecting non-social memory, anxiety-like behavior, or locomotor activity.

Chemogenetic activation of the medial cingulate cortex (mCg) to prelimbic cortex (PrL) pathway

in offspring of stressed dams rescued social deficits, whereas inhibition of this pathway in

control mice recapitulated the impairments. Notably, co-housing stressed dams with

experienced parous females during the early postpartum period restored pup-directed maternal

behaviors and normalized offspring social outcomes, demonstrating that social experience can

buffer the intergenerational consequences of ELA. Circuit analyses identified the mCg→PrL

projections as a predominantly excitatory pathway critical for social behavior.

Electrophysiological recordings revealed hypofunction of this circuit in offspring of stressed

dams, which was ameliorated by postpartum co-housing with parous females. Together, these

findings define a neural circuit linking adolescent psychosocial adversity to impaired maternal

caregiving and intergenerational social dysfunction, and highlight social support from

experienced mothers as a potent modulator of maternal and offspring behavioral outcomes.

Significance Statement

Adverse social experiences before motherhood can compromise maternal caregiving and

transmit social vulnerability across generations, yet the underlying mechanisms remain poorly

understood. Using a mouse model of subtle adolescent social isolation, we demonstrate that

even mild isolation selectively impairs postpartum caregiving behaviors while sparing other

behavioral domains, resulting in social deficits in adult offspring. We identify the midcingulate-

prelimbic cortex circuit as a critical neural substrate mediating these effects and show that social

support from experienced mothers during the early postpartum period restores both maternal

behavior and offspring social outcomes. Together, these findings define a neural and social

mechanism for the intergenerational effects of adolescent psychosocial adversity.

Introduction

Maternal behavior is a fundamental social behavior that supports offspring survival and long-term health across species (1, 2). In mammals, early mother-infant attachment shapes both maternal mental health and offspring socioemotional development (3). Disruption of maternal care due to stress, psychiatric conditions, or limited social support can compromise offspring development (4-7). Early life adversity (ELA), including maltreatment, family dysfunction, trauma, and social deprivation, increases long-term risk for psychiatric and physical disorders and is associated with postpartum mood disturbances and impaired maternal care (8-13). However, the neural mechanisms by which ELA affects postpartum maternal care remain poorly understood, in part because most animal models rely on hormonal manipulations during pregnancy (14-18), which do not capture the lasting neurodevelopmental consequences of ELA on neural circuit function.

Adolescence is a sensitive period for brain and social development, marked by hormonal changes and refinement of circuits that regulate emotion and social behavior (19). Social isolation in rodents is a widely used model for psychosocial ELA (20-22), but most paradigms involve prolonged or severe isolation that produce immediate behavioral and endocrine deficits (23, 24). In contrast, we developed a model of mild adolescent social isolation that produces no detectable changes but generates a latent vulnerability that emerges during the postpartum period as impairments in maternal mood- and social-related behaviors (25, 26). This model enables investigation of how subclinical adolescent adversity programs maternal behavior and affects the next generation. Consistent with this framework, mild adolescent isolation is associated with prolonged dysregulation of the hypothalamic-pituitary-adrenal (HPA) axis and sustained corticosterone elevation specifically during the postpartum period, paralleling observations in women with postpartum depression following early adversity (26). Together,

these findings define a developmental trajectory in which adolescent psychosocial adversity

produces latent, context-dependent postpartum dysfunction, with stress-axis alterations serving

as a downstream correlate rather that a sufficient explanatory mechanism.

Pregnancy induces lasting adaptations in the maternal brain, including structural remodeling of

the anterior cingulate cortex (ACC) (27). In humans, the dorsal ACC (dACC), homologous to the

rodent prelimbic cortex (PrL), is activated when healthy mothers respond to infant cues but

shows reduced activation in mothers with caregiving difficulties (28, 29). The medial prefrontal

cortex (mPFC), including the PrL, plays a central role in social cognition, emotional responses,

and the expression of maternal behaviors (30-33). Within the rodent cingulate cortex, the

anterior cingulate cortex [aCg; antero-posterior (AP) +0.0 mm] and mid-cingulate cortex (mCg;

AP +1.0 mm) are structurally and functionally distinct, with the mCg implicated in attention,

conflict monitoring, and social cognition (34-38). Although the role of the rodent mCg in maternal

and social behaviors remains largely unexplored, its involvement in behavioral domains that are

consistently disrupted following ELA suggests that it may represent a neural locus vulnerable to

adversity-related dysfunction. Based on these observations, we hypothesize that disruption of

cortico-cortical communication, particularly the mCg→PrL pathway, contributes to impaired

maternal care and the intergenerational transmission of social deficits.

Maternal circuits remain plastic in adulthood, allowing social experience to shape caregiving

behaviors. Mothers can learn maternal skills by observing experienced caregivers (39-44),

highlighting the role of social learning in parenting. Social support may be particularly important

for mothers with a history of ELA, who are at increased risk for postpartum mood disturbances

that can compromise caregiving (45, 46). However, the mechanisms through which social

support from experienced caregivers mitigates these deficits remain largely unknown.

In the present study, we show that mild adolescent social isolation creates a latent vulnerability

in female mice that manifests during the postpartum period as impaired maternal care, mediated

by reduced function of the mCg→PrL circuit. Dysfunction of this pathway drives social deficits in

offspring, establishing a neural substrate for intergenerational transmission of ELA effects.

Importantly, co-housing with parous females, defined here as females that have experienced at

least one pregnancy and delivery, during the postpartum period ameliorates maternal behavioral

deficits and rescues offspring social outcomes, revealing a mechanism through which social

support can buffer the long-term consequences of early adversity.

Results

Adolescent social isolation impairs maternal caring behavior in the postpartum period

We first examined whether females exposed to social isolation during late adolescence

exhibited alterations in maternal behavior after delivery. At postnatal day (P) 7, stressed dams

spent significantly less time nursing (**Figure 1A**) and licking their pups (**Figure 1B**) compared to

unstressed controls, resulting in a marked reduction in overall maternal caring behavior, defined

as the combined duration of nursing and licking (**Figure 1C**). In contrast, the time spent climbing

and digging (**Figure 1D**) as well as self-grooming (**Figure 1E**) did not differ between groups.

Accordingly, total self-care behavior, defined as the combined duration of climbing/digging and

self-grooming, was unchanged by adolescent isolation (**Figure 1F**). These findings indicate that

adolescent social isolation selectively impairs maternal care toward offspring without broadly

affecting self-directed maintenance behaviors. Consistent with this interpretation, nests built by

stressed dams were smaller and less organized than those of unstressed controls (**Figure 1G**).

Moreover, stressed dams required significantly more time to retrieve displaced pups to the nest

(**Figure 1H**), further supporting impaired maternal responsiveness at one week postpartum.

To determine whether these maternal deficits were evident immediately after birth, we assessed

maternal behavior at P1. At this early stage, stressed dams showed only mild or partial

impairments, limited to licking and pup retrieval behaviors (**Supplementary Figures 1A-H**). This

pattern suggests that deficits in maternal care develop progressively over the first postpartum

week, possibly reflecting delayed adaptation to the maternal care state.

Because maternal caregiving disruptions can influence offspring health, we evaluated the

general well-being of the litters. Despite reduced maternal care, offspring reared by stressed

dams (stressed offspring) exhibited body weight gain comparable to that of offspring reared by

unstressed dams (unstressed offspring) (**Supplementary Figure 2A**). However, mortality rates

were higher in litters reared by stressed dams, particularly during the early postnatal days

(**Supplementary Figure 2B**). The total number of surviving pups per litter tended to be lower in

stressed dams than in unstressed controls, and by P8, more stressed offspring had died

compared with unstressed offspring; however, these differences were not statistically significant

(**Supplementary Figure 2C**). These results indicate that adolescent social isolation leads to

gradually emerging deficits in maternal caregiving during the first postpartum week, which

contribute to increased early pup mortality despite apparently normal growth among surviving

offspring.

Adolescent isolation-induced maternal deficits lead to impaired social behavior in adult

offspring

We next examined whether postpartum maternal caregiving deficits induced by adolescent

social isolation affected behavioral outcomes in offspring during adulthood (>P56). In the social

interaction test (SIT), stressed offspring exhibited reduced sociability and impaired social

novelty recognition compared with unstressed controls (**Figures 2A and 2B**). In the social

olfactory recognition test (SORT), stressed offspring showed markedly reduced investigation

toward both familiar social cues from the test mouse's home cage and novel social cues from

the stranger's cage (**Figure 2C**), indicating broad impairments in social cue recognition. In

contrast, stressed offspring showed no differences from unstressed offspring in the novel object

recognition test (NORT), open field test (OFT), elevated plus maze test (EPM), or light-dark box

test (LDB) (**Figures 2D-G**). Locomotor activity in the OFT and discrimination index during the

NORT training session were also comparable between groups (**Supplementary Figures 3A**

and 3B). Consistent with heightened stress sensitivity, serum corticosterone levels were

significantly elevated in adult stressed offspring relative to unstressed controls (**Supplementary**

Figure 4).

These findings suggest that adolescent isolation-induced deficits in maternal caregiving

selectively impair social behavior in adult offspring without affecting general locomotion, anxiety-

related behavior, or non-social memory. The persistent elevation in corticosterone levels

indicates long-lasting dysregulation of HPA axis function that may contribute to the observed

social deficits. Collectively, these results support the idea that disrupted maternal care during

the early postnatal period can produce long-lasting, domain-specific alterations in offspring

social behavior and stress physiology.

The mCg→PrL pathway regulates social behavior and mediates the effects of poor

maternal care

Given the established role of the PrL in regulating social behavior (28) and our hypothesis that

cortico-cortical connectivity serves as a key circuit substrate for social behavior, we next

investigated whether the mCg→PrL pathway plays a critical role in social behaviors in offspring.

To manipulate this pathway, we employed Designer Receptors Exclusively Activated by

Designer Drugs (DREADDs), expressing engineered G-protein coupled receptors [M3

muscarinic receptor (hM3Dq; excitatory) and hM4Di (inhibitory)] selectively activated by the

synthetic ligand clozapine-N-oxide (CNO) (47).

Excitation of the mCg→PrL pathway in stressed offspring restored sociability and social novelty

recognition in the SIT (**Figures 3A-C**). Conversely, inhibition of the pathway in unstressed

offspring induced behavioral deficits similar to those observed in stressed offspring (**Figures**

3D-F). These results indicate that hypofunction of the mCg→PrL pathway contributes to social

behavioral deficits in stressed offspring and that the pathway is functionally required for normal

social behavior.

Co-housing with parous females rescues pup-directed maternal behaviors and offspring

social deficits

Parous mice exhibited significantly higher levels of maternal care than virgin mice, even when

caring for foster pups (**Supplemental Figures 5A-G**). Specifically, parous females spent more

time nursing (**Supplemental Figure 5A**) and licking pups (**Supplemental Figure 5B**), resulting

in greater overall caring behavior, defined as the combined duration of nursing and licking

(**Supplemental Figure 5C**). In contrast, self-directed behaviors, including climbing and digging

(**Supplemental Figure 5D**) and self-grooming (**Supplemental Figure 5E**), did not differ

between groups. Accordingly, total self-care behavior, defined as the combined duration of

climbing/digging and self-grooming, was comparable between parous and virgin females

(**Supplemental Figure 5F**). Pup retrieval behavior was significantly increased in parous

females relative to virgins (**Supplemental Figure 5G**). Together, these findings indicate that

pregnancy and delivery selectively enhance offspring-directed maternal behaviors without

broadly affecting self-directed maintenance activities.

Based on these observations, we next investigated whether social exposure to experienced

mothers could rescue maternal deficits. Stressed dams were co-housed with parous females

from P0 to P7, the day following completion of maternal behavioral testing. During co-housing,

pups remained with the stressed dams and were accessible to both stressed and parous

females, allowing observation and potential social learning. Immediately prior to maternal

behavioral testing, the parous females were temporarily removed so that measured behaviors

reflected the stressed dams' responses to their own pups rather that direct interactions with the

parous females. This experimental design is summarized in **Figure 4A**. Co-housing from P0 to

P7 improved nest-building and rescued deficits in pup-directed caring and retrieval behaviors in

stressed dams, while self-directed behaviors remained unchanged (**Figures 4B-I**). In contrast,

co-housing had no effect on maternal behavior in unstressed dams (**Figures 4B-I**).

Importantly, this maternal behavioral rescue translated into improved offspring behavioral

outcomes. Stressed offspring reared by dams co-housed with parous females during the

postpartum period exhibited restored sociability and social novelty recognition in the SIT

(**Figures 5A and 5B**). These findings indicate that social support from experienced mothers

during the early postpartum period selectively mitigates pup-directed maternal deficits and

ameliorates intergenerational social behavioral impairments.

Excitatory mCg→PrL firing is reduced in stressed offspring and restored by co-housing

with parous females

Building on evidence of cortico-cortical connectivity between the mCg and PrL in mice, we

performed retrograde viral tracing to identify upstream cortical inputs to the PrL. Consistent with

our prior identification of mCg→PrL projections (36), we focused on further characterizing this

pathway. Following retrograde viral injection into the PrL, the majority of labeled mCg cells co-

expressed CaMKII, a marker of excitatory neurons, whereas only a small fraction co-localized

with GABA, an inhibitory marker (**Figures 6A-C**). Furthermore, injection of a CaMKII-promoter-

driven retrograde virus into the PrL labeled numerous mCg cells (**Figure 6D**), confirming that

the mCg→PrL pathway is predominantly excitatory.

We next assessed the electrophysiological properties of mCg→PrL neurons in adult offspring.

Because cortico-cortical projection neurons from the mCg to the PrL are predominantly located

in layer V (48, 49), we focused our recordings on neurons in this layer. Whole-cell patch-clamp

recordings revealed a significant reduction in firing frequency in stressed offspring compared

with unstressed controls (**Figure 6E**), indicating hypofunction of the mCg→PrL pathway. In

contrast, rheobase and resting membrane potential were unchanged (**Supplemental Figures**

6A and 6B), suggesting that intrinsic membrane excitability was not altered. Analysis of

spontaneous excitatory and inhibitory postsynaptic currents showed no significant differences in

amplitude or frequency between groups (**Supplemental Figures 6C-F**), indicating that

presynaptic input to mCg→PrL neurons remains intact. Collectively, these data suggest that

impaired maternal care reduces the output activity of mCg→PrL neurons without affecting their

intrinsic excitability or upstream synaptic inputs. Notably, co-housing with parous females

restored firing activity in these neurons (**Figure 6E**), corresponding with normalized social

behaviors in adult offspring (**Figure 5**).

Discussion

Social isolation during adolescence establishes a latent vulnerability in female mice, selectively

impairing postpartum pup-directed behaviors, including licking, nursing, nest-building, and pup

retrieval. These specific maternal deficit traits are transmitted to the next generation, causing

persistent social deficits in adult offspring, while sparing other behavioral domains. Our research

revealed that the mCg→PrL projection is a critical excitatory pathway underlying these

intergenerational effects. Crucially, social contact with experienced parous mothers during the

early postpartum period selectively restores maternal care and rescues offspring's social

behavior. This demonstrates a causal link between maternal experience, circuit function, and

offspring development. These findings underscore the dual importance of adolescent

psychosocial experiences and the postpartum social environment in shaping maternal behavior

and offspring social trajectories.

The prefrontal cortex constitutes a core substrate for maternal and social behavior. Our data

indicate that adolescent social isolation reduces the firing output of an excitatory cortico-cortical

pathway in the prefrontal cortex (mCg→PrL), while presynaptic inputs remain largely intact. This

is consistent with prior reports that ELA selectively remodels excitatory cortico-cortical circuits,

with behavioral specificity reflecting vulnerability of defined projections (25, 32, 50, 51).

Candidate mechanisms include altered ion channel expression and modulation of conductance-

regulating signaling pathways, reminiscent of TRPM3-dependent regulation in social hierarchy

circuits (50). Functional manipulation sufficiently rescued social deficits by restoring activity

within this pathway, while its inhibition was confirmed to mimic adversity-induced impairments,

emphasizing a causal role for excitatory cortico-cortical dynamics in integrating maternal cues

and guiding offspring social outcomes.

The selectivity of behavioral changes in offspring highlights the critical influence of early

maternal care on social development (52, 53). Offspring exposed to suboptimal maternal care

exhibit elevated corticosterone levels while maintaining normal locomotion, anxiety responses,

and non-social memory, suggesting persistent stress axis dysregulation (6, 54). This pattern

aligns with prior evidence linking deficits in tactile and nursing stimulation to impaired HPA axis

negative feedback and abnormal stress responsivity (26, 55, 56). While oxytocinergic signaling

may also contribute to social vulnerability, its precise regulation by maternal care remains

unexplored (57-60). Our findings suggest that maternal deficits shape the development of social

circuits without broadly impairing other behavioral domains, providing a framework for

understanding how ELA confers selective vulnerability to social dysfunction across generations.

Communal rearing with experienced parous dams selectively improves maternal behavior in

stressed females, demonstrating the plasticity of maternal circuits (61, 62). Mechanistically, this

effect may arise through three complementary pathways: (1) Social support $-$ affiliative

presence reduces psychological stress and enhances caregiving motivation (63); (2) Behavioral

modeling $-$ stressed dams learn maternal behaviors by observing parous partners, leading to

sustained improvements even when the model is absent (40); (3) Stress buffering $-$ enhancing

caregiving capacity through normalization of corticosterone levels and restoration of mCg$\rightarrow$PrL

neuronal activity (64, 65). These results align with prior work demonstrating that maternal

behavior can be socially transmitted in mice and other mammals (39, 40, 66), emphasizing the

translational potential of supportive social environments to buffer adversity.

Collectively, our study elucidated a mechanistic framework linking adolescent psychosocial

adversity to postpartum maternal dysfunction and selective intergenerational social deficits. By

identifying the mCg$\rightarrow$PrL pathway as a critical substrate for experience-dependent regulation of

social behavior, these findings provide a basis for interventions to mitigate the persistent effects

of ELA. Future studies should elucidate the molecular and synaptic mechanisms underlying

circuit hypofunction, investigate the potential for transgenerational epigenetic modifications, and

explore the efficacy of structured social support interventions across diverse mammalian

models. Understanding these processes offers insight into the pathogenesis of social

dysfunction in humans and contributes to developing strategies that promote resilience to ELA.

Materials and Methods

Mice

C57BL/6J (B6J; JAX 000664) and B6;129S-Slc17a7tm1.1(cre)Hze/J (Vglut1-Cre; 023527) mice

were obtained from the Jackson Laboratory. All mice were backcrossed for more than ten

generations onto the C57BL/6J genetic background. Mice were maintained under controlled

environmental conditions (23 ± 3 °C; 40 ± 5% humidity) with a 12-hour light/dark cycle (lights on

at 6 a.m., lights off at 6 p.m.). Animals were housed in wire-topped plastic cages ($20 \times 31 \times 13$

cm) with ad libitum access to food and water. All experimental procedures adhered to the

National Institutes of Health Guidelines for the Care and Use of Laboratory Animals and were

approved by the Institutional Animal Care and Use Committee at the University of Alabama at

Birmingham (IACUC-21547). Both sexes were included to ensure generalizability, but sex

differences were not analyzed because the study was primarily designed to test maternal and

offspring outcomes and was not powered for sex-specific comparisons.

Unstressed and Stressed Dams; Unstressed and Stressed Offspring

Virgin female mice were either group-housed or socially isolated from five to eight weeks of age

to induce psychosocial adversity. At eight weeks of age, all females were mated with healthy

B6J males. Pregnancy was monitored daily, and the day of delivery was designated as P0.

These mice were designated as stressed dams. Age-matched controls (unstressed dams) were

group-housed (≥3 per cage) in wire-topped plastic cages during the same period. Pups raised

by unstressed dams were designated unstressed offspring, and pups raised by stressed dams

were designated stressed offspring. All offspring were weaned at P23 and group-housed

thereafter until adulthood (>P56), when experimental procedures were performed. Both sexes

were used in all offspring experiments.

Parous Mice

Female nulliparous mice were group-housed and mated, and following weaning of their litters,

they remained group-housed until use. These dams, with only one prior delivery and maternal

experience, are referred to as "parous females." To assess the effects of co-housing on

maternal behavior, either a virgin or parous female was introduced into the cages of unstressed

or stressed dams with their pups immediately after delivery (P0) (**Fig. 4A**). Co-housed females

remained together for the first postpartum week, during which maternal behaviors were

evaluated in both the co-housed females and the experimental dams. To examine the long-

lasting impact of co-housing with parous females on offspring social behavior, virgin or parous

females were co-housed with unstressed or stressed dams and their pups from birth until

weaning (P23). The introduced females were unfamiliar with the focal dams prior to co-housing.

All co-housed females were age-matched and had comparable social histories, except for prior

pregnancy and delivery. For parous females, at least three to five weeks elapsed between

weaning of their previous litter and introduction to the focal dams, ensuring recovery and

stabilization of maternal experience.

Behavioral testing overview

Basal maternal behavior and pup retrieval tests were conducted on P1 and P7, with both tests

performed at each time point. Unstressed and stressed dams co-housed with either parous or

virgin females were evaluated on P7. Separate cohorts of dams were used for P1 and P7, and

data from each timepoint were analyzed independently.

Behavioral assessments for adult offspring were conducted according to the experimental

timeline. The SIT included a habituation phase on P53-55 and the test phase on P56, followed

by the open field test on P57, the elevated plus maze test on P58, the light-dark box test on

P59, and brain/blood sampling on P60. Separate cohorts of mice were used for the novel object

recognition test on P56 and the social olfactory recognition test on P56.

Behavioral data were analyzed using EthoVision XT (versions 15 and 17; Noldus Information

Technology) for automated tracking, and manual measurements were used for the basal

maternal behavior and pup retrieval tests.

Basal Maternal Behavior Test

The basal maternal behavior test was conducted as previously described with minor

modifications (67, 68). Basal maternal behaviors were assessed in the home cage to evaluate

spontaneous caregiving activity in dams under undisturbed conditions. Each dam was observed

for 30 min, and all behaviors were recorded for subsequent manual scoring by an observer

blinded to experimental conditions. Nest quality was scored on a 4-point scale (0–3) based on

structural completeness and pup coverage, as follows: 0 = no nest; scattered bedding with no

structure; 1 = shallow or incomplete nest structure; 2 = well-defined nest with partial enclosure

of pups; 3 = fully enclosed nest with pups gathered and covered. The duration (in seconds) of

pup-directed and non-pup-directed behaviors was quantified as follows: Pup-directed behaviors:

Licking: the dam licks or grooms any part of the pup's body, primarily the anogenital region.

Nursing: the dam displays a crouching or arched-back posture with legs extended to cover the

litter. *Caring behavior:* total duration of licking and nursing combined. Non-pup-directed

behaviors: *Climbing and digging:* pushing or displacing bedding material with the snout or forepaws. *Self-grooming:* licking or scratching any part of the dam's own body. *Self-care:* total duration of climbing/digging and self-grooming combined. All behaviors were manually scored from video recordings by an experimenter blind to group assignments.

Pup Retrieval Test

Maternal responsiveness was evaluated using the pup retrieval test, as previously described with minor modifications (40, 59, 69). Pups were temporarily removed from the home cage, and dams were allowed to habituate for 30 minutes prior to testing. Three pups from the dam's own litter were placed individually in separate corners of the home cage, equidistant from the nest site. In the P7 condition, to prevent spontaneous pup movement, transparent cups were placed in three corners of the cage, and one pup from the dam's own litter was placed in each cup. The latency to retrieve each pup and return it to the nest was recorded. The dam was given up to 30 minutes to complete the retrieval of all three pups. The primary measure was the latency to retrieve the first pup (in seconds). If the dam failed to retrieve any pup within 30 min, a latency of 1800 s was assigned for statistical analysis. All trials were video recorded and scored by an observer blind to the experimental conditions.

Social Interaction Test (SIT)

The SIT was performed in adult offspring as previously described with minor modifications (25, 26, 70-72). The test consisted of two sequential trials: the Sociability Trial (S-trial) and the Social Novelty Trial (SN-trial), which were designed to assess progressively complex social behaviors. The testing apparatus consisted of a three-chamber box (60 cm × 40 cm × 35 cm) divided into left, middle, and right chambers, with small, barred cages positioned in the left and right compartments. The subject mouse was allowed to habituate to the three-chamber environment

for 10 minutes daily over three consecutive days before testing. On the test day, each mouse was placed in the middle chamber and allowed to habituate for 10 minutes before being confined to the middle chamber. The doors were then closed to create three separate chambers, and the subject mouse returned to the middle chamber for 5 minutes. During this period, a novel mouse was introduced into one of the barred cages, while a mouse-shaped object was placed in the other barred cage. In the S-trial, the subject mouse encountered the mouse in the barred cage and the object in the other barred cage for 10 minutes. At the 10-minute mark, the test mice were again restricted from entering the right and left chambers for 5 minutes, and the mouse-shaped object was replaced with another novel mouse. During the SN-trial, the subject mouse encountered a familiar mouse in one barred cage and a novel mouse in the other barred cage for 10 minutes. Familiar and novel mice were chosen to be age- and sex-matched healthy mice. Behavioral data were recorded and analyzed using Ethovision XT 17 software (Noldus). Nose contact and proximity within a 1cm boundary zone surrounding each cage were automatically detected, and the time spent sniffing each cage were quantified. Sociability and social novelty indexes were calculated as follows: (time interacting with mouse or novel mouse cage – time interacting with an object or familiar mouse cage) / (time interacting with a mouse or novel mouse cage + time interacting with an object or familiar mouse cage). An index of 0 indicated no preference. Positive indexes indicate increased sociability or social novelty behavior, whereas negative indexes indicated social avoidance or deficits in social novelty behavior.

Social Olfactory Recognition Test (SORT)

The SORT was conducted in adult offspring to assess social odor recognition and discrimination, as previously described with minor modifications (73). Each mouse was habituated to a clean testing cage, identical in dimensions to its home cage, for 30 min before testing. Odor stimuli

were presented using cotton balls soaked with odor solutions and placed in open 15 mL conical

tubes secured at one corner of the cage using a custom plastic holder. Three odor conditions

were used: (1) MilliQ water, (2) familiar scent, generated by agitating soiled bedding from the

subject's own home cage in MilliQ water, and (3) stranger scent, prepared from soiled bedding

collected from an age- and sex-matched unfamiliar mouse housed in a separate cage with no

prior contact. Each odor type was presented in three consecutive 2-min trials with no inter-trial

intervals. Cotton balls were replaced between trials to prevent cross-contamination and ensure

consistent odor presentation. The testing sequence was water → three sessions of familiar

scent → three sessions of stranger scent. Olfactory investigation was defined as direct nasal

contact or proximity (≤ 2.54 cm) to the cotton ball. Investigation time was recorded for each trial

and quantified as total exploration duration by an observer blinded to the experimental group.

Animals displaying little exploratory behavior (<10 s cumulative investigation across all trials)

were excluded from subsequent analysis.

Novel Object Recognition Test (NORT)

The NORT was conducted in adult offspring as previously described with minor modifications

(74, 75). Each trial consisted of three phases: habituation, training, and testing, each lasting 10

minutes. During the habituation phase on postnatal days 52-54, the subject mouse was

introduced into a chamber apparatus (40 cm × 40 cm × 42 cm) and allowed to habituate for 10

minutes per day over the course of three days. On postnatal day 55, during the training phase,

the mouse was once again habituated for 10 minutes in the chamber apparatus. Following this,

the mouse was briefly removed and then reintroduced into the chamber, which contained two

identical cone objects, for 2 minutes. Data during the training phase were collected over the

subsequent 10 minutes. On postnatal day 56, the test phase began with a 10-minute habituation

in the chamber apparatus. Subsequently, the mouse was briefly removed from the chamber.

During this brief removal, one of the two familiar cone objects was replaced with a novel

pyramid object. The mouse was then returned to the chamber for 2 minutes, after which data

collection took place over the following 10 minutes. All data were manually analyzed by

reviewing video recordings to identify instances of nose contact or physical interactions within a

1cm boundary zone around the objects. Stopwatch timers were used to measure the time spent

sniffing. The discrimination index was calculated as follows: (time sniffing the novel object – time

sniffing the familiar object / (time sniffing the novel object + time sniffing the familiar object). An

index of 0 indicated no recognition of the novel object, positive values indicated increased

recognition of the novel object, and negative values indicated deficits in recognizing the novel

object.

Open Field Test (OFT)

The OFT was conducted in adult offspring to assess general locomotor activity, as previously

described with minor modifications (76). Each mouse was placed in the center of a square

open-field arena (40 cm × 40 cm × 42 cm) constructed from opaque material to prevent external

visual distractions. The mice were allowed to explore the arena freely for 10 min under

standardized room lighting and ambient noise conditions. The total distance traveled was

automatically recorded and analyzed using EthoVision XT software.

Elevated Plus Maze Test (EPM)

Anxiety-like behavior was evaluated in adult offspring using the EPM, as previously described

with minor modifications (77-79). The maze consisted of two open arms (40 × 10 cm) and two

closed arms (40 × 10 cm, enclosed by 20-cm-high opaque gray walls) elevated 50 cm above

the floor. The apparatus was positioned in a quiet, dimly lit room to minimize environmental

stressors. Each mouse was placed in the central platform facing an open arm and allowed to

explore the maze freely for 10 min. Behavioral parameters measured included time spent in

open versus closed arms, number of entries into each arm, and total distance traveled. All data

were automatically recorded and analyzed using EthoVision XT software. Time spent in closed

arms was used as the primary index of anxiety-like behavior. The experimenter was positioned

outside the line of sight of the animal to avoid introducing stress-related artifacts.

Light-Dark Box test (LDB)

The LDB was conducted in adult offspring to assess anxiety-like behavior, as previously

described with minor modifications (80). The apparatus consisted of two compartments: a dark

chamber (opaque black walls and lid, impervious to visible light) and a light chamber (opaque

walls, illuminated under standard room lighting). The chambers were connected by a small

opening, allowing free movement between compartments. Each mouse was placed in the center

of either the light or dark compartment (facing away from the opening) and allowed to explore

for 10 min. Behavioral measures included total time spent in each compartment, number of

transitions between compartments, and latency to first enter the dark chamber. Time spent in

the dark chamber was considered a measure of anxiety-like behavior, with greater time

indicating increased anxiety. All behavior was automatically tracked and analyzed using

EthoVision XT software. Lighting and ambient noise were controlled to minimize environmental

variability.

DREADD manipulation of the mCg→PrL pathway

To selectively manipulate the mCg→PrL pathway, we employed DREADDs, which are

engineered G-protein coupled receptors activated exclusively by synthetic ligands such as CNO

(47). At P35, mice were anesthetized with isoflurane (5% induction, 1 - 2.5% maintenance) and

placed in a stereotaxic apparatus. Bilateral stereotaxic injections of 400 nL AAV-hSyn-

hM3D(Gq)-mCherry (4.3×10^{12} genomic copies/mL; #50474, Addgene) or AAV-hSyn-hM4D(Gi)-

mCherry (4.0×10^{12} genomic copies/mL; #50475, Addgene) were delivered into the mCg

[coordinates from bregma: AP +0.2 mm, medial-lateral (ML) ±0.3 mm, dorsal-ventral (DV) −1.7

mm] at a rate of 0.1 µL/min. The injection cannula was left in place for 5 minutes post-injection

to allow diffusion and prevent backflow. hM3Dq and hM4Di were used to induce neuronal

excitation and inhibition, respectively (81). During the same surgery, bilateral guide cannulas

(C235G-1.0/SPC, 26-gauge, Protech International) were implanted targeting the PrL (AP +2.34

mm, DV -1.45 mm, 0.5 mm above the intended microinjection site at DV -1.95 mm). Dummy

cannulas were inserted to maintain guide patency until P56. At P56, internal cannula extending

0.5 mm beyond the guides were used for microinjections. Neuronal activity was manipulated 30

minutes prior to SIT via bilateral microinjection of CNO (300 µM, 300 nL) or vehicle [1% dimethyl

sulfoxide (DMSO) in artificial cerebrospinal fluid (aCSF)] into the PrL (AP +2.34 mm, ML ±0.4

mm, DV −1.95 mm) using microsyringe pumps. Viral expression in the mCg and cannula

placement were verified via mCherry fluorescence, and only mice with confirmed expression in

the mCg→PrL pathway were included in behavioral analyses.

Brain slice electrophysiology of mCg→PrL projections

At P35, mice were stereotactically injected with 400 nL of rgAAV-hSyn-EGFP (1.93×10^{13}

genomic copies/mL; #50465, Addgene), rgAAV-CaMKIIa-mCherry (2.2×10^{13} genomic

copies/mL; #114469, Addgene), or rgAAV-CaMKIIa-EGFP (3.8×10^{12} genomic copies/mL;

#50469, Addgene) into the PrL (coordinates from bregma: AP +2.34 mm, ML ±0.4 mm, DV

−1.95 mm) at a rate of 40 nL/min. Three weeks were allowed for viral expression. At P56, mice

were anesthetized with isoflurane and decapitated. Brains were quickly removed, and 300-µm

coronal slices of the medial prefrontal cortex were prepared using a Leica VT1200S vibratome

in ice-cold, oxygenated slicing buffer (pH 7.3, 305 mOsm) containing 206 mM sucrose, 25 mM

sodium bicarbonate, 2.5 mM potassium chloride, 10 mM magnesium sulfate, 1.45 mM sodium

dihydrogen phosphate, 0.5 mM calcium chloride, and 11 mM D-glucose. The buffer was

continuously oxygenated with 5% CO_2 and 95% O_2. Slices were transferred to a holding

chamber containing oxygenated aCSF (126 mM NaCl, 26 mM $NaHCO_3$, 2.5 mM KCl, 1.45 mM

NaH_2PO_4, 1 mM $MgCl_2$, 2 mM $CaCl_2$, 9 mM D-glucose) at room temperature (22-25 °C) for at

least one hour. For recording, slices were placed in a submersion-type chamber on a

microscope stage and perfused with oxygenated aCSF at 30 ± 1.5 °C using an inline heater.

Recording pipettes were pulled from borosilicate glass (4-7 MΩ). Whole-cell patch-clamp

recordings were performed under differential interference contrast and fluorescence microscopy

targeting EGFP-positive neurons in layer V of the PrL. Spontaneous postsynaptic currents were

recorded in voltage-clamp mode using SutterPatch, held at -70 mV, with compensation for

electrode and cell capacitance. Recordings started five minutes after achieving whole-cell

configuration and lasted five minutes. Only neurons with stable access resistance (<20%

change) and baseline currents were included. Synaptic events were detected and analyzed with

SutterPatch software; mean amplitude and frequency for each neuron were used for statistics.

For spontaneous excitatory postsynaptic current experiments, 20 µM bicuculline was added to

the bath. Intracellular solution contained 117 mM K-gluconate, 13 mM KCl, 10 mM HEPES, 2

mM Na_2ATP, 0.4 mM Na_2GTP, 1 mM $MgCl_2$, 0.1 mM EGTA, 0.07 mM $CaCl_2$, pH 7.3, 290

mOsm. Action potential firing was recorded using current injection from -50 to +490 pA in 20 pA

steps; rheobase was the minimal current eliciting action potentials. For spontaneous inhibitory

postsynaptic current experiments, 20 µM DNQX and 50 µM D-APV were added to isolate

GABAergic currents. Intracellular solution contained 140 mM CsCl, 10 mM HEPES, 2 mM

Na_2ATP, 0.4 mM Na_2GTP, 1 mM $MgCl_2$, 0.05 mM EGTA, 3.6 mM NaCl, pH 7.3, 290 mOsm.

Blood and brain collection

Blood collection and brain perfusion were performed with minor modifications to previously

described methods (26, 82). To minimize the effects of circadian rhythms and acute stress from

behavioral testing, blood was collected at the same time of day, 24 hours after the last

behavioral session. One day following the light-dark box test, mice were anesthetized with

isoflurane. One mL of blood was collected from the right atrium between 8:00 am and 11:00 am

and transferred to 1.5 mL low-binding tubes. After blood collection, mice were perfused

intracardially with phosphate-buffered saline (PBS) followed by 4% paraformaldehyde. Brains

were extracted and post-fixed overnight in 4% paraformaldehyde at 4 °C and then transferred to

a 15% sucrose solution in PBS for one day, followed by 30% sucrose for an additional day.

Brains were subsequently embedded in optimal cutting temperature compound and stored at -

80 °C until sectioning.

Immunohistochemistry

Free-floating coronal brain sections (30 µm thickness) were prepared using a Leica CM3050S

cryostat and collected in PBS. Sections were blocked and permeabilized in PBS containing 5%

normal goat serum (NGS) and 0.1% Triton X-100 for 1 hour at room temperature to reduce

nonspecific binding. Sections were then incubated overnight at room temperature with one of

the following primary antibodies: Mouse anti-CaMKIIα (1:500, Millipore, 05-532) and rabbit anti-

GABA (1:1000, Sigma, A2052-25UL). After three 10-minute washes in PBS, sections were

incubated with the appropriate Alexa Fluor 647-conjugated secondary antibodies (1:400,

Thermo Fisher) for 2 hours at room temperature in the dark. Nuclear staining was performed

with DAPI (1:50,000, Roche, 10236276001) for 3 minutes at room temperature. Sections were

mounted on glass slides using Dako® fluorescent mounting medium and stored at 4 °C

protected from light until imaging. Images were acquired with either a Zeiss LSM800 confocal

microscope using Zen software or an Olympus BX61 epifluorescence microscope. The number

of $EGFP^+$, $EGFP^+CaMKII\alpha^+$, or $EGFP^+GABA^+$ cells was counted using Fiji/ImageJ software

package (83).

Serum Corticosterone Measurement

Blood samples were allowed to clot at room temperature for 1 hour and then centrifuged at

3,000 rpm for 15 minutes at 4 °C. The resulting serum was carefully collected and transferred

into 1.5 mL low-binding microcentrifuge tubes, then stored at −80 °C until analysis. Serum

corticosterone concentrations were measured using a commercially available ELISA kit

(Cayman Chemical, #501320) according to the manufacturer's instructions with minor

modifications (26, 82).

Statistical Analysis

All statistical analyses were performed using SPSS 28 (IBM) and GraphPad Prism 8.

Experiments were randomized whenever possible. For continuous data, assumptions of

normality and homogeneity of variances were assessed using the Shapiro-Wilk test and

Levene's test, respectively. For experiments involving multiple groups or repeated measures,

one-way analysis of variance (ANOVA), two-way ANOVA, or mixed-design ANOVA (including at

least one paired measure) was performed, followed by relevant pairwise comparisons with

Bonferroni correction. Data that violated only the assumption of homogeneity of variances were

analyzed using Welch's t-test or Welch ANOVA. For data that violated assumptions of normality

and/or homogeneity, non-parametric tests were applied, including the Mann-Whitney U test or

Kruskal-Wallis test. Statistical significance was defined as $p < 0.05$. Comparisons between two

groups were conducted using either Student's t-test or paired Student's t-test, as appropriate.

Effect sizes are reported for all analyses, and additional statistical parameters are provided in

Supplementary Table 1. Data are presented as mean ± standard error of the mean (SEM).

Acknowledgments

This work was supported by the National Institutes of Health (NIH) R01 MH-116869 (M.N.), the

UAB Comprehensive Neuroscience Center Pilot Award (M.N. and S.-i.K.), the Hill Crest

Foundation (M.N.), JST PRESTO grant JPMJPR14M6 (M.N.), and UAB Department of

Psychiatry startup funds (M.N.). R.T. is supported by a scholarship from the Uehara Memorial

Foundation. We thank members of the Niwa and Kano laboratories for their technical

assistance, data analysis support, and insightful discussions. We also thank Dr. Akira Sawa and

members of his laboratory for helpful discussions during the initial phase of the project. Figures

3A, 3D, 4A, 6A, and 6D were created with BioRender.com.

Figure 1. Adolescent social isolation impairs pup-directed maternal behavior without affecting

self-directed behavior at 1-week postpartum. (A-B) Stressed dams showed reduced nursing (A)

and pup-licking (B) behaviors. (C) Combined pup-directed care (nursing + licking) was

significantly decreased in stressed dams. (D-E) Self-directed behaviors, including

climbing/digging (D) and self-grooming (E), were unchanged. (F) Combined self-care behavior

(climbing/digging + self-grooming) did not differ between groups. (G) Stressed dams built

poorer-quality nests, reflected by lower nest scores. (H) Pup retrieval latency was significantly

prolonged in stressed dams, indicating diminished maternal responsiveness. Data are shown as

means ± SEM, N = 11-12 per group. *p < 0.05. Statistical tests: Student's t-test (A, C), Welch's

t-test (B), and Mann-Whitney U (D-H) were applied as appropriate. Detailed statistical analyses

are provided in **Supplementary Table 1**.

Figure 2. Offspring reared by stressed dams show selective social behavior deficits. (A) In the three-chamber social interaction test (SIT), stressed offspring showed reduced sociability compared with unstressed offspring. (B) Stressed offspring also displayed impaired social novelty recognition in the SIT. (C) In the social olfactory recognition test (SORT), stressed offspring exhibited diminished investigation toward both familiar and novel social odors. (D) Performance in the novel object recognition test (NORT) was unaffected by maternal adversity. (E) In the open field test (ORT), time spent in the center did not differ between groups. (F and G) Anxiety-related behaviors in the elevated plus maze (EPM) and light-dark box (LDB) showed no significant differences between groups. Together, these results indicate that impaired maternal behavior in stressed dams selectively disrupts offspring social behavior while sparing cognitive and anxiety-related domains. Data are shown as means ± SEM, N = 18-32 per group. *$p < 0.05$. Statistical tests: Welch's t-test (A, B), two-way mixed ANOVA (C), Mann-Whitney U (D, G), and Student's t-test (E, F), and were applied as appropriate. Detailed statistics are provided in **Supplementary Table 1**.

Figure 3

Figure 3. Chemogenetic modulation of mCg→PrL pathway rescues or induces social deficits. (A) Schematic illustrating virus delivery of AAV-hSyn-hM3D(Gq)-mCherry into the mCg and placement of guide cannulas into the PrL. (B and C) Activation of M3Dq-expressiong mCg→PrL neurons via CNO in stressed offspring rescued sociability and social novelty recognition deficits. (D) Schematic illustrating virus delivery of AAV-hSyn-hM4D(Gi)-mCherry into the mCg and placement of guide cannulas into the PrL. (E and F) Inhibition of the same circuit in unstressed offspring using M4Di receptors induced social deficits. Data are shown as means ± SEM, N = 6 per group. *p < 0.05. Statistical tests: Two-way ANOVA (B-C) and Student's t-test (E-F) were applied. Detailed statistics are provided in **Supplementary Table 1**.

Figure 4

Figure 4. Co-housing with parous females rescues impaired maternal caregiving in stressed

dams. (A) Experimental design of co-housing with parous females. (B-D) Pup-directed

behaviors were increased in stressed dams co-housed with parous, but not virgin, females. (E-

G) Self-care behaviors were not affected by co-housing conditions. (H and I) Nest quality and

pup retrieval performance were improved following co-housing with parous females. Data are

shown as means ± SEM, N = 12 per group. *p < 0.05. Statistical tests: All non-normal data were

analyzed using pairwise planned comparisons in Mann-Whitney U tests. Detailed statistics are

provided in **Supplementary Table 1**.

Figure 5. Co-housing with parous mice rescues social deficits in stressed offspring. (A)

Sociability index was decreased in stressed offspring but rescued by co-housing with parous

females from birth to weaning. (B) Social novelty recognition followed a similar pattern and was

rescued by parous co-housing. Data are shown as means ± SEM. Sample size for sociability

index is N = 7-18 per group. *p < 0.05. Statistical tests: Kruskal-Wallis test (A) and Welch's one-

way ANOVA (B) were applied. Detailed statistics are provided in **Supplementary Table 1**.

Figure 6

Figure 6. The mCg→PrL pathway is predominantly excitatory and shows reduced firing in stressed offspring that is restored by co-housing. (A) Left: Schematic illustrating virus delivery of rgAAV-hSyn-EGFP into the PrL. Right: Quantification of EGFP-labeled mCg neurons retrogradely labeled from PrL co-localized with CaMKII or GABA. Data are shown as means ± SEM, N = 3 per group. Each dot represents an average of three slices per animal. (B and C) Representative immunohistochemistry images of EGFP-labeled mCg neurons retrogradely labeled from PrL co-localized with CaMKII or GABA. Scale bars: 50 μm. (D) Schematic image and representative mCg image showing mCherry-labeled neurons following injection of a CaMKII promoter-driven retrograde virus into the PrL, confirming the excitatory nature of the mCg→PrL circuit. Scale bars: 100 μm. (E) Firing frequency of action potentials in layer V mCg→PrL neurons was reduced in stressed offspring, and rescued in stressed offspring co-housed with parous mice. Sample size for firing frequency is 10-13 cells per group, obtained from 4 different mice in each group. *p < 0.05. Statistical tests: Two-way mixed ANOVA (C) was used. Detailed statistics are provided in **Supplementary Table 1**.

References

1. R. Orso *et al.*, How Early Life Stress Impact Maternal Care: A Systematic Review of
Rodent Studies. *Front Behav Neurosci* **13**, 197-197 (2019).

2. J. G. Green *et al.*, Childhood adversities and adult psychiatric disorders in the national
comorbidity survey replication I: associations with first onset of DSM-IV disorders. *Arch*
Gen Psychiatry **67**, 113-123 (2010).

3. M. Numan, L. J. Young, Neural mechanisms of mother-infant bonding and pair bonding:
Similarities, differences, and broader implications. *Horm Behav* **77**, 98-112 (2016).

4. K. A. O'Donnell *et al.*, The maternal adversity, vulnerability and neurodevelopment
project: theory and methodology. *Can J Psychiatry* **59**, 497-508 (2014).

5. L. Murray, P. J. Cooper, Postpartum depression and child development. *Psychol Med*
27, 253-260 (1997).

6. F. A. Champagne, M. J. Meaney, Stress during gestation alters postpartum maternal
care and the development of the offspring in a rodent model. *Biol Psychiatry* **59**, 1227-
1235 (2006).

7. C. Heim *et al.*, Pituitary-adrenal and autonomic responses to stress in women after
sexual and physical abuse in childhood. *JAMA* **284**, 592-597 (2000).

8. J. Belsky, R. Conger, D. M. Capaldi, The intergenerational transmission of parenting:
introduction to the special section. *Dev Psychol* **45**, 1201-1204 (2009).

9. J. S. Merrick *et al.*, The Child Life Challenges Scale (CLCS): Associations of a Single-
Item Rating of Global Child Adversity with Children's Total Life Stressors and Parent's
Childhood Adversity. *Children (Basel)* **7** (2020).

10.   M. H. Labella, K. L. Raby, J. Martin, G. I. Roisman, Romantic functioning mediates

692   prospective associations between childhood abuse and neglect and parenting outcomes

693   in adulthood. *Dev Psychopathol* **31**, 95-111 (2019).

694   11.   S. Lockhart, A. Sawa, M. Niwa, Developmental trajectories of brain maturation and

695   behavior: Relevance to major mental illnesses. *Journal of pharmacological sciences*

696   **137**, 1-4 (2018).

697   12.   A. C. Viguera *et al.*, Episodes of mood disorders in 2,252 pregnancies and postpartum

698   periods. *Am J Psychiatry* **168**, 1179-1185 (2011).

699   13.   C. Fu *et al.*, The Relationship Between Adverse Childhood Experiences and Postpartum

700   Depression: A Systematic Review and Meta-Analysis. *Trauma Violence Abuse* **25**, 3066-

701   3081 (2024).

702   14.   S. Brummelte, L. A. Galea, Depression during pregnancy and postpartum: contribution

703   of stress and ovarian hormones. *Prog Neuropsychopharmacol Biol Psychiatry* **34**, 766-

704   776 (2010).

705   15.   S. Brummelte, J. L. Pawluski, L. A. Galea, High post-partum levels of corticosterone

706   given to dams influence postnatal hippocampal cell proliferation and behavior of

707   offspring: A model of post-partum stress and possible depression. *Horm Behav* **50**, 370-

708   382 (2006).

709   16.   J. Maguire, I. Mody, GABA(A)R plasticity during pregnancy: relevance to postpartum

710   depression. *Neuron* **59**, 207-213 (2008).

711   17.   J. W. Smith, J. R. Seckl, A. T. Evans, B. Costall, J. W. Smythe, Gestational stress

712   induces post-partum depression-like behaviour and alters maternal care in rats.

713   *Psychoneuroendocrinology* **29**, 227-244 (2004).

714   18.   M. L. Boccia *et al.*, Repeated long separations from pups produce depression-like

715   behavior in rat mothers. *Psychoneuroendocrinology* **32**, 65-71 (2007).

716   19.   S. J. Blakemore, The social brain in adolescence. *Nat Rev Neurosci* **9**, 267-277 (2008).

717   20.   A. Orben, L. Tomova, S. J. Blakemore, The effects of social deprivation on adolescent

718   development and mental health. *Lancet Child Adolesc Health* **4**, 634-640 (2020).

719   21.   V. Begni *et al.*, Social isolation in rats: Effects on animal welfare and molecular markers

720   for neuroplasticity. *PLoS One* **15**, e0240439 (2020).

721   22.   A. Borsini, J. Giacobbe, G. Mandal, M. Boldrini, Acute and long-term effects of

722   adolescence stress exposure on rodent adult hippocampal neurogenesis, cognition, and

723   behaviour. *Mol Psychiatry* **28**, 4124-4137 (2023).

724   23.   K. C. Fone, M. V. Porkess, Behavioural and neurochemical effects of post-weaning

725   social isolation in rodents-relevance to developmental neuropsychiatric disorders.

726   *Neurosci Biobehav Rev* **32**, 1087-1102 (2008).

727   24.   E. J. Nestler, S. E. Hyman, Animal models of neuropsychiatric disorders. *Nat Neurosci*

728   **13**, 1161-1169 (2010).

729   25.   K. Kin, J. Francis-Oliveira, S. I. Kano, M. Niwa, Adolescent stress impairs postpartum

730   social behavior via anterior insula-prelimbic pathway in mice. *Nat Commun* **14**, 2975

731   (2023).

732   26.   M. Niwa *et al.*, Prolonged HPA axis dysregulation in postpartum depression associated

733   with adverse early life experiences: A cross-species translational study. *Nat Ment Health*

734   **2**, 593-604 (2024).

735   27.   E. Hoekzema *et al.*, Pregnancy leads to long-lasting changes in human brain structure.

736   *Nat Neurosci* 10.1038/nn.4458 (2016).

737   28.   J. L. Pawluski, J. S. Lonstein, A. S. Fleming, The Neurobiology of Postpartum Anxiety

738   and Depression. *Trends Neurosci* **40**, 106-120 (2017).

739   29.   J. K. Rilling, L. J. Young, The biology of mammalian parenting and its effect on offspring

740   social development. *Science* **345**, 771-776 (2014).

30. L. K. Bicks, H. Koike, S. Akbarian, H. Morishita, Prefrontal Cortex and Social Cognition
in Mouse and Man. *Front Psychol* **6**, 1805 (2015).

31. C. E. Forbes, J. Grafman, The role of the human prefrontal cortex in social cognition and
moral judgment. *Annu Rev Neurosci* **33**, 299-324 (2010).

32. M. Febo, A. C. Felix-Ortiz, T. R. Johnson, Inactivation or inhibition of neuronal activity in
the medial prefrontal cortex largely reduces pup retrieval and grouping in maternal rats.
Brain Res **1325**, 77-88 (2010).

33. V. M. Afonso, M. Sison, V. Lovic, A. S. Fleming, Medial prefrontal cortex lesions in the
female rat affect sexual and maternal behavior and their sequential organization. *Behav*
Neurosci **121**, 515-526 (2007).

34. S. van Heukelum *et al.*, Where is Cingulate Cortex? A Cross-Species View. *Trends*
Neurosci **43**, 285-299 (2020).

35. B. A. Vogt, Midcingulate cortex: Structure, connections, homologies, functions and
diseases. *J Chem Neuroanat* **74**, 28-46 (2016).

36. J. Francis-Oliveira, O. Leitzel, M. Niwa, Are the Anterior and Mid-Cingulate Cortices
Distinct in Rodents? *Front Neuroanat* **16**, 914359 (2022).

37. A. Etkin, T. Egner, R. Kalisch, Emotional processing in anterior cingulate and medial
prefrontal cortex. *Trends Cogn Sci* **15**, 85-93 (2011).

38. G. Bush, P. Luu, M. I. Posner, Cognitive and emotional influences in anterior cingulate
cortex. *Trends Cogn Sci* **4**, 215-222 (2000).

39. T. R. Insel, L. J. Young, The neurobiology of attachment. *Nat Rev Neurosci* **2**, 129-136
(2001).

40. I. Carcea *et al.*, Oxytocin neurons enable social transmission of maternal behaviour.
Nature **596**, 553-557 (2021).

41.   C. Dulac, L. A. O'Connell, Z. Wu, Neural control of maternal and paternal behaviors.

766   *Science* **345**, 765-770 (2014).

767   42.   R. Feldman, The Neurobiology of Human Attachments. *Trends Cogn Sci* **21**, 80-99

768   (2017).

769   43.   M. Opendak, E. Gould, R. Sullivan, Early life adversity during the infant sensitive period

770   for attachment: Programming of behavioral neurobiology of threat processing and social

771   behavior. *Dev Cogn Neurosci* **25**, 145-159 (2017).

772   44.   B. A. Hotelling, Newborn capabilities: parent teaching is a necessity. *J Perinat Educ* **13**,

773   43-49 (2004).

774   45.   J. Guintivano *et al.*, Adverse life events, psychiatric history, and biological predictors of

775   postpartum depression in an ethnically diverse sample of postpartum women. *Psychol*

776   *Med* **48**, 1190-1200 (2018).

777   46.   S. E. Mathisen, K. Glavin, L. Lien, P. Lagerlov, Prevalence and risk factors for

778   postpartum depressive symptoms in Argentina: a cross-sectional study. *Int J Womens*

779   *Health* **5**, 787-793 (2013).

780   47.   B. L. Roth, DREADDs for Neuroscientists. *Neuron* **89**, 683-694 (2016).

781   48.   S. Moberg, N. Takahashi, Neocortical layer 5 subclasses: From cellular properties to

782   roles in behavior. *Front Synaptic Neurosci* **14**, 1006773 (2022).

783   49.   C. Song, J. R. Moyer, Jr., Layer- and subregion-specific differences in the

784   neurophysiological properties of rat medial prefrontal cortex pyramidal neurons. *J*

785   *Neurophysiol* **119**, 177-191 (2018).

786   50.   A. C. Nelson *et al.*, Molecular and neural control of social hierarchy by a forebrain-

787   thalamocortical circuit. *Cell* **188**, 5535-5554 e5523 (2025).

788   51.   S. Tzanoulinou, O. Riccio, M. W. de Boer, C. Sandi, Peripubertal stress-induced
789   behavioral changes are associated with altered expression of genes involved in
790   excitation and inhibition in the amygdala. *Transl Psychiatry* **4**, e410 (2014).

791   52.   C. Caldji *et al.*, Maternal care during infancy regulates the development of neural
792   systems mediating the expression of fearfulness in the rat. *Proc Natl Acad Sci U S A* **95**,
793   5335-5340 (1998).

794   53.   C. M. Lindeyer, M. J. Meaney, S. M. Reader, Early maternal care predicts reliance on
795   social learning about food in adult rats. *Dev Psychobiol* **55**, 168-175 (2013).

796   54.   M. Nishi, N. Horii-Hayashi, T. Sasagawa, Effects of early life adverse experiences on the
797   brain: implications from maternal separation models in rodents. *Front Neurosci* **8**, 166
798   (2014).

799   55.   D. Liu *et al.*, Maternal care, hippocampal glucocorticoid receptors, and hypothalamic-
800   pituitary-adrenal responses to stress. *Science* **277**, 1659-1662 (1997).

801   56.   A. S. Ivy, K. L. Brunson, C. Sandman, T. Z. Baram, Dysfunctional nurturing behavior in
802   rat dams with limited access to nesting material: a clinically relevant model for early-life
803   stress. *Neuroscience* **154**, 1132-1142 (2008).

804   57.   S. Kojima, R. A. Stewart, G. E. Demas, J. R. Alberts, Maternal contact differentially
805   modulates central and peripheral oxytocin in rat pups during a brief regime of mother-
806   pup interaction that induces a filial huddling preference. *J Neuroendocrinol* **24**, 831-840
807   (2012).

808   58.   A. S. Todeschin *et al.*, Effects of neonatal handling on social memory, social interaction,
809   and number of oxytocin and vasopressin neurons in rats. *Horm Behav* **56**, 93-100
810   (2009).

811   59.   P. Toepfer *et al.*, Oxytocin pathways in the intergenerational transmission of maternal
812   early life stress. *Neurosci Biobehav Rev* **73**, 293-308 (2017).

813   60.   S. C. Walker, P. D. Trotter, W. T. Swaney, A. Marshall, F. P. McGlone, C-tactile

814   afferents: Cutaneous mediators of oxytocin release during affiliative tactile interactions?

815   *Neuropeptides* **64**, 27-38 (2017).

816   61.   J. P. Curley, F. A. Champagne, Influence of maternal care on the developing brain:

817   Mechanisms, temporal dynamics and sensitive periods. *Front Neuroendocrinol* **40**, 52-66

818   (2016).

819   62.   I. Branchi, The mouse communal nest: investigating the epigenetic influences of the

820   early social environment on brain and behavior development. *Neurosci Biobehav Rev*

821   **33**, 551-559 (2009).

822   63.   M. R. Denomme, G. J. Mason, Social Buffering as a Tool for Improving Rodent Welfare.

823   *J Am Assoc Lab Anim Sci* **61**, 5-14 (2022).

824   64.   J. Varela, J. Wang, A. Varnell, D. Cooper, Control over stress induces plasticity of

825   individual prefrontal cortical neurons: A conductance-based neural simulation. *Nature*

826   *Precedings* 10.1038/npre.2011.6267.1 (2011).

827   65.   J. A. Varela, J. Wang, J. P. Christianson, S. F. Maier, D. C. Cooper, Control over stress,

828   but not stress per se increases prefrontal cortical pyramidal neuron excitability. *J*

829   *Neurosci* **32**, 12848-12853 (2012).

830   66.   B. J. Marlin, M. Mitre, A. D'Amour J, M. V. Chao, R. C. Froemke, Oxytocin enables

831   maternal behaviour by balancing cortical inhibition. *Nature* **520**, 499-504 (2015).

832   67.   K. O. Kuroda, K. Tachikawa, S. Yoshida, Y. Tsuneoka, M. Numan, Neuromolecular

833   basis of parental behavior in laboratory mice and rats: with special emphasis on

834   technical issues of using mouse genetics. *Prog Neuropsychopharmacol Biol Psychiatry*

835   **35**, 1205-1231 (2011).

836   68.   H. J. Kim, J. Jang, H. Y. Koh, Abnormal maternal behavior in mice lacking

837   phospholipase Cbeta1. *Anim Cells Syst (Seoul)* **26**, 291-299 (2022).

69.   G. Esposito *et al.*, Infant calming responses during maternal carrying in humans and
mice. *Curr Biol* **23**, 739-745 (2013).

70.   O. Kaidanovich-Beilin, T. Lipina, I. Vukobradovic, J. Roder, J. R. Woodgett, Assessment
of social interaction behaviors. *J Vis Exp* 10.3791/2473 (2011).

71.   S. S. Moy *et al.*, Sociability and preference for social novelty in five inbred strains: an
approach to assess autistic-like behavior in mice. *Genes Brain Behav* **3**, 287-302 (2004).

72.   C. Sandi, J. Haller, Stress and the social brain: behavioural effects and neurobiological
mechanisms. *Nature reviews. Neuroscience* **16**, 290-304 (2015).

73.   M. Yang, J. N. Crawley, Simple behavioral assessment of mouse olfaction. *Curr Protoc*
Neurosci **Chapter 8**, Unit 8 24 (2009).

74.   Q. Sun *et al.*, Ventral Hippocampal-Prefrontal Interaction Affects Social Behavior via
Parvalbumin Positive Neurons in the Medial Prefrontal Cortex. *iScience* **23**, 100894
(2020).

75.   J. H. Lee *et al.*, Astrocytes phagocytose adult hippocampal synapses for circuit
homeostasis. *Nature* **590**, 612-617 (2021).

76.   A. K. Kraeuter, P. C. Guest, Z. Sarnyai, The Open Field Test for Measuring Locomotor
Activity and Anxiety-Like Behavior. *Methods Mol Biol* **1916**, 99-103 (2019).

77.   A. K. Kraeuter, P. C. Guest, Z. Sarnyai, The Elevated Plus Maze Test for Measuring
Anxiety-Like Behavior in Rodents. *Methods Mol Biol* **1916**, 69-74 (2019).

78.   K. Yoshizaki, M. Asai, T. Hara, High-Fat Diet Enhances Working Memory in the Y-Maze
Test in Male C57BL/6J Mice with Less Anxiety in the Elevated Plus Maze Test. *Nutrients*
12 (2020).

79.   E. Dohi *et al.*, Behavioral Changes in Mice Lacking Interleukin-33. *eNeuro* **4** (2017).

80.   A. C. Campos, M. V. Fogaca, D. C. Aguiar, F. S. Guimaraes, Animal models of anxiety
disorders and stress. *Braz J Psychiatry* **35 Suppl 2**, S101-111 (2013).

81.   S. Zhang *et al.*, Molecular basis for selective activation of DREADD-based

864   chemogenetics. *Nature* **612**, 354-362 (2022).

865   82.   M. Niwa *et al.*, Adolescent stress-induced epigenetic control of dopaminergic neurons

866   via glucocorticoids. *Science* **339**, 335-339 (2013).

867   83.   J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat*

868   *Methods* **9**, 676-682 (2012).