Modeling VEGF and GLUT1 Expression as Coadapted Foraging Strategies in Cancer

Ranjini Bhattacharya [1,2,3], Robert A. Gatenby [1,2,4], Joel S. Brown [1,2]

1 Cancer Biology and Evolution Program, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, USA

2 Department of Integrated Mathematical Oncology, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, USA

3 Department of Cancer Biology, University of South Florida, Tampa, FL, USA

4 Department of Radiology, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, USA

Corresponding Author: Ranjini Bhattacharya, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, USA

Corresponding Author Email: Ranjini.bhattacharya@moffitt.org

Abstract

Natural selection acting on cancer cells within their tumor microenvironment should favor cells with fast or efficient nutrient uptake strategies. Here, we develop and analyze a game-theoretic model focusing on the coadaptation between two foraging traits: vascular endothelial growth factor (VEGF) and glucose transporter 1 (GLUT1). Studies show that VEGF and GLUT1 are often co-expressed and are associated with more aggressive tumor phenotypes and poor clinical prognosis. VEGF is a diffusible paracrine factor that recruits blood vessels towards neighborhoods of cancer cells (angiogenesis). GLUT1 is a cell-surface transporter that enables the uptake of nutrients, especially glucose. We model these strategies operating at different scales: VEGF influences resource availability at the neighborhood level, while GLUT1 determines resource uptake at the cellular level. For VEGF, we introduce a resource-sharing continuum. With no resource sharing, cells access resources in proportion to their VEGF contribution. With uniform sharing, cells have equal access to resources, regardless of their VEGF contribution. The former leads to a tragedy of the commons and overproduction of VEGF. The latter yields a public goods game with moderate VEGF expression matching a group optimum. GLUT1 expression mediates uptake of resources recruited by VEGF and is largely independent of the degree of resource sharing. Therapeutically, both VEGF and GLUT1 inhibitors are more effective in high resource-sharing neighborhoods and less so as resource sharing declines. Overall,

34 inhibition of GLUT1-mediated uptake emerges as more effective. The model, perhaps
35 the first to consider VEGF and GLUT1 as coadaptations, emphasizes the need to
36 consider cancer cell traits jointly.

37 **Keywords**: VEGF, GLUT1, tumor angiogenesis, resource sharing, evolutionary game
38 theory, glucose uptake, cancer foraging, coadapted syndromes, therapy

1. <u>Introduction</u>

Cancer cells exist in a resource-limited tumor microenvironment. This scarcity arises
from both inadequate vascular supply and rapid nutrient consumption by proliferating
cells [1,2]. In response, cancer cells modulate a range of signaling pathways and
transport mechanisms, including the expression of vascular endothelial growth factor
(VEGF), which promotes angiogenesis, and glucose transporter 1 (GLUT1), which
facilitates glucose uptake [3–6]. While VEGF expression increases blood supply within a
region, GLUT1 expression increases a cell's capacity to take up glucose. Together,
these responses act in concert as a coadaptation [7,8] for optimizing resource
acquisition in cancer cells. Evolutionary pressures within the tumor should shape the
cell's investment in these two strategies and determine how expression of one
influences the expression of the other. In this study, we will explore how these
complementary strategies evolve together under shared constraints.

In response to hypoxia and resource deprivation in the tumor, the master transcriptional
regulator, hypoxia inducible factor-1 (HIF-1), is activated [9,10]. Among its downstream
targets are VEGF and GLUT1. VEGF is a paracrine signaling factor released into the
tumor microenvironment that instigates the angiogenic cascade. It binds to VEGF
receptors (primarily VEGFR-2) on endothelial cells lining existing blood vessels,
activating signaling pathways that promote endothelial cell proliferation, migration, and
sprouting toward regions of high VEGF concentration [11,12]. The local concentration of
VEGF in a region influences the extent of neovascular growth. In addition to its
paracrine effects, VEGF can also bind to VEGF receptors expressed on cancer cells,
promoting autocrine signaling that aids cellular proliferation [13,14].

Glucose transporters (GLUTs) belong to the SLC2A family of facilitative membrane
transporters and mediate glucose uptake [6]. Among the fourteen isoforms, GLUT1 is
the most consistently upregulated in cancers and is used clinically as the basis for
fluorodeoxyglucose positron emission tomography (FDG-PET) imaging [5,15–18].
GLUT1 localizes to the plasma membrane and increases cellular glucose influx,
supporting the elevated glycolytic demands of cancer cells. Other isoforms, such as
GLUT3, are also upregulated in tumors with high metabolic demand [19–21]. Although
glucose remains the primary substrate driving tumor metabolism, certain GLUT isoforms
can transport other hexoses and related molecules, such as dehydroascorbic acid,

reflecting the metabolic plasticity of cancer cells under resource-limited conditions
[22,23].

Co-expression of VEGF and GLUT1 is frequently associated with increased microvessel
density, elevated metabolic activity, and more aggressive tumor phenotypes [24–26].
For example, studies in esophageal and ovarian cancers have reported associations
between GLUT1 expression, VEGF expression, and measures of vascularization and
glucose uptake [27]. In esophageal squamous cell carcinoma, GLUT1 expression has
been shown to correlate with increased VEGF expression and FDG uptake [28]. These
findings suggest a close relationship between these two traits.

These markers have also been evaluated for their prognostic significance. Elevated
expression of VEGF and GLUT1, either individually or in combination with other
markers such as carbonic anhydrase IX (CAIX), has been associated with poor survival,
increased tumor aggressiveness, and resistance to therapy across multiple cancer
types [29–32]. In some cases, a combined assessment of VEGF and GLUT1 provides
greater prognostic value than the individual markers [31]. Together, these observations
highlight the importance of considering these two adaptations as co-evolved strategies,
rather than in isolation.

Tumor angiogenesis has been extensively modeled. Early continuum and hybrid models
developed by Anderson and Chaplain [33,34], as well as Byrne and collaborators
[35,36] laid the foundation for describing endothelial cell migration, chemotaxis, and
vascular sprouting in response to angiogenic factors. Subsequent work has
incorporated increasingly detailed mechanistic descriptions of VEGF signaling and
transport. For example, Finley et al. developed compartmental models of VEGF
distribution that account for VEGF isoforms, receptor binding, and interactions across
tumor compartments [37,38]. Hybrid models such as those by Phillips et al. [39] and
Nikmaneshi et al. [40] have coupled tumor growth with angiogenesis through feedback
between hypoxia, VEGF production, and vascular supply. In addition to mechanistic and
multiscale models, evolutionary and game-theoretic approaches have examined
angiogenesis as a collective behavior, framing VEGF production as a public goods
game, where individual cells balance the costs of VEGF production against shared
benefits from VEGF expression [41,42].

In contrast to the extensive modeling of angiogenesis, mathematical models of tumor
metabolism have largely focused on glucose as a diffusible resource, with less attention
given to the regulatory mechanisms governing glucose uptake, such as GLUT1
expression. Early mathematical models of tumor growth, such as those developed by
Greenspan [43], incorporated nutrient limitation through diffusion-based frameworks to
describe how the availability of substrates such as oxygen and glucose constrain tumor
expansion. Subsequent work by Casciari et al. [44] and Ward and King [45] extended
these approaches by modeling the kinetics of glucose and oxygen consumption using

reaction–diffusion equations, linking nutrient transport to cell proliferation, quiescence,
and necrosis. Gatenby et al. [46] modeled glucose metabolism and acidosis using
cellular automaton frameworks, to examine how gradients of oxygen, glucose, and
acidity drive the selection of glycolytic phenotypes, providing a theoretical basis for the
emergence of the Warburg effect in cancer [47–49]. Subsequent work has treated
glycolytic and oxidative cells as distinct subpopulations. For example, Kareva et al. [50]
modeled tumor–immune–glucose interactions using predator–prey dynamics. More
recent approaches have incorporated glucose uptake into models of fluorodeoxyglucose
(FDG) transport and uptake used to interpret PET imaging data [51,52]. Mechanistic
models have also examined the role of glucose transport in tumor proliferation, including
work by Ampatzoglou et al. [53], which explicitly incorporates GLUT1 expression into a
multiscale tumor growth model.

Despite extensive empirical and theoretical work on angiogenesis and glucose
metabolism, existing models have largely treated VEGF and GLUT1 expression as
independent processes. To our knowledge, no modeling framework has examined
these traits as coordinated, co-adapted strategies. Here, we address this gap.

In our earlier work, we developed an evolutionary game-theoretic model to study how
cancer cells evolve VEGF expression as a function of local resource sharing [21]. We
showed that in tumor neighborhoods with high resource sharing, where resources are
distributed uniformly among cells, cells engage in a cooperative public goods game. In
contrast, when resource sharing is limited, cells are incentivized to increase VEGF
expression to secure a larger share of resources, giving rise to a tragedy of the
commons [54,55]. In this regime, excessive individual investment in VEGF ultimately
reduces overall tumor fitness. These results demonstrated the influence of cell-cell
interactions on the evolution of VEGF as a foraging strategy.

Here, we build on this perspective and study the coadaptation of VEGF and GLUT1
expression as foraging strategies [56]. VEGF acts as a diffusible, paracrine signal that
modulates resource availability at the neighborhood level, such that a cell's access to
nutrients depends on both its own expression and that of its neighbors [21]. In contrast,
GLUT1 functions at the level of individual cells, with glucose uptake determined
primarily by a cell's own expression. These traits therefore, operate at different scales
and impose distinct costs [57,58]: VEGF production requires sustained biosynthetic
investment and generates shared benefits, whereas GLUT1 overexpression increases
metabolic demand and glycolytic flux. Natural selection within the tumor
microenvironment acts on the expression of these traits, shaping how cells allocate
investment between these strategies under varying ecological and resource-sharing
conditions.

Here, we model the "game" that emerges as cancer cells allocate investment to these
two complementary foraging strategies within a tumor. Specifically, we 1) develop an

evolutionary game-theoretic framework [59,60] in which the VEGF and GLUT1 are
coadapted traits, 2) analyze how investment in these traits varies along a continuum of
resource-sharing conditions, 3) analyze the effects of key parameters, and 4) explore
the implications of these dynamics for therapeutic strategies targeting VEGF and
GLUT1.

2. <u>Methods</u>

To investigate how cancer cells allocate investment between VEGF and GLUT1
expression, we consider a population of cancer cells organized into local neighborhoods
within the tumor microenvironment. The neighborhood is described by cells that directly
interact and influence each other's access to nutrients supplied by shared blood
vasculature [21,61]. The collective VEGF production of the neighborhood diverts blood
flow towards it, determining the total resource supply. A part of this supply is available to
each of the cancer cells in the neighborhood [62]. For a cancer cell to obtain its share, it
must invest in GLUT1, which enables resource uptake from the blood. We model VEGF
expression and GLUT1 expression as continuous quantitative traits representing
investment in resource supply and uptake, respectively. VEGF acts at the neighborhood
level by modulating resource availability, whereas GLUT1 acts at the cellular level by
determining a cell's ability to capture available resources.

We formalize these interactions using a fitness-generating function (G-function) [63–65],
which describes the per capita growth rate of a focal cell within its neighborhood. A focal
cell's fitness is influenced by the number of cancer cells within its neighborhood (N), the
average VEGF expression of the other $N-1$ cells of the neighborhood (u), the focal
cell's GLUT1 expression (y), and the focal cell's VEGF expression (v): $G(N, u, v, y)$.

We assume that the total VEGF production of all cells, $x = v + u(N-1)$, influences the
overall vasculature and resource supply to the neighborhood, and that both v and u can
influence how much of this supply is available to the focal cell. Given a share of the
resource supply, it is only the focal individual's investment in GLUT1, y, and not that of
others, which influences the uptake of its share. We develop these assumptions into the
following model:

Resource uptake via GLUT1

Resource redirection by total VEGF in the neighborhood

$$G(N, u, v, y) = b\left((1 - e^{-ky})\left(\frac{q}{N} + \frac{(1-q)v}{x}\right)R(1 - e^{-ax}) - cv - hy - f\right) \qquad (1)$$

Resource distribution between focal cell and its neighbors

Cost

where R is the maximum possible resource influx, a is the efficacy of VEGF mediated
resource redirection to the neighborhood, k is the efficacy of GLUT1 mediated resource
uptake, c and h are the per unit costs of VEGF and GLUT1 expression, respectively,
and f is the fixed cost of survival. In Eq. (1), $R(1 - e^{-ax})$ determines the total amount of
resources entering a neighborhood as a function of x, the neighborhood's total VEGF
expression. $q \in [0,1]$ is the degree of resource sharing in the neighborhood. When the
degree of resource sharing is maximum ($q = 1$), resources are evenly distributed
regardless of each cell's VEGF contribution. When there is no resource sharing ($q = 0$),
resources are allocated in proportion to each cell's VEGF contribution. The term,
$(1 - e^{-ky})$, represents GLUT1 mediated resource import, with the assumption that
glucose uptake increases with GLUT1 expression but eventually saturates (as the
function approaches 1).

The *G*-function models both cancer population dynamics and trait dynamics. The rate of
evolution of the two traits, VEGF and GLUT1 expression, is a function of the selection
gradient and the cell's phenotypic plasticity in these traits. The cancer population (N),
VEGF expression (u), and GLUT1 expression (y) dynamics are governed by Eq. (2),
(3), (4).

$$\frac{dN}{dt} = NG \tag{2}$$

$$\frac{du}{dt} = \sigma_u^2 \left.\frac{\partial G}{\partial v}\right|_{v=u} \tag{3}$$

$$\frac{dy}{dt} = \sigma_y^2 \frac{\partial G}{\partial y} \tag{4}$$

195 where $\frac{\partial G}{\partial v}$ and $\frac{\partial G}{\partial y}$ are the selection gradients with respect to the focal cell's VEGF
196 expression and GLUT1 expression respectively. The parameters σ_u^2 and σ_y^2 are the
197 speed of phenotypic plasticity in VEGF and GLUT1 expression, respectively, in
198 response to the strength of selection [23,66].

2.1 Modeling Therapy

To model therapy, we consider two drug modalities. The first are VEGF inhibitors or
neutralizing agents that reduce the effectiveness of VEGF-mediated resource
recruitment to the tumor (a). This class includes clinically approved drugs such as
bevacizumab and sunitinib, which target VEGF signaling and angiogenesis [67–69]. The
second are GLUT1 inhibitors, which reduce the efficacy of GLUT1-mediated glucose
import into cells (k). While fewer GLUT1 inhibitors are currently in clinical use, several
compounds are under development or study, including STF-31, WZB117, BAY-876, and

ritonavir, which act through competitive or noncompetitive mechanisms to block glucose
uptake [70–73]. We modify Eq. (1) to incorporate these therapies as follows-

In Eq. (5), m_G is the efficacy of the GLUT1 inhibitor, and m_A is the efficacy of VEGF

$$G(N, u, v, y) = b\left(\left(1 - e^{-\frac{ky}{1+em_G}}\right)\left(\frac{q}{N} + \frac{(1-q)v}{x}\right)R\left(1 - e^{-\frac{ax}{1+dm_A}}\right) - cv - hy - f\right) \tag{5}$$

211 inhibitor.

2.2 Simulations

All simulations were performed in Python. To identify the evolutionarily stable strategy
(ESS) and the team optimum, the system was iterated until equilibrium was reached,
and the resulting steady-state values were recorded. Details of the sensitivity analysis
are provided in the Supplementary Section. Variables and parameters used in the
model are summarized in Tables 1 and 2.

In this theoretical body of work, we are interested in the emergent model dynamics in
different parameter ranges. The scaling and units of the parameters can be adjusted to
align with experimental measurements. For example, VEGF may be quantified using
tissue-level RNA expression or circulating protein concentration, while GLUT1
expression may be measured via RNA expression or immunofluorescence intensity
[72,74–77].

Symbol	Meaning
v	VEGF produced by focal cell
u	Average VEGF produced by all cells in a neighborhood
x	Total VEGF produced by all cells in a neighborhood
N	Number of cells in a neighborhood

	Average GLUT1 expression by cancer population
y	

Parameter	Units	Meaning	Value
q	dimensionless	Degree of resource sharing	0-1
R	1/time	Rate of resource influx into the neighborhood	100
a	1/(VEGF units)	Effectiveness of VEGF mediated resource redirection	0.05
c	1/(VEGF units × time)	Cost of VEGF expression	0.01
f	1/time	Constant cost of survival	5
m_G	dimensionless	Efficacy of GLUT1 inhibitor	0-20
m_A	dimensionless	Efficacy of VEGF inhibitor	0-20
σ_u^2	VEGF units	Evolvability of VEGF expression strategy	0.1
σ_y^2	GLUT1 units	Evolvability of GLUT1 expression strategy	0.1
h	1/(GLUT1 units × time)	Cost of GLUT1 expression	0.01
k	1/(GLUT1 units)	Effectiveness of GLUT1 mediated resource uptake	0.05
b	dimensionless	Conversion of net resource profit into fitness	0.5
d	dimensionless	Scaling factor for VEGF inhibitor	1
e	dimensionless	Scaling factor for GLUT1 inhibitor	1

225   Table 1: Summary of variables and their symbols used in the model.

226   Table 2: Summary of parameters and parameter values used in the model.

3. <u>Results</u>

We use an evolutionary game theoretic framework [59,64] to examine how cancer cells
allocate investment between VEGF and GLUT1 expression. We examine and compare
the team optimum and the evolutionarily stable strategy (ESS) [21,55]. The team
optimum is the strategy that maximizes the collective fitness, whereas the evolutionary
stable strategy (ESS) is the strategy that, when common among the cancer cells cannot
be invaded by rare alternative strategies. In our model, the ESS will be expected
outcome of natural selection.

3.1 Fixed Neighborhood

237 We begin by assuming that the number of cancer cells within a neighborhood (N) is
238 fixed, and analyze how VEGF and GLUT1 expression evolve across neighborhoods of
239 different sizes.

3.1.1 Team Optimum

For the team optimum, we determine the collective value of VEGF (u) and the individual value of GLUT1 (y) that maximizes fitness, G, for a certain neighborhood size (N). We start by setting $v = u$ in Eq. (1):

$$G(N, u, y) = b\left(\frac{R}{N}(1 - e^{-ky})(1 - e^{-aNu}) - cu - hy - f\right) \tag{6}$$

At the fitness-maximizing strategies, the partial derivatives of G with respect to each trait u and y must equal zero:

$$\frac{\partial G}{\partial u} = b(Ra(1 - e^{-ky})e^{-aNu} - c) \tag{7}$$

$$\frac{\partial G}{\partial y} = b\left(\frac{Rk(1 - e^{-aNu})e^{-ky}}{N} - h\right) \tag{8}$$

Yielding the following necessary conditions, respectively:

$$Ra(1 - e^{-ky})e^{-aNu} - c = 0 \tag{9}$$

$$\frac{Rk(1 - e^{-aNu})e^{-ky}}{N} - h = 0 \tag{10}$$

The overall team optimum (u_{team}, y_{team}) is obtained *numerically* by solving Eq. (9) and (10).

As we have noted before [21], the team optimum is independent of the resource-sharing parameter q. Also, the team optimum is independent of the rate at which net profit can be converted into survivorship and proliferation, b, and independent of the cell's fixed cost, f. We see how the two traits become co-adapted in that increasing GLUT1 expression increases the marginal value of VEGF production and vice-versa.

Fig. S1 shows how the team-optimum varies with model parameters in fixed neighborhoods. Both per cell VEGF and GLUT1 expression (u_{team} and y_{team}, respectively) increase with greater resource availability, R, and decrease with increasing neighborhood size, N, and higher expression costs, c and h. Interestingly, increasing the effectiveness of VEGF-mediated resource redirection, a, decreases the team-optimum VEGF expression while increasing GLUT1 expression, whereas the opposite pattern is observed when GLUT1-mediated uptake efficiency, k, increases. These trends arise because greater functional effectiveness reduces the level of investment required in that strategy to achieve the same benefit. Fig. S1B shows that VEGF expression at the team

Fig. 1: Effect of neighborhood size (N) on VEGF and GLUT1 expression at team optimum.

optimum is most strongly influenced by neighborhood size, N, followed by VEGF effectiveness, a, whereas GLUT1 expression is most sensitive to resource availability, R, neighborhood size, N, and GLUT1 efficiency, k.

Fig. 1 shows the temporal dynamics of per cell VEGF (u) and GLUT1 (y) expression as the system adapts towards the team optimum for five different neighborhood sizes, increasing from left to right. Since we are interested in relative changes in VEGF and GLUT1 expression in response to parameter variation, we chose parameters such that, at $N = 1$, VEGF and GLUT1 expressions are equal. Accordingly, we set the costs and efficacies of VEGF and GLUT1 to be the same, i.e., $c = h$ and $a = k$. Thus, in the absence of neighbors, the focal cell invests equally in VEGF and GLUT1. As neighborhood size increases, investment in both traits declines. This occurs because, as the number of cells in the neighborhood increases, VEGF concentration increases, reducing the marginal benefit of further investment in VEGF. At the same time, the amount of resources available per cell decreases, lowering the payoff from investing in GLUT1. Note that across all neighborhood sizes, the team optimum consistently favors higher investment in GLUT1 relative to VEGF. Although individual investment in VEGF and GLUT1 decreases with increasing neighborhood size, total VEGF expression always increases with neighborhood size (Fig. S2).

3.1.2 Evolutionary Stable Strategy

The ESS VEGF expression, u_{ESS}, occurs when $v = u_{ESS}$ maximizes an individual's fitness given the VEGF production of others. At the ESS, when all cells adopt the same strategy, no alternative strategy can invade. The first order necessary condition for ESS values of u and v occur when the selection gradients in v and y equal zero when evaluated at $v = u_{ESS}$ [78]:

Fig. 2: Evolution of VEGF and GLUT1 expression towards the ESS in two neighborhood sizes ($N = 2$ and $N = 10$) for different degrees of resource sharing (q).

$$\frac{\partial G}{\partial v}\bigg|_{v=u} = b\left(\frac{Ra}{N}(1 - e^{-ky})e^{-aNu} + R(1 - e^{-ky})(1 - q)\frac{(N-1)}{N^2}\frac{(1 - e^{-aNu})}{u} - c\right) \quad (11)$$
$$= 0$$

$$\frac{\partial G}{\partial y}\bigg|_{v=u} = b\left(\frac{Rk}{N}(1 - e^{-aNu})e^{-ky} - h\right) = 0 \quad (12)$$

Fig. 3: Dynamics of a cell's VEGF and GLUT1 expression towards the ESS for high ($q = 0.9$), intermediate ($q = 0.5$), and low ($q = 0.1$) resource sharing in neighborhoods of different sizes ($N = 1$ to 50).

For GLUT1, the ESS condition, Eq. (12), is identical to the team optimum condition, Eq. (10), and both are independent of q. This is because GLUT1 expression is purely density dependent and does not depend on the strategies of neighboring cells. For VEGF, Eq. (11) has more terms than the condition for the team optimum, Eq. (9). But when there is complete equal sharing ($q = 1$) or when the neighborhood size is just one ($N = 1$) then the conditions for the ESS become the same as the team optimum. Thus when $q = 1$ or $N = 1$: $u_{ESS} = u_{team}$ and $y_{ESS} = y_{team}$.

Consistent with this, Fig. S3A shows that increasing the degree of resource sharing leads to a decrease in VEGF expression at the ESS, while GLUT1 expression remains relatively unchanged. This pattern is also evident in Fig. 2 and S4: for a fixed neighborhood size, varying q has an imperceptible effect on GLUT1 expression, whereas increased resource sharing consistently reduces VEGF expression at the ESS.

Fig. S5 depicts the adaptive landscape of a focal cell introduced into neighborhoods of varying sizes (1-50). In each landscape, the red star marks the evolutionary stable strategy (ESS), and the black dots trace the trajectory of the population towards that point. When resource sharing is low, equilibrium VEGF expression is substantially higher than when sharing is high. However, in both regimes, increasing neighborhood size consistently reduces the equilibrium values of both VEGF and GLUT1. As neighborhood size (N) increases, the entire landscape progressively sinks toward zero

fitness and becomes increasingly flat. This indicates that large neighborhoods confer
little selective advantage to any single cell. In other words, when many cells share the
same resource pool, fitness differences across strategies vanish, as both VEGF and
GLUT1 production go to 0.

In Fig. 3 and S3 we examine the effect of neighborhood size on the ESS. We find that
with increasing neighborhood size, VEGF expression at the ESS initially increases. This
is because for smaller neighborhoods, cells have more to gain by co-opting resources
from their neighbors. This incentive vanishes as the neighborhood size increases. Once
there are too many cells in the neighborhood, there is little fitness to gain by producing
more VEGF. Thus, after a certain point, VEGF expression decreases with increasing
neighborhood size. GLUT1 expression decreases smoothly with increasing
neighborhood size (in all three resource-sharing environments, see Fig. S3B, C, D),
reflecting the actual harvestable resources available to a cell. As before, total VEGF and
GLUT1 expressions increase with neighborhood size (Fig. S6).

As neighborhood size increases from 1 to 2 cells, VEGF expression increases across all
three resource-sharing conditions (Fig. 3). This reflects the introduction of direct
competition, which leads to an "arms race" as cells invest more in VEGF to secure
resources [79]. With further increases in neighborhood size, VEGF expression declines,
as the benefits of additional investment in VEGF are shared among neighbors. The
neighborhood size at which per-cell VEGF expression (u_{ESS}) begins to decline with N
occurs at lower N under high resource sharing. Regardless u_{ESS} is substantially higher
under low resource sharing than under high resource sharing.

GLUT1 expression also decreases with increasing neighborhood size, although this
decline is more gradual than for VEGF. This reduction becomes more pronounced
under higher resource sharing. Overall, VEGF expression is greater in smaller
neighborhoods with low resource sharing, whereas GLUT1 expression dominates in
larger neighborhoods with high resource sharing. The interplay between neighborhood
size and resource sharing is illustrated in Fig. 3. In large neighborhoods ($N = 50$),
VEGF expression is more dominant under low resource sharing ($q = 0.1$), whereas
under high resource sharing ($q = 0.9$), GLUT1 expression becomes the strategy in
which cells invest more heavily. In contrast, in very small neighborhoods ($N = 2$), VEGF
expression remains higher even under high resource sharing conditions, reflecting the
strong incentive for cells to invest in VEGF in small neighborhoods. These patterns
reinforce that VEGF and GLUT1 are not independent strategies, but coadapted
responses to local cellular interactions and resource distribution.

To analyze how these dynamics change with model parameters, we performed
parameter sweeps (Fig. S3) and sensitivity analyses (Fig. S7). Both VEGF and GLUT1
expression increase with resource abundance, R. Over the range of values considered,

variation in VEGF efficacy, a, has minimal effects on the ESS, whereas GLUT1 efficacy,
k, strongly influences system behavior. Increasing k leads to reduced GLUT1
expression at the ESS, accompanied by increased VEGF expression. Increasing the
costs of VEGF, c, or GLUT1, h, reduces expression of both traits; these reductions are
more pronounced under high resource-sharing conditions.

Sensitivity analysis further reveals that GLUT1 expression at the ESS is most sensitive
to changes in the efficacy of transporters, k, whereas VEGF expression is most
sensitive to neighborhood size, N, resource abundance, R, and the cost of VEGF, c,
particularly under low resource sharing. In high resource-sharing conditions, VEGF
expression is most strongly influenced by the degree of sharing, q.

3.2 Integrating Ecological and Evolutionary Dynamics

In the previous section, the number of cells in a neighborhood (N) was fixed. Here, we
let the number of cells in the number increase or decrease based on whether fitness is
positive ($G > 0$) or negative ($G < 0$). The ESS will now include VEGF and GLUT1
expression and an equilibrium neighborhood population size, N_{ESS}. The value u_{ESS}, y_{ESS}
and N_{ESS} now represent eco-evolutionary dynamics.

We can visualize the eco-evolutionary dynamics on adaptive landscapes (Fig. S8).
Each graph is a snapshot in time in the trajectory of the population towards eco-
evolutionary stability. The condition for eco-evolutionary stability is attained when Eq.
(2), (3), (4) are all satisfied. The black dots show the trajectory and the red star is the
ESS strategy itself. As in the fixed-neighborhood case, higher resource sharing leads to
lower investment in both VEGF and GLUT1 at the ESS, and to a higher equilibrium
population size.

Fig. 4A shows that population size, N_{ESS}, increases monotonically with the degree of
resource sharing (q), while VEGF and GLUT1 expression, u_{ESS} and y_{ESS}, are inversely
related to q (also see Fig. S9). Higher resource sharing reduces the escalating VEGF-
expression arms race that occurs at low resource sharing as cancer cells attempt to
secure a higher share of the bloodborne resources. Figure 4B further illustrates how
q determines the relative balance between VEGF and GLUT1 expression within a
neighborhood. Under low resource-sharing conditions, the ratio of VEGF to GLUT1
expression is high, reflecting the strong incentive for individual cells to invest in VEGF-
mediated signaling. In contrast, high resource-sharing environments reduce this ratio,
favoring relatively greater investment in GLUT1 expression (also see Fig. S10).

Fig. 4: A) Population Size (N), VEGF Expression (u), and GLUT1 Expression (y) dynamics for different values of q. Darker the shade, higher the resource sharing (q). B) Comparison of ESS dynamics in low ($q = 0.1$), intermediate ($q = 0.5$), and high ($q = 0.9$) resource sharing neighborhoods.

To characterize how ecological and evolutionary interactions shape neighborhood dynamics, we examine how key model parameters influence both the ESS and team optimum through parameter sweeps and sensitivity analyses (Fig. S11-S14). Across all regimes, increasing resource abundance, R, increases population size. At the ESS, VEGF and GLUT1 expression increase with R under low and intermediate resource sharing, whereas under high resource sharing, both traits exhibit non-monotonic behavior, initially decreasing before increasing at higher resource levels. In contrast, at the team optimum, increasing R leads to monotonic reductions in VEGF and GLUT1 expression, reflecting reduced need for costly resource acquisition.

Increasing the efficacy of GLUT1-mediated uptake, k, increases population size while reducing both VEGF and GLUT1 expression at the ESS, reflecting more efficient resource acquisition and utilization. This effect is consistent across low, intermediate, and high resource-sharing conditions, as well as at the team optimum. In contrast, variation in VEGF efficacy, a, has little effect on the ESS across all resource-sharing regimes. But at the team optimum, increasing a increases population size while reducing investment in both VEGF and GLUT1 expression.

The effect of increasing the cost of VEGF, c, depends on the degree of resource sharing. Under low resource sharing, VEGF expression decreases sharply with increasing c, with little effect on the population size or GLUT1 expression. At intermediate resource sharing, both VEGF and GLUT1 expression decline with increasing c, while population size remains nearly constant. Under high resource sharing, increasing c reduces both population size and VEGF expression, with only little change in GLUT1. At the team optimum, increasing c reduces population size and VEGF expression while increasing GLUT1 expression, consistent with compensatory allocation toward GLUT1.

Increasing the cost of GLUT1 expression, h, has a consistent effect across all resource-sharing regimes: population size and GLUT1 expression decline, while VEGF expression increases, indicating a compensatory shift away from costly GLUT1 investment and toward VEGF. The same qualitative pattern is also observed at the team optimum. Finally, increasing the cost of survival, f, reduces population size while increasing investment in both VEGF and GLUT1 across all resource sharing values and at the team optimum.

Normalized sensitivity analysis (Fig. S15) identifies the dominant drivers of system behavior. At the team optimum, population size is most strongly influenced by resource abundance, R, and GLUT1 efficacy, k. VEGF expression is primarily governed by the cost of GLUT1 expression, h. At the ESS, the sensitivities depend on the degree of resource sharing. Under low and intermediate sharing, VEGF expression is most sensitive to VEGF cost, c, whereas GLUT1 expression is primarily driven by GLUT1 efficacy, k. In high resource sharing, VEGF expression becomes overwhelmingly sensitive to the degree of sharing, q, and other parameters become less impactful.

3.3 The Effect of Therapy

In this section, we examine the effects of two potential therapeutic strategies: one targeting VEGF signaling and one targeting GLUT1-mediated glucose uptake [67,72,80]. The VEGF-targeting therapy, m_A, acts by reducing the efficacy of VEGF-mediated resource redirection, a, while the GLUT1-targeting therapy, m_G, reduces the efficacy of glucose uptake, k. We expect either therapy to induce predictable coadapted phenotypic responses in both VEGF and GLUT1 expression. As in Sections 3.1 and 3.2, we analyze both the team optimum and evolutionarily stable strategy (ESS) under these interventions. Details of the derivations are provided in the Supplementary material.

Considering the both therapies, m_A and m_G, the necessary conditions for the team optimum VEGF and GLUT1 expression, u_{team} and y_{team}, respectively are

(13)

Fig. 5: Effect of GLUT1 inhibitor (top) and anti-angiogenic therapy (bottom) on population size (N), VEGF expression (u), GLUT1 expression (y) for different degrees of resource sharing (q).

For the top row: $m_G = 10$ and $m_A = 0$. For the bottom row, $m_G = 0$ and $m_A = 10$. Therapy is administered at $t = 10000$ (dashed line).

$$b\left(\frac{Ra\left(1 - e^{-\frac{ky}{1+em_G}}\right)e^{-\frac{aNu}{1+dm_A}}}{1 + dm_A} - c\right) = 0$$

$$b\left(\frac{Rk\left(1 - e^{-\frac{aNu}{1+dm_A}}\right)e^{-\frac{ky}{1+em_G}}}{N(1 + em_G)} - h\right) = 0$$

441  Under therapy, the ESS conditions for VEGF and GLUT1, u_{ESS} and y_{ESS} are:

$$b\left(\frac{Ra}{N}\left(1-e^{-\frac{ky}{1+em_G}}\right)e^{-\frac{aNu}{1+dm_A}}+R\left(1-e^{-\frac{ky}{1+em_G}}\right)(1-q)\frac{(N-1)}{N^2}\frac{\left(1-e^{-\frac{aNu}{1+dm_A}}\right)}{u}-c\right)$$
$$=0 \tag{15}$$

$$\tag{16}$$

$$b\left(\frac{Rk}{N}\left(1-e^{-\frac{aNu}{1+dm_A}}\right)e^{-\frac{ky}{1+em_G}}-h\right)=0$$

443   For either the team optimum or the ESS, the condition for the neighborhood's
444   equilibrium population size, N_{team} and N_{ESS}, respectively, is:

$$G(N,u,v,y)=b\left(\left(1-e^{-\frac{ky}{1+em_G}}\right)\left(\frac{R}{N}\right)\left(1-e^{-\frac{aNu}{1+dm_A}}\right)-cu-hy-f\right)=0 \tag{17}$$

Eco-evolutionary stability is attained when Eq. (15), (16), and (17) are satisfied.

As shown in Fig. 5, inhibition of GLUT1 consistently reduces tumor size more effectively
than anti-angiogenic therapy across all degrees of resource sharing. In our framework,
this happen because GLUT1-mediated uptake operates at cellular level and is largely
independent of the degree of resource sharing (q). In contrast, the effectiveness of anti-
angiogenic therapy is heavily influenced by q. VEGF inhibition is most effective under
high resource-sharing conditions ($q=0.9$) and has minimal impact under low resource-
sharing regimes. While one might expect VEGF inhibitors to be more effective in tumors
with higher VEGF expression, our results suggest that the tragedy of the commons that
emerges under low resource sharing buffers the impact of anti-angiogenic therapy. This
could mean that therapies targeting shared paracrine factors may be less effective than
those targeting cell-autonomous mechanisms such as glucose uptake.

We further examine these differences in Fig. 6, which shows the effect of varying
therapy doses across low, intermediate, and high resource-sharing environments. As
expected, increasing the dose generally reduces tumor size. However, anti-angiogenic
therapy remains largely ineffective in low and intermediate resource-sharing regimes,
whereas both therapies are more effective under high resource sharing. These results
underscore that therapeutic response is shaped not only by drug potency, but also by
the interactions between cells in a tumor.

Fig. S16 summarizes the percentage decrease in population size following
monotherapy across the three resource-sharing regimes. Fig. 7 examines the combined
effect of VEGF

Fig. 6: Effect of different doses of (A) GLUT1 inhibitor (m_G) and VEGF inhibitor (m_A) (B) on population size (N), VEGF expression (u), GLUT1 expression (y) at ESS. For the top row: $q = 0.1$ (low resource sharing), middle row: $q = 0.5$ (intermediate resource sharing), bottom row: $q = 0.9$ (high resource sharing). Therapy is administered at $t = 10000$ (dashed line).

Fig. 7: Effect of combination therapy with GLUT1 inhibitor (m_G) and VEGF inhibitor (m_A) on population size (N), VEGF expression (u), GLUT1 expression (y) at ESS. For the top row: $q = 0.1$ (low resource sharing), middle row: $q = 0.5$ (intermediate resource sharing), bottom row: $q = 0.9$ (high resource sharing).

and GLUT1 inhibition. Relatively low doses of GLUT1 inhibition ($m_G = 5$) achieve reductions in tumor size that are not attainable even with high doses of VEGF inhibition ($m_A = 50$) alone. Moreover, combining moderate anti-angiogenic therapy with lower doses of GLUT1 inhibition yields comparable reductions in tumor size. These results suggest that combination strategies can achieve substantial therapeutic benefit at reduced dosing, which may be particularly important given the systemic constraints associated with targeting glucose uptake.

Our analysis suggests that therapeutic response is heavily influenced by the degree of resource sharing within the tumor microenvironment. The degree of resource sharing is

an empirical question. This parameter may vary by disease type, patient to patient, or
even among different regions of the same tumor, leading to spatial variation in the
evolutionary incentives governing VEGF and GLUT1 co-expression. Consequently, the
effectiveness of a given therapy may differ across tumor neighborhoods, even within a
single lesion. By framing VEGF and GLUT1 expression as coadaptations shaped by
neighborhood structure, our model provides a foundation for designing therapies that
anticipate evolutionary responses rather than react to them [81,82].

4. <u>Discussion</u>

In this study, we examined the coadaptation [83,84] of VEGF-mediated angiogenesis
and GLUT1-mediated glucose uptake as complementary foraging strategies in cancer
cells [56]. Both traits are downstream effectors of HIF-1, which is activated in response
to resource deprivation and hypoxia, linking the two coadaptations with a shared
regulatory program [9–11,24,26]. Clinical and translational studies have consistently
documented coordinated expression of hypoxia-related markers, including VEGF,
GLUT1, CA-IX, and HIF-1α, across multiple malignancies [29,31]. In cancer, both VEGF
and GLUT1 are frequently overexpressed relative to normal tissue.
Immunohistochemical analyses in cervical, ovarian, gastric, and bladder cancers
demonstrate frequent co-expression and spatial co-localization of these markers, often
correlating with tumor progression, microvessel density, and poor prognosis
[25,27,28,85–87]. Although these studies do not explicitly address evolutionary
dynamics, they provide strong biological evidence that angiogenic and metabolic
responses are activated together in human tumors.

A substantial mathematical oncology literature has modeled angiogenesis and tumor
metabolism, but typically in isolation. Mechanistic models of angiogenesis have focused
on VEGF dynamics, endothelial sprouting, vascular remodeling, and nutrient transport
using reaction–diffusion, continuum, and multiscale frameworks to predict tumor growth
and vascular architecture [34,35,88]. Separately, models of tumor metabolism have
examined glucose uptake, glycolytic flux, and metabolic competition, and recent
evolutionary game-theoretic work has treated GLUT1 expression as a tragedy of the
commons [41,44–46,55]. Our framework explicitly models VEGF and GLUT1 as
coadapted traits whose joint evolutionary dynamics are shaped by neighborhood
interactions via resource supply and degree of resource sharing.

VEGF influences resource availability at the neighborhood scale, whereas GLUT1
governs resource uptake at the cellular scale. Modeling these traits jointly reveals that
their evolutionary dynamics are coupled and shaped by cell–cell interactions,
specifically the degree of resource sharing. Here, resource sharing refers to how
resources supplied by blood vessels are distributed among cells within a local habitat.

We show that the degree of resource sharing within tumor neighborhoods is a key
determinant of strategy selection [21]. Under low-resource sharing, cells are
incentivized to overinvest in VEGF, leading to a tragedy of the commons that reduces
collective tumor fitness [54,55]. This mirrors ecological systems in which shared
resources are overexploited when access is unequal [89–93]. In contrast, when
resources are more equitably distributed, the selective pressure to overproduce VEGF
diminishes, and cellular investment shifts toward uptake efficiency through GLUT1
expression. Thus, resource sharing dynamics influences the dominant foraging strategy
in a neighborhood.

GLUT1 reflects a cell's capacity for resource uptake, whereas VEGF reflects the 'effort'
per resource recruited into the neighborhood. The ratio of VEGF to GLUT1 expression,
thus, provides a proxy for how much effort is expended on resource recruitment relative
to uptake. A high VEGF-to-GLUT1 ratio suggests environments where cells must invest
heavily in recruiting resources that are not readily shared, consistent with low resource-
sharing conditions. Conversely, a lower ratio indicates that resources are more
accessible, allowing cells to prioritize uptake over blood vessel recruitment.

Different tumor regions are likely to exhibit distinct degrees of resource sharing
depending on the resource supply, vascular density, hypoxia, and cell packing [94–96].
Consequently, some neighborhoods may favor VEGF and others GLUT1 expression. To
empirically estimate resource-sharing dynamics in cancer, multiplex imaging platforms
such as immunofluorescence and spatial transcriptomics can be used to simultaneously
quantify multiple tumor markers at single-cell resolution while preserving spatial context
[97,98]. This spatial information enables the identification of local neighborhoods, within
which marker intensities can be quantified for each cell. From these measurements, the
VEGF-to-GLUT1 ratio can be computed at the neighborhood level and used as a proxy
for resource-sharing dynamics. Model predictions can be tested by examining how this
ratio varies with features such as vascular density and local cell packing. This spatial
heterogeneity suggests that therapeutic responses may vary not only between tumors in
different patients but also across regions of a single lesion.

Our simulations suggest that targeting glucose uptake exerts a stronger suppressive
effect on tumor size than anti-angiogenic therapy across different degrees of resource
sharing. Clinically, anti-angiogenic agents such as bevacizumab and other VEGF
pathway inhibitors are widely used, yet their benefits are often transient, with tumors
frequently developing resistance [99–102]. Several studies have reported that
resistance to VEGF-targeted therapies is accompanied by enhanced GLUT1 expression
and metabolic reprogramming toward glycolysis [94,103], suggesting a compensatory
shift from vascular recruitment to cellular resource acquisition.

In contrast, systemic GLUT1 inhibition is currently not a viable strategy due to the
essential role of glucose transport in normal tissues [6,104,105]. Several GLUT
inhibitors have been explored in preclinical stages, including small molecules such as
WZB117, BAY-876, and STF-31 [70,71,73], which inhibit GLUT1-mediated glucose
transport and have demonstrated anti-tumor effects in cell lines and xenograft models.
Metabolic analogues such as 2-deoxy-D-glucose (2-DG) have also been explored as
glycolytic inhibitors [106], though systemic toxicity and limited efficacy have constrained
clinical translation. The widespread expression of GLUT1 in erythrocytes and the blood–
brain barrier raises significant concerns about on-target toxicity, limiting the therapeutic
window of direct GLUT1 blockade [6,107].

Emerging precision-delivery approaches offer potential strategies to overcome these
limitations by selectively targeting tumor metabolism. Recent preclinical studies have
demonstrated tumor-biased delivery of GLUT1-targeting payloads, including Glut1
siRNA delivered via engineered nanoparticles [108,109], as well as nanoformulations
carrying GLUT1 inhibitors such as BAY-876 [110,111]. More broadly, advances in
nanomedicine and targeted delivery systems have enabled selective modulation of
tumor metabolic pathways while sparing normal tissues [112]. If such tumor-specific
targeting can be achieved, our model predicts that GLUT1 inhibition would be highly
effective across a wide range of resource-sharing conditions.

An alternative therapeutic approach is to reduce systemic glucose availability rather
than inhibiting glucose transport. Dietary interventions such as ketogenic diets, which
lower circulating glucose and shift metabolism toward ketone utilization, have shown
potential in preclinical models and early clinical studies to impair tumor growth and
enhance response to therapy, although their efficacy remains context-dependent and
requires further validation [113–115]. Our previous modeling work [21] demonstrated
that reducing resource supply is more effective at limiting tumor growth than reducing
the efficacy of VEGF-mediated resource recruitment. Potentially, combining reduced
resource inflow with inhibition of GLUT1-mediated uptake could be an even more
effective therapeutic approach.

Several limitations must be addressed. Our framework is not spatially explicit. Although
we consider resource distribution between cells through the parameter q, we do not
explicitly represent spatial diffusion of oxygen or glucose, or vascular topology. In our
model, q is treated as a fixed parameter; a natural extension would allow resource
sharing itself to evolve or vary dynamically over time.

In addition, we assume that both VEGF and GLUT1 expression incur fitness costs.
These costs are biologically plausible. VEGF production requires sustained biosynthetic
investment and contributes to structurally abnormal vasculature that may reduce
perfusion efficiency and increase metabolic stress. Elevated GLUT1 expression is

associated with high glycolytic flux, increased metabolic demand, and altered redox balance, imposing energetic constraints on the cell [57,116–119]. While these costs are represented abstractly in our framework, experimental quantification of the metabolic and physiological trade-offs associated with VEGF and GLUT1 expression would further strengthen evolutionary models of tumor resource acquisition.

In summary, our study frames angiogenesis and glucose uptake as coadapted strategies shaped by ecological interactions within the tumor microenvironment. By modeling VEGF and GLUT1 as coadaptations, we demonstrate that the balance between these two foraging strategies depends on resource sharing and neighborhood structure. These results highlight that tumor behavior and therapeutic response cannot be fully understood without considering how adapted traits interact with each other. More broadly, our findings underscore the importance of integrating ecological structure and evolutionary dynamics into cancer modeling [81]. Viewing tumors as systems of interacting, coevolving strategies may provide a foundation for designing therapies that anticipate evolutionary responses rather than inadvertently selecting for them.

5. Conflicts of Interest

The authors declare no conflicts of interest.

6. Funding

We appreciate grant support from NIH/NCI 1R01CA258089 and 1U54 CA274507-01A1, and DoD W81XWH2210680.

7. Author Contributions

RB and JB conceptualized the model and study. RB created the simulations and wrote the manuscript. RB, JB, and RG reviewed and edited the manuscript.

8. Acknowledgments

The authors thank Anuraag Bukkuri for his valuable discussions and input in the design of the model.

9. Declaration of generative AI and AI-assisted technologies in the writing process

The authors used Grammarly and ChatGPT for copyediting the manuscript. After using the tools, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

10. Data Availability

Codes used to generate simulations in the paper are available at
https://github.com/rbhattacharya49/vegf-glut1-coadaptation

11. References

[1] Sullivan MR, Heiden MGV. Determinants of nutrient limitation in cancer. Crit Rev Biochem Mol Biol 2019;54:193–207. https://doi.org/10.1080/10409238.2019.1611733.

[2] García-Jiménez C, Goding CR. Starvation and pseudo-starvation as drivers of cancer metastasis through translation reprogramming. Cell Metab 2019;29:254–67. https://doi.org/10.1016/j.cmet.2018.11.018.

[3] Ferrara N, Gerber H-P, LeCouter J. The biology of VEGF and its receptors. Nat Med 2003;9:669–76. https://doi.org/10.1038/nm0603-669.

[4] Shibuya M. Vascular Endothelial Growth Factor (VEGF) and Its Receptor (VEGFR) Signaling in Angiogenesis. Genes Cancer 2011;2:1097–105. https://doi.org/10.1177/1947601911423031.

[5] Medina RA, Owen GI. Glucose transporters: expression, regulation and cancer. Biol Res 2002;35:9–26. https://doi.org/10.4067/s0716-97602002000100004.

[6] Mueckler M, Thorens B. The SLC2 (GLUT) family of membrane transporters. Mol Aspects Med 2013;34:121–38. https://doi.org/10.1016/j.mam.2012.07.001.

[7] Wallace B. Coadaptation revisited. J Hered 1991;82:89–95. https://doi.org/10.1093/oxfordjournals.jhered.a111061.

[8] Pazos F, Valencia A. Protein co-evolution, co-adaptation and interactions. EMBO J 2008;27:2648–55. https://doi.org/10.1038/emboj.2008.189.

[9] Semenza GL. Targeting HIF-1 for cancer therapy. Nat Rev Cancer 2003;3:721–32. https://doi.org/10.1038/nrc1187.

[10] Lee J-W, Bae S-H, Jeong J-W, Kim S-H, Kim K-W. Hypoxia-inducible factor (HIF-1)α: its protein stability and biological functions. Exp Mol Med 2004;36:1–12. https://doi.org/10.1038/emm.2004.1.

[11] Yang Y, Sun M, Wang L, Jiao B. HIFs, angiogenesis, and cancer. J Cell Biochem 2013;114:967–74. https://doi.org/10.1002/jcb.24438.

[12] Carmeliet P. VEGF as a key mediator of angiogenesis in cancer. Oncology 2005;69 Suppl 3:4–10. https://doi.org/10.1159/000088478.

[13] Perrot-Applanat M, Di Benedetto M. Autocrine functions of VEGF in breast tumor cells. Cell Adhes Migr 2012;6:547–53. https://doi.org/10.4161/cam.23332.

[14] Goggins E, Mironchik Y, Kakkad S, Jacob D, Wildes F, Bhujwalla ZM, et al. Reprogramming of VEGF-mediated extracellular matrix changes through autocrine signaling. Cancer Biol Ther 2023;24:2184145. https://doi.org/10.1080/15384047.2023.2184145.

[15] Adekola K, Rosen ST, Shanmugam M. Glucose transporters in cancer metabolism. Curr Opin Oncol 2012;24:650–4. https://doi.org/10.1097/CCO.0b013e328356da72.

[16] Ancey P-B, Contat C, Meylan E. Glucose transporters in cancer – from tumor cells to the tumor microenvironment. FEBS J 2018;285:2926–43. https://doi.org/10.1111/febs.14577.

[17] Castell F, Cook GJR. Quantitative techniques in 18FDG PET scanning in oncology. Br J Cancer 2008;98:1597–601. https://doi.org/10.1038/sj.bjc.6604330.

[18] Meyer H-J, Wienke A, Surov A. Associations between GLUT expression and SUV
values derived from FDG-PET in different tumors—A systematic review and meta
analysis. PLoS ONE 2019;14:e0217781.
https://doi.org/10.1371/journal.pone.0217781.
[19] Amann T, Maegdefrau U, Hartmann A, Agaimy A, Marienhagen J, Weiss TS, et al.
GLUT1 Expression Is Increased in Hepatocellular Carcinoma and Promotes
Tumorigenesis. Am J Pathol 2009;174:1544–52.
https://doi.org/10.2353/ajpath.2009.080596.
[20] Younes M, Brown RW, Stephenson M, Gondo M, Cagle PT. Overexpression of
Glut1 and Glut3 in stage I nonsmall cell lung carcinoma is associated with poor
survival. Cancer 1997;80:1046–51. https://doi.org/10.1002/(sici)1097-
0142(19970915)80:6%253C1046::aid-cncr6%253E3.0.co;2-7.
[21] Bhattacharya R, Bukkuri A, Gatenby RA, Brown JS. Tumor angiogenesis may be a
tragedy of the commons. SSRN [Preprint]. 2026.
http://dx.doi.org/10.2139/ssrn.5441855
[22] Rumsey SC, Kwon O, Xu GW, Burant CF, Simpson I, Levine M. Glucose
transporter isoforms GLUT1 and GLUT3 transport dehydroascorbic acid. J Biol
Chem 1997;272:18982–9. https://doi.org/10.1074/jbc.272.30.18982.
[23] Fendt S-M, Frezza C, Erez A. Targeting metabolic plasticity and flexibility dynamics
for cancer therapy. Cancer Discov 2020;10:1797–807.
https://doi.org/10.1158/2159-8290.CD-20-0844.
[24] Song I-S, Wang A-G, Yoon SY, Kim J-M, Kim JH, Lee D-S, et al. Regulation of
glucose metabolism-related genes and VEGF by HIF-1α and HIF-1β, but not HIF-
2α, in gastric cancer. Exp Mol Med 2009;41:51–8.
https://doi.org/10.3858/emm.2009.41.1.007.
[25] Des Guetz G, Uzzan B, Nicolas P, Cucherat M, Morere J-F, Benamouzig R, et al.
Microvessel density and VEGF expression are prognostic factors in colorectal
cancer. Meta-analysis of the literature. Br J Cancer 2006;94:1823–32.
https://doi.org/10.1038/sj.bjc.6603176.
[26] Melstrom LG, Salabat MR, Ding X-Z, Strouch MJ, Grippo PJ, Mirzoeva S, et al.
Apigenin down-regulates the hypoxia response genes: HIF-1α, GLUT-1, and VEGF
in human pancreatic cancer cells. J Surg Res 2011;167:173–81.
https://doi.org/10.1016/j.jss.2010.10.041.
[27] Semaan A, Munkarah AR, Arabi H, Bandyopadhyay S, Seward S, Kumar S, et al.
Expression of GLUT-1 in epithelial ovarian carcinoma: correlation with tumor cell
proliferation, angiogenesis, survival and ability to predict optimal cytoreduction.
Gynecol Oncol 2011;121:181–6. https://doi.org/10.1016/j.ygyno.2010.11.019.
[28] Kobayashi M, Kaida H, Kawahara A, Hattori S, Kurata S, Hayakawa M, et al. The
relationship between GLUT-1 and vascular endothelial growth factor expression
and 18F-FDG uptake in esophageal squamous cell cancer patients. Clin Nucl Med
2012;37:447–52. https://doi.org/10.1097/RLU.0b013e31823924bb.
[29] De Schutter H, Landuyt W, Verbeken E, Goethals L, Hermans R, Nuyts S. The
prognostic value of the hypoxia markers CA IX and GLUT 1 and the cytokines
VEGF and IL 6 in head and neck squamous cell carcinoma treated by radiotherapy
+/- chemotherapy. BMC Cancer 2005;5:42. https://doi.org/10.1186/1471-2407-5-42.

[30] Qiang B, Kang Y, Yang J, Su H, Wang Z, Zhang C, et al. The prognostic value of
vascular endothelial growth factor in endometrial cancer: A protocol for systematic
review and meta-analysis. Medicine (Baltimore) 2024;103:e40933.
https://doi.org/10.1097/MD.0000000000040933.
[31] Kim K, Park WY, Kim JY, Sol MY, Shin DH, Park DY, et al. Prognostic Relevance of
the Expression of CA IX, GLUT-1, and VEGF in Ovarian Epithelial Cancers. Korean
J Pathol 2012;46:532–40. https://doi.org/10.4132/KoreanJPathol.2012.46.6.532.
[32] Minami K, Saito Y, Imamura H, Okamura A. Prognostic significance of p53, Ki-67,
VEGF and Glut-1 in resected stage I adenocarcinoma of the lung. Lung Cancer
2002;38:51–7. https://doi.org/10.1016/S0169-5002(02)00108-3.
[33] Anderson AR, Chaplain MA. Continuous and discrete mathematical models of
tumor-induced angiogenesis. Bull Math Biol 1998;60:857–99.
https://doi.org/10.1006/bulm.1998.0042.
[34] Chaplain M, Anderson A. Mathematical modelling of tumour-induced angiogenesis:
network growth and structure. Cancer Treat Res 2004;117:51–75.
https://doi.org/10.1007/978-1-4419-8871-3_3.
[35] Byrne HM, Chaplain MA. Mathematical models for tumour angiogenesis: numerical
simulations and nonlinear wave solutions. Bull Math Biol 1995;57:461–86.
https://doi.org/10.1007/BF02460635.
[36] Owen MR, Alarcón T, Maini PK, Byrne HM. Angiogenesis and vascular remodelling
in normal and cancerous tissues. J Math Biol 2009;58:689–721.
https://doi.org/10.1007/s00285-008-0213-z.
[37] Finley SD, Dhar M, Popel AS. Compartment Model Predicts VEGF Secretion and
Investigates the Effects of VEGF Trap in Tumor-Bearing Mice. Front Oncol 2013;3.
https://doi.org/10.3389/fonc.2013.00196.
[38] Finley SD, Engel-Stefanini MO, Imoukhuede PI, Popel AS. Pharmacokinetics and
pharmacodynamics of VEGF-neutralizing antibodies. BMC Syst Biol 2011;5:193.
https://doi.org/10.1186/1752-0509-5-193.
[39] Phillips CM, Lima EABF, Woodall RT, Brock A, Yankeelov TE. A hybrid model of
tumor growth and angiogenesis: In silico experiments. PLOS ONE
2020;15:e0231137. https://doi.org/10.1371/journal.pone.0231137.
[40] Nikmaneshi MR, Firoozabadi B, Mozafari A, Munn LL. A multi-scale model for
determining the effects of pathophysiology and metabolic disorders on tumor
growth. Sci Rep 2020;10:3025. https://doi.org/10.1038/s41598-020-59658-0.
[41] Sartakhti JS, Manshaei MH, Basanta D, Sadeghi M. Evolutionary emergence of
angiogenesis in avascular tumors using a spatial public goods game. PLOS ONE
2017;12:e0175063. https://doi.org/10.1371/journal.pone.0175063.
[42] Archetti M, Ferraro DA, Christofori G. Heterogeneity for IGF-II production
maintained by public goods dynamics in neuroendocrine pancreatic cancer. Proc
Natl Acad Sci U S A 2015;112:1833–8. https://doi.org/10.1073/pnas.1414653112.
[43] Greenspan HP. Models for the Growth of a Solid Tumor by Diffusion. Stud Appl
Math 1972;51:317–40. https://doi.org/10.1002/sapm1972514317.
[44] Casciari JJ, Sotirchos SV, Sutherland RM. Variations in tumor cell growth rates and
metabolism with oxygen concentration, glucose concentration, and extracellular
pH. J Cell Physiol 1992;151:386–94. https://doi.org/10.1002/jcp.1041510220.

[45] Ward JP, King JR. Mathematical modelling of avascular-tumour growth. IMA J Math
Appl Med Biol 1997;14:39–69. https://doi.org/10.1093/imammb/14.1.39.
[46] Gatenby RA, Smallbone K, Maini PK, Rose F, Averill J, Nagle RB, et al. Cellular
adaptations to hypoxia and acidosis during somatic evolution of breast cancer. Br J
Cancer 2007;97:646–53. https://doi.org/10.1038/sj.bjc.6603922.
[47] Gatenby RA, Gillies RJ. Why do cancers have high aerobic glycolysis? Nat Rev
Cancer 2004;4:891–9. https://doi.org/10.1038/nrc1478.
[48] Epstein T, Gatenby RA, Brown JS. The Warburg effect as an adaptation of cancer
cells to rapid fluctuations in energy demand. PLOS ONE 2017;12:e0185085.
https://doi.org/10.1371/journal.pone.0185085.
[49] Shamsi M, Saghafian M, Dejam M, Sanati-Nezhad A. Mathematical Modeling of the
Function of Warburg Effect in Tumor Microenvironment. Sci Rep 2018;8:8903.
https://doi.org/10.1038/s41598-018-27303-6.
[50] Kareva I, Berezovskaya F. Cancer immunoediting: A process driven by metabolic
competition as a predator-prey-shared resource type model. J Theor Biol
2015;380:463–72. https://doi.org/10.1016/j.jtbi.2015.06.007.
[51] Williams S-P, Flores-Mercado JE, Port RE, Bengtsson T. Quantitation of glucose
uptake in tumors by dynamic FDG-PET has less glucose bias and lower variability
when adjusted for partial saturation of glucose transport. EJNMMI Res 2012;2:6.
https://doi.org/10.1186/2191-219X-2-6.
[52] Kashkooli FM, Abazari MA, Soltani M, Ghazani MA, Rahmim A. A spatiotemporal
multi-scale computational model for FDG PET imaging at different stages of tumor
growth and angiogenesis. Sci Rep 2022;12:10062. https://doi.org/10.1038/s41598-
022-13345-4.
[53] Ampatzoglou P, Kariotou F, Hadjinicolaou M. A Multiscale Mathematical Model for
Tumor Growth, Incorporating the GLUT1 Expression. Adv Exp Med Biol
2021;1338:273–81. https://doi.org/10.1007/978-3-030-78775-2_32.
[54] Hardin G. The Tragedy of the Commons. Sci New Ser 1968;162:1243–8.
[55] Bukkuri A, Gatenby RA, Brown JS. GLUT1 production in cancer cells: a tragedy of
the commons. Npj Syst Biol Appl 2022;8:1–13. https://doi.org/10.1038/s41540-022-
00229-6.
[56] Amend SR, Gatenby RA, Pienta KJ, Brown JS. Cancer Foraging Ecology: Diet
Choice, Patch Use, and Habitat Selection of Cancer Cells. Curr Pathobiol Rep
2018;6:209–18. https://doi.org/10.1007/s40139-018-0185-7.
[57] Zhang H, Wang Y, Li J, Chen H, He X, Zhang H, et al. Biosynthetic energy cost for
amino acids decreases in cancer evolution. Nat Commun 2018;9:4124.
https://doi.org/10.1038/s41467-018-06461-1.
[58] Pfeiffer T, Schuster S, Bonhoeffer S. Cooperation and competition in the evolution
of ATP-producing pathways. Science 2001;292:504–7.
https://doi.org/10.1126/science.1058079.
[59] Smith JM. Evolution and the Theory of Games. Cambridge: Cambridge University
Press; 1982. https://doi.org/10.1017/CBO9780511806292.
[60] Smith JM. Evolutionary game theory. Phys Nonlinear Phenom 1986;22:43–9.
https://doi.org/10.1016/0167-2789(86)90232-0.

[61] Kimmel GJ, Gerlee P, Brown JS, Altrock PM. Neighborhood size-effects shape
growing population dynamics in evolutionary public goods games. Commun Biol
2019;2:1–10. https://doi.org/10.1038/s42003-019-0299-4.
[62] Willer R. Groups Reward Individual Sacrifice: The Status Solution to the Collective
Action Problem. Am Sociol Rev 2009;74:23–43.
https://doi.org/10.1177/000312240907400102.
[63] Bukkuri A, Brown JS. Evolutionary Game Theory: Darwinian Dynamics and the G
Function Approach. Games 2021;12:72. https://doi.org/10.3390/g12040072.
[64] Vincent TL, Brown JS. Evolutionary Game Theory, Natural Selection, and
Darwinian Dynamics. Cambridge: Cambridge University Press; 2005.
https://doi.org/10.1017/CBO9780511542633.
[65] Brown JS. Why Darwin would have loved evolutionary game theory. Proc R Soc B
Biol Sci 2016;283:20160847. https://doi.org/10.1098/rspb.2016.0847.
[66] Tong M, Han BB, Holpuch AS, Pei P, He L, Mallery SR. Inherent phenotypic
plasticity facilitates progression of head and neck cancer: endotheliod
characteristics enable angiogenesis and invasion. Exp Cell Res 2013;319:1028–
42. https://doi.org/10.1016/j.yexcr.2013.01.013.
[67] Cook KM, Figg WD. Angiogenesis Inhibitors – Current Strategies and Future
Prospects. CA Cancer J Clin 2010;60:222–43. https://doi.org/10.3322/caac.20075.
[68] Jain RK. Antiangiogenesis strategies revisited: from starving tumors to alleviating
hypoxia. Cancer Cell 2014;26:605–22. https://doi.org/10.1016/j.ccell.2014.10.006.
[69] Lopes-Coelho F, Martins F, Pereira SA, Serpa J. Anti-Angiogenic Therapy: Current
Challenges and Future Perspectives. Int J Mol Sci 2021;22:3765.
https://doi.org/10.3390/ijms22073765.
[70] Dalva-Aydemir S, Bajpai R, Martinez M, Adekola KUA, Kandela I, Wei C, et al.
Targeting the Metabolic Plasticity of Multiple Myeloma with FDA-Approved
Ritonavir and Metformin. Clin Cancer Res Off J Am Assoc Cancer Res
2015;21:1161–71. https://doi.org/10.1158/1078-0432.CCR-14-1088.
[71] Siebeneicher H, Cleve A, Rehwinkel H, Neuhaus R, Heisler I, Müller T, et al.
Identification and Optimization of the First Highly Selective GLUT1 Inhibitor BAY-
876. ChemMedChem 2016;11:2261–71. https://doi.org/10.1002/cmdc.201600276.
[72] Chan DA, Sutphin PD, Nguyen P, Turcotte S, Lai EW, Banh A, et al. Targeting
GLUT1 and the Warburg effect in renal cell carcinoma by chemical synthetic
lethality. Sci Transl Med 2011;3:94ra70.
https://doi.org/10.1126/scitranslmed.3002394.
[73] Liu Y, Cao Y, Zhang W, Bergmeier S, Qian Y, Akbar H, et al. A small-molecule
inhibitor of glucose transporter 1 downregulates glycolysis, induces cell-cycle
arrest, and inhibits cancer cell growth in vitro and in vivo. Mol Cancer Ther
2012;11:1672–82. https://doi.org/10.1158/1535-7163.MCT-12-0131.
[74] Hurwitz H, Fehrenbacher L, Novotny W, Cartwright T, Hainsworth J, Heim W, et al.
Bevacizumab plus Irinotecan, Fluorouracil, and Leucovorin for Metastatic
Colorectal Cancer. N Engl J Med 2004;350:2335–42.
https://doi.org/10.1056/NEJMoa032691.
[75] Wang J, Ye C, Chen C, Xiong H, Xie B, Zhou J, et al. Glucose transporter GLUT1
expression and clinical outcome in solid tumors: a systematic review and meta-
analysis. Oncotarget 2017;8:16875–86. https://doi.org/10.18632/oncotarget.15171.

[76] Maae E, Nielsen M, Steffensen KD, Jakobsen EH, Jakobsen A, Sørensen FB.
Estimation of Immunohistochemical Expression of VEGF in Ductal Carcinomas of
the Breast. J Histochem Cytochem 2011;59:750–60.
https://doi.org/10.1369/0022155411412599.
[77] Kut C, Mac Gabhann F, Popel AS. Where is VEGF in the body? A meta-analysis of
VEGF distribution in cancer. Br J Cancer 2007;97:978–85.
https://doi.org/10.1038/sj.bjc.6603923.
[78] Apaloo J, Brown JS, Vincent T. Evolutionary game theory: ESS, convergence
stability, and NIS. Evol Ecol Res 2009;11:489–515.
[79] Dawkins R, Krebs JR. Arms races between and within species. Proc R Soc Lond B
Biol Sci 1979. https://doi.org/10.1098/rspb.1979.0081.
[80] Liu Z-L, Chen H-H, Zheng L-L, Sun L-P, Shi L. Angiogenic signaling pathways and
anti-angiogenic therapy for cancer. Signal Transduct Target Ther 2023;8:1–39.
https://doi.org/10.1038/s41392-023-01460-1.
[81] Gatenby RA, Brown JS. Integrating evolutionary dynamics into cancer therapy. Nat
Rev Clin Oncol 2020;17:675–86. https://doi.org/10.1038/s41571-020-0411-1.
[82] Gatenby RA, Brown J, Vincent T. Lessons from applied ecology: cancer control
using an evolutionary double bind. Cancer Res 2009;69:7499–502.
https://doi.org/10.1158/0008-5472.CAN-09-1354.
[83] Thompson JN. The Geographic Mosaic of Coevolution. Chicago, IL: University of
Chicago Press; 2005.
[84] Thompson JN. The Coevolutionary Process. Chicago, IL: University of Chicago
Press; 1994.
[85] Abd El-Azeem MA, Ali MA, El-Shorbagy SH. Expression of GLUT4 and FAP in
urothelial bladder carcinoma: correlation with angiogenesis and clinicopathological
characteristics. J Egypt Natl Cancer Inst 2022;34:40.
https://doi.org/10.1186/s43046-022-00145-0.
[86] Macedo F, Ladeira K, Longatto-Filho A, Martins SF. Gastric Cancer and
Angiogenesis: Is VEGF a Useful Biomarker to Assess Progression and Remission?
J Gastric Cancer 2017;17:1–10. https://doi.org/10.5230/jgc.2017.17.e1.
[87] Yin C, Gao B, Yang J, Wu J. Glucose Transporter-1 (GLUT-1) Expression is
Associated with Tumor Size and Poor Prognosis in Locally Advanced Gastric
Cancer. Med Sci Monit Basic Res 2020;26:e920778-1-e920778-10.
https://doi.org/10.12659/MSMBR.920778.
[88] Alarcon T. Modelling tumour-induced angiogenesis: a review of individual-based
models and multiscale approaches. Contemp Math 2009;494:47–73.
[89] Gersani M, Brown JS., O'Brien EE, Maina GM, Abramsky Z. Tragedy of the
commons as a result of root competition. J Ecol 2001;89:660–9.
https://doi.org/10.1046/j.0022-0477.2001.00609.x.
[90] McNickle GG, Gonzalez-Meler MA, Lynch DJ, Baltzer JL, Brown JS. The world's
biomes and primary production as a triple tragedy of the commons foraging game
played among plants. Proc R Soc B Biol Sci 2016;283:20161993.
https://doi.org/10.1098/rspb.2016.1993.
[91] Paavola J. Climate change: the ultimate 'tragedy of the commons'? In: Cole DH,
Ostrom E, editors. Property in Land and Other Resources. Cambridge, MA: Lincoln
Institute of Land Policy; 2011. p. 417–34.

[92] Allen E, Miéville P, Warren CM, Saghafinia S, Li L, Peng M-W, et al. Metabolic
Symbiosis Enables Adaptive Resistance to Anti-angiogenic Therapy that Is
Dependent on mTOR Signaling. Cell Rep 2016;15:1144–60.
https://doi.org/10.1016/j.celrep.2016.04.029.
[93] Patkulkar PA, Subbalakshmi AR, Jolly MK, Sinharay S. Mapping Spatiotemporal
Heterogeneity in Tumor Profiles by Integrating High-Throughput Imaging and
Omics Analysis. ACS Omega 2023;8:6126–38.
https://doi.org/10.1021/acsomega.2c06659.
[94] Chai X, Tao Q, Li L. Spatiotemporal Heterogeneity of Tumor Glucose Metabolism
Reprogramming: From Single-Cell Mechanisms to Precision Interventions. Int J
Mol Sci 2025;26:6901. https://doi.org/10.3390/ijms26146901.
[95] Andreou C, Weissleder R, Kircher MF. Multiplexed imaging in oncology. Nat
Biomed Eng 2022;6:527–40. https://doi.org/10.1038/s41551-022-00891-5.
[96] Cilento MA, Sweeney CJ, Butler LM. Spatial transcriptomics in cancer research
and potential clinical impact: a narrative review. J Cancer Res Clin Oncol
2024;150:296. https://doi.org/10.1007/s00432-024-05816-0.
[97] Bergers G, Hanahan D. Modes of resistance to anti-angiogenic therapy. Nat Rev
Cancer 2008;8:592–603. https://doi.org/10.1038/nrc2442.
[98] Baraniskin A, Buchberger B, Pox C, Graeven U, Holch JW, Schmiegel W, et al.
Efficacy of bevacizumab in first-line treatment of metastatic colorectal cancer: A
systematic review and meta-analysis. Eur J Cancer Oxf Engl 1990 2019;106:37–
44. https://doi.org/10.1016/j.ejca.2018.10.009.
[99] Carlisle B, Demko N, Freeman G, Hakala A, MacKinnon N, Ramsay T, et al.
Benefit, Risk, and Outcomes in Drug Development: A Systematic Review of
Sunitinib. JNCI J Natl Cancer Inst 2015;108:djv292.
https://doi.org/10.1093/jnci/djv292.
[100] Haibe Y, Kreidieh M, El Hajj H, Khalifeh I, Mukherji D, Temraz S, et al. Resistance
Mechanisms to Anti-angiogenic Therapies in Cancer. Front Oncol 2020;10:221.
https://doi.org/10.3389/fonc.2020.00221.
[101] McIntyre A, Harris AL. Metabolic and hypoxic adaptation to anti-angiogenic
therapy: a target for induced essentiality. EMBO Mol Med 2015;7:368–79.
https://doi.org/10.15252/emmm.201404271.
[102] Ren Z, Zhao J, Li S, Yuan H. Targeting Glucose Transporter 1 (GLUT1) in
Cancer: Molecular Mechanisms and Nanomedicine Applications. Int J
Nanomedicine 2025;20:11859–79. https://doi.org/10.2147/IJN.S534976.
[103] Temre MK, Kumar A, Singh SM. An appraisal of the current status of inhibition of
glucose transporters as an emerging antineoplastic approach: Promising potential
of new pan-GLUT inhibitors. Front Pharmacol 2022;13:1035510.
https://doi.org/10.3389/fphar.2022.1035510.
[104] Raez LE, Papadopoulos K, Ricart AD, Chiorean EG, Dipaola RS, Stein MN, et al.
A phase I dose-escalation trial of 2-deoxy-D-glucose alone or combined with
docetaxel in patients with advanced solid tumors. Cancer Chemother Pharmacol
2013;71:523–30. https://doi.org/10.1007/s00280-012-2045-1.
[105] Devraj K, Klinger ME, Myers RL, Mokashi A, Hawkins RA, Simpson IA. GLUT-1
GLUCOSE TRANSPORTERS IN THE BLOOD-BRAIN BARRIER: DIFFERENTIAL

PHOSPHORYLATION. J Neurosci Res 2011;89:10.1002/jnr.22738. https://doi.org/10.1002/jnr.22738.

[106] Li J, Yu J, Fang Q, Du Y, Zhang X. Gold Nanoparticle Delivery of Glut1 SiRNA Facilitates Glucose Starvation Therapy in Lung Cancer. Chembiochem Eur J Chem Biol 2024;25:e202400239. https://doi.org/10.1002/cbic.202400239.

[107] Wu B, Wang Z, Liu J, Li N, Wang X, Bai H, et al. Dual rectification of metabolism abnormality in pancreatic cancer by a programmed nanomedicine. Nat Commun 2024;15:10526. https://doi.org/10.1038/s41467-024-54963-y.

[108] Wang J-W, Chen Q-W, Luo G-F, Ji P, Han Z-Y, Song W-F, et al. Interference of Glucose Bioavailability of Tumor by Engineered Biohybrids for Potentiating Targeting and Uptake of Antitumor Nanodrugs. Nano Lett 2022;22:8735–43. https://doi.org/10.1021/acs.nanolett.2c03608.

[109] Ren X, Cheng Z, He J, Yao X, Liu Y, Cai K, et al. Inhibition of glycolysis-driven immunosuppression with a nano-assembly enhances response to immune checkpoint blockade therapy in triple negative breast cancer. Nat Commun 2023;14:7021. https://doi.org/10.1038/s41467-023-42883-2.

[110] Wang X, Wang L, Hao Q, Cai M, Wang X, An W. Harnessing glucose metabolism with nanomedicine for cancer treatment. Theranostics 2024;14:6831–82. https://doi.org/10.7150/thno.100036.

[111] Kasumi E, Sato N. A ketogenic diet improves the prognosis in a mouse model of peritoneal dissemination without tumor regression. J Clin Biochem Nutr 2019;64:201–8. https://doi.org/10.3164/jcbn.18-103.

[112] Seyfried TN, Flores RE, Poff AM, D'Agostino DP. Cancer as a metabolic disease: implications for novel therapeutics. Carcinogenesis 2014;35:515–27. https://doi.org/10.1093/carcin/bgt480.

[113] Weber DD, Aminazdeh-Gohari S, Kofler B. Ketogenic diet in cancer therapy. Aging 2018;10:164–5. https://doi.org/10.18632/aging.101382.

[114] DeBerardinis RJ, Chandel NS. Fundamentals of cancer metabolism. Sci Adv 2016;2:e1600200. https://doi.org/10.1126/sciadv.1600200.

[115] Vander Heiden MG, Cantley LC, Thompson CB. Understanding the Warburg Effect: The Metabolic Requirements of Cell Proliferation. Science 2009;324:1029–33. https://doi.org/10.1126/science.1160809.

[116] Jain RK. Normalization of tumor vasculature: an emerging concept in antiangiogenic therapy. Science 2005;307:58–62. https://doi.org/10.1126/science.1104819.

[117] Dekel E, Alon U. Optimality and evolutionary tuning of the expression level of a protein. Nature 2005;436:588–92. https://doi.org/10.1038/nature03842.

[118] Gatenby RA, Brown JS. The Evolution and Ecology of Resistance in Cancer Therapy. Cold Spring Harb Perspect Med 2020;10:a040972. https://doi.org/10.1101/cshperspect.a040972.

[119] Wölfl B, Te Rietmole H, Salvioli M, Kaznatcheev A, Thuijsman F, Brown JS, et al. The Contribution of Evolutionary Game Theory to Understanding and Treating Cancer. Dyn Games Appl 2022;12:313–42. https://doi.org/10.1007/s13235-021-00397-w.