1 **Evolution and adaptations of the seminal proteome in an insect with traumatic**
2 **insemination**
3

4 Martin D. Garlovsky[1], Oliver Otti[1], Klaus Reinhardt[1], Timothy L. Karr[2,3]
5
AFFILIATIONS
[1]Applied Zoology, Faculty Biology, Technische Universität Dresden, Dresden,
Germany
[2]Biosciences Mass Spectrometry Core Research Facility, Knowledge Enterprise,
Arizona State University, USA
[3]Neurodegenerative Disease Research Center, The Biodesign Institute, Arizona
State University, USA
13
14 **CORRESPONDING AUTHOR**
15 Martin D. Garlovsky, Faculty of Biology, Technische Universität Dresden, 01217
16 Dresden, Germany, +49 351 463-39451, martin.garlovsky@tu-dresden.de

17
AUTHOR CONTRIBUTIONS
Conceptualisation: MDG, OO, KR, TLK; Data curation: MDG, TLK; Formal analysis:
MDG, TLK; Funding acquisition: OO, KR; Investigation: MDG, OO; Methodology:
MDG, OO, TLK; Project administration: MDG, OO, KR, TLK; Resources: MDG, OO,
KR, TLK; Software: MDG; Validation: MDG, TLK; Visualization: MDG; Writing –
original draft preparation: MDG; Writing – review and editing: MDG, OO, KR, TLK.
24
25 **KEYWORDS**
26 Sperm | seminal fluid proteins | bedbugs | leucyl-aminopeptidases | AlphaFold3
27
28 **COMPETING INTERESTS**
29 The authors declare no competing interests
30
31 **DATA AVAILABILITY**

ABSTRACT
The protein composition of sperm and seminal fluid are key to male fitness.
However, we currently lack an understanding of the factors that shape seminal
proteome composition. The common bedbug (*Cimex lectularius*) mates by traumatic
insemination, subjecting the ejaculate to a unique selective environment as sperm
traverse the female genital and paragenital system. We provide the first high-
throughput proteomic characterisation of the sperm and seminal fluid proteome in a
hemimetabolous insect and the first in-depth proteomic characterisation of the male
bedbug reproductive system. Our analysis revealed conserved and unique features
of the sperm and seminal fluid proteome with possible links to features of sperm
behaviour linked to traumatic insemination. The sperm proteome showed elevated
rates of molecular evolution, unlike most other studied species. Conversely, the
sperm proteome also contained many conserved proteins. Notably, we found an
expansion of Sperm-leucylaminopeptidases (S-Laps) in bedbugs and other
hemimetabolous insects, suggesting the origin of S-Laps is perhaps even more
ancient than previously thought. Using *in silico* protein-ligand binding predictions, we
show that S-Laps have likely retained catalytic activity. Our results provide a list of
candidate proteins involved in reproduction and a foundation for future studies of this
expanding global pest.

INTRODUCTION

The proteomic composition of sperm and seminal fluid is central to fertility and fitness (Snook 2005; Perry et al. 2013). Proteins are a major component of gametes, in large part dictating spermatozoa form and fertilisation capacity (Dorus and Karr 2009). Likewise, seminal fluid proteins transferred along with sperm in the ejaculate aid sperm transport and storage and affect female physiology and morphology after mating (Wigby et al. 2020). However, until recently the tools needed to study the molecular composition of the ejaculate were lacking (Karr 2019). Consequently, our understanding of the factors shaping seminal proteome composition and evolution remain poorly understood.

High-throughput shotgun proteomics studies have revealed the sperm and seminal fluid proteome contain hundreds or thousands of proteins (Wigby et al. 2020; Pini et al. 2025). Recent studies have highlighted a nuanced landscape of the seminal proteome. For instance, some seminal fluid proteins may have a testis origin (Garlovsky et al. 2022; McCullough et al. 2022). Many reproductive proteins show elevated rates of molecular evolution (Swanson and Vacquier 2002; Dapper and Wade 2020; Patlar et al. 2021). However, the sperm and seminal fluid proteome also show conservation across taxa (Bayram et al. 2016; Pini et al. 2025). For instance, Sperm-leucylaminopeptidases (S-Laps), first identified in the *Drosophila melanogaster* sperm proteome, have been identified in many insects (Dorus et al. 2011; Whittington et al. 2015; Whittington et al. 2017; Bayram et al. 2019; Degner et al. 2019; Laurinyecz et al. 2019; Kumar et al. 2022; Gomez et al. 2024). Similarly, peptidases and peptidase inhibitors are common features of seminal fluid proteomes that initiate cascades of downstream interactions between female and male proteins to ensure long-term postmating responses (LaFlamme et al. 2014; Avila et al. 2015; Singh et al. 2018; Plakke et al. 2019; Wainwright et al. 2021). Studying the seminal proteome in non-model organisms allows testing commonly held assumptions about reproductive proteins (Zuk et al. 2014).

The common bedbug (*Cimex lectularius*) presents an attractive model to study the factors shaping the seminal proteome, with unique selective pressures potentially acting on the male ejaculate. First, in species where males are the heterogametic sex (i.e., XY), genes with male-biased expression are expected to be underrepresented on the X chromosome which spends two-thirds of time in females (Vibranovski et al. 2009). The karyotype in bedbugs is $2n = 26 + X_1X_2Y$ in males (Slack 1939; Ueshima 1967; Grozeva et al. 2010), which may reduce sex biased selection and alter the landscape of sex biased genes. Second, bedbugs obligately mate by traumatic insemination, where the male intromittent organ (paramere) pierces the female cuticle, bypassing the female reproductive tract, which is used only during oviposition (Siva-Jothy 2006). Insemination takes place in the mesospermalege, a novel organ that has evolved in the Cimicidae (Roth et al. 2024), which reduces mating costs to females (Morrow and Arnqvist 2003; Reinhardt et al. 2003). As the site of insemination, sperm first interact with the female in the

mesospermalege before sperm exit into the female haemolymph (Usinger 1966;

Martens et al. 2026). Sperm then travel through the haemolymph and cross into the

female reproductive tract to enter the oviduct via an as yet unknown mechanism,

where some sperm are stored in a unique storage organ (Usinger 1966). Other

sperm move up the oviduct and, unlike in other insects, fertilise eggs in the ovaries

(Usinger 1966). Sperm must therefore traverse multiple tissues, membranes and

substrates to reach the egg (Usinger 1966).

To explore features of the bedbug seminal proteome associated with traumatic

insemination, we provide a detailed proteomic investigation of the male bedbug

reproductive organs, including the sperm and seminal proteomes. This is the first

high-throughput seminal proteome of a hemimetabolous insect, greatly increasing

the number of ejaculate proteins in bedbugs (Reinhardt, Wong, et al. 2009). Recently

published genomic resources (Law et al. 2025; Miles et al. 2025) allow us to further

investigate the evolutionary dynamics of bedbug reproductive proteins, including

rates of molecular evolution and chromosomal distribution. Finally, we compare the

bedbug seminal proteome to *Drosophila* and mosquitoes (another blood-feeding

insect). In doing so, we identified an expansion of the *S-Lap* gene family in the

Cimicomorpha, showing retention of catalytic activity based on sequence similarity

and *in silico* protein-ligand interactions.

RESULTS

Label-free quantitation using LC-MS/MS of the *Cimex lectularius* male reproductive
tract tissues and sperm provide a unique and quantitative view of the bedbug
seminal proteome. In total, we identified 4,651 proteins, of which we retained 3,348
(71.7%; Table 1; Fig. S2) after filtering proteins identified by fewer than 2 unique
peptides and default filtering steps in *MSstats* (Kohler et al. 2023).

Table 1. Total numbers of proteins identified in each tissue/sperm and the number of
tissue/sperm-biased proteins (as a percentage of total in brackets. SFCs, seminal
fluid containers.

	Testes	Sperm	Mesadenes	SFCs
No. proteins identified	3244	2494	3017	2272
No. tissue-biased (%)	535 (16.5%)	487 (19.5%)	708 (23.5%)	239 (10.5%)

Unlike most other insects, true bugs have separate organs which are the source of
seminal fluid production; the mesadenes, which are physically separated from the
seminal fluid storage organ, the seminal fluid containers (Fig. 1A). A principal
component analysis (PCA) of normalised protein abundances showed the first two
PCs explained more than 83% of variation, separating all four tissues (Fig. 1B). PC1
(47.9% variance explained) separated the sperm and testes from the seminal fluid
containers and mesadenes (Fig. 1B). PC2 (35.2% variance explained) further
separated the sperm and seminal fluid containers from the testes and mesadenes.
This suggests some gross similarities between the organs that are the sources of
protein production (testes and mesadenes), separate from the sinks (sperm and
seminal fluid containers).

Figure 1. (A) *Dorsal view of male* Cimex lectularius *reproductive system highlighting
organs dissected for proteomics (adapted from* (Usinger 1966)*). Sperm were isolated
from the seminal vesicles. Bacteriomes (aka mycetomes) were included with the
testes for LC-MS/MS. **B)** Principal component analysis of protein abundances for the*

*500 most variable proteins. **C)** Reproduce protein classification workflow. Numbers*
in Venn diagrams show number of proteins biased towards each tissue (log2FC > 1
*& adjusted p-value < 0.05). **D)** Enrichment of proteins containing a signal peptide*
sequence in the bedbug reproductive proteome. Numbers below bars are the
observed and expected number of proteins in each group, respectively. The dashed
line indicates the null expectation, i.e., the expected proportion of proteins in the
*whole proteome. P-values obtained from X^2 tests. **E)** Chromosomal distribution of*
male-biased proteins on the autosomes and each of the two C. lectularius *X*
chromosomes. Asterisks represent results from comparing the observed to expected
number of proteins on each chromosome after multiple testing correction using X^2
*tests. The dashed line indicates the null expectation. **F)** Pairwise nonsynonymous*
(dN) to synonymous (dS) nucleotide substitution rate, dN/dS (mean ± standard error)
between C. lectularius and C. hemipterus. Numbers below points indicate numbers
of genes in each category. Asterisks represent results from Mann-Whitney U tests
comparing each gene set to the genome average shown by dashed line (dN/dS =
0.167 ± 0.001, n = 17580). * p < 0.05, ** p < 0.01, *** p < 0.001. Sfp, seminal fluid
proteins. SFCs, seminal fluid containers.
164
165   **Characterisation of the male reproductive proteome**
166   We defined proteomes for each tissue/sperm based on proteins having higher
167   abundance (or the exclusive presence) compared to all other tissues (log2FC > 1 &
168   adjusted *p*-value < 0.05; Fig. 1C; Table 1; Fig. S2&S3).
169
170   **Testis and sperm proteome.** The combined testis and sperm proteome comprised
171   1,226 proteins and showed underrepresentation of proteins containing a signal
172   peptide sequence (X^2 = 30.67, df = 1, $p < 0.001$; Fig. 1D). Sperm- and testes-
173   biased proteins were underrepresented on chromosome X_2 (sperm: X^2 = 10.0, df =
174   1, $p = 0.046$; testes: X^2 = 9.14, df = 1, $p = 0.05$; Fig. 1E & Fig. S5).
175
The putative sperm proteome, comprising proteins biased towards sperm or sperm
and testes (n = 691; Fig. 1C) was evolving at an elevated rate compared to the
genome average (mean pairwise dN/dS ± standard error = 0.20 ± 0.01, n = 580;
pairwise Wilcoxon test, $p < 0.001$; Fig. 1F). The sperm proteome showed GO
enrichment of biological processes involving metabolic and glycolytic processes,
proteolysis, cilium assembly and microtubule-based process; cellular components
typically associated with sperm cells, such as mitochondria, cilium, axoneme, dynein
complex, cytoplasm and microtubules; and molecular functions including ion binding
and serine-type endopeptidase activity (Table S3).
185
Proteins exclusively biased towards testes (n = 535) showed GO enrichment of
biological processes including protein folding, cellular transport and organisation and
translation; cellular components including proteasomes, cytoplasm, microtubules and
mitochondria; molecular functions including protein folding, ATP binding and heat
shock protein binding (Table S4). Endosymbiotic bacteria (e.g., *Wolbachia*)
(Thongprem et al. 2020) are localised in bacteriomes (aka mycetomes) attached to
the *vas deferens* at the base of the testes (Usinger 1966; Hosokawa et al. 2010). As

expected, *Wolbachia* proteins showed the highest abundance in the
testes/bacteriome samples (Fig. S4).
195

Mesadene and seminal fluid proteome. The seminal fluid producing proteome,
including proteins biased towards mesadenes and/or seminal fluid containers
comprised 1022 proteins (Fig. 1C). Proteins containing a signal peptide sequence
were overrepresented in the mesadenes and seminal fluid containers (X^2 = 123.45,
df = 1, *p* < 0.001; Fig. 1D). We classified the 249 proteins containing a signal peptide
sequence from the seminal fluid producing organs as our putative seminal fluid
proteome (Fig. 1C); a number similar to the number of "high confidence" Sfps
identified in *Drosophila melanogaster* (Wigby et al. 2020). While the presence of a
signal peptide sequence is indicative of a protein being secreted (Teufel et al. 2022)
and a hallmark of seminal fluid proteins, not all secreted proteins contain signal
peptides. We found a subset of putative seminal fluid proteins expressed highly (in
the top 20% most abundant proteins) in sperm (3.6%; 9/249) and testes (8%;
20/249), in line with recent studies suggesting a more complex landscape of seminal
fluid biology (Garlovsky et al. 2022; McCullough et al. 2022).
210

Seminal fluid proteins were evolving at a similar rate to sperm (*p* = 0.106) but were
not evolving faster than the genome average (dN/dS = 0.18 ± 0.01, n = 224; *p* =
0.135; Fig. 1F). The seminal fluid proteome showed GO enrichment for biological
processes including protein folding and proteolysis, glycosylation and negative
regulation of peptidase activity; cellular components including endoplasmic reticulum
and extracellular space; and molecular functions including ion and molecule binding,
and (serin-type) endopeptidase inhibitor activity (Table S5).
218

The remaining 773 proteins classified as the seminal fluid producing mesadenal
category were overrepresented on X_2 (X^2 = 11.8, df = 1, *p* = 0.035; Fig. 1E).
Mesdadenal proteins showed GO enrichments often associated with accessory
gland/secretory functions; biological processes such as translation, protein transport
and Golgi/vesicle transport; cellular components including ribosome, endoplasmic
reticulum and Golgi apparatus; molecular functions including RNA- and GTP- binding
(Table S6).
226

Comparison with other seminal proteomes
In the bedbug sperm proteome (n = 691), 189 orthologs of *D. melanogaster* sperm
proteins were identified (Garlovsky et al. 2022; McCullough et al. 2022) and 58
Aedes aegypti sperm proteins (Degner et al. 2019). Among the most abundant
proteins in the bedbug sperm proteome were orthologs of *D. melanogaster* sperm
proteins, including proteins involved in spermatid development (e.g., betaTub56D,
bb8), S-Laps, mitochondrial proteins (e.g., P5CDh1, Cs1, Ant2, mAcon, Mitofilin) and
Y-linked fertility factors (e.g., ORY, WDY, Ppr-Y, kl-3, kl-5) (Table 2; Table S7). The
bedbug seminal fluid proteome (n = 249) contained fewer orthologs; just 15 *D.*

melanogaster seminal fluid proteins (Wigby et al. 2020), and 5 *Ae. aegypti* seminal
fluid proteins (Degner et al. 2019).

Table 2. *The most abundant proteins in the bedbug sperm- and seminal fluid-*
proteomes with D. melanogaster *orthologs (rank ordered by abundance). Bedbug*
proteins matching more than one Drosophila *gene are separated by a comma. S-*
Laps are labelled based on one-to-one orthology with D. melanogaster *or*
membership to S-Lap cluster 1 (g1) or 2 (g2) (Dorus et al. 2011) *followed by a*
unique ID. D. melanogaster *Sfps are noted with an asterisk (all found in the Sfp*
category).

Sperm proteome			Seminal fluid proteome		
Protein	*Drosophila* name(s)	Rank	Protein	*Drosophila* name(s)	Rank
A0A8I6R8Q4	betaTub56D	100	A0A8I6TE74	Pdi	99.8
A0A8I6S6N3	S-Lap3	99.9	A0A8I6S6T0	Gp93	99.6
A0A8I6THS3	S-Lap-g1_2	99.7	A0A8I6R8G9	OstDelta	99.1
A0A8I6RD41	S-Lap4	99.6	A0A8I6S4A5	Grp170	98.6
A0A8I6RDX4	bb8	99.5	A0A8I6SFV2	mfas	95.0
A0A8I6R8J1	S-Lap-g2_4	99.1	A0A8I6SCN7	Swim	84.4
A0A8I6RZZ4	P5CDh1	99.0	A0A8I6RVV7	NUCB1	81.0
A0A8I6TEZ7	S-Lap-g1_5	98.9	A0A8I6RXP7	GCS2alpha	88.6
A0A8I6TD95	S-Lap-g2_6	98.7	A0A8I6SQD0	Prx4	85.3
A0A8I6TJT9	Cs1	98.5	A0A8I6RJB5	Spn88Ea, Spn88Eb	65.4
A0A8I6RI95	Ant2	98.4	A0A8I6RPW4	CG17271	74.1
A0A8I6TIM8	mAcon1, mAcon2	98.4	A0A8I6RND8	Phm	46.7
A0A8I6RBN0	Mitofilin	98.0	A0A8I6RWL8	CG8420	67.3
A0A8I6RL88	Alat	97.6	A0A8I6RAJ9	CG12093	23.0
A0A8I6SB96	S-Lap-g1_7	97.5	A0A8I6RFN7	Qsox1, Qsox3, Qsox2, Qsox4	33.9
A0A8I6SDI3	CG3213, CG13405	97.3			
A0A8I6RRZ2	CG3339, kl-5	97.2			
A0A8I6TFY4	SdhA	96.6			
A0A8I6S5H6	kl-3	96.4			
A0A8I6SIB1	AIF	96.3			

247
Figure 2. *Ranked abundance of the top 5% most abundant proteins in each*
tissue/sperm. Points are coloured by proteome category (mesadenes, dark blue;

seminal fluid proteins, light blue; sperm, yellow; testes, orange). Orthologs of
Drosophila melanogaster *sperm (Garlovsky et al. 2022) and seminal fluid proteins*
(Wigby et al. 2020) are labelled. S-Laps are labelled based on one-to-one orthology
with D. melanogaster *or membership to S-Lap cluster 1 or 2 (Dorus et al. 2011)*
followed by a unique ID.
255

Sperm-leucylaminopeptidase evolution

Notably, 6 of the 10 most abundant bedbug sperm proteins with *D. melanogaster*
orthologs were S-Laps (Fig. 2; Table 2) (Dorus et al. 2011). We identified 13 *C.*
lectularius S-Lap orthologs in total; five belonging to S-Lap cluster 1, eight in cluster
2 (Dorus et al. 2011), all of which were most abundant in sperm (Fig. S6). To further
explore orthology between the bedbug and *Drosophila* S-Laps we used BLASTp
(Camacho et al. 2009) which showed matches to all 8 *Drosophila* S-Laps. A more
stringent reciprocal best BLAST search (e-value = 1e-2, min. ident = 30, min.
coverage = 50) returned four one-to-one S-Lap orthologs: S-Laps 2, 3, 4 and 7 (Fig.
3B).
266

There appear to be no S-Lap orthologs in more distantly related Hemiptera
(*Acyrthosiphon pisum*) or Orthoptera (*Gryllus bimaculatus*), but S-Lap orthologs were
present in the Coleoptera (*Callosobruchus maculatus*) (see also (Gomez et al. 2024)
(Fig. 3A). However, orthologs of *granny-smith*, the closest S-Lap paralog (Dorus et
al. 2011), were identified in all species studied (Fig. 3A & Fig. S7). Across the 73 S-
Lap orthologs from 7 species studied (including *D. melanogaster*), proteins first
cluster by S-Lap cluster (1 or 2) rather than species based on sequence similarity,
suggesting the origin of S-Laps predates the separation of the Holometabola from
the Hemimetabola (Misof et al. 2014).

*Figure 3. Sperm-leucylaminopeptidase (S-Lap) enzymatic activity in bedbugs. **A)***
Unrooted gene tree of S-Lap and granny-smith orthologs in insects. Protein
sequences were aligned using MAFFT (Madeira et al. 2024) and bootstrapped 1,000
times. Tip colour indicates Order and shape indicates species. Drosophila *and*
*bedbug S-Laps shown with open circles. **B)** Alignment of bedbug S-Lap orthologs*
focused on amino acid residues against the M17 amino peptidase metal binding
motif indicated above (Burley et al. 1992). Amino acids are coloured by alignment to
consensus (conserved, red; compatible, pink; non-functional, white). Zn^{2+} ion binding
at the "loose" binding site (Zn1) and "tight" binding site (Zn2) is scored as canonical
(red), uncertain (pink) or non-canonical (white). Overall activity is scored as likely
active (blue), uncertain (green), or likely inactive (yellow) based on coordinating
residue geometry, number and inter-ligand angles and inter-zinc distances measured
*from AlphaFold3 models in ChimeraX. **C)** AlphaFold predictions for structures of the*
canonical M17 LAP, cytosol aminopeptidase (LAP3; UniProt ID: P00727), and
bedbug S-Laps focussed on zinc binding sites. Residues containing atoms within 4Å
of Zn^{2+} ions (purple spheres) are highlighted: N, blue; O, red; S, yellow; C, blue.
293
S-Laps belong to the M17 family of leucyl aminopeptidases (Dorus et al. 2011) which
bind two Zn^{2+} ions (Maric et al. 2009). However, in *D. melanogaster*, S-Laps
belonging to "cluster 1" have mutations at most of the active site residues and sperm
display very weak enzymatic activity (Dorus et al. 2011; Laurinyecz et al. 2019).
Most (11 out of 13) bedbug S-Laps, including those in "cluster 1", retain at least 4 out
of 5 metal ion binding residues including all the residues coordinating tight Zn^{2+}
binding and cofactor binding sites (Fig. 3B).
301
Structure-based predictions to determine 3D coordination of Zn^{2+} ions also showed
that most bedbug S-Laps (at least 8 out of 13) retain the M17 LAP zinc binding
coordination geometry (Fig. 3B,C). However, five bedbug S-Laps (two in cluster 1,
three in cluster 2) are likely inactive, with inter-zinc distances exceeding the ~3Å
observed in the conserved M17 structure (Burley et al. 1992; McGowan et al. 2010)
(Fig. 3B,C).
308
This pattern of (i) retention of metal binding active site residues, and (ii) 3D
coordination of Zn^{2+} was observed across the species we studied outside of
Drosophila including other hemipterans and hemimetabolous insects as well as in
Diptera and Coleoptera (Fig. S7). Notably, *C. maculatus* S-Laps in "cluster 1" do not
have the conserved Asp residue coordinating site 2 tight zinc binding and lack zinc
binding geometry (Fig. S7).
315

DISCUSSION

Bedbugs are unusual among insects in obligately mating by traumatic insemination. After insemination, sperm move in aggregations out of the mesospermalege towards the ovaries or female sperm storage organs (Usinger 1966; Martens et al. 2026). Sperm must therefore cross multiple tissue membranes, likely shaping key adaptations in the bedbug seminal proteome. Our analysis revealed conserved and unique features of the sperm and seminal fluid proteome with possible links to this unusual sperm behaviour.

Sperm and testes proteins were underrepresented on chromosome X_2, as expected for proteins with male-biased expression (Vibranovski et al. 2009). However, the same pattern was not found for chromosome X_1. The two X chromosomes in *Cimex* are thought to be derived from fission of an ancestral X (Grozeva et al. 2010; Poggio et al. 2014). Mesadenal proteins, on the other hand, were overrepresented on X_2, which may result if new mesadenal genes have arisen on the X (Levine et al. 2006). The multiple sex chromosomes present interesting opportunities for unusual evolutionary dynamics that warrant further investigation.

We did not observe the commonly found pattern of rapidly evolving seminal fluid proteins (Swanson and Vacquier 2002), but found few seminal fluid protein orthologs in *Drosophila* or *Aedes* mosquitoes. Surprisingly, elevated rates of molecular evolution were observed in the bedbug sperm proteome. In many taxa, the sperm proteome evolves slowly, thought to result from purifying selection (Garlovsky et al. 2022). Elevated rates of sperm evolution might be driven by sexually antagonistic co-evolution in the bedbug associated with traumatic insemination and movement of sperm through the haemolymph. The mating rate in bedbugs is at least ten times above the female optimum (Stutt and Siva-Jothy 2001; Reinhardt et al. 2011), setting the stage for sexual conflict affecting ejaculate × female interactions (Morrow and Arnqvist 2003; Reinhardt et al. 2003; Sirot et al. 2015). Sperm traversing the female haemolymph and female reproductive tract might entail novel and unusual selective pressures acting on sperm and seminal fluid proteins. As such, it is unclear if or for how long seminal fluid associates with sperm after insemination, as found in other species (Misra and Wolfner 2020). We also found a subset of putative seminal fluid proteins were abundant in sperm and testes, in line with recent studies which point to a possible testis origin of some seminal fluid proteins (Garlovsky et al. 2022; McCullough et al. 2022).

Despite overall elevated rate of evolution in the bedbug sperm proteome, many of the most abundant sperm proteins had orthologs in other insects. Most notably, we identified 13 Sperm-leucylaminopeptidases (S-Laps) in bedbugs and other insects including other hemimetabolous insects. The S-Laps in *Drosophila* show evidence of purifying selection (Dorus et al. 2011) and indeed were evolving more slowly than the rest of the genome in bedbugs (Fig. S8). Other S-Lap orthologs have been identified in Lepidoptera, Coleoptera and Diptera, suggesting an ancient origin of

360 these sperm proteins after more than 380 million years since the last common
361 ancestor (Dorus et al. 2011; Whittington et al. 2015; Whittington et al. 2017; Bayram
362 et al. 2019; Degner et al. 2019; Laurinyecz et al. 2019; Garlovsky et al. 2022; Kumar
363 et al. 2022; Gomez et al. 2024).
364
S-Laps belong to the M17 family of metal-dependent leucyl aminopeptidases, but in
Drosophila have mostly lost enzymatic activity (Dorus et al. 2011; Laurinyecz et al.
2019). Our functional analysis revealed that S-Laps in *Cimex* retain the Zn^{2+} binding
motif, suggesting that S-Laps retained enzymatic properties. Additional direct
biochemical assay of *Cimex* S-Laps will help support the idea that enzymatic activity
has been secondarily lost in *Drosophila*. Our analysis suggests neofunctionalisation
of this class of M17 leucyl-aminopeptidases while retaining an active site in bedbugs
and other insects. S-Laps have been identified in whirligig beetles and also retain the
metal binding motif (Gomez et al. 2024). Whirligig sperm move in conjugates
attached to spermatostyles (Breland and Simmons 1970). Bedbug sperm also move
in conjugates, lending support to the notion that S-Laps play a role in sperm
conjugation and cooperation (Gomez et al. 2024).
377
Conclusions
In conclusion, our study provides a substantially expanded database of male
reproductive proteins in bedbugs using high-throughput proteomics, increasing
coverage of the bedbug seminal proteome by two orders of magnitude (Reinhardt,
Wong, et al. 2009). Our analysis provides insights into the landscape of sperm and
seminal fluid dynamics in an insect with unique features of reproductive anatomy and
physiology. The peculiarities of traumatic insemination and multiple sex
chromosomes in the bedbug have led to unique evolutionary characteristics of the
seminal proteome. Sperm genes are underrepresented on the second X
chromosome and evolve rapidly. S-Laps have retained catalytic activity, which may
aid in sperm conjugation. A better understanding of the proteins important for
reproduction will aid the development of effective strategies to control and suppress
populations of insect pests and disease vectors (Doggett et al. 2012). Unlike many
other blood-feeding insects, bedbugs do not transmit any known human pathogens
(Doggett et al. 2012). Nevertheless, bedbugs are an expanding global pest, causing
sleeplessness, skin irritation, severe allergic reactions and growing economic costs
(Reinhardt and Siva-Jothy 2007; Reinhardt, Kempke, et al. 2009; Doggett et al.
2012). Our results provide a useful resource for future studies and a list of candidate
proteins involved in reproduction in this economically important insect.

MATERIALS AND METHODS

Bedbug maintenance

We used bedbugs originally collected in London, UK (F4 population) in 2006 and maintained at TU Dresden, Germany, as a large, outbred population with overlapping generations. Bedbugs were maintained at 26ºC, 70% relative humidity on a 12:12 hour light:dark cycle. Bedbugs were fed on human blood once each week during the juvenile stages following long-term experimental protocols (Reinhardt et al. 2003). We used males aged 3 weeks post-final moult. Males were housed in groups of 10-15 and fed once a week for 2 weeks after eclosion as adults and left to mature for a further week.

Tissue collection and protein extraction

All dissections were performed in freshly prepared ice-cold phosphate-buffered saline (1X PBS) with 1X protease inhibitors (HALT) and 5mM EDTA (Thermo Fisher). We anaesthetised males on ice and removed reproductive tracts with forceps, insect pins, and dissecting scissors under a stereo dissecting microscope. We had four replicates consisting of 4-6 males each.

To collect sperm samples, seminal vesicles were separated from the remaining reproductive tract tissue and transferred to a fresh drop of PBS/HALT. We then made a small incision at the distal end of the seminal vesicle and exuded the sperm mass using insect pins. The sperm mass was then transferred to an Eppendorf containing 1mL PBS/HALT kept on ice. Seminal fluid containers and mesadenes were moved to a fresh drop of PBS/HALT and separated using insect pins. Mesadenes were transferred to an Eppendorf containing 1mL PBS/HALT kept on ice. Seminal fluid containers were transferred in 2 µL PBS/HALT per gland to a separate Eppendorf. Testes were separated from the remaining tissue using insect pins and transferred to an Eppendorf containing 1mL PBS/HALT kept on ice. Sperm, testis, and mesadene samples were pelleted at 15,000 × G for 2 minutes at 4ºC, supernatant carefully removed, and the pellet resuspended by addition of 1 mL PBS/HALT followed by an additional 2 mins centrifugation at 15,000 × G. The washing procedure was repeated twice, and the final pellet resuspended in 1 mL PBS/HALT and stored at -80ºC. Seminal fluid containers were freeze/thawed and vortexed for 30 seconds three times and then spun at 15,000 × G for 5 minutes at 4ºC and stored at -80ºC suspended in 10 µL of PBS/HALT.

Liquid chromatography tandem mass spectrometry

All LC-MS analyses were performed at the Biosciences Mass Spectrometry Core Facility (https://cores.research.asu.edu/mass-spec/) at Arizona State University. For LC–MS/MS, solubilised proteins were quantified (Thermo Fisher EZQ Protein Quantitation Kit or the Pierce BCA). Proteins were reduced with 50 mM dithiothreitol (Sigma-Aldrich) at 95 °C for 10 min and alkylated for 30 min with 40 mM iodoacetamide (Pierce). Proteins were digested using 2.0 µg of MS-grade porcine trypsin (Pierce), and peptides were recovered using S-trap Micro Columns (Protifi)

per manufacturer directions. Recovered peptides were dried via speed vac and
resuspended in 30 µL of 0.1% formic acid. All data-dependent mass spectra were
collected in positive mode using an Orbitrap Fusion Lumos mass spectrometer
(Thermo Scientific) coupled with a Dionex UltiMate 3000 system equipped with an
NCS-3500 nano capillary pump and UltiMate 3000 RS autosampler. One µL of the
peptide was fractionated using an Easy-Spray LC column (50 cm Å ~75 µm ID,
PepMap C18, 2 µm particles, 100 Å pore size, Thermo Scientific). Electrospray
potential was set to 1.9 kV and the ion transfer tube temperature to 275 °C. The
mass spectra were collected using the method optimised for peptide analysis
provided by Thermo Scientific. Full MS scans (350–1500 m/z range) were acquired
in profile mode with the following settings: Orbitrap resolution 120,000, cycle time
3 s and mass range "Normal;" RF lens at 35%, and the AGC set to "Standard".
Monoisotopic peak determination (MIPS) at "peptide" and included charge states 2–
7; dynamic exclusion at 60 s, mass tolerance 10 ppm, intensity threshold at 5.0e4;
MS/MS spectra acquired in a centroid mode using quadrupole isolation window at
1.6 (m/z); Higher-energy collisional dissociation (HCD) energies at 35%. Spectra
were acquired over a 180-min gradient, flow rate 0.250 µL/min as follows: 0–10 min
at 0-2%, 10–100 min at 2–8%, 100–110 min at 8–32%, 110-120 min at 32–35%,
120–125 min at 35–45%, 125–155 min at 45-98%,155–165 min at 98%, 165 –
180 min at 2%,followed by post run wash step.
461
Protein identification and label-free quantification
We analysed raw files searched against the UniProt (www.uniprot.org) *Cimex*
lectularius database that included two bacterial predicted proteomes
(CIMLE_UP000494040; wCle UP000031663; Rickettsia UP000826664) using
Proteome Discover 2.5 (Thermo Scientific). Input data included Trypsin as enzyme,
maximum missed cleavage site 3, min/max peptide length 6/144, and precursor ion
(MS1) mass tolerance set to 20 ppm and fragment mass tolerance set to 0.5 Da.
Carbamidomethyl (C) was specified as fixed modification, and dynamic modifications
set to Aceytl and Met-loss at the N-terminus, and oxidation of Met. SequestHT and
Percolator ID validation was performed using a minimum of 1 peptide identified using
a concatenated target/decoy strategy and false-discovery rate (FDR) set to 1.0%.
The Minora feature detector identified between run peak matches and quantification
intensities, normalised and scaled to a maximum of 100. Intensity values were
averaged across all six runs and ranked from highest to lowest to obtain a relative
abundance profile. A combined list of predicted signal peptide sequences from the
entire *C. lectularius* proteome was obtained using signal-P 6 (Teufel et al. 2022) and
Phobius (Käll et al. 2004).
479
Ortholog identification
We downloaded proteomes for *Cimex hemipterus*, assassin bug (*Triatoma*
infestans), kissing bug (*Rhodnius prolixus*), pea aphid (*Acyrthosiphon pisum*), seed
beetle (*Callosobruchus maculatus*), yellow fever mosquito (*Aedes aegypti*) and fruit
fly (*Drosophila melanogaster*) from uniprot.org and two-spotted cricket (*Gryllus*

bimaculatus) from NCBI (https://www.ncbi.nlm.nih.gov/) (Fig. S1; Table S1). Species
were chosen because of their evolutionary relationship to *Cimex lectularius*, i.e., the
true bugs and hemimetabolous insects (aphid, cricket), because of similar biology,
such as blood feeding (mosquito), or because of a wealth of data on reproductive
biology, including the occurrence of copulatory wounding (*D. melanogaster* and seed
beetle). Hierarchical orthologs were identified in each species with *OrthoFinder* using
default settings (Emms and Kelly 2015; Emms and Kelly 2019).
492

Evolutionary dynamics
We obtained previously calculated pairwise rates of nonsynonymous (dN) and
synonymous (dS) nucleotide substitution rates (dN/dS) and chromosomal positions
of genes (Martens et al. 2026). Briefly, rates of molecular evolution (dN/dS) were
calculated between *C. lectularius* and *C. hemipterius* using PAML (Yang 2007).
Chromosomal distribution of genes was determined by aligning coding sequences
from the *C. lectularius* reference genome Clec_2.1 (Benoit et al. 2016) to the
recently published chromosome-level *C. lectularius* genome assembly (Miles et al.
2025) using BLASTn (Camacho et al. 2009).
502

Gene ontology enrichment analysis
We performed gene ontology enrichment analysis using the topGO package
(Rahnenfuhrer 2022) in R (R Core Team 2024) using the GO annotations from
uniprot.org. The entire bedbug proteome was used as background for enrichment
tests. Statistical analyses in topGO used Fisher's exact tests with topnodes = 10,
and corrected p-values for multiple testing using the FDR method (Benjamini and
Hochberg 1995). All GO terms reported had an FDR-corrected *p*-value < 0.05.
510

511   ***Statistical analysis***
512   All statistical analysis was performed in R v4.5.0 (R Core Team 2024). Output from
513   Proteome Discoverer was imported into R and formatted for use in the *MSstats*
514   pipeline (Kohler et al. 2023) using the following options: normalisation = quantile,
515   imputed missing values using the TMP method, MBimpute = TRUE.
516

To identify proteins biased towards each tissue (seminal fluid containers,
mesadenes, testes) or sperm, we performed pairwise differential abundance analysis
between each tissue/sperm. We considered a protein to be biased towards a
tissue/sperm based on a log2-fold-change > 1 and false discovery corrected *p*-value
< 0.05 compared to all other tissues. Proteins had to be identified in two or more
replicates of a tissue/sperm to count for that tissue/sperm. We then classified
proteins into 1 of 4 categories: (1) the putative seminal fluid proteome was defined
as proteins biased towards the mesadenes and/or seminal fluid containers
containing a signal peptide sequence; (2) the remaining proteins from the seminal
fluid producing organs were classified as the mesadenal (seminal fluid producing)
proteome; (3) the putative *C. lectularius* sperm proteome was defined as proteins

more abundant in sperm or sperm and testes; (4) the remaining proteins biased
towards testes we classified as the testes proteome (Fig. 1C).
530
Predicting protein-interactions
We characterised catalytic activity for S-Laps based on conservation of the M17
aminopeptidase metal-binding and catalytic residue motif (Maric et al. 2009).
Orthologs of S-Laps and granny-smith (UniProt ID: Q9V3D8) identified using
OrthoFinder and cytosol aminopeptidase (LAP3; UniProt ID: P00727) were aligned
using *MUSCLE* (Edgar 2022). The catalytic domain of each sequence was identified
by mapping a ±150 residue window around the active site centre of *granny-smith*.
The position equivalent to the start of the motif (Lys327) was pinned in each
sequence and the 7 active site positions scored at fixed offsets from the anchor.
Each position was classified as (i) conserved, exact match to the M17 LAP
consensus; (ii) compatible, a residue with structural or mutagenesis evidence for
metal coordination in M17 LAPs (Burley et al. 1992; Gu and Walling 2002; Dorus et
al. 2011); or (iii) non-functional (Table S2).
544
Zinc coordination sites were also assessed using mmCIF results from modelling
complexes of each protein (after removing signal peptide sequences identified in
Phobius) with two Zn^{2+} ions using Alphafold3 (Abramson et al. 2024). Zinc
coordination geometry was assessed with ChimeraX v.1.11.1 (Meng et al. 2023) in
headless mode using a custom Python script. Atoms within 3.5Å of each zinc ion
were identified and contacts classified as canonical if they involved the sidechain
nitrogen of lysine (NZ) or carboxylate oxygens of aspartate (OD1/OD2) or glutamate
(OE1/OE2) consistent with the experimentally derived coordination chemistry of M17
leucyl aminopeptidases (PDB: 1LAP) (Burley et al. 1992). Where both carboxylate
oxygens of a single Asp or Glu residue fell within the distance cutoff (bidentate
geometry) only the closer oxygen was retained.
556
Each zinc ion was classified as occupying either the exchangeable site (Zn1,
characterised by lysine coordination) or the tight site (Zn2, characterised by two
aspartate contacts and absence of lysine), based on coordination chemistry of LAP3
(P00727). We applied a scoring rubric (from 1 to 10) for coordination environments
of Zn1 and Zn2 that incorporated presence of canonical coordinating residues (6
points), coordination number (2 points), bond distance (1 point, penalised for
contacts > 3.2Å) and mean pLDDT confidence at the binding site (≥ 70, 1 point).
Each binding site was classified as canonical (score ≥ 6), uncertain (score 3-5), or
non-canonical (score < 3). Distance thresholds were set to account for the
systematic overestimation of metal-ligand bond lengths in AF3 predictions relative to
experimental crystal structures.
568
After scoring each site, structures were assessed for presence of a shared bridging
ligand/residue providing sidechain contacts to both zinc ions and Euclidean distance
between the two zinc ions was calculated to determine if zinc ions resided in the

same zinc pocket (≤4Å) (Burley et al. 1992). Proteins were classified as; "likely
active": both binding sites canonical and zinc ions residing within the same zinc
pocket; "uncertain": both canonical sites and both zincs within the inter-zinc distance
but no bridging ligand; or "likely inactive", i.e., non-functional: one or two non-
canonical binding sites or separate binding pockets (inter-zinc distances > 7Å).
577

ACKNOWLEDGEMENTS
We would like to thank Christin Froschauer and Susanne Broschk for help in the lab
and Caitlin McDonough-Goldstein for comments on an earlier version of the
manuscript. We thank members of the ASU Biosciences Core Mass Spectrometry
facility, Kyle Tucker, Sydney Canning and Yishai Gilron for the acquisition of mass
spectra and downstream computational analyses. Yan Ge and the ZIH team at TU
Dresden provided valuable assistance in implementing AlphaFold3. This work was
funded by DFG grants to OO 521/4-1 and KR 1666/4-1. We are grateful for high-
performance computing at the NHR Center of TU Dresden, jointly supported by the
Federal Ministry of Education and Research and the state governments participating
in the NHR (www.nhr-verein.de/unsere-partner).

REFERENCES

Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard AJ, Bambrick J, et al. 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* 630:493–500.

Avila FW, Mattei AL, Wolfner MF. 2015. Sex peptide receptor is required for the release of stored sperm by mated *Drosophila melanogaster* females. *J. Insect Physiol.* 76:1–6.

Bayram H, Claydon AJ, Brownridge PJ, Hurst JL, Mileham A, Stockley P, Beynon RJ, Hammond DE. 2016. Cross-species proteomics in analysis of mammalian sperm proteins. *J. Proteomics* 135:38–50.

Bayram H, Sayadi A, Immonen E, Arnqvist G. 2019. Identification of novel ejaculate proteins in a seed beetle and division of labour across male accessory reproductive glands. *Insect Biochem. Mol. Biol.* 104:50–57.

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* 57:289–300.

Benoit JB, Adelman ZN, Reinhardt K, Dolan A, Poelchau M, Jennings EC, Szuter EM, Hagan RW, Gujar H, Shukla JN, et al. 2016. Unique features of a global human ectoparasite identified through sequencing of the bed bug genome. *Nat. Commun.* 7:10165.

Breland OP, Simmons E. 1970. Preliminary Studies Of The Spermatozoa And The Male Reproductive System Of Some Whirligig Beetles Coleoptera Gyrinidae. *Entomol. News* 81:101–110.

Burley SK, David PR, Sweet RM, Taylor A, Lipscomb WN. 1992. Structure determination and refinement of bovine lens leucine aminopeptidase and its complex with bestatin. *J. Mol. Biol.* 224:113–140.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421.

Dapper AL, Wade MJ. 2020. Relaxed selection and the rapid evolution of reproductive genes. *Trends Genet.* 36:640–649.

Degner EC, Ahmed-Braimah YH, Borziak K, Wolfner MF, Harrington LC, Dorus S. 2019. Proteins, Transcripts, and Genetic Architecture of Seminal Fluid and Sperm in the Mosquito Aedes aegypti. *Mol. Cell. Proteomics* 18:S6–S22.

Doggett SL, Dwyer DE, Peñas PF, Russell RC. 2012. Bed Bugs: Clinical Relevance and Control Options. *Clin. Microbiol. Rev.* 25:164–192.

Dorus S, Karr TL. 2009. Sperm proteomics and genomics. In: Sperm biology: an evolutionary perspective. 1st ed. Burlington, MA: Academic press. p. 435–469.

Dorus S, Wilkin EC, Karr TL. 2011. Expansion and functional diversification of a
leucyl aminopeptidase family that encodes the major protein constituents of
Drosophila sperm. *BMC Genomics* 12:177.

Edgar RC. 2022. Muscle5: High-accuracy alignment ensembles enable unbiased
assessments of sequence homology and phylogeny. *Nat. Commun.* 13:6968.

Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. *Genome*
Biol. 16:157.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for
comparative genomics. *Genome Biol.* 20:238.

Garlovsky MD, Sandler JA, Karr TL. 2022. Functional Diversity and Evolution of the
Drosophila Sperm Proteome. *Mol. Cell. Proteomics* [Internet] 21. Available
from: https://www.mcponline.org/article/S1535-9476(22)00089-5/abstract

Gomez RA, Dallai R, Sims-West DJ, Mercati D, Sinka R, Ahmed-Braimah Y, Pitnick
S, Dorus S. 2024. Proteomic diversification of spermatostyles among six
species of whirligig beetles. *Mol. Reprod. Dev.* 91:e23745.

Grozeva S, Kuznetsova VG, Anokhin B. 2010. Bed bug cytogenetics: karyotype, sex
chromosome system, FISH mapping of 18S rDNA, and male meiosis in Cimex
lectularius Linnaeus, 1758 (Heteroptera: Cimicidae). *Comp. Cytogenet.*
4:151–160.

Gu Y-Q, Walling LL. 2002. Identification of residues critical for activity of the wound-
induced leucine aminopeptidase (LAP-A) of tomato. *Eur. J. Biochem.*
269:1630–1640.

Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. 2010. Wolbachia as a
bacteriocyte-associated nutritional mutualist. *Proc. Natl. Acad. Sci.* 107:769–
774.

Käll L, Krogh A, Sonnhammer ELL. 2004. A Combined Transmembrane Topology
and Signal Peptide Prediction Method. *J. Mol. Biol.* 338:1027–1036.

Karr TL. 2019. Reproductive proteomics comes of age. *Mol. Cell. Proteomics* 18:S1–
S5.

Kohler D, Staniak M, Tsai T-H, Huang T, Shulman N, Bernhardt OM, MacLean BX,
Nesvizhskii AI, Reiter L, Sabido E, et al. 2023. MSstats Version 4.0: Statistical
Analyses of Quantitative Mass Spectrometry-Based Proteomic Experiments
with Chromatography-Based Quantification at Scale. *J. Proteome Res.*
22:1466–1482.

Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, Stecher G,
Hedges SB. 2022. TimeTree 5: An Expanded Resource for Species
Divergence Times. *Mol. Biol. Evol.* 39:msac174.

LaFlamme BA, Avila FW, Michalski K, Wolfner MF. 2014. A *Drosophila* protease
cascade member, Seminal Metalloprotease-1, is activated stepwise by male
factors and requires female factors for full activity. *Genetics* 196:1117–1129.

Laurinyecz B, Vedelek V, Kovács AL, Szilasi K, Lipinszki Z, Slezák C, Darula Z,
Juhász G, Sinka R. 2019. Sperm-Leucylaminopeptidases are required for
male fertility as structural components of mitochondrial paracrystalline
material in Drosophila melanogaster sperm. *PLOS Genet.* 15:e1007987.

Law STS, Nong W, Li C, Chong TK, Yip HY, Swale T, Chiu SW, Chung RY-N, Lam H-
M, Wong SYS, et al. 2025. Genome of tropical bed bug Cimex hemipterus
(Cimicidae, Hemiptera) reveals tetraspanin expanded in bed bug ancestor.
Insect Sci. 32:42–54.

Levine MT, Jones CD, Kern AD, Lindfors HA, Begun DJ. 2006. Novel genes derived
from noncoding DNA in Drosophila melanogaster are frequently X-linked and
exhibit testis-biased expression. *Proc. Natl. Acad. Sci.* 103:9935–9939.

Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, Lopez R,
Butcher S. 2024. The EMBL-EBI Job Dispatcher sequence analysis tools
framework in 2024. *Nucleic Acids Res.* 52:W521–W525.

Maric S, Donnelly SM, Robinson MW, Skinner-Adams T, Trenholme KR, Gardiner
DL, Dalton JP, Stack CM, Lowther J. 2009. The M17 Leucine Aminopeptidase
of the Malaria Parasite Plasmodium falciparum: Importance of Active Site
Metal Ions in the Binding of Substrates and Inhibitors. *Biochemistry* 48:5435–
5439.

Martens BM, McDonough-Goldstein CE, Otti O, Broschk S, Kullmann L, Reinhardt K,
Garlovsky MD. 2026. Novel female reproductive organ differentiates
postmating transcriptional response to insemination versus arrival of sperm in
bedbugs. :2026.03.17.707905. Available from:
https://www.biorxiv.org/content/10.64898/2026.03.17.707905v1

McCullough EL, Whittington E, Singh A, Pitnick S, Wolfner MF, Dorus S. 2022. The
life history of *Drosophila* sperm involves molecular continuity between male
and female reproductive tracts. *Proc. Natl. Acad. Sci.* 119:e2119899119.

McGowan S, Oellig CA, Birru WA, Caradoc-Davies TT, Stack CM, Lowther J,
Skinner-Adams T, Mucha A, Kafarski P, Grembecka J, et al. 2010. Structure of
the Plasmodium falciparum M17 aminopeptidase and significance for the
design of drugs targeting the neutral exopeptidases. *Proc. Natl. Acad. Sci.*
107:2449–2454.

Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE.
2023. UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci.*
32:e4792.

Miles LS, Adams R, Francioli YZ, Card DC, Castoe TA, Booth W. 2025. A
chromosome-level reference genome for the common bed bug, Cimex
lectularius, with identification of sex chromosomes. *J. Hered.* 116:382–388.

Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware
J, Flouri T, Beutel RG, et al. 2014. Phylogenomics resolves the timing and
pattern of insect evolution. *Science* 346:763–767.

Misra S, Wolfner MF. 2020. *Drosophila* seminal sex peptide associates with rival as
well as own sperm, providing SP function in polyandrous females. *eLife* 9.

Morrow EH, Arnqvist G. 2003. Costly traumatic insemination and a female counter-
adaptation in bed bugs. *Proc. R. Soc. Lond. B Biol. Sci.* 270:2377–2381.

Patlar B, Jayaswal V, Ranz JM, Civetta A. 2021. Non-adaptive molecular evolution of
seminal fluid proteins in Drosophila. *Evolution* 75:2102–2113.

Perry JC, Sirot L, Wigby S. 2013. The seminal symphony: how to compose an
ejaculate. *Trends Ecol. Evol.* 28:414–422.

Pini T, Nixon B, Karr TL, Teperino R, Sanz-Moreno A, Silva-Buttkus P da, Tüttelmann
F, Kliesch S, Gailus-Durner V, Fuchs H, et al. 2025. Towards a kingdom of
reproductive life - the core sperm proteome. *Reproduction* 169:e250105.

Plakke MS, Walker JL, Lombardo JB, Goetz BJ, Pacella GN, Durrant JD, Clark NL,
Morehouse NI. 2019. Characterization of female reproductive proteases in a
butterfly from functional and evolutionary perspectives. *Physiol. Biochem.
Zool.* 92:579–590.

Poggio MG, Iorio OD, Turienzo P, Papeschi AG, Bressa MJ. 2014. Heterochromatin
characterization and ribosomal gene location in two monotypic genera of
bloodsucker bugs (Cimicidae, Heteroptera) with holokinetic chromosomes and
achiasmatic male meiosis. *Bull. Entomol. Res.* 104:788–793.

R Core Team. 2024. R: A language and environment for statistical computing.
Available from: http://www.R-project.org

Rahnenfuhrer AA. 2022. topGO: Enrichment Analysis for Gene Ontology.

Reinhardt K, Kempke D, Naylor RA, Siva-Jothy MT. 2009. Sensitivity to bites by the
bedbug, Cimex lectularius. *Med. Vet. Entomol.* 23:163–166.

Reinhardt K, Naylor R, Siva–Jothy MT. 2003. Reducing a cost of traumatic
insemination: female bedbugs evolve a unique organ. *Proc. R. Soc. Lond. B
Biol. Sci.* 270:2371–2375.

Reinhardt K, Naylor R, Siva-Jothy MT. 2011. Male mating rate is constrained by
seminal fluid availability in bedbugs, *Cimex lectularius. PLOS ONE* 6:e22082.

Reinhardt K, Siva-Jothy MT. 2007. Biology of the Bed Bugs (Cimicidae). *Annu. Rev.
Entomol.* 52:351–374.

Reinhardt K, Wong CH, Georgiou AS. 2009. Detection of seminal fluid proteins in the
bed bug, Cimex lectularius, using two-dimensional gel electrophoresis and
mass spectrometry. *Parasitology* 136:283–292.

Roth S, Siva-Jothy MT, Balvín O, Morrow EH, Willassen E, Reinhardt K. 2024. The
evolution of female-biased genital diversity in bedbugs (Cimicidae). *Evolution*
78:329–341.

Singh A, Buehner NA, Lin H, Baranowski KJ, Findlay GD, Wolfner MF. 2018. Long-
term interaction between *Drosophila* sperm and sex peptide is mediated by
other seminal proteins that bind only transiently to sperm. *Insect Biochem.*
Mol. Biol. 102:43–51.

Sirot LK, Wong A, Chapman T, Wolfner MF. 2015. Sexual conflict and seminal fluid
proteins: a dynamic landscape of sexual interactions. *Cold Spring Harb.*
Perspect. Biol. 7:a017533.

Siva-Jothy MT. 2006. Trauma, disease and collateral damage: conflict in cimicids.
Philos. Trans. R. Soc. B Biol. Sci. 361:269–275.

Slack HD. 1939. Chromosomes of Cimex. *Nature* 143:78–78.

Snook RR. 2005. Sperm in competition: not playing by the numbers. *Trends Ecol.*
Evol. 20:46–53.

Stutt AD, Siva-Jothy MT. 2001. Traumatic insemination and sexual conflict in the bed
bug *Cimex lectularius*. *Proc. Natl. Acad. Sci.* 98:5683–5687.

Swanson WJ, Vacquier VD. 2002. The rapid evolution of reproductive proteins. *Nat.*
Rev. Genet. 3:137–144.

Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD,
Winther O, Brunak S, von Heijne G, Nielsen H. 2022. SignalP 6.0 predicts all
five types of signal peptides using protein language models. *Nat. Biotechnol.*
40:1023–1025.

Thongprem P, Evison SEF, Hurst GDD, Otti O. 2020. Transmission, Tropism, and
Biological Impacts of Torix Rickettsia in the Common Bed Bug Cimex
lectularius (Hemiptera: Cimicidae). *Front. Microbiol.* [Internet] 11. Available
from:
https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2020.6
08763/full

Ueshima N. 1967. Supernumerary chromosomes in the human bed bug, Cimex
lectularius Linn. (Cimicidae:Hemiptera). *Chromosoma* 20:311–331.

Usinger RL. 1966. Monograph of Cimicidae (Hemiptera-Heteroptera). Available from:
https://www.jstor.org/stable/3276661?origin=crossref

Vibranovski MD, Lopes HF, Karr TL, Long M. 2009. Stage-specific expression
profiling of Drosophila spermatogenesis suggests that meiotic sex
chromosome inactivation drives genomic relocation of testis-expressed
genes. *PLOS Genet.* 5:e1000731.

Wainwright SM, Hopkins BR, Mendes CC, Sekar A, Kroeger B, Hellberg JEEU, Fan
S-J, Pavey A, Marie PP, Leiblich A, et al. 2021. *Drosophila* Sex Peptide

controls the assembly of lipid microcarriers in seminal fluid. *Proc. Natl. Acad.*
Sci. 118:e2019622118.

Whittington E, Forsythe D, Borziak K, Karr TL, Walters JR, Dorus S. 2017.
Contrasting patterns of evolutionary constraint and novelty revealed by
comparative sperm proteomic analysis in Lepidoptera. *BMC Genomics*
18:931.

Whittington E, Zhao Q, Borziak K, Walters JR, Dorus S. 2015. Characterisation of
the Manduca sexta sperm proteome: Genetic novelty underlying sperm
composition in Lepidoptera. *Insect Biochem. Mol. Biol.* 62:183–193.

Wigby S, Brown NC, Allen SE, Misra S, Sitnik JL, Sepil I, Clark AG, Wolfner MF.
2020. The *Drosophila* seminal proteome and its role in postcopulatory sexual
selection. *Philos. Trans. R. Soc. B Biol. Sci.* 375:20200072.

Yang Z. 2007. PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Mol. Biol.*
Evol. 24:1586–1591.

Zuk M, Garcia-Gonzalez F, Herberstein ME, Simmons LW. 2014. Model systems,
taxonomic bias, and sexual selection: beyond *Drosophila*. *Annu. Rev.*
Entomol. 59:321–338.

799