Chemical Genetic Screen Identifies PSD3 as a Direct Substrate of NUAK1 that Regulates Dendritic Spine Maturation

Josilyn R. Sejd[1], Daphnée M. Marciniak[1], Moira A. Cornell[1], Angel Sondhi[1], Shao-En Ong[1] and Smita Yadav[1,2*]

[1]Department of Pharmacology, University of Washington, Seattle, United States.

[2]Institute of Stem Cell and Regenerative Medicine, Seattle, United States

* Corresponding author

Abstract

Novel *(nua)* Kinase 1 *(NUAK1)* encodes a serine-threonine protein kinase, mutations in which are associated with neurodevelopmental disorder. Direct phosphorylation targets of NUAK1 have been elusive hindering mechanistic understanding of its role in brain development. Here, we characterize autism-associated NUAK1 variants and shows their differential impact on catalytic activity and subcellular distribution. Next, we employed a chemical-genetic strategy with engineered ATP analog-sensitive kinase to identify over 30 direct phosphorylation targets of NUAK1. We demonstrate that Pleckstrin Homology and Sec7-domain containing protein 3 (PSD3) is a *bona fide* phosphorylation target of NUAK1. A guanine exchange factor (GEF) for ARF6 GTPase, PSD3 is phosphorylated by NUAK1 at S476. Expression of phosphodeficient PSD3 leads to aberrant activation of ARF6 and generation of $PI(4,5)P_2$ that accumulates in intracellular vesicles. In neurons, phosphomutant PSD3 leads to enhanced spine maturation in an ARF6 dependent fashion. This study reveals direct neuronal substrates of an autism risk gene NUAK1, and delineates a mechanism by which PSD3 phosphorylation regulates ARF6 activation and spine maturation.

Introduction

NUAK1 is a serine threonine protein kinase that belongs to the AMP-activated protein kinase (AMPK) family[1]. Despite the association of NUAK1 with pathophysiology of autism spectrum disorder[2,3], neurodegenerative disorder[4–6], fibrosis[7,8], immune dysfunction[9], and cancer[10–12], our understanding of this enigmatic kinase has been limited. Several factors have hindered progress in understanding the physiological function of NUAK1 in normal and disease states. NUAK1 knockout mice are embryonically lethal[13], precluding investigation of its physiological function. Moreover, both overexpression and loss of function of NUAK1 are associated with pathogenesis suggesting the importance of its regulated activity. For example, high NUAK1 expression is associated with reduced survival in pancreatic ductal adenocarcinoma and colorectal cancers[9,12]. It is also upregulated in multiple fibrotic tissues in humans, and NUAK1 inhibition in mice attenuates fibrosis[7,8,14]. NUAK1 stabilizes microtubule binding protein tau, and in a tauopathy mouse model its deletion protects against neurodegeneration[5]. In contrast, loss of NUAK1 in the brain causes defects in axonal development and heterozygous NUAK1 knockout mice exhibit social behavior deficits[15,16]. The fact that physiological phosphorylation targets of NUAK1 with a few exceptions have not been identified, has constrained understanding of NUAK1 function in normal and disease states.

The canonical isoform of human NUAK1 contains 661 amino acids with an N-terminal kinase domain and a largely disordered C-terminal region. Most studied in context of tumor suppression and cell proliferation[10,17–19], NUAK1 activity is regulated by LKB1 and NDR2 phosphorylation at the activation loop residue T211[20,21]. Phosphorylation by AKT at S600 has been shown to further increase the catalytic activity of NUAK1[22]. The LKB1-NUAK1 mediated signaling controls cell adhesion through regulation of myosin phosphatase complexes[17]. Three Gly-Iso-Leu-Lys (GILK) motifs in NUAK1 mediate its interaction with phosphatase complex MYPT1-PP1β that controls cell adhesion through regulation of myosin light chain[17]. An alternative pathway for NUAK1 phosphoregulation mediated by calcium-dependent activation of PKCα has been implicated in cancer cells lacking LKB1[23,24], highlighting several routes for achieving NUAK1 activation in distinct cellular context. Greater complexity to NUAK1 signaling arises from distinct subcellular compartments in which NUAK1 has been reported to function. A bipartite nuclear localization signal shuttles NUAK1 into the nucleus, with oxidative stress increasing its nuclear transport[25]. In cancer cell lines, NUAK1 appears to be predominantly nuclear, localized within nuclear speckles where it plays an important role in regulating splicing[24]. Therefore, in the cytoplasm, NUAK1 controls the function of PP1 complex by phosphorylating the MYPT1

regulatory subunit, while in the nucleus NUAK1 interacts with PNUTS/PPP1CB complexes

through phosphorylation of PNUTS to regulate cotranscriptional splicing[17,24].

NUAK1 function and signaling in the brain is less understood. Highly expressed in the

central nervous system, NUAK1 has been shown to be important for axonal and dendrite

development[15,16,26,27]. It is clear that catalytic activity of NUAK1 is essential for its function in the

neuronal development. For example, heterozygous NUAK1 knockout mice exhibit impaired axon

elongation which can be rescued by expression of WT NUAK1, but not by kinase dead NUAK1[16].

Further, small molecule kinase inhibitors against NUAK1 impair both axon and dendritic

development[26]. However, mechanisms through which NUAK1 mediated phosphorylation

promotes neuronal development are unknown largely because its direct phosphorylation targets

in the brain have not been identified. Further, whether autism associated NUAK1 variants impact

its catalytic activity have not been tested.

In our study, we biochemically characterized NUAK1 autism-associated variants and show

their differential impact on catalytic function and subcellular localization. Given the importance of

its kinase activity and dearth of mapped NUAK1 neuronal substrates, we employed an unbiased

chemical-genetic strategy to identify direct phosphorylation targets of NUAK1 in the mouse brain.

We reveal neuronal substrates of NUAK1 kinase that span cellular processes of endocytosis,

splicing, axon development, dendritic growth and synaptogenesis. From these candidate

substrate hits, we demonstrate that PSD3, a nucleotide exchange factor for the small GTPase

ARF6, is a *bona fide* substrate of NUAK1. Through a combination of in vitro biochemical assays,

phosphomutant analyses in neurons and super resolution imaging, we have uncovered an

essential role of NUAK1 mediated phosphorylation of PSD3 that controls ARF6 and PI(4,5)P2

dependent endocytic trafficking and dendritic spine maturation.

Results

Autism-Associated NUAK1 Variants Perturb Kinase Activity and Subcellular Distribution

NUAK1 has previously been reported to have differential subcellular localization to either the

nucleus and/or cytosol depending on cell/tissue type and nutrient conditions[8]. To study the role of

NUAK1 in brain development, we first investigated its localization in neuronal cells. Rat embryonic

hippocampal neurons were fixed at 14 days *in vitro* (DIV) and then immunostained with an

antibody against NUAK1 to visualize neuronal localization of endogenous NUAK1, along with an

antibody against the dendrite marker MAP2 *(Fig.1a)*. We found that endogenous NUAK1 is

localized to the cytosol within the soma, dendrites, and axon with no observable nuclear

enrichment within neurons (Fig.1a). In human brain, the Human Brain Transcriptome

database[28] shows NUAK1 is also highly expressed in early development across all measured

brain regions (NCX, HIP, AMY, STR, MD, CBC), and its expression persists into adulthood

(Fig.1b). Similarly, NUAK1 is highly expressed in neurons within the cortex, hippocampus, and

cerebellum as seen in ISH staining data by the Allen Brain Atlas (*Fig.1c*). Similar to endogenous

NUAK1, exogenously expressed GFP-tagged NUAK1 was also largely cytosolic. We next

interrogated the impact of NUAK1 autism-associated variants on its catalytic activity and neuronal

localization. Three NUAK1 variants have been reported to date; missense mutation Q161E within

the kinase domain, premature truncation mutant Q433*, and missense mutation A653V in the C-

terminal tail[29–31] (Fig.1d). Evolutionary analyses with ConSurf[32] revealed conservation of the

residues mutated in missense mutations. The Q161 residue shows 100% conservation across

150 species and there was 98% conservation of residue A653, with only the small amino acids

glycine or serine observed at this position (Fig.1e). Alphafold3[33,34] model of NUAK1 predicts A653

residue to be located in an alpha helix. We used HeliQuest[35] to probe the nature of this alpha

helix and found that residues 640-661 formed an amphipathic helix (Fig.S1a-b), with A653 located

in the hydrophobic face (Fig.1d and 1f).

Next, we tested whether autism-associated NUAK1 variants impacted its catalytic activity

by performing an *in vitro* kinase activity assay[36]. Briefly, GFP-tagged NUAK1 variants were

expressed in HEK293T cells, followed by lysis, immunoprecipitation, and incubation with Mg^{2+}

and ATP. We observed robust autophosphorylation with WT NUAK1, which was lost with the

catalytic lysine mutation K84M kinase dead (KD) mutation (Fig.1g-h). We observed a greater that

80-fold reduction in catalytic activity with the Q161E mutant, while the Q433* and A653V

mutations had unchanged kinase activity relative to WT NUAK1 (Fig.1g-h). To evaluate the impact

of NUAK1 variants on its subcellular distribution in neurons, we expressed GFP-tagged NUAK1

mutants in hippocampal neurons and found that while the Q161E and A653V variants had similar

localization to WT, expression of NUAK1 truncation variant Q433* dramatically induced nuclear

localization (Fig.1i). While the A653V mutation did not impact catalytic activity or nuclear

localization in neurons, we sought to ascertain if any other functions of this protein were impacted

by this mutation. While neuronal NUAK1 is excluded from the nucleus, NUAK1 in mitotically active

cells has well-established kinase activity-dependent roles in regulation of splicing[24,37]. WT NUAK1

has been shown to colocalize with SC35, a marker of spliceosomes, in the nucleus in various

contexts[24,37]. We expressed GFP-tagged NUAK1 (WT, Q161E, Q433* and A653V) in human

WTC11 (Corielle) iPSC-derived neural progenitor cells (NPCs) (Fig.S1b). In NPCs, WT NUAK1

localizes to the cytosol and nucleus, and is also enriched in nuclear speckles. In contrast,

truncation mutant Q433* NUAK1 is entirely absent from the cytosol and instead enriched in

nuclear speckles positive for SC35. Remarkably, the kinase deficient mutant Q161E NUAK1 does

not show colocalization with SC35 positive nuclear speckles. While catalytically active, in NPCs

the mutant A653V NUAK1 shows nuclear and cytosolic localization, but is absent from SC35

positive nuclear speckles (Fig.S1a). These experiments suggest that NUAK1 mutations might

lead to decreased phosphorylation of its neuronal substrates either through deficits in its catalytic

function (Q161E) or aberrant shift in its distribution between the cytoplasm to the nuclear speckles

(A653V and Q433stop).

Chemical-Genetic Screen for Identification of NUAK1 direct neuronal targets

NUAK1 phosphorylation targets in the developing brain have not been identified. We

employed a chemical-genetics approach[38–41] where we engineered a bulky ATP analog sensitive

(AS) NUAK1 that uniquely thiophosphorylates substrates that are covalently captured by

iodoacetamide beads and then identified by mass spectrometry. The advantages of this method

are that first, it uniquely identifies direct phosphorylation targets of the kinase of interest, and

second, the phosphorylation site on the substrate is revealed that can be then modified to study

physiological impact of phosphorylation. To engineer the AS-NUAK1 kinase, we mutated the

conserved gatekeeper residue M132 in the kinase domain of NUAK1 (Fig.2a) to a smaller amino

acid Alanine or Glycine. With the enlarged hydrophobic nucleotide binding pocket, the engineered

AS-NUAK1 can utilize N^6-substituted bulky ATPɣS molecules to thiophosphorylate its substrates.

As a negative control, we also generated kinase dead (KD) NUAK1 with a Lys84Met (K84M)

mutation, abolishing the conserved catalytic lysine residue that is responsible for coordinating the

ɣ-phosphate of ATP. We bacterially purified the kinase domain of NUAK1 (residues 1-433) with

either active wildtype (WT), kinase-dead (K84M), and analog sensitive (M132G) mutations

(Fig.S2a-c). We next combinatorially tested different NUAK1 AS mutants and bulky ATP analogs

and determined that N^6-furfuryl-ATPɣS is optimally utilized by M132G NUAK1 for phosphorylation

but not by WT NUAK1 (Fig.2b-c). We confirmed that full-length NUAK1 M132G shows preferential

utilization of N^6-furfuryl-ATPɣS using full-length HA-NUAK1 immunoprecipitated from HEK293T

cells (Fig.S2d-e).

Since motifs outside the NUAK1 kinase domain such as the "Gly-Iso-Leu-Lys" (GILK) motif

are important for substrate binding[4,16], we utilized full-length AS-NUAK1 (M132G) to identify

thiophosphorylated substrates from mouse brain lysate. We affinity purified full-length AS-
NUAK1(M132G) and kinase-dead NUAK1(K84M) from HEK293T cells using HA-conjugated
beads and incubated them with mouse brain lysate in the presence of furfuryl ATPɣS (Fig.2d).
Compared to the lysate only or kinase dead controls, we observed robust signal for
thiophosphorylated substrates identified by a phosphothioester antibody in the AS samples but
not in the KD or lysate only samples. After validating that full-length HA-NUAK1 AS is able to
effectively phosphorylate substrates using furfuryl ATPɣS, we scaled up our system for mass
spectrometry analysis. Immunoprecipitated NUAK1 was incubated with mouse brain lysate and
furfuryl ATPɣS to allow AS-NUAK1 to thiophosphorylate its substrates. Samples were then
denatured with urea, reduced with TCEP, and digested to generate peptides. The resulting
thiopeptides were then captured on iodoacetamide resin and treated with oxone to specifically
elute peptides containing a thiophosphate moiety, while cysteine (thiol) containing peptides were
retained on beads. Eluted phosphopeptides were separated using liquid chromatography and
analyzed using mass spectrometry (Fig.2e).
177
Identification of NUAK1 Consensus Sequence and Direct Neuronal Substrates
Mass spectrometry analyses of five replicate samples revealed 972 total phosphopeptides. Our
criteria for narrowing candidate hits to be further tested as a *bona fide* substrate of NUAK1
following the chemical genetic screen was that the substrate appears in at least three out of five
AS replicates and was absent in (0/5) all kinase dead (KD) replicates. From our list of 972 total
phospho-serine and phospho-threonine peptides, 31 occurred in at least 3/5 AS replicates and
were absent in all 5 KD samples. These 31 potential NUAK1 substrate proteins were analyzed
for their cellular role(s) which binned into the following categories: endocytic recycling, splicing,
cytoskeletal reorganization, as well as axonal, dendritic, and synaptic development (Fig.3a).
Proteins were assigned to one or more categories based on their previously established roles in
each cellular process and color coded with either a single color or a gradient of 2-4 colors to
indicate their pleiotropic functions (Fig 3a). We used the sequence of these 31 peptides to
generate a sequence logo using Phosphosite Plus to show the preferred phosphorylation motif of
NUAK1 in our mass spectrometry experiment (Fig.3b). Next, we further shortlisted candidates
where the phosphosite sequence fit the experimentally determined NUAK1 stringent consensus
sequence of a basic residue (R/K) at the -3 position[42]. Out of the 31 phosphopeptides, 17 fit the
stringent consensus of a basic residue at the -3 position (Fig.3c). The substrate phosphopeptides
are color coded based on how well the residue fits our optimal NUAK1 phosphorylation motif at
positions -5 and +4 (purple), basic residues at positions -4, -3 (cyan), and preferred residue serine

at -2 position (chartreuse). The central phosphoacceptor at position 0 is colored in dark green for

serine which is preferred over threonine (light green) (Fig.3c). The function and subcellular

localization of the 17 putative NUAK1 substrates is listed (Fig.3c) and both nuclear and cytosolic

proteins were identified in the chemical genetic screen.

PSD3 is a Bona Fide Substrate of NUAK1 Kinase

We next focused on our top candidate in the screen for NUAK1 substrates which was Pleckstrin

Homology Domain and SEC7 domain-containing protein 3 (PSD3). A multidomain protein, PSD3

consists of a Pleckstrin Homology (PH) domain which binds specifically to phospholipid PI(4,5)P$_2$

and a SEC7 domain that acts as an guanine exchange factor (GEF) for small GTPase ARF6[43,44].

A coiled coil (CC) region following the PH domain enhances membrane association of the PH

domain and has been shown to bind actin[45,46]. These 3 functional domains are present in the

alternatively spliced human isoforms of PSD3 (Fig.4a), the canonical isoform 1 with 1048 amino

acids (Uniprot #Q9NYI0), isoform 2 with a single amino acid deletion with 1047 amino acids

(Uniprot #Q9NYI0-2), and isoform 3 which has 513 amino acids (Uniprot #Q9NYI0-3). The

putative NUAK1 phosphorylation site is present in the CC region shared by all three isoforms

(Fig.4a). Previous studies have shown that the predominant isoform of PSD3 expressed in the

mouse brain is 56kD, which corresponds to isoform 3 [45,47]. PSD3 isoform 3 amino acids 12-513

are identical to the C terminal region of PSD3 isoform 1, with only the first 11 amino acids of

isoform 3 differing. We utilized ConSurf[32] to perform evolutionary conservation analysis, which

revealed that the C-terminus of PSD3 is highly conserved, specifically around the PH, Sec7, and

CC domains. Further the phosphosite S476 and the motif surrounding it are extremely conserved

(Fig.4b and Fig.S3a). Because our MS-identified phosphosite is located within the evolutionarily

conserved C-terminus of both isoforms, we elected to utilize the more abundant shorter isoform

3 for further analyses.

To assess whether affinity purified human PSD3 was phosphorylated by NUAK1, we

immunoprecipitated V5-tagged PSD3-WT or phosphomutant V5-PSD3-S476A from HEK293T

cells and incubated them with purified NUAK1-WT or NUAK1-KD in an in vitro kinase reaction

with ATP. We observed robust phosphorylation of WT PSD3 by WT NUAK1, which was completely

lost in the PSD3 S476A mutant (Fig.4c-d). We next determined whether NUAK1 and PSD3

interacted with each other. In an affinity pulldown assay with HA-beads, PSD3 was co-

immunoprecipitated by HA-tagged NUAK1 but not in the control condition (Fig.4e-f). Similarly,

HA-NUAK1 was co-immunoprecipitated by V5-PSD3 (Fig.4g-h), demonstrating that PSD3 and

NUAK1 are interacting proteins. To ensure that a co-immunoprecipitating kinase was not

responsible for phosphorylating PSD3, we confirmed that bacterially expressed and purified

NUAK1 and PSD3 would be sufficient to reconstitute the phosphorylation. We expressed and

purified the GST-tagged C-terminal tail of PSD3 which is identical in isoforms 1 and 3 (amino

acids 378-513, isoform 3/amino acids 912-1048, isoform 1) encompassing the coiled coil domain

and C-terminal tail, and henceforth referred to as 'PSD3-CC-Tail' (Fig.S2a-c). Incubation of

purified NUAK1 WT with PSD3-CC Tail in a kinase activity buffer led to phosphorylation of PSD3-

CC-Tail which was abolished in the PSD3 S476A mutant. No phosphorylation was observed as

expected with the kinase dead NUAK1 (Fig.4i and 4j).

Next, we used Alphafold3 structure prediction to understand the mechanism through which

NUAK1 binds its substrates. First, we performed a prediction of NUAK1 in complex with its known

substrate MYPT1, along with the requisite adaptor Protein Phosphatase 1 beta (PP1β)[48]. In this

structure, the region of MYPT1 from amino acids 435-455, centered around the phosphosite at

S445, docks into the kinase domain of NUAK1, which is shown as an electrostatic surface model

(Fig.4k). The phospho-acceptor S445 is located adjacent to the binding site for ATP, while basic

residues at the -3 and -4 position bind to acidic patches in the NUAK1 kinase domain. The

aliphatic isoleucine residues at -5 and +4 positions in MYPT1 dock into small pockets within the

NUAK1 kinase domain. To determine whether NUAK1 had a similar binding mode towards the

PSD3 phosphosite, we next modeled NUAK1 in complex with full length PSD3 isoform 1 (Fig.4l).

Here, the region surrounding the PSD3 S1011(isoform1)/S476(isoform3) phosphosite docks into

the catalytic cleft of NUAK1, in a manner highly similar to MYPT1. The phospho-acceptor

S1011/S476 of PSD3 is docked directly next to the ATP-binding site, and the basic residues at

positions -4 and -3 are aligned with the acidic patches of the NUAK1 kinase domain. Finally, the

aliphatic leucine residues at positions -5 and +4 fit into the pockets in the NUAK1 kinase domain

in a similar manner to the respective residues in MYPT1. Altogether, our computational modeling

highlights the similarity of the binding conformation of NUAK1 towards known substrate MYPT1

and PSD3 discovered in our chemical genetic screen.

NUAK1 Phosphorylation of PSD3 Regulates Dendritic Spine Maturation

PSD3 is widely expressed in the central nervous system both during and after developmental

time periods. In mouse brain, the highest expression is within the hippocampus and cerebral

cortex (Fig.5a), while in human brain PSD3 expression is broad including hippocampus, cortex,

striatum and cerebellum. PSD3 expression increases during embryonic brain development,

peaks at birth and is maintained at high levels throughout the human lifespan (Fig.5b). PSD3 is a

guanine exchange factor specifically for small GTPase ARF6[44], which has important roles in

dendritic spine development[49,50]. To determine how NUAK1-mediated PSD3 phosphorylation

impacts neuronal development, we transfected DIV12 rat hippocampal neurons with membrane

marker myr-tdTomato with either GFP-tagged wildtype PSD3 or the S476A phosphomutant and

imaged at DIV14. WT-PSD3 localized throughout the neuron including soma, dendrites, axons

and was particularly enriched in mushroom-shaped spines (Fig.5c-d). Notably, we observed that

neurons expressing the phosphomutant S476A-PSD3 showed significantly increased enrichment

of PSD3 within dendritic spines concomitant with increased number of mushroom spines (Fig.5c

and 5d). Percent dendritic spines enriched in PSD3 increased from a mean of 33.8% (n=16 from

three experiments) in WT-PSD3 expressing neurons to 63.8% (n=17 from three experiments,

p<0.0001) in neurons expressing the S476A phosphomutant (Fig.5e). To further analyze the

impact of PSD3 phosphorylation on dendritic protrusions, we binned protrusions into four classes:

stubby, thin, filopodia and mushroom. While there were no significant differences in the percent

of stubby and thin protrusions, the filopodial protrusions decreased from 13.9% to 7.5% in

neurons expressing the phosphomutant PSD3 S476A (n=56 neurons from three experiments,

SEM=1.33, p<0.05) and mushroom spines increased from 36.1% in WT (n=65, SEM=2.08) to

48% in S476A mutant expressing neurons (n=56, SEM=2.0, p<0.0001) (Fig.5f). No significant

differences were observed in total protrusions between the wildtype and S476A mutant PSD3

expressing neurons (Fig.5g). These results suggest that NUAK1 mediated phosphorylation of

PSD3 at S476 does not impact formation of new protrusions but rather controls the maturation of

filopodial protrusions into mature mushroom shaped spines enriched in PSD3.

Phosphomutant PSD3 S476A Disrupts ARF6 Mediated Endocytosis

Localization of PSD3 to the plasma membrane is required for conversion of GDP-bound ARF6 to

active GTP-bound ARF6. The C-terminal tail of PSD3 containing both the PH domain and the

coiled-coils is required for its plasma membrane association[45]. Since S476, the NUAK1

phosphosite on PSD3 lies in the C terminal tail, we tested whether PSD3 phosphorylation

regulates ARF6 localization or activation. Expression of GFP-tagged wildtype PSD3 along with

mScarlet-tagged wildtype ARF6, led to the recruitment to active ARF6 to the membrane (Fig.6a,

top row), while expression of catalytically dead PSD3 mutant E114K, as expected, was unable to

recruit and activate ARF6 (Fig.6a, second row). Expression of phosphomutant PSD3 recruited

ARF6 to the plasma membrane but led to an aberrant accumulation of large vacuole-like

intracellular vesicles positive for both ARF6 and PSD3 (Fig.6a, third row). Since constitutively

active ARF6 mutant Q67L leads to similar accumulation of GTP-bound ARF6 that is unable to
recycle back, we hypothesized that S476A phosphomutant PSD3 causes aberrant ARF6
activation or prevented the conversion of ARF6 GTP back to its GDP form. To test this hypothesis,
we tested whether dominant negative ARF6 mutant T44N would rescue the effect of PSD3 S476A
mutant. Indeed, co-expression of PSD3 S476A along with ARF6 T44N did not cause internal
accumulation of PSD3 and ARF6 positive vesicles (Fig.6a, bottom row). Area of internalized
PSD3/ARF6 positive vesicles increased from 9.6% in WT-PSD3/WT-ARF6 expressing cells to
20.9% in S476A-PSD3/WT-ARF6 expressing cells (n=26, p<0.005) (Fig.6b). Expression of
S476A-PSD3 with DN-ARF6 lead to almost complete rescue to 2.2% (n=20, p<0.0001) similar to
the inactive E114K PSD3 expression. GTP-bound ARF6 activates phosphatidylinositol 4-
phosphate 5-kinase (PIP 5-kinase) that generates phosphatidylinositol 4,5-bisphosphate
(PI(4,5)P2) which is important for ARF6 dependent endocytosis[51] . Expression of a GTP
hydrolysis-resistant ARF6 mutant or overexpression of PIP 5-kinase alpha induces the formation
of PI(4,5)P2-positive actin-coated vacuoles that are unable to recycle membrane back to the
plasma membrane[51] . These vesicles have been shown to contain ARF6-dependent cargo that
remains trapped as they are unable to recruit early endosome markers RAB5 and EEA1 to recycle
back[52]. To further test our hypothesis that loss of C-terminal PSD3 phosphorylation leads to
aberrant ARF6 activation, we determined if levels of PI(4,5)P2 and recruitment of RAB5 to
internalized membrane would be altered in cells expressing S476A mutant. We expressed WT or
S476A phosphomutant PSD3 in HEK293T cells along with the PI(4,5)P2 sensor GFP-C1-
PLCdelta-PH[51] and endosomal marker BFP-tagged RAB5. In cells expressing WT-PSD3,
PI(4,5)P2 was normally localized to the cell membrane, however, cells expressing phosphomutant
PSD3 developed numerous large internal vacuoles that were positive for PSD3 and PI(4,5)P2 but
devoid of early endosome marker RAB5 (Fig.6c). Total area of internalized vesicles positive for
PI(4,5)P2 increased from 4.7% in WT-PSD3 expressing cells to 25.8% in cells expressing the
S476A mutant PSD3 (n=23 cells from three experiments, p<0.0001) (Fig.6d). Visualization of
single z frames images of internalized PIP2 vesicles through super resolution
microscopy[53] indicates clear absence of RAB5 on the PSD3 / PI(4,5)P2 positive vesicles (Fig.6e).
These data together indicate the phosphorylation of PSD3 at S476 is important for ARF6
mediated endosomal trafficking.
329
Discussion
In this study, we utilize a chemical-genetic screen to reveal phosphorylation targets of the
autism-associated kinase NUAK1 in the brain. This unbiased approach allowed us to map the

basophilic consensus motif preferred by NUAK1. We uncovered direct substrates of NUAK1 in

cellular processes NUAK1 has been previously implicated in including splicing, axon growth, and

dendrite development, as well as discovered new functions of NUAK1 in regulation of endocytosis

and spine maturation (Fig.7a). Among the identified substrates, we focused on the ARF6-specific

guanine exchange factor PSD3 and delineated mechanistically how its phosphorylation by

NUAK1 negatively controls ARF6-dependent endocytosis and spine maturation (Fig.7b and 7c).

We report the biochemical characterization of autism-associated NUAK1 variants and show

that NUAK1 mutations have differential impact on catalytic activity and subcellular localization.

The residue Q161 in the αE-helix in the C-lobe is under high evolutionary constraint and makes

key contacts with residues N191 and I192 in the β8 strand that stabilize the catalytic loop of

NUAK1. We found that the autism variant Q161E led to a significant decrease in the catalytic

activity of NUAK1. The corresponding Q residue in protein kinase A (PKA) is Q149 which has

been shown to allosterically regulate the DFG loop during transitions between catalytic active and

inactive states[54,55]. The premature termination variant Q433* is predicted to escape nonsense

mediated decay, and our data shows that it has intact catalytic activity. However, on expression

in neurons and dividing cells, we found that the truncated NUAK1 Q433* is trapped in nuclear

speckles positive for SC35. The third NUAK1 disease variant A653V also does not impact

catalytic activity but decreases NUAK1 association with nuclear speckles. Our structural analyses

suggests that the residue A653 lies in an amphipathic helix within the C-terminal tail. Mechanisms

through which the latter two variants have opposing impact on NUAK1 association with nuclear

speckles remains to be determined. Collectively, our data suggest that heterozygous autism-

associated mutations in NUAK1 are predicted to either decrease substrate phosphorylation or

alter substrate availability by shifting the localization of the active kinase between cytoplasm and

nuclear speckles, two distinct locations where NUAK1 likely phosphorylates non-overlapping set

of substrates. This is highlighted in our dataset of identified NUAK1 substrates that include both

nuclear and cytoplasm enriched proteins.

To shortlist the putative substrate candidates identified in our chemical-genetic screen, we

used the following criteria. We reasoned that a true NUAK1 substrate would: (1) be

phosphorylated by NUAK1 *in vitro,* (2) show abrogated phosphorylation upon mutation of the

identified phospho-site, (3) biochemically interact with the purified kinase, and that (4) purified

NUAK1 and substrate in vitro would be sufficient to reconstitute phosphorylation at the identified

phosphosite. PSD3, an ARF6 specific GEF, identified in our screen meets all four of the above

criteria consistent with a bona fide NUAK1 substrate. PSD3 isoform 3 is phosphorylated at S476

(S1001 isoform 1), an evolutionarily conserved site that fits the NUAK1 consensus motif. Notably,

mutation of the S476 residue in PSD3 to alanine abolishes all phosphorylation by NUAK1,

indicating it is the only site within PSD3 that is phosphorylated by NUAK1. The phosphosite is

situated in the C-terminal tail of PSD3 following the PH domain and the coiled coil region which

binds cortical actin. The C-terminus of PSD3 mediates its plasma membrane association which

is required for activation of ARF6 at the membrane via the catalytic SEC7 domain. While it is

unlikely that phosphorylation at S476 directly impacts SEC7 domain mediated GEF activity, it can

indirectly increase ARF6 activation by promoting membrane or actin association of PSD3.

Accumulation of PI(4,5)P2 positive vesicles that fail to acquire endosomal markers such as RAB5

in cells expressing phosphomutant PSD3 are phenotypically similar to expression of ARF6 Q67L

mutant which cannot hydrolyze GTP. Another possibility is that phosphorylation of the PSD3 tail

does not impact ARF6 activity but regulates recruitment of ARF6 GAPs that convert GTP-bound

ARF6 to GDP.

It is important to note that copy number variation and missense mutations in PSD3 have been

identified in patients with autism spectrum disorder[29,30,56]. ARF6 GTPase is an important regulator

of actin dynamics and membrane trafficking, and plays an important role in neurite development,

axon growth, and synaptic plasticity[57]. PSD3 is enriched within dendritic spines, and our data

shows that abolishing NUAK1 mediated phosphorylation at S476 further increases PSD3 spine

enrichment, as well as causes loss of filopodial protrusions and stabilization of mature mushroom

dendritic spines. This is consistent with the effect of expression of fast cycling ARF6 mutant T157A

which increases the conversion of filopodia into mature spines[50]. The impact of PSD3 variants

associated with autism spectrum disorder on spine stability and ARF6 activity has not been tested

and will be subject of future investigation. In addition to autism, both NUAK1 and PSD3/ARF6

have been implicated in cancer progression, fatty liver and metabolic disease[19,58–62]. Our study

raises the interesting possibility that NUAK1 mediated phosphorylation of PSD3 underlies diverse

physiological effects of ARF6 dependent endocytic trafficking.

Acknowledgement

We are grateful for funding support by the National Institutes of Health to SY (R01MH121674,

R01MH130336), JS (T32 GM007750) and SEO (R01GM129090, R21CA288806). Funding for

the purchase of ELYRA7 Lattice SIM microscope was generously provided by the Murdock

Foundation Equipment Grant to SY with matching funds from the University of Washington. This

study used an EASY-nLC1200 UHPLC and Thermo Scientific Orbitrap Fusion Lumos Tribrid mass
spectrometer purchased with funding from a National Institutes of Health SIG grant
S10OD021502.
404

Author Contributions
J.S. designed and performed all experiments under the supervision of S.Y. D.M. ran the
proteomics samples on the LC-MS/MS and performed data analysis under the supervision of
S.E.O. A.S. performed dendritic spine morphology and vesicle area quantifications. M.C.
performed Alphafold predictions and Pymol modeling. Manuscript was written by J.S. and S.Y.,
and edited by all authors.
411

Methods
Expression Constructs, Cloning and Reagents
Full-length human NUAK1 was PCR amplified from K84M kinase-DEAD Nuak1-3x Flag
(Addgene #97219) and inserted into vector sfGFP-C1 (Addgene #54579) or PRK5-HA (PMID:
28065648) using PCR amplification followed by Gibson Assembly (NEB). The K84M mutation
was corrected using primers to generate WT NUAK1. All mutants of NUAK1 (K84M, M132A,
M132G, Q161E, Q433*, A653V) were generated using PCR amplification followed by Gibson
assembly (NEB). NUAK1 Q433* was cloned into the bacterial expression vector 8xHis-MsyB
(HM) (Addgene #112570) to generate HM-NUAK1 1-433 using PCR amplification followed by
Gibson assembly (NEB). The K84M or M132G mutations were introduced in this plasmid using
PCR amplification followed by Gibson assembly (NEB). Full-length human PSD3 isoform 1
(canonical) was obtained from DNASU and inserted into an N-terminal V5 mammalian expression
using Gateway cloning (Thermo). PSD3 isoform 3, which is 513 amino acids long, shares all
except the first 11 amino acids with isoform 1. PSD3 isoform 3 was cloned using PCR
amplification to insert the 11 amino acids at the N-terminus, followed by Gibson assembly (NEB).
PSD3-isoform 3 E114K and S476A point mutations were cloning using PCR amplification followed
by Gibson assembly (NEB). V5-PSD3-iso3-sfGFP was cloned by inserting sfGFP from sfGFP-C1
vector (Addgene #54579) at the C-terminus of PSD3 followed by a stop codon using PCR
amplification and Gibson Assembly. mCherry-V5-PSD3-iso3 was cloned by inserting the mCherry
from mCherry-Rab5 (Addgene #49201) upstream of V5-PSD3-iso3 using PCR amplification
followed by Gibson Assembly (NEB). GST-PSD3-coiled coil tail was generated by PCR
amplification of amino acids 378-513 of PSD3-isoform 3 followed by Gibson Assembly (NEB) with
PCR-amplified pGex-4T3 vector (Addgene #31669).

Arf6-mNeonGreen was obtained from Addgene, and the mNeonGreen was exchanged for
mScarlet from mScarlet-Rab7 (Addgene #169068) using PCR amplification and Gibson Assembly
(NEB). Arf6-mScarlet T44N (DN) mutation was introduced using PCR amplification followed by
Gibson Assembly (NEB).

Construct name	Source
Myristoyl-tdtomato	Yadav et al, *Neuron*, 2017 (PMID 28065648)
PRK5-HA vector	Yadav et al, *Neuron*, 2017 (PMID 28065648)
Rab5b-BFP	Addgene #49147
Arf6-mNeonGreen WT	Addgene #162027
Arf6-mScarlet (WT, T44N)	Cloned in this study
PSD3 in pDONR221	DNASU #HsCD00296934
GFP-C1-PLC-delta-PH	Addgene # 21179
K84M kinase-DEAD Nuak1-3x Flag	Addgene #97219
pExp-MsyB	Addgene #112570
GST-PSD3-CC-tail	Cloned in this study
PRK5-HA-NUAK1 (WT, K84M, M132G, M132A)	Cloned in this study
sfGFP-C1-NUAK1 (WT, K84M, Q161E, Q433*, A653V)	Cloned in this study
pGEX-PPP1R12A wt	Addgene #31669
His-MsyB-NUAK1 1-433 (WT, K84M, M132G)	Cloned in this study
sfGFP-C1	Addgene # 54579

Cell Line Maintenance and Transfection

HEK293T cells were grown in Dulbecco's modified Eagle's medium (Thermo Fisher
Scientific, Gibco) with 10% fetal bovine serum (Axenia BioLogix) and 1% penicillin-streptomycin
(Invitrogen). Cells were maintained at 5% CO_2 and 37°C and passaged every 3 to 4 days. Cells
were tested for mycoplasma using the Abm PCR-based kit. For biochemical experiments,
HEK293T cells were transfected with 0.5-1 ug plasmid DNA using the JetOptimus system
(Genesee Scientific) following the manufacturer's instructions. For confocal imaging, HEK293T
cells were transfected with 100-500 ng plasmid DNA using Lipofectamine 2000 and Optimem
(Thermo Scientific) according to the manufacturer's instructions.

Primary neurons were obtained from dissociating hippocampi obtained from embryonic
day 18 (E18) Sprague-Dawley rat (Envigo). Animal work was performed under University of

Washington Institutional Animal Care and Use Committee approved protocol 4421-01. Rat

embryos were trypsin-dissociated and plated at a density of 150,000 neurons per 18-mm glass

coverslip (Thermo Fisher Scientific) or 450,000 neurons per 35-mm glass-bottom dish (MaTek),

which were coated with poly-D-lysine (0.06 mg/ml; Sigma-Aldrich) and laminin (0.0025 mg/ml;

Sigma-Aldrich). Plating medium containing heat-inactivated 10% fetal bovine serum (HyClone),

0.45% dextrose, sodium pyruvate (0.11 mg/ml), and 2 mM glutamine in minimum essential

medium with Earle's Balanced Salt Solution was used. Medium was changed 4 hours after

seeding neurons into maintenance medium consisting of Neurobasal medium (Invitrogen), 2%

B27 (Invitrogen), penicillin (50 U/ml) and streptomycin (50 µg/ml), and 0.5 mM Glutamax

(Thermo). Half of the medium was replaced with new maintenance medium every 3 to 4 days.

For exogenous gene expression, neurons were transfected with 0.5 to 1.0 µg of plasmid DNA per

coverslip in a 12-well plate (or 1 to 2 µg of DNA per 35-mm dish) using Lipofectamine 2000

(Invitrogen) following the manufacturer's guide.

WTC11 iPSCs (Corielle) were differentiated into neuroprogenitor cells (NPCs) as reported

previously[63] . Stem cells were treated with 1U/mL Dispase, lifted using a cell scraper,

resuspended in mTeSR media (STEMCELL Technologies) containing 10µm ROCK Inhibitor

(Selleck Chemical), and then transferred to uncoated T25 flasks for embryoid body (EB)

formation. After 24–48 hours (Day 0), EBs were gravity settled and transferred to new uncoated

T25 flasks in neural media (DMEM/F12 (Gibco), 1% N-2 supplement (Gibco), 1% Non-essential

amino acids (Gibco), 2µg/mL Heparin, 1% Penicillin/Streptomycin [Gibco]) containing small

molecule TGF-β/SMAD inhibitors, SB431542 (5µM; Peprotech) and LDN193189 (0.25µM;

Peprotech). On Day 3, EBs were transferred to Matrigel (1:50 dilution; Corning)-coated 6 well

plates in fresh neural media with SB and LDN. Media was changed with neural media without SB

and LDN every other day or as needed until Day 11 for formation of neural rosettes. On Day 11,

neuroepithelia were mechanically lifted using a cell scraper and transferred to uncoated T25

flasks in neural media. Neurospheres were fed with neural media every other day, or as needed,

until Day 25. On Day 25, neurospheres were plated onto Matrigel-coated plates in STEMDiff™

Neural Progenitor Media (STEMCELL Technologies) and NPCs were obtained. NPCs were tested

for neural progenitor markers Nestin (Invitrogen) and PAX6 (Sigma).

Protein Purification

HM-NUAK1 purification

NUAK1 (WT, KD, or M132G) were inserted into 8x-His-tagged pExp-MsyB (HM) vector for

bacterial expression and purification. Due to poor expression of full-length NUAK1 1-661, we

elected to use NUAK1 1-433, which is kinase active but contains less unstructured domains than

full-length NUAK1. Expression of NUAK1 1-433 was dramatically improved over full length, so we

moved forward with purification of HM-NUAK1 1-433 WT, K84M, and M132G. HM-NUAK1 1-433

WT, K84M, or M132G were transformed into BL21 *E. coli* and sequenced using full-plasmid

sequencing (Plasmidsaurus) to confirm the sequences were correct. 50 mL cultures of HM-

NUAK1 were grown overnight with Carb and used to inoculate 1L cultures, which were grown at

37C with 220 rpm shaking until Optical Density (OD) of 0.6-0.8 was reached. At this point, the

incubator was cooled to 20C (room temperature), and IPTG was added to all cultures for a final

concentration of 0.4mM. Protein induction was performed overnight at 20C with 220 rpm shaking.

Bacteria were pelleted the next day by spinning at 3,200 rcf at 4C for 15 minutes, followed by

removal of LB and addition of 30 mL 1x DPBS. The cultures suspended in PBS were transferred

to 50 mL conical tubes and spun at 3,200g at 4C for 15 minutes. The PBS was then removed,

and 25 mL lysis buffer (20 mM Tris pH 7.5, 200 mM NaCl, 0.4 mM TCEP, 20 mM imidazole, 1x

cOmplete protease inhibitor (Sigma), 0.1% PMSF, and 10% glycerol) was added to the tube.

Bacteria were lysed using sonication at 25% power for 3.5 minutes, followed by centrifugation at

24,500g for 60 minutes at 4C to pellet debris. Following centrifugation, the supernatant was added

to 1 mL HisPur Ni-NTA resin (Thermo) for 60 min at 4C with rotation to bind HM-NUAK1 to the

beads. After binding, beads were added to a fritted column and the supernatant allowed to drain

by gravity. The beads were then washed with 20 mL lysis buffer, followed by the addition of lysis

buffer + 200 mM imidazole. This high imidazole buffer was incubated with the beads for 5 minutes,

after which eluted protein was collected every 10 minutes for a total of 6 elutions, and quantified

using Pierce 600nm protein assay (Thermo). The eluted protein was then incubated with

sulphopropyl (SP) beads (GE healthcare) to enrich for acidic proteins for 1 hour rotating at 4C,

followed by elution with 450mM NaCl. Finally, proteins were concentrated to a final concentration

of ~10 ug/mL using concentrators with a 30kD molecular weight cutoff (Thermo). Following

concentration, protein aliquots were flash frozen in liquid nitrogen and stored at -80C. All

purification samples were assayed for purity by SDS-Page, Coomassie staining, and Western

blotting with anti-His-HRP antibody (Proteintech).

GST-PSD3 purification

PSD3-CC tail (WT or S>A) was cloned into the pGEX-4T3 bacterial expression vector,

transformed into BL21 *E. coli*, and sequenced using full-plasmid sequencing (Plasmidsaurus). 25

mL overnight cultures of GST-PSD3 were grown overnight with Carb and used to inoculate 1L

cultures, which were grown at 37C with 220 rpm shaking until Optical Density (OD) of 0.6-0.8 was

reached. At this point, the incubator was cooled to 30C, and IPTG was added to all cultures for a

final concentration of 0.3mM. Protein induction was performed for 2 hours at 30C with 220 rpm
shaking. Bacteria were pelleted by spinning at 3,200 rcf at 4C for 15 minutes, followed by removal
of LB and addition of 30 mL 1x DPBS. The cultures suspended in PBS were transferred to 50 mL
conical tubes and spun at 3,200g at 4C for 15 minutes. The PBS was then removed, and samples
were resuspended in 7 mL lysis buffer (50mM Tris pH 8.0, 200mM NaCl, 5mM EDTA, 10%
glycerol, 5mM DTT, 1x cOmplete protease inhibitor (Sigma), 0.1% PMSF). 4 mg of lysozyme was
added to each sample, followed by incubation for 30 minutes on ice with occasional mixing. After
this step, Triton was added to the sample for a final concentration of 1%, followed by lysis with
sonication as described above. After lysis, samples were centrifuged at 24,500g for 60 minutes
at 4C to pellet debris. Columns for batch binding were prepared using 500 uL of glutathione resin
(Thermo) and 10 mL binding buffer (50mM Tris pH 8.0, 200mM KCl, 1mM DTT). The buffer was
removed prior to the addition of the post-centrifugation bacterial supernatant to the beads. The
binding of GST-PSD3 to the beads was performed overnight at 4C with rotation. The next day,
columns were washed 2x with 15 mL wash buffer (50mM Tris pH 8.0, 200mM KCl, 0.1% Tween-
20 1mM DTT), followed by 2x washes with 1 mL binding buffer to remove detergent. Samples
were then eluted in binding buffer + 25 mM glutathione in 500 uL fractions every 10 minutes, for
a total of 6 elutions. Protein concentration was quantified using Pierce 600nm protein assay
(Thermo), and purification samples were assayed for purity by SDS-Page, Coomassie staining,
and Western blotting with anti-GST-HRP antibody (Thermo). Following protein quantification,
GST-PSD3 was diluted to 1 mg/mL final concentration with binding buffer, and aliquots were flash
frozen in liquid nitrogen and stored at -80C.
540
Immunoprecipitation and Kinase Assay
HEK293T cells were transfected with 0.5-1 ug plasmid DNA using the JetOptimus system
according to the manufacturer's instructions (Genesee Scientific). After overnight transfections,
cells were lysed in HKT lysis buffer (25 mM HEPES pH 7.2, 150 mM KCl, 1% Triton X-100, 1mM
DTT) with 1x cOmplete protease inhibitor (Sigma). Lysate was then spun for 10 minutes at
15,000g at 4C to pellet out cell debris. For pre-conjugated beads (HA, V5), beads were
equilibrated with 2x HKT washes, followed by addition of 1 mL cell lysate for 30 uL equilibrated
beads. For non-conjugated beads (Protein G), beads were equilibrated with 2x HKT washes,
followed by pre-clearing of 1 mL cell lysate with 15 uL equilibrated beads for 30-60 min at 4C
rotating. 1 ug of mouse primary antibody (anti-GFP, Roche) was conjugated to 30 uL of Protein
G beads during this pre-clear step, followed by addition of pre-cleared lysate to the conjugated
beads for immunoprecipitation. Proteins were immunoprecipitated for 3-16 hours at 4C with

rotation. Following IP, samples were washed 2x with HKT, 1x with high salt HKT (25 mM HEPES

pH 7.2, 1 M NaCl, 1% Triton X-100, 1mM DTT) for 10 minutes, 2x with HK buffer (25 mM HEPES

pH 7.2, 1 M NaCl, 1mM DTT), and 2x with kinase buffer (20 mM Tris HCl pH 7.5, 10 mM MgCl2,

1 mM DTT, 1x cOmplete protease inhibitor). Following bead washes, kinase labeling reaction was

performed for 45 minutes at 30C on a shaking heat block (Thermo) in kinase buffer with 0.5mM

ATP (Sigma) or ATPɣS (Thermo) and 2x Halt Phosphatase inhibitor (Thermo). Following the

kinase reaction, samples with normal ATP were quenched by the addition of 4X Lammeli-SDS

sample buffer (Thermo) with 200mM DTT. Labeling was quenched in samples with ATPgS by the

addition of EDTA to a final of concentration of 50mM, and thiophosphates were then alkylated

with p-nitrobenzyl mesylate (pNBM, Sigma) at a final concentration of 6.67mM for 45 minutes at

room temperature. After alkylation of the ATPgS-containing samples, the reaction was quenched

by the addition of 4X Lammeli-SDS sample buffer (Thermo) with 200mM DTT. For co-IP

experiments, 0.5μg of V5-PSD3 was co-transfected with 0.5μg of HA-NUAK1 WT or HA-empty

vector with JetOptimus (Genesee Scientific) as above. This was followed by lysis, equilibration of

beads, and immunoprecipitation with either HA or V5-conjugated beads for 3 hours. Following IP,

samples were washed 3x with HKT and 3x with HK, followed by addition of 4X Lammeli-SDS

sample buffer (Thermo) with 200mM DTT.

Antibodies and Reagents

Antibody/Reagent	Company	Catalog number
HA beads	Sigma	E6779-1ML
V5 beads	Sigma	A7345-1ML
HisPur Ni-NTA beads	ThermoFisher	88221
Glutathione beads	ThermoFisher	16100
Pierce Protein G beads	ThermoFisher	20398
Anti-NUAK1	MRC PPU	Sheep 628B
Anti-GFP (Roche)	Sigma	11814460001
Anti-p-S/T	Phosphosolutions	PP2551
Anti-pTE	Abcam	ab92570
Anti-HA	Proteintech	66006-2-Ig
His-HRP	Proteintech	HRP-66005
GST-HRP	ThermoFisher	MA4-004-HRP
Anti-V5	ThermoFisher	R960-25

Anti-MAP2	Abcam	ab5392
Phenyl-ATPɣS	Axxora	BLG-P044
Furfuryl-ATPɣS	Axxora	BLG-F008-05
Benzyl-ATPɣS	Axxora	BLG-B072
ATPɣS	Thermo	40-801-0
pNBM	Sigma	SML3810-25MG

Confocal Microscopy

Live and fixed cell confocal imaging was performed on a Nikon Ti2 Eclipse-CSU-X1 confocal spinning disk microscope equipped with four laser lines—405, 488, 561, and 670 nm—and the Nikon Elements software. An Andor Zyla4.2 Plus sCMOS camera was used for image acquisition. The microscope was caged within the OkoLab environmental control setup, enabling temperature and CO_2 control during live imaging. Imaging was performed using Nikon 1.49 100×, Apo 60×, or 40× oil objectives. All image analyses were done using the open access Fiji software.

Super Resolution Microscopy

The Zeiss ELYRA7 Lattice SIM2 structural illumination microscope was used for super resolution imaging. Live cell imaging was performed on a Zeiss Axio Observer 7 - Lattice SIM inverted microscope for super resolution microscopy equipped with four laser lines – 405 nm (50mW), 488 nm (100mW), 561 nm (100mW) and 642 nm (150mW) – and the ZEN imaging software. Two Hamamatsu ORCA-Fusion BT sCMOS cameras were used for simultaneous image acquisition. The microscope was equipped with a Zeiss Stage Top incubation and gas system for Lattice SIM inverted microscopes, enabling temperature, humidity, and CO2 control during live imaging. Imaging was performed using Plan-Apochromat 63x/1.40 oil DIC objective.

Immunofluorescence and Western Blotting

All cells, including neurons, were fixed using warm 4% paraformaldehyde with 4% sucrose for 20 min at room temperature followed by three washes with 1x phosphate-buffered saline (PBS). One-hour incubation with blocking buffer [4% normal donkey serum, 0.2 M glycine (pH 7.4), and 0.1% Triton X-100, in 1x PBS] was followed by overnight incubation with primary antibody at 1:100-1000 dilution in blocking buffer at 4°C. After six 5-min washes in PBS, cells were incubated with secondary antibody at 1:1000 dilution in blocking buffer for overnight at 4°C. Coverslips were washed 6 times in PBS and then mounted onto microscope glass slides with Fluoromount-G (EMS) and sealed with clear nail polish (EMS).

Samples for Western blot analysis were treated with NuPAGE LDS sample buffer (4×)
(Invitrogen) with Dithiothreitol (DTT) at 200mM and subsequently heated for 10 min at 95°C.
Samples were electrophoresed on NuPAGE 4 to 12% Bis-Tris polyacrylamide gels (Invitrogen)
with NuPAGE MOPS or MES running buffer (Invitrogen). Western blots were transferred to
polyvinylidene difluoride membrane Immobilon-P (MilliporeSigma) with transfer buffer [25 mM
Tris, 192 mM glycine, 0.1% SDS and 20% (v/v) methanol]. Transferred blots were blocked in 5%
nonfat dry milk or 5% Bovine Serum Album (Equitech Bio) and subjected to primary antibody
overnight, washed 3x with blocking buffer, and incubated 3 hours at room temperature with
horseradish peroxidase–conjugated secondary antibody before visualization with Pierce
SuperSignal West Pico PLUS Chemiluminescent Substrate (Thermo Fisher Scientific). Western
blot images were obtained using the ChemiDoc Imaging System (Bio-Rad). Western blot images
were analyzed and quantified using Image Lab Software for Mac Version 6.1 (Bio-Rad).

Mouse Brain Lysate
To prepare mouse brain lysate, P13-P15 C57BL/6 mice pups of both sexes were euthanized by
CO2 (5psi for 10min) followed by decapitation. The brains were dissected out and incubated in
high detergent lysis buffer (20mM Tris pH7.5, 100mM NaCl, 10mM MgCl2, 0.5mM DTT, 1%
TritonX-100, 1% deoxycholic acid, and 1x cOmplete protease inhibitor) on ice for 30min.
Homogenization was achieved by sonication. Supernatant was collected after centrifugation at
14,000g at 4°C for 10 min, aliquots were prepared by diluting with lysis buffer without detergent
to 20µg/µl, flash frozen in liquid nitrogen and stored at -80°C.

NUAK1 Substrate Labeling and Covalent Capture
Covalent capture to identify NUAK1 kinase substrates using mass spectrometry was done as
previously described[14,15,22,23]. Briefly, HA-tagged NUAK1 M132G was purified from HEK293T cells
by affinity purification using HA antibody-coated beads. Cells expressing HA-tagged kinase-dead
NUAK1-K84M were used as negative controls. Cells were lysed with 25mM HEPES pH 7.2,
150mM KCl, 1x cOmplete protease inhibitor cocktail (Sigma), 1x Halt phosphatase inhibitor
(Thermo), 1 mM DTT, 1 mM EDTA, and 1% Triton X-100. Lysate was incubated on ice for 30 min
with frequent vortexing to lyse cells, followed by centrifugation at 20,000g for 15 min. Supernatant
was incubated with 30µL anti-HA resin for 3hr rotating at 4C. 100µl brain lysate at 20µg/µl protein
concentration was used for labeling by NUAK1-AS or NUAK1-KD. A total of 5 experimental
replicates each for NUAK1-AS or NUAK1-KD were completed with p15 mouse brain lysate.

The labeling reaction comprised of 1μM cAMP-dependent protein kinase inhibitor (PKI 5-24,
Tocris), 0.2μM protein kinase C inhibitor (Bisindolylmaleimide-I, Calbiochem), 3mM GTP (Sigma),
100μM ATP (Sigma), 1μM okadaic acid (Sigma), 2x Halt Phosphatase inhibitor (ThermoFisher)
and 0.5mM Furfuryl-ATP-γ-S (Axxora).

Labeling of the brain lysate was quenched by addition of EDTA to 50mM final concentration. The
labeled reaction was denatured by adding solid urea (60% w/v) and TCEP (10mM final
concentration) and incubating at 55°C for an hour. Samples were then diluted with two volumes
of ammonium bicarbonate (Sigma) and two volumes of TCEP (to maintain 10 mM final
concentration) and the pH adjusted to 8.0, then digested with Trypsin (Promega, sequencing
grade #V5113) with overnight rotation at 37C. Trypsin was quenched the following day by adding
0.5% TFA (Trifluoroacetic acid, Sigma).

Peptides were ascertained to be at pH 3.0, then desalted and cleaned using C18 cartridges
(Waters Sep-Pak Classic C18 Cartridges), and finally eluted by 70% Acetonitrile (ACN) and 0.1%
TFA. Peptide solution was reduced in volume by SpeedVac and final solution adjusted to 50%
ACN and 50mM HEPES pH 7.3.

Samples were evenly adjusted to pH7.0 using 5% NaOH, incubated with SulfoLink Coupling
Resin (ThermoFisher, 20401) and mixed overnight on a nutator at room temperature protected
from light. Addition of 25μg Bovine Serum Albumin (BSA) to the beads prior to sample addition
ensured low background binding. The following day, the beads were washed over equilibrated
fritted columns (Sigma) sequentially using Ultrapure H2O (Thermo), 5M NaCl, 50% ACN, 5%
Formic Acid and 1mM DTT. Bound thio-phospho-peptides were eluted through hydrolysis reaction
by addition of Oxone (1mg/ml Oxone in ddH2O freshly prepared, Sigma 228036). The resulting
phosphopeptides were extracted and desalted using C18 StageTips according to the published
protocol[24].

Reversed-Phase LC-MS/MS Analysis

Peptide samples were separated on an EASY-nLC 1200 System (Thermo Fisher Scientific) using
20 cm long fused silica capillary columns (100 μm ID, laser pulled in-house with Sutter P-2000,
Novato CA) packed with 3 μm 120 Å reversed phase C18 beads (Dr. Maisch, Ammerbuch,
DE). The LC gradient was 90 min long with 5–35% B at 300 nL/min. LC solvent A was 0.5% (v/v)
aq. acetic acid and LC solvent B was 80% acetonitrile in 0.5% (v/v) acetic acid. MS data was
collected with a Thermo Fisher Scientific Orbitrap Fusion Lumos using a data-independent
acquisition (DIA) method with a 120K resolution Orbitrap MS1 scan and 12 m/z isolation window,
30K resolution Orbitrap HCD MS2 scans for precursors from 400-1000 m/z.

Protein and Phosphosite Identification

Data .raw files were converted to .mzML using MSConvert 3.0.24285 and searched using FragPipe version 21.1 and MSFragger version 4.0, with quantification through DIA-NN version 1.8.2[64,65]. The database search was against the UniProt mouse protein sequence database (downloaded 02-14-2024) with supplemental spike-in of common contaminants, containing 17642 sequences and 17642 reverse-sequence decoys, as well as the sequence for human NUAK1. For the MSFragger analysis, both precursor and initial fragment mass tolerances were set to 20 ppm. Enzyme specificity was set to "stricttrypsin" and up to two missed cleavages were allowed. Oxidized Met, acetylated N-termini, and phosphorylated Ser/Thr/Tyr were set as variable modifications and carbamidomethylated Cys was a fixed modification. The maximum number of variable modifications per peptide was set to 3. FragPipe/DIA-NN output files were processed using a Python script to extract +/- 7 AA sequence flanking identified phosphosites from the FASTA file.

Database Usage

Developing and adult human brain transcriptomics data were obtained and plotted using HBT database (https://hbatlas.org/). These data were generated using Affymetrix GeneChip Human Exon 1.0 ST Arrays from over 1340 tissue samples sampled from both hemispheres of postmortem human brains, with specimens ranging from embryonic development to adulthood and representative of both males and females from multiple ethnicities. Allen Brain Atlas was used to search for NUAK1 mRNA expression in the mouse brain (https://mouse.brain-map.org/experiment/show/73497621) as well as PSD3 mRNA expression (https://mouse.brain-map.org/experiment/show/69095967). Evolutionary conservation analysis for both NUAK1 and PSD3 was performed using the ConSurf-DB web server[32]. Amphipathic helix prediction was performed using the NUAK1 protein sequence from Uniprot and the Heliquest[35]. NUAK1 consensus sequence and phosphorylation preference for substrates was determined using the publicly available Kinase Library database which was generated using the data from Johnson et al, 2023[42].

Molecular Modeling of NUAK1 and its Substrates

Prediction of NUAK1(O60285) with PSD3(Q9NYI0) and NUAK1(O60285) with PP1beta(P62140) and MYPT1(O14974) was obtained using Alphafold3[33]. Alphafold3 was installed via GitHub (https://github.com/google-deepmind/alphafold3) on a machine running Ubuntu 24.04 LTS with

two Nvidia A6000 48GB GPUs. Structural figures were prepared in PyMOL (Schrodinger).

Predictions were performed with 20 recycles with 5 diffusion samples per seed. Multiple seeds

(217, 275, 428, 551, and 924) were used for random number generation to increase robustness

of prediction.

Statistics and Reproducibility

Statistical analyses were performed with GraphPad Prism (version 10.0). No statistical

method was used to pre- determine sample size, and no data were excluded from the analyses.

The experiments were not randomized and investigators were not blinded during experiments

and data analysis. All experiments were performed in at least three independent replicates.

References

1. Manning, G., Whyte, D. B., Martinez, R., Hunter, T. & Sudarsanam, S. The protein kinase complement
of the human genome. *Science* **298**, 1912–1934 (2002).

2. Courchet, V. *et al.* Haploinsufficiency of autism spectrum disorder candidate gene NUAK1 impairs
cortical development and behavior in mice. *Nat. Commun.* **9**, 4289 (2018).

3. Bennison, S. A., Liu, X. & Toyo-oka, K. Nuak kinase signaling in development and disease of the
central nervous system. *Cell. Signal.* **100**, 110472 (2022).

4. Ferrarelli, L. K. A tauopathy kinase. *Sci. Signal.* **9**, (2016).

5. Lasagna-Reeves, C. A. *et al.* Reduction of Nuak1 Decreases Tau and Reverses Phenotypes in a
Tauopathy Mouse Model. *Neuron* **92**, 407–418 (2016).

6. Taylor, L. W. *et al.* p-tau Ser356 is associated with Alzheimer's disease pathology and is lowered in
brain slice cultures using the NUAK inhibitor WZ4003. *Acta Neuropathol.* **147**, 7 (2024).

7. Zhang, T. *et al.* NUAK1 promotes organ fibrosis via YAP and TGF-β/SMAD signaling. *Sci. Transl.*
Med. **14**, eaaz4028 (2022).

8. Carney, E. F. NUAK1 promotes kidney fibrosis. *Nat. Rev. Nephrol.* **18**, 345–345 (2022).

9. Ni, X., Pan, F., Lang, Y. K. & Zhang, W. Prognostic significance of NUAK1 and its association with
immune infiltration in stomach adenocarcinoma. *Discov. Oncol.* **15**, (2024).

10. Golkowski, M. *et al.* Pharmacoproteomics Identifies Kinase Pathways that Drive the Epithelial-
Mesenchymal Transition and Drug Resistance in Hepatocellular Carcinoma. *Cell Syst* **11**, 196-207.e7
(2020).

11. Yao, C.-Y. *et al.* NUAK1 acts as a novel regulator of PD-L1 via activating GSK-3β/β-catenin
pathway in hepatocellular carcinoma. *Mol. Med.* **31**, 38 (2025).

12. Whyte, D. *et al.* NUAK1 governs centrosome replication in pancreatic cancer via MYPT1/PP1β and
GSK3β-dependent regulation of PLK4. *Mol. Oncol.* **17**, 1212–1227 (2023).

13. Inazuka, F. *et al.* Muscle-specific knock-out of NUAK family SNF1-like kinase 1 (NUAK1) prevents
high fat diet-induced glucose intolerance. *J. Biol. Chem.* **287**, 16379–89 (2012).

14. Crunkhorn, S. NUAK1 inhibition prevents fibrosis. *Nat. Rev. Drug Discov.* **21**, 338–338 (2022).

15. Courchet, J. *et al.* Terminal Axon Branching Is Regulated by the LKB1-NUAK1 Kinase Pathway via
Presynaptic Mitochondrial Capture. *Cell* **153**, 1510–1525 (2013).

16. Lanfranchi, M. *et al.* The AMPK-related kinase NUAK1 controls cortical axons branching by locally
modulating mitochondrial metabolic functions. *Nat. Commun.* **15**, 2487 (2024).

17. Zagórska, A. *et al.* New Roles for the LKB1-NUAK Pathway in Controlling Myosin Phosphatase
Complexes and Cell Adhesion. *Sci Signal* **3**, ra25 (2010).

18. Hou, X. *et al.* A new role of NUAK1: directly phosphorylating p53 and regulating cell proliferation.
Oncogene **30**, 2933–2942 (2011).

19. Cullen, N. M. *et al.* NUAK kinases: Signaling mechanisms and therapeutic applications. *J. Biol.*
Chem. **301**, 110565 (2025).

20. Lizcano, J. M. *et al.* LKB1 is a master kinase that activates 13 kinases of the AMPK subfamily,
including MARK/PAR-1. *EMBO J.* **23**, 833–843 (2004).

21. Suzuki, A., Ogura, T. & Esumi, H. NDR2 Acts as the Upstream Kinase of ARK5 during Insulin-like
Growth Factor-1 Signaling*. *J. Biol. Chem.* **281**, 13915–13921 (2006).

22. Suzuki, A. *et al.* Identification of a Novel Protein Kinase Mediating Akt Survival Signaling to the
ATM Protein*. *J. Biol. Chem.* **278**, 48–53 (2003).

23. Monteverde, T. *et al.* Calcium signalling links MYC to NUAK1. *Oncogene* **37**, 982–992 (2018).

24. Cossa, G. *et al.* Localized Inhibition of Protein Phosphatase 1 by NUAK1 Promotes Spliceosome
Activity and Reveals a MYC-Sensitive Feedback Control of Transcription. *Mol. Cell* **77**, 1322-1339.e11
(2020).

25. Palma, M. *et al.* Identification of a nuclear localization signal and importin beta members mediating
NUAK1 nuclear import inhibited by oxidative stress. *J. Cell. Biochem.* **120**, 16088–16107 (2019).

26. Blazejewski, S. M., Bennison, S. A., Liu, X. & Toyo-oka, K. High-throughput kinase inhibitor
screening reveals roles for Aurora and Nuak kinases in neurite initiation and dendritic branching. *Sci.*
Rep. **11**, 8156 (2021).

27. Satoh, D. & Arber, S. Carving Axon Arbors to Fit: Master Directs One Kinase at a Time. *Cell* **153**,
1425–1426 (2013).

28. Kang, H. J. *et al.* Spatio-temporal transcriptome of the human brain. *Nature* **478**, 483–489 (2011).

29. Iossifov, I. *et al.* De Novo Gene Disruptions in Children on the Autistic Spectrum. *Neuron* **74**, 285–
299 (2012).

30. Iossifov, I. *et al.* The contribution of de novo coding mutations to autism spectrum disorder. *Nature*
515, 216–221 (2014).

31. Wang, T. *et al.* De novo genic mutations among a Chinese autism spectrum disorder cohort. *Nat.*
Commun. **7**, 13316 (2016).

32. Chorin, A. B. *et al.* ConSurf-DB: An accessible repository for the evolutionary conservation patterns
of the majority of PDB proteins. *Protein Sci.* **29**, 258–267 (2020).

33. Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with AlphaFold 3.
Nature **630**, 493–500 (2024).

34. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* 1–7 (2021)
doi:10.1038/s41586-021-03819-2.

35. Gautier, R., Douguet, D., Antonny, B. & Drin, G. HELIQUEST: a web server to screen sequences
with specific alpha-helical properties. *Bioinformatics* **24**, 2101–2102 (2008).

36. Beeman, N., Sapre, T., Ong, S.-E. & Yadav, S. Neurodevelopmental disorder–associated mutations in
TAOK1 reveal its function as a plasma membrane remodeling kinase. *Sci Signal* **16**, eadd3269 (2023).

37. Kerkhofs, M. *et al.* The AMPK-related kinase NUAK1 regulates neuronal morphogenesis through the
RNA splicing co-factor SON. *bioRxiv* 2025.10.10.681550 (2025) doi:10.1101/2025.10.10.681550.

38. Hertz, N. T. *et al.* Chemical genetic approach for kinase-substrate mapping by covalent capture of
thiophosphopeptides and analysis by mass spectrometry. *Curr Protoc Chem Biol* **2**, 15–36 (2010).

39. Blethrow, J. D., Glavy, J. S., Morgan, D. O. & Shokat, K. M. Covalent capture of kinase-specific
phosphopeptides reveals Cdk1-cyclin B substrates. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 1442–1447 (2008).

40. Blethrow, J., Zhang, C., Shokat, K. M. & Weiss, E. L. Design and use of analog-sensitive protein
kinases. *Curr Protoc Mol Biol* **Chapter 18**, Unit 18.11-18.11.19 (2004).

41. Yadav, S. *et al.* TAOK2 Kinase Mediates PSD95 Stability and Dendritic Spine Maturation through
Septin7 Phosphorylation. *Neuron* **93**, 379–393 (2017).

42. Johnson, J. L. *et al.* An atlas of substrate specificities for the human serine/threonine kinome. *Nature*
613, 759–766 (2023).

43. Fukaya, M., Ohta, S., Hara, Y., Tamaki, H. & Sakagami, H. Distinct subcellular localization of
alternative splicing variants of EFA6D, a guanine nucleotide exchange factor for Arf6, in the mouse
brain. *J. Comp. Neurol.* **524**, 2531–2552 (2016).

44. Gonzalez, D. A. *et al.* The Arf6 activator Efa6/PSD3 confers regional specificity and modulates
ethanol consumption in Drosophila and humans. *Mol. Psychiatry* **23**, 621–628 (2018).

45. Kanamarlapudi, V. Exchange Factor EFA6R Requires C-terminal Targeting to the Plasma Membrane
to Promote Cytoskeletal Rearrangement through the Activation of ADP-ribosylation Factor 6 (ARF6)*. *J.
Biol. Chem.* **289**, 33378–33390 (2014).

46. Macia, E. *et al.* The C-terminal domain of EFA6A interacts directly with F-actin and assembles F-
actin bundles. *Sci. Rep.* **9**, 19209 (2019).

47. Fukaya, M., Ohta, S., Hara, Y., Tamaki, H. & Sakagami, H. Distinct subcellular localization of
alternative splicing variants of EFA6D, a guanine nucleotide exchange factor for Arf6, in the mouse
brain. *J. Comp. Neurol.* **524**, 2531–2552 (2016).

48. Banerjee, S. *et al.* Interplay between Polo kinase, LKB1-activated NUAK1 kinase, PP1βMYPT1
phosphatase complex and the SCFβTrCP E3 ubiquitin ligase. *Biochem. J.* **461**, 233–245 (2014).

49. Kim, Y. *et al.* ADP-ribosylation Factor 6 (ARF6) Bidirectionally Regulates Dendritic Spine
Formation Depending on Neuronal Maturation and Activity*. *J. Biol. Chem.* **290**, 7323–7335 (2015).

50. Choi, S. *et al.* ARF6 and EFA6A Regulate the Development and Maintenance of Dendritic Spines. *J.*
Neurosci. **26**, 4811–4819 (2006).

51. Brown, F. D., Rozelle, A. L., Yin, H. L., Balla, T. & Donaldson, J. G. Phosphatidylinositol 4,5-
bisphosphate and Arf6-regulated membrane traffic. *J. Cell Biol.* **154**, 1007–1018 (2001).

52. Naslavsky, N., Weigert, R. & Donaldson, J. G. Convergence of Non-clathrin- and Clathrin-derived
Endosomes Involves Arf6 Inactivation and Changes in Phosphoinositides. *Mol. Biol. Cell* **14**, 417–431
(2003).

53. Mezache, L. & Leterrier, C. Advancing Super-Resolution Microscopy: Recent Innovations in
Commercial Instruments. *Microsc. Microanal.* **31**, ozaf004 (2025).

54. Eyck, L. F. T., Taylor, S. S. & Kornev, A. P. Conserved spatial patterns across the protein kinase
family. *Biochim. Biophys. Acta (BBA) - Proteins Proteom.* **1784**, 238–243 (2008).

55. Kennedy, E. J., Yang, J., Pillus, L., Taylor, S. S. & Ghosh, G. Identifying Critical Non-Catalytic
Residues that Modulate Protein Kinase A Activity. *PLoS ONE* **4**, e4746 (2009).

56. Zhou, X. *et al.* Integrating de novo and inherited variants in 42,607 autism cases identifies mutations
in new moderate-risk genes. *Nat. Genet.* **54**, 1305–1319 (2022).

57. Jaworski, J. ARF6 in the nervous system. *Eur. J. Cell Biol.* **86**, 513–524 (2007).

58. Mancina, R. M. *et al.* PSD3 downregulation confers protection against fatty liver disease. *Nat. Metab.*
4, 60–75 (2022).

59. Sheng, M. *et al.* NUAK1 promotes metabolic dysfunction-associated steatohepatitis progression by
activating Caspase 6–driven pyroptosis and inflammation. *Hepatol. Commun.* **8**, e0479 (2024).

60. Chen, P., Li, K., Liang, Y., Li, L. & Zhu, X. High NUAK1 expression correlates with poor prognosis
and involved in NSCLC cells migration and invasion. *Exp. Lung Res.* **39**, 9–17 (2013).

61. Jin, L. *et al.* PSD3 is an oncogene that promotes proliferation, migration, invasion, and G1/S
transition while inhibits apoptotic in papillary thyroid cancer. *J. Cancer* **12**, 5413–5422 (2021).

62. Luo, S. *et al.* PSD3 as a context-dependent modulator of immune landscape and tumor aggressiveness
in esophageal squamous cell carcinoma. *Front. Immunol.* **16**, 1641254 (2025).

63. Byeon, S. *et al.* TAOK2 drives opposing cilia length deficits in 16p11.2 deletion and duplication
carriers. *Stem Cell Rep.* **20**, 102608 (2025).

64. Kong, A. T., Leprevost, F. V., Avtonomov, D. M., Mellacheruvu, D. & Nesvizhskii, A. I. MSFragger:
ultrafast and comprehensive peptide identification in mass spectrometry–based proteomics. *Nat. Methods*
14, 513–520 (2017).

65. Demichev, V., Messner, C. B., Vernardis, S. I., Lilley, K. S. & Ralser, M. DIA-NN: Neural networks
and interference correction enable deep proteome coverage in high throughput. *Nat. methods* **17**, 41–44
(2020).

FIGURE LEGENDS

Figure 1. Autism-associated NUAK1 mutations impact catalytic activity and

subcellular localization

(a) Day *in vitro* (div) 14 rat hippocampal neurons stained with antibodies against NUAK1

and MAP2, with nuclear marker DAPI. Right panels show magnified soma with anti-

NUAK1 and either MAP2 (top panel) or DAPI (bottom panel). Error bars represent 20μm

in full images and 10μm in zoomed images at right. **(b)** Human brain transcriptomic data

for NUAK1 obtained from Human Brain Transcriptome (HBT) database across different

stages of human brain development. Genome-wide, exon-level transcriptome data in HBT

was generated using the Affymetrix GeneChip Human Exon 1.0 ST Arrays from over

1,340 tissue samples sampled from both hemispheres of postmortem human brains.

Specimens range from embryonic development to adulthood (periods 1-15) and represent

both males and females. Brain regions plotted are the neocortex (NCX), hippocampus

(HIP), amygdala (AMY), striatum (STR), mediodorsal nucleus of the thalamus (MD), and

cerebellar cortex (CBC). **(c)** *In situ* hybridization data from the Allen Brain Atlas using

antisense probe against NUAK1 on sagittal sections of postnatal day 56 (P56) C57BL6/J

male mouse brain. Scale bar represents 839μm. **(d)** Two-dimensional schematic of

NUAK1 protein domains, with the N-terminal kinase domain, 3 GILK motifs, and C-

terminal amphipathic helix indicated with colored boxes. The three NUAK1 ASD-

associated mutations are shown with lollipops above the protein domains: the two

missense mutations Q161E and A653V, and the early truncation mutation Q433ter

(Q433*). **(e)** ConSurf evolutionary conservation analysis of the two ASD-associated

missense mutations in NUAK1: Q161E and A653V. Residues are colored according to

their evolutionary conservation, with warm colors indicating conserved residues and cool

colors indicating variable residues. **(f)** Heliquest-predicted amphipathic helix at the C-

terminus of NUAK1, which spans amino acids 644-661. Alanine 653, the site of the A653V

ASD-associated mutation, is indicated. The arrow indicates the hydrophobic face of the

amphipathic helix. **(g)** Western blot of immunoprecipitated GFP-NUAK1 mutants

incubated plus/minus ATP to allow NUAK1 to autophosphorylate. Immunoblots show anti-

phospho-serine/threonine and anti-GFP (loading control). **(h)** Quantification of kinase

activity of NUAK1 mutants, with the phospho-serine/threonine signal divided by the GFP

loading control and all samples normalized as a percentage of WT + ATP per experiment

(n=3 independent experiments, ** denotes $p<0.005$; *** denotes $p<0.001$, one way

ANOVA with Tukey's multiple comparisons). All error bars represent standard error of the

mean. **(i)** DIV 14 rat embryonic hippocampal neurons expressing GFP-tagged NUAK1

mutants and membrane marker myristoyl-tdtomato (myr-tdtomato). Neurons were

transfected with GFP-NUAK1 and myr-tdTomato on DIV 12 and fixed 48 hours later. The

arrowhead in Q433* image indicates the predominantly nuclear localization of this

mutation. Scale bar represents 20μm in all images.

Figure 2. Engineering ATP-analog sensitive NUAK1 kinase through the gatekeeper

residue mutation M132G

(a) Multiple sequence alignment of the kinase domains of serine/threonine kinases,

including NUAK1, NUAK1, AMPKα2, MST3, and TAOK2. The conserved methionine

gatekeeper residue is colored red and underlined, while the flanking residues are colored

blue. The four highly conserved residues surrounding the gatekeeper are indicated with

a box. **(b)** Top: bulky side groups (benzyl, phenyl, and furfuryl) that are N6-substitutents

of ATPɣS (right) in kinase assays. Bottom: Western blots showing utilization of bulky

ATPɣS molecules by purified HM-tagged NUAK1 (amino acid 1-433). Antibody against

phosphothioester (pTE) is used as a readout of autophosphorylation, while anti-His is

used as a loading control. The optimal gatekeeper mutant/bulky ATPɣS combination of

M132G NUAK1 and furfuryl-ATPɣS is indicated with a red box. **(c)** Quantification of

utilization of bulky ATPɣS analogs by HM-NUAK1, with pTE signal divided by loading

control His for each sample. Data were normalized to WT for each analog, with KD and

MG represented as a ratio of WT. N=3 independent experiments. Error bars represent

standard error of the mean. **(d)** HA-tagged full-length NUAK1 KD or MG was expressed

in HEK293T cells followed by lysis and immunoprecipitation for the indicated time.

Immunoprecipitated NUAK1 was incubated with mouse brain lysate and furfuryl ATPɣS,

followed by alkylation to generate phosphothioester for WB detection. HA is used as a

loading control for NUAK1, while phosphothioester (pTE) is used as a readout of

substrate phosphorylation in mouse brain lysate. **(e)** Schematic of labeling of mouse brain

lysate by analog-sensitive (AS) NUAK1 and N6-furfuryl-ATPɣS (reaction 1), or negative

control K84M (KD) NUAK1 and N6-furfuryl-ATPɣS (reaction 2). Thiophosphorylated

substrate proteins were digested with trypsin, and thiophosphorylated peptides were then

covalently captured using iodoacetamide (IAA) beads. Beads were treated with oxone to

specifically elute off thiophosphorylated peptides as phosphopeptides, while peptides

with other thiol groups (e.g., cysteines) are retained on the beads. The resulting

phosphopeptides are then analyzed by LC-MS/MS.

Figure 3. NUAK1 phosphorylation targets identified by chemical-genetic screen

(a) Putative NUAK1 substrate proteins (31 total) that occurred in at least 3/5 analog

sensitive samples and 0/5 kinase dead samples were analyzed for their roles in

endocytosis, splicing, cytoskeletal reorganization, as well as development of dendrites,

axons, or synapses. Each substrate is color-coded according to its role(s) in each of these

processes, with light green for endocytosis, teal for splicing, yellow for cytoskeleton, pink

for axon, and blue for dendrites, with the legend in the lower right panel showing each

individual color. Substrates with roles in synaptic development are located in the "synapse"

box on the middle right. Substrates with multiple functions are indicated with gradients

with 2-4 colors. **(b)** The 9 amino acids surrounding the phosphoacceptor (from residues -

5 to +4) for the 31 potential substrates of NUAK1 that occurred in at least 3/5 analog

sensitive samples and 0/5 kinase dead samples were analyzed using the Phosphosite.org

Weblogo tool to generate a preference map for NUAK1 substrate phosphorylation. **(c)**

Putative NUAK1 substrate proteins that occurred in at least 3/5 analog sensitive samples

and 0/5 kinase dead samples were analyzed for how well they fit the established NUAK1

consensus sequence. Most stringently, NUAK1 requires a basic residue (K/R) at the -3

position. After filtering the 31 unique potential substrates by this stringent consensus

sequence, 17 putative substrates remained. The mouse Uniprot ID and protein name for

each protein is listed, as are the human and mouse phosphosites. The amino acids

surrounding the phosphosite are listed in this table, from the -5 position to the +4 position

relative to the phosphoacceptor. Based on the established NUAK1 consensus sequence,

along with the data from our MS runs, we were able to establish the optimal motif for

substrate phosphorylation by NUAK1. This optimal motif: (1) contains a basic residue at

-3 position; (2) contains a serine at -2 position; (3) contains a serine phosphoacceptor;

and (4) contains an aliphatic branched chain amino acid at -5 and +4 positions. Residues

are color coded based on whether the residue fits the optimal NUAK1 motif at the given

site, with positions -5 and +4 in purple, positions -4, -3, and -1 in cyan, and positions -2,

+1, +2, and +3 in chartreuse. The central phosphoacceptor at position 0 is colored in dark

green for serine (more preferred) and light green for threonine (less preferred).

Figure 4. PSD3 is a bona fide direct phosphorylation target of NUAK1 kinase.

(a) Two-dimensional schematic representation of the function domains of PSD3 isoform

1 (Uniprot # Q9NYI0), isoform 3 (Uniprot # Q9NYI0-3), and the shortened PSD3 construct

used for biochemical experiments, PSD3 CC tail. PSD3 isoforms 1 and 3 share the same

3 conserved functional domains, including a Sec7 GEF domain (green) which activates

ARF6, Pleckstrin Homology (PH) domain with specificity for PI(4,5)P2, and a Coiled Coil

(CC) domain which binds to actin. The MS-identified phosphosite on PSD3, S1011 in

isoform 1 or S476 in isoform 3, is indicated with a lollipop. The shortened CC tail domain

used for protein purification contains an N-terminal GST tag followed by the C-terminal

region of PSD3, from the CC domain to the end of the protein containing the S476/S1011

phosphosite. **(b)** ConSurf evolutionary conservation analysis of the MS-identified

phosphosite of PSD3, S1011 in isoform 1 or S476 in isoform 3. The S1011

phosphoacceptor residue is boxed in violet. Residues are colored according to their

evolutionary conservation, with warmer colors indicating conserved residues and cool

colors indicating variable residues. **(c)** V5-tagged PSD3 isoform 3 WT or S476A was

immunoprecipitated from HEK293T cells, followed by incubation with purified HM-NUAK1

WT and ATP. Immunoblots show V5 as a loading control for PSD3 and His as a control

for NUAK1, and phospho-serine/threonine as a readout of phosphorylation for both

NUAK1 and PSD3 at their respective molecular weights. **(d)** Quantification of

phosphorylation of PSD3 by NUAK1, as measured by phospho-serine/threonine divided

by V5 signal. Each experiment is normalized to WT PSD3, with S>A PSD3 as a ratio of

WT signal (n=3 independent experiments; *** denotes $p<0.001$, student's t-test). Error

bars represent standard error of the mean. **(e)** HEK293T cells were co-transfected with

mCherry-V5-PSD3 isoform 3 and either HA-empty vector or HA-NUAK1 WT. Cells were

then lysed, and protein was immunoprecipitated using HA beads. Immunoblots show HA

signal for HA-empty vector or HA-NUAK1, and V5 signal for mCherry-V5-PSD3. **(f)**

Quantification of V5 signal for input and HA IP for n=3 independent experiments. Error

bars represent standard error of the mean; ** denotes $p<0.005$, one way ANOVA with

Tukey's multiple comparisons. **(g)** HEK293T cells were co-transfected with mCherry-V5-

PSD3 isoform 3 and either HA-empty vector or HA-NUAK1 WT. Cells were then lysed,

and protein was immunoprecipitated using V5 beads. Immunoblots show HA signal for

HA-empty vector or HA-NUAK1, and V5 signal for mCherry-V5-PSD3. **(h)** Quantification

of HA signal for input and V5 IP, for n=3 independent experiments. Error bars represent

standard error of the mean; * denotes $p<0.05$, one way ANOVA with Tukey's multiple

comparisons. **(i)** HM-NUAK1 WT or KD was incubated with GST-PSD3-CC tail WT or

S>A in the presence of Mg^{2+} and ATPɣS, followed by alkylation with PNBM to generate

thiophosphate esters as a readout of thiophosphorylation by NUAK1. **(j)** Quantification of

phosphorylation of GST-PSD3-CC tail WT or S>A by HM-NUAK1 WT or KD. Data are

normalized to WT PSD3 + WT HM-NUAK1 for each experiment, with all other samples

as a percentage of WT/WT (n=3 independent experiments; **** indicates p<0.0001, one

way ANOVA with Tukey's multiple comparisons). Error bars represent standard error of

the mean **(k)** Alphafold-predicted structure of NUAK1 WT (Uniprot # O60285) with the

previously identified and validated substrate MYPT1 (Uniprot ID #O14974), along with

adaptor protein Protein Phosphatase 1 beta (Uniprot ID #P62140). NUAK1 kinase domain

is displayed as an electrostatic surface representation, with red-toned colors representing

a negative charge, and blue-toned colors representing a positive charge. MYPT1 is

colored in light blue, and amino acids within the NUAK1 substrate binding groove are

color coded according to their position and side chain, with Leu at -5 and +4 in limon,

Lys/His at -4, -3, and -1 in blue, and central phospho-acceptor Ser in green. Amino acid

side chains are shown for positions -5 to +4 surrounding the MYPT1 S445 phosphosite.

(l) Alphafold3 predicted structure of NUAK1 WT (Uniprot # O60285) with PSD3 isoform 1

WT (Uniprot # Q9NYI0), following the same color scheme as figure 4k. Amino acid side

chains are shown for positions -5 to +4 surrounding the PSD3 S476 phosphosite.

Figure 5. Phosphorylation of PSD3 at S476 regulates ARF6 dependent trafficking

(a) HEK393T cells co-expressing mCherry-tagged WT or dominant negative T44N ARF6

(magenta) and sfGFP-tagged WT-PSD3, Sec7 mutant E114K or phosphomutant S476A

(green). Scale bar is 5μm. **(b)** Area of intracellular vesicles positive for ARF6 and PSD3

normalized to the total cell area is plotted for each condition (n>20 cells from 3

experiments, ** denotes $p<0.005$ and **** indicates $p<0.0001$, one-way ANOVA with

Tukey's multiple comparisons). **(c)** HEK393T cells co-expressing the PI(4,5)P2 sensor

EGFP-PLCD1 (gray) along with either mCherry-tagged WT or phosphomutant PSD3

S476A (red). Scale bar is 5μm. **(d)** Area of intracellular vesicles positive for PI(4,5)P2

sensor PH-PLCD1 normalized to the total cell area is plotted for each condition (n>23

cells from 3 experiments, **** indicates $p<0.0001$, unpaired t-test with Welch's correction).

All error bars are standard error of the mean.

Figure 6. NUAK1 mediated PSD3 phosphorylation at S476 restricts maturation of

filopodia into dendritic spines

(a) Representative images of DIV13 rat embryonic hippocampal neurons co-expressing

membrane marker myristoylated-tdTomato (red) along with either sfGFP-tagged wildtype

(WT) PSD3 or phosphomutant S476A (cyan). Scale bar is 10mm. **(b)** Magnified insets

are shown to highlight the morphology of dendritic spines and the enrichment for PSD3

within spines. Grayscale and pseudocolored images show intensity of PSD3 distribution

in WT and S476 conditions. Scale bar is 1μm. **(c)** Quantification of percent spines

enriched in PSD3 in hippocampal neurons expressing WT-PSD3 and S476A

phosphomutant (n>15 neurons for each condition from three experiments, **** denotes

p<0.0001, t-test with Welch's correction). **(d)** Protrusion density per micron dendrite

length is plotted for neurons expressing WT-PSD3 and S476A phosphomutant (n>55

neurons for each condition from three experiments, ns denotes p>0.05, t-test with Welch's

correction). **(e)** Percent protrusions classified into stubby, thin, filopodia and mushrooms

spines are plotted neurons expressing WT-PSD3 and S476A phosphomutant (n>55

neurons for each condition from three experiments, **** denotes p<0.0001 and * denotes

p<0.05, 2-way ANOVA with multiple comparisons). All error bars are standard error of the

mean. **(f)** Schematic depicts working model based on the results showing role of NUAK1

phosphorylation of PSD3 in regulation of ARF6 mediated membrane internalization and

PI(4,5)P2 production. Role of PSD3 phosphorylation in maturation of dendritic spines

through ARF6 is shown.

Figure 7. NUAK1 phosphorylation targets in neuronal development and disease

(a) Schematic shows substrates of NUAK1 identified by the chemical genetic screen

which are binned by their roles in axonal, dendritic, and synaptic development, as well as

splicing. Each substrate is color-coded according to its known function(s), with light green

for endocytosis, teal for splicing, yellow for cytoskeleton, pink for axon, and blue for

dendrites, with the legend in the lower left panel showing each individual color. Substrates

with multiple functions are indicated with gradients of 2-4 colors. **(b)** Schematic

representation of the role of PSD3 and NUAK1 in ARF6 activation. At the plasma

membrane, PSD3 recruits inactive, GDP-bound ARF6. PSD3 exchanges GDP for GTP,

generating active ARF6, which promotes the formation of PI(4,5)P2. This increase in PIP2

causes internalization of PSD3 and ARF6 on PIP2-enriched vesicles. In normal conditions,

Rab5, ARF6 GAPs, and lipid phosphatases are recruited to these vesicles, leading to a

cessation of ARF6 activity and depletion of PIP2. These vesicles can then continue in the

endolysosomal pathway, or be recycled to the cell surface. When NUAK1 is not present,

or when the PSD3 S476 phosphosite is mutated to alanine, ARF6 remains active and

bound to GTP, accumulating in large, PIP2-enriched vesicles. **(c)** Schematic of the role

of PSD3 in dendritic spine maturation. PSD3 activation of ARF6 leads to an increased

proportion of mature, mushroom-shaped dendritic spines and a corresponding loss of

immature filopodia. In normal conditions, PSD3 is phosphorylated by NUAK1, keeping

ARF6 activation in check. When NUAK1 is inactive or when the PSD3 phosphosite is

mutated to alanine, PSD3 phosphorylation is lost, leading to an aberrant increase in the

proportion of active ARF6.

Chemical Genetic Screen Identifies PSD3 as a Direct Substrate of NUAK1 that Regulates Dendritic Spine Maturation

Josilyn R. Sejd[1], Daphnée M. Marciniak[1], Moira A. Cornell[1], Angel Sondhi[1], Shao-En Ong[1] and Smita Yadav[1,2*]

[1]Department of Pharmacology, University of Washington, Seattle, United States.

[2]Institute of Stem Cell and Regenerative Medicine, Seattle, United States

* Corresponding author

Supplementary materials

Figure S1: NUAK1 structure and localization in mitotically active cells

Figure S2: NUAK1 chemical genetic method development

Figure S3: PSD3 evolutionary conservation and purification from *E. coli*

Table S4: Excel spreadsheet with 3 sheets. Sheet 1 contains an index and definitions of column titles. Sheet 2 contains curated proteins which occurred in 3/5 M132G replicates and 0/5 KD replicates and fit stringent consensus of K/R at -3 position. Sheet 3 contains all proteins identified in LC-MS/MS-based chemical genetic screen

Supplemental Figure Legends

Figure S1: NUAK1 structure and localization in mitotically active cells

(a) Alphafold-predicted structure for NUAK1 (AF-O60285-F1-v6), visualized in Pymol viewer. Color scheme reflects pLDDT value per residue, with dark blue representing very high confidence (>90%), cyan representing high confidence (70-90%), yellow representing low confidence (50-70%), and orange representing very low confidence

(<50%). The C-terminal helix containing Alanine 653 is labeled as "A653". Dashed box

represents the region that is magnified in S1b.

(b) Left: Zoom on the kinase domain of NUAK1 with unstructured domains hidden,

showing the position of the A653-containing helix relative to the kinase domain. Dashed

box shows area which is magnified in the right panel. Right: Zoom on the NUAK1 C-

terminal alpha helix containing Alanine 653, with A653 shown in licorice stick format.

(c)GFP-tagged NUAK1 mutants were expressed in human neural progenitor cells

(NPCs), followed by fixation and staining with anti-SC35 antibody and DAPI. Panels

show, from top to bottom, GFP-NUAK1, SC35 staining, merge of GFP-NUAK1, SC35,

and DAPI with a dashed box around the nucleus, and magnified nuclear area with merge

of GFP-NUAK1 and SC35.

Figure S2: NUAK1 chemical genetic method development

(a) Purification of HM-NUAK1 1-433 WT from *E. coli*. HM-NUAK1 1-433 WT was

expressed in *E. coli*, followed by induction with IPTG and lysis. Coomassie stain shows

protein gel with ladder of indicated molecular weights, with HM-NUAK1 1-433 WT

purification. Samples show pre-induction, whole cell lysate post sonication, supernatant

and pellet following centrifugation post sonication, flow through from bead binding, elution

1, HisPur beads following elution, and elution after incubation with SP beads. **(b)**

Purification of HM-NUAK1 1-433 WT from *E. coli* with the same samples loading as in (c),

with IB against His tag. **(c)** Purification of HM-NUAK1 1-433 KD and M132G from *E. coli*,

with samples from the first elution (E1), flow through after binding to SP beads (FT), wash

1 after binding to SP beads, and purified protein after elution from SP beads (final), with

purification process shown for both KD and M132G HM-NUAK1 1-433, with final WT HM-

NUAK1 1-433 (lane 9) shown for reference. A red dashed box indicates the relevant

bands. **(d)** Western blot of full-length HA-NUAK1 WT, K84M (KD), M132A, and M132G

incubated with the indicated ATPγS analogs and Mg2+ to allow autophosphorylation,

followed by alkylation of thiophosphates with pNBM to generate thiophosphate esters

(pTE). Anti-HA is used as a loading control, while pTE is used as a readout of

autophosphorylation. **(e)** Quantification of utilization of N6-bulky ATPγS analogs by HA-

NUAK1, with pTE signal divided by loading control His for each sample. Error bars

represent standard error of the mean.

Figure S3: PSD3 evolutionary conservation and purification from *E. coli*

(a) The sequences for human PSD3 (NP_056125.3) and 457 orthologs were obtained

from the NCBI gene browser, followed by alignment using Cobalt. Multiple sequence

alignments were visualized using SnapGene viewer. Amino acids L995 to D1017 in

human PSD3 are shown, with the height and color of each bar corresponding to

percentage conservation of that residue in the 457 orthologs analyzed. The consensus

amino acid sequence is listed, with X indicating a lack of conservation of one specific

amino acid at a given site. The phosphosite on PSD3 that was identified by MS is

indicated with an asterisk, and the flanking residues from -5 to +4 are surrounded by a

dashed box. The legend shows the color scheme based on percentage conservation of

the indicated amino acid, with >88% conservation in dark red, 80-87% in red, 63-79% in

orange, 50-64% in yellow-orange, and <49% in light blue. **(b)** GST-PSD3-CC tail (WT or

S476A) was expressed in *E. coli*, followed by induction with IPTG and lysis. Coomassie

stain shows protein gel with ladder of indicated molecular weights, with GST-PSD3-CC

tail purification. Samples show pre-induction, 1 and 2 hours post IPTG induction, whole

cell lysate post sonication, supernatant and pellet following centrifugation post sonication,

flow through from bead binding, elutions 1-6 (E1-6), and glutathione beads following 6

elutions with 25mM glutathione. The relevant bands are indicated with a dashed red box.

Figure 1

Figure 2

a

Endocytic Recycling

CLIP4, Aftiphilin, NHSL2, CLAP2, Erbin, MRCKα, RIMS1, NAV1, SPAR1, PSD3, Dynamin-1, TCPG

Splicing

CHERP, MPRIP, CHTOP, LS14B, PRRC2B, RBM14, RL13

Cytoskeleton

CEP170B, GAB1, Aftiphilin, CLIP2, MAP7D1, CLAP2, TBB6, MYPT2, MRCKα, CLIP4, DCLK1, NAV1, NHSL2, TNIK, PSD3, MPRIP, Dynamin-1, TCPG

Axon

GAB1, PCY1a, MAP7D1, LS14B, MYPT2, DCLK1, Aftiphilin, TNIK, CLAP2, Erbin, MRCKα, RIMS1, NAV1, SPAR1, PSD3, Dynamin-1, TCPG

Dendrite

CNK2, PCY1a, RBM14, Aftiphilin, Erbin, CLAP2, RIMS1, MRCKα, SPAR1, NAV1, DCLK1, PSD3, TNIK, TCPG

Synapse

CNK2, DCLK1, Erbin, CLAP2, TNIK, RIMS1, PSD3, RL13, SPAR1, Dynamin-1

LEGEND Endocytosis, Axon, Splicing, Cytoskeleton, Dendrite

b NUAK1 Substrate Preference

31 input sequences

c

Site	-5	-4	-3	-2	-1	0	+1	+2	+3	+4
Optimal motif	P	K	R	S	K/R/H	S	A	S/T	S	L/V
Secondary	L/I	R	K	X	P	S	S	X	X	S
Additional	R	S	X	X	T	T	X	X	X	X

Mouse ID	Protein name	human site	mouse site	-5	-4	-3	-2	-1	S/T	+1	+2	+3	+4	Localization (Uniprot)
Q2PFD7	PSD3	S1011	S1000	L	K	K	S	H	S	S	P	S	L	Membrane, cytosol
A2AJI0	MAP7D1	S313	S315	L	A	R	S	R	S	A	V	T	L	Cytosol
Q80U49	CEP170B	S1364	S1349	P	T	R	S	P	S	L	G	N	V	Cytosol
P83510	TNIK	S570	S541	P	P	R	S	E	S	F	S	I	S	Nucleus, cytosol
Q0PHV7	DACT3	S426	S409	A	A	R	S	Q	S	E	T	S	L	Cytosol
Q3UU96	MRCKa	S1629	S1616	P	G	R	S	M	S	A	S	S	G	Cytosol
Q8CH77	NAV1	S364	S367	P	M	R	S	P	S	K	L	S	H	Cytosol
P97434	MPRIP	S993	S992	I	M	K	S	K	S	N	P	D	F	Cytosol
Q8C0T5	SPAR1	T9	T9	L	K	R	S	Q	T	E	R	P	V	Membrane, cytosol
Q9QYY0	GAB1	T387	T388	P	S	R	S	N	T	I	S	T	V	Cytosol
Q9CY57	CHTOP	T15	T15	V	L	K	S	T	T	K	M	S	L	Nucleus
Q80YA9	CNKR2	T327	T327	I	P	R	S	P	T	S	S	V	A	Membrane, cytosol
Q8CGC4	LSM14B	S181	S181	P	S	K	V	T	S	A	T	Q	L	Cytosol
P47963	RL13	S106	S106	R	R	R	N	K	S	T	E	S	L	Cytosol
Q8CGZ0	CHERP	S902	S911	Y	R	R	N	K	S	Y	S	F	I	Nucleus, cytosol
P80318	TCPG	S252	S252	Y	K	K	G	E	S	Q	T	D	I	Cytosol
Q80TH2	Erbin	S1158	S1150	P	E	R	T	M	S	V	S	D	F	Nucleus, cytosol

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure S1

a

Coomassie: HM-NUAK1 1-433 WT

b

IB: anti-His: HM-NUAK1 1-433 WT

c

Coomassie: HM-NUAK1 1-433 KD, M132G, WT

d

| HA-NUAK1: | WT | KD | MA | MG | WT | KD | MA | MG | WT | KD | MA | MG |
| ATPγS: | | Benzyl | | | | Furfuryl | | | | Phenyl | | |

IB: pTE — 80 kD

IB: HA — 80 kD

e

HA-NUAK1 utilization of N6-ATPγS analogs

pTE over HA

| HA-NUAK1: | WT | KD | MA | MG | WT | KD | MA | MG | WT | KD | MA | MG |
| ATPγS: | | Benzyl | | | | Furfuryl | | | | Phenyl | | |

Figure S2

a

PSD3 Phosphosite Conservation

b

Coomassie: GST-PSD3 CC tail purification

Gel Loading:
1. Pre-ind
2. 1 hour ind.
3. 2 hour ind.
4. Whole cell lysate
5. Pellet
6. Supernatant
7. Flow thru
8. E1
9. E2
10. E3
11. E4
12. E5
13. E6
14. Beads

Figure S3