Metatranscriptome data support the existence of two distinct morphotypes in a single parmalean species in natural environments

Hiroto Sasaki[1], Hisashi Endo[1], Eric Pelletier[2,3], Shiya Yoshikawa[4], Akira Kuwata[5], Hiroyuki Ogata[1,*]

[1]Bioinformatics Center, Institute for Chemical Research, Kyoto University, Gokasho, Uji, Kyoto, 611-0011, Japan

[2]Génomique Métabolique, Genoscope, Institut François-Jacob, CEA, CNRS, Univ Evry, Université Paris-Saclay, F-91000 Evry, France

[3]Research Federation for the Study of Global Ocean Systems Ecology and Evolution, FRA2022/Tara GOSEE, F-75016 Paris, France

[4]Faculty of Marine Science and Technology, Fukui Prefectural University, Fukui, 917-0003, Japan

[5]Shiogama Field Station, Fisheries Resources Institute, Japan Fisheries Research and Education Agency, 3-27-5 Shinhama-cho, Shiogama, Miyagi, 985-0001, Japan

*Corresponding author: Hiroyuki Ogata, E-mail: ogata@kuicr.kyoto-u.ac.jp, Phone: +81-774-38-3270

Topics: Genomics, metagenomics, and bioinformatics for microbiology

Running headline: Two morphotypes for a single parmalean

Abstract

Parmales (Bolidophyceae), the closest relatives of diatoms, includes isolates exhibiting one of two distinct morphotypes: a silicified non-flagellated form (S-type) and a naked flagellated form (F-type). Although alternation of these two forms for a single isolate has not been formally established, previous studies hypothesized that these morphotypes represent alternating stages of a single parmalean. In this study, we investigated the global expression patterns of S- and F-type marker genes by integrating parmalean Metagenome-Assembled Genomes (MAGs) and the metatranscriptomic dataset from *Tara* Oceans. We detected the expression of both S- and F-type marker genes from individual environmental MAGs. This finding provides the first metatranscriptomic evidence that natural parmalean genomes possess the potential to manifest both morphotypes. Furthermore, our analysis revealed different geographical expression patterns between the two forms. The expression of F-type markers showed a broad distribution, whereas that of S-type markers was more restricted, suggesting distinct niches for the two morpho-phases. Moreover, S-type gene expression appears to require specific environmental triggers that lead to a higher population density, whereas F-type expression is rather constitutively maintained. Overall, our results support the hypothesis of a life cycle involving morphological switching and reconcile the long-standing discrepancy between the ubiquity of parmaleans in molecular surveys and the limited geographical range for the observation of silicified cells. Based on these patterns, we propose a threshold-based model in which F-type-dominated populations persist under conditions unfavorable for growth and a morphological switch to the S-type is triggered once environmental conditions exceed a critical threshold for growth.

Keywords: Parmales, Metatranscriptome, Morphotypes

Introduction

Diatoms are one of the most ecologically successful groups of phytoplankton, accounting for approximately 20% of primary production in modern oceans (Armbrust, 2009). However, the drivers of their ecological success remain elusive, and comparative studies involving their close relatives are increasingly recognized as promising approaches to address this question (Ichinomiya et al., 2016).

The order Parmales comprises pico-eukaryotic algae that form the sister group of diatoms (Guillou et al., 1999; Ichinomiya et al., 2011). Originally, two morphological forms have been described: a silicified form surrounded by silica plates (S-type) (Booth & Marchant, 1987), and a naked flagellated form possessing two unequal flagella (F-type) (Guillou et al., 1999). Recently, a novel silicified morphotype covered with distinct silica scales, referred to as 'Scaly parma', was identified within the basal lineage of Parmales (Ban et al., 2023). This discovery led to the recent establishment of the order Lepidoparmales to accommodate this lineage (Kamakura et al., 2025). Although these two morphological forms were historically regarded as distinct lineages, molecular phylogenetic analyses revealed their close relationship (Ichinomiya et al., 2011). Consequently, both S- and F-type algae—originally classified separately—were reclassified into the single order Parmales within the class Bolidophyceae (2). Ecologically, S-type cells are most abundant in polar and subarctic regions (Booth et al., 1980; Ichinomiya & Kuwata, 2015), whereas F-type populations were originally isolated from the equatorial Pacific Ocean and the Mediterranean Sea (Guillou et al., 1999). Despite these habitat preferences based on isolation, microscopic observations and molecular studies indicate that the algae in Bolidophyceae are distributed globally (Ban et al., 2024; Silver et al., 1980) and the morphological forms display different distributions across depth and seasons (Kuwata et al., 2018). These algae are referred to as parmaleans in this study.

The close evolutionary relationship between the S-type and F-type parmaleans raised the possibility that these morphotypes correspond to different life stages of a single organism. Mann and Marchant hypothesized that the life cycle of ancestral diatoms consisted of haploid flagellated cells and diploid silicified zygotes akin to that of the early branching centric diatoms (Mann & Marchant, 1989). This type of life cycle is also known for other algae such as the haptophyte *Gephyrocapsa huxleyi* (Green et al., 1996). Given the evolutionary relatedness between S-type and F-type parmalean strains and their sister relationship to diatoms, Ichinomiya et al. (2011) hypothesized that Parmales possess a life cycle that switches between a diploid silicified form and a haploid flagellated form (Ichinomiya et al., 2011). This hypothesis was supported by several lines of evidence. First, a study of a flagellated strain revealed a low level of heterozygosity, suggesting the haploid status of the genome for this strain (Kessenich et al., 2014). Second, the flagellated strain *Triparma* sp. RCC1657 was found to belong to a clade of silicified forms based on SSU rRNA gene sequences, despite its morphology (Ichinomiya et al., 2016). Third, a genome sequencing study of silicified strains revealed the presence of genes related to the flagellated form in their genomes led to the hypothesis that the hypothesized life cycle switching corresponds to an alternation between a silicified photoautotrophic stage and a naked flagellated phago-mixotrophic stage (Ban et al., 2023). Specifically, Ban et al. (2024) identified genes associated with phagocytosis in parmalean genomes—genes that have been lost in diatoms—and proposed that this mixotrophic strategy facilitates their adaptation to varying nutrient environments. Furthermore, a recent global metabarcoding study consistently revealed cosmopolitan distributions of parmaleans, which may be explained by their life cycle strategy (Ban et al., 2024). However, to our knowledge, the life cycle switches between the two morphological forms in a single parmalean has not been formally established, although Ichinomiya et al. (2016) mentioned the appearance of flagellated cells in cultures of silicified parmaleans (Ichinomiya et al., 2016). Therefore, the hypothesis on the life cycle and genome state of

parmaleans remains to be tested.

To assess the hypothesis, large-scale environmental omics resources provide a powerful
framework. The *Tara* Oceans expedition generated global-scale metagenomic and metatranscriptomic
datasets for surface and mesopelagic plankton (Sunagawa et al., 2020). Among its key outputs, the
Marine Atlas of *Tara* Oceans Unigenes (MATOU) provides a comprehensive catalog of eukaryotic gene
expression in the marine environment (Carradec et al., 2018).

In this study, we examine the hypothesis that the S-type and F-type forms represent distinct life-
cycle stages of a single parmalean. Under this hypothesis, genes specific to S-type or F-type forms
would be expressed from a single parmalean, possibly in different geographic locations or
environmental conditions. Specifically, we aim to clarify whether the genes related to silicified or
flagellated forms are expressed from individual parmalean populations (at the species level) in the ocean
and whether the geographic distributions of these gene expressions are environmentally regulated. To
achieve this, we first select marker genes representing the two distinct morphological forms based on
differential gene expression between cultured strains based on publicly available transcriptome data. We
then examine the presence of these genes in two parmalean Metagenome-Assembled Genomes (MAGs)
derived from *Tara* Oceans metagenomic data (Delmont et al., 2022). By integrating these MAGs with
the *Tara* Oceans metagenome and MATOU dataset, we characterize the global expression patterns of
these marker genes and correlate them with the environmental parameters.

Materials and Methods

Transcriptome data

We analyzed publicly available RNA-seq data from S-type and F-type parmalean strains. For the
S-type strain, raw reads of *Triparma laevis* f. *inornata* strain NIES-2565 [six samples: three silicate-rich
(concentration of Si, 50 μM) and three silicate-deficient (0.23 μM) replicates] were retrieved from NCBI
Sequence Read Archive (SRA) (BioProject PRJDB4267). Both conditions were treated as S-type in this
study because the overall expression profiles were quiasi-indistinguishable between conditions (see **Fig.
S1**), though a previous study reported that the strain tend to lose their silica plates under silicate-
deficient conditions (Yamada et al., 2014). For the F-type strains, data for four strains (*Triparma
eleuthera* CCMP1866, *Triparma* sp. RCC1657, RCC208, and RCC2347) were obtained from the
MMETSP database (Keeling et al., 2014).

Reads were quality-filtered with fastp v0.24.0 (Chen et al., 2018), rRNA-depleted with
SortMeRNA v4.3.7 (Kopylova et al., 2012) using the PR2 database v5.1.0 (Guillou et al., 2013),
assembled de novo with Trinity v2.15.2 (Grabherr et al., 2011), and redundancy-reduced using the
EvidentialGene pipeline (https://sourceforge.net/projects/evidentialgene/) to obtain non-redundant CDS
and protein sequences for each strain.

Gene expression quantification and completeness assessment

To quantify gene expression levels, the quality-filtered RNA-seq reads were mapped back to the
constructed non-redundant reference gene catalogs using Salmon (version 1.10.3) (Patro et al. 2017) in
quasi-mapping mode with default parameters. This provided transcript abundance estimates for each
sample, which were subsequently used for downstream differential expression analysis.

To assess the completeness of the assembled transcriptomes, we performed BUSCO v5.8.3
analysis (Manni et al., 2021). The non-redundant predicted protein sequences from the five

transcriptomes were evaluated against the stramenopile_odb12 lineage dataset. For comparative
purposes, the same BUSCO assessment was also performed on the coding sequences of eight publicly
available parmalean genomes.

Orthology inference

To investigate evolutionary relationships and identify core gene sets, we performed an orthology
analysis using OrthoFinder v2.3.8 (Emms & Kelly, 2019). The input dataset included the predicted
protein sequences from (1) the five assembled transcriptomes (one S-type and four F-types described
above), (2) eight publicly available parmalean genomes, (3) two parmalean Metagenome-Assembled
Genomes (MAGs) retrieved from the *Tara* Oceans dataset: TARA_ARC_108_MAG_00221 (ARC-
MAG) and TARA_PON_109_MAG_00217 (PON-MAG), and (4) three diatom genomes (*Thalassiosira
pseudonana*, *Pseudo-nitzschia multistriata*, *Fragilariopsis cylindrus*) included as outgroups.

Phylogenetic analysis

144 Based on the OrthoFinder results, single-copy orthologous genes conserved across all genomes
145 and transcriptomes were extracted to reconstruct the phylogeny. The sequences were concatenated and
146 aligned, and a maximum likelihood phylogenetic tree was constructed using IQ-TREE 3.0.1 (Wong et
147 al., 2025). ModelFinder (Kalyaanamoorthy et al., 2017) was used to determine the best-fit substitution
148 model, and branch support was assessed using ultrafast bootstrapping (Hoang et al., 2018).

Identification of morphotype-specific marker genes

150 From the OrthoFinder output, we extracted orthogroups conserved across all eight Parmales
151 reference genomes and filtered for genes with high expression levels (logCPM (Counts per million) > 5).
152 Marker genes were then classified based on their exclusive expression in either S-type or F-type
153 cultures. This conservative classification strategy was adopted to minimize false positives arising from
154 inter-strain differences.

Metagenomic and metatranscriptomic mapping to MAGs

Morphotype-specific marker gene sequences were extracted from the two parmalean MAGs
(ARC-MAG and PON-MAG) based on orthogroup assignments. Metagenomic and metatranscriptomic
reads were mapped to these MAG-derived marker genes using bwa-mem2 v2.2.1 (Vasimuddin et al.,
2019) with default parameters. The resulting alignments were sorted and deduplicated using samtools
v1.18 (Danecek et al., 2021). To ensure high-confidence mapping (Carradec et al., 2018), reads were
filtered using bamFilters v1.10.2 (https://github.com/institut-de-genomique/bamFilters) with a minimum
sequence identity of 95% and a minimum alignment length coverage of 90%. Finally, mapped reads
were extracted, and raw read counts were aggregated per sample to construct a count matrix.

Clustering and visualization of expression patterns

165 A Bray-Curtis dissimilarity matrix was calculated from the marker gene expression matrix using
166 the vegan package v2.7-1 in R v4.5.1, and hierarchical clustering was performed using the Ward.D2
167 method. Results were visualized as heatmaps with clustering dendrograms using the pheatmap package
168 v1.0.13. Expression data were also integrated with *Tara* Oceans environmental metadata.

Environmental modeling

170 To investigate the relationships between morphotype expression and environmental factors, we
171 employed Generalized Additive Models (GAMs) using the mgcv package v1.9.3. In this modeling
172 framework, the response variable was defined as a normalized expression index calculated using log10-

173 transformed reads with a pseudo count of 1. Specifically, the index for each type (F or S) was calculated
174 as:

$$\text{Index} = \frac{\log_{10}(T_{type} + 1)}{\log_{10}(G_{total} + 1)}$$

where T_{type} represents metatranscriptomic read counts for type-specific marker genes (F- or S-type),
and G_{total} represents the sum of metagenomic read counts for both F- and S-type genes. Samples with
low gene abundance ($G_{total} < 10$) were excluded from the analysis. The explanatory variables included
nutrient concentrations (total carbon, PO_4, Si, and NO_2+NO_3) and physical parameters (temperature,
salinity, and density). Sampling depth (Surface [SRF] or Deep Chlorophyll Maximum [DCM]) was
included as a categorical factor. Prior to analysis, we confirmed that the Variance Inflation Factor (VIF)
for all continuous predictors was less than 10. The model assumed a Tweedie error distribution with a
log link function, and parameter estimation was performed using REML with the "select = TRUE"
option enabled. This applies additional shrinkage, allowing the model to shrink non-contributing smooth
terms toward zero, effectively removing them.

Results

De novo transcriptome assembly of five cultured Parmales strains

To explore possible morphotype-specific genes in parmaleans, we generated de novo RNA-seq
assembly for five strains: the silicified strain NIES-2565 (n=6) and four flagellated strains (CCMP1866,
RCC1657, RCC208, and RCC2347; n=1 each), yielding 19,000–30,000 transcripts per strain (**(6)**).
BUSCO analysis (stramenopiles_odb12) confirmed completeness scores exceeding 60% for all
assemblies, which is comparable to the reference complete genomes.

Identification of morphology-specific markers

Prior to marker gene selection, we assessed whether the silicate-rich and silicate-deficient culture
conditions of NIES-2565 produced distinct expression profiles. A Principal Coordinates Analysis
(PCoA) based on Bray-Curtis dissimilarity of ortholog-mapped read counts across all ten samples
revealed that the first principal coordinate (PCoA1), accounting for 63.3% of the total variance, clearly
separated the four F-type strains from all six NIES-2565 samples. The silicate-rich and silicate-deficient
replicates of NIES-2565 were intermingled along this axis, indicating that culture condition had a
negligible effect on global expression patterns relative to the morphotype difference. Therefore, all six
NIES-2565 samples were treated as S-type replicates in subsequent marker gene selection.

To identify genes specific to S-type or F-type forms, we conducted an ortholog analysis of 18
strains (five transcriptomes, eight parmalean genomes, and five diatom outgroups) and yielded 25,752
orthogroups. Of 5,638 orthogroups conserved across all eight reference genomes ("Parmales core gene
set"), 39 were expressed exclusively in silicified cultures and 652 exclusively in flagellated cultures.
Applying a stringent expression threshold (logCPM > 5) resulted in 30 S-type and 180 F-type marker
genes (**Fig. 1**). Functional annotation confirmed that S-type markers included the silicon transporter
(SIT) and F-type markers contained intraflagellar transport (IFT) genes, validating their correspondence
to the silica cell wall and flagella, respectively. ARC-MAG and PON-MAG, the two parmalean MAGs
that we analyze in this study, encoded 92% (166/181) of F-type and 73% (22/30) of S-type markers,
while PON-MAG encoded 81% (139/181) and 90% (26/30), respectively.

Phylogenetic relationships and MAG placement

To ensure our MAGs were representatives of the Parrmales known diversity, we inferred A phylogenetic tree based on 75 concatenated single-copy orthologous genes, which reproduced the established classification of Parmales with full bootstrap support (**Fig. 2**). The two MAGs were placed in distinct clades: ARC-MAG in the Lepidoparmales (Clade I) and PON-MAG in the *Tetraparma* group (Clade IV), representing two phylogenetically distant lineages. Average Nucleotide Identity (ANI) analysis further confirmed the ARC-MAG's close affinity to the *Lepidoparma frigida* reference genome (99.4% identity).

In situ expression of both morphotype markers from individual MAGs

To investigate the geographic distributions of S-type and F-type markers, we mapped *Tara* Oceans metagenomic and metatranscriptomic reads to the two MAGs. Metagenomic abundances (i.e., "presence") of S-type and F-type markers were highly correlated within each MAG, confirming stable genomic copy number ratios (**Fig. S2**, **Table 2**). In contrast, metatranscriptomic abundances (i.e., "activity") varied substantially across samples, suggesting that marker expression is dynamically regulated by local environmental conditions.

Mapping *Tara* Oceans metatranscriptomic reads to the two MAGs revealed that all of the S-type and F-type marker genes encoded in the MAGs were transcriptionally active at least in one sample. Hierarchical clustering of expression profiles showed that S-type and F-type markers formed distinct clusters within each MAG (**Fig. 3**), indicating that the two marker sets are independently regulated.

Spatially, metagenomic distributions were synchronized between markers, whereas transcriptomic patterns diverged (**Fig. 4**). ARC-MAG transcription was restricted to polar regions, with F-type markers notably upregulated in the Atlantic inflow at the DCM layer (**Fig. 4d**). Conversely, PON-MAG transcripts were concentrated in low-to-middle latitudes, showing higher abundance in the DCM (Fig. 4h) than in surface waters (Fig. 4g). While F-type transcripts remained prevalent, S-type expression prominently increased within the DCM, especially in the Eastern Pacific.

Distinct scaling regimes of morphotype-specific gene expression

The relationship between total gene abundance and S-type transcript abundance revealed two distinct scaling regimes (**Fig. 5**). In the high-expression regime (Group 1), S-type transcripts scaled near-linearly with gene abundance (slopes: 1.18 ± 0.12 for ARC-MAG, 0.90 ± 0.08 for PON-MAG; $R_2 = 0.82$ and 0.86, respectively), indicating silicification gene expression proportional to relative genome abundance within the plankton community. In the basal regime (Group 2), scaling was markedly weaker (slopes: 0.54 ± 0.07 and 0.28 ± 0.07; $R_2 = 0.24$ and 0.08), with S-type transcript levels remaining low and stochastic regardless of gene abundance. In contrast, F-type transcript abundance scaled uniformly with gene abundance without exhibiting such regime separation (slopes: 0.80 ± 0.09 and 0.87 ± 0.06; $R_2 = 0.28$ and 0.54). These results suggest that S-type gene expression requires specific environmental triggers that lead to a higher population density, whereas F-type expression is constitutively maintained.

Environmental drivers of morphology-specific gene expression

We further explored the environmental parameters that may affect expression profiles of morphotypes marker genes. GAMs revealed contrasting environmental dependencies between the two marker types (**Fig. 6, Fig. S3**). Notably, S-type models explained substantially more deviance (48.8% for ARC-MAG, 26.0% for PON-MAG) than F-type models (19.7% and 5.8%), indicating that S-type expression is more tightly coupled to abiotic variables. In the ARC-MAG (n=146), S-type expression was significantly associated with PO_4, Si, temperature, salinity, and total carbon, whereas F-type expression responded primarily to temperature and NO_2+NO_3. In the PON-MAG (n=142), temperature

256 was the dominant driver for both types, with PO_4 additionally influencing S-type and NO_2+NO_3
257 influencing F-type expression. Sampling depth was not significant in any model.

Discussion

In this study, we detected the expression of marker genes for both S- and F-type forms from
individual parmalean MAGs. While Ban et al. (2023) previously demonstrated that parmaleans (S-type)
possess the genomic repertoire for these distinctive functions (Ban et al., 2023), our findings indicate,
for the first time, that environmental parmalean populations (at the species level) manifest gene
expression for both morphotypes. Previously, the hypothesis that these two forms belong to the same
species was primarily inferred from molecular phylogenies (Ichinomiya et al., 2016). Our results support
the view that these dual genetic traits are functional in actual marine environments, at least within the
lineages represented by the two analyzed MAGs.

The global distribution patterns of the two MAGs align well with the biogeography of parmalean
clades elucidated by Ban et al. (2024): ARC-MAG was predominantly found in polar and subarctic
regions, mirroring Clade I (Lepidoparmales), while PON-MAG was detected in tropical and temperate
regions, corresponding to the warmer-water populations within the ubiquitous Clade IV.

A notable observation was the contrasting distribution patterns between S-type and F-type
marker genes at the transcriptomic level, despite their stable genomic ratios across all sites. This
genomic stability indicates that the geographically distinct expression patterns are driven by differential
transcriptional activity rather than differences in intra-species genetic variations. This finding offers a
coherent explanation for the long-standing discrepancy between the global ubiquity reported in
Amplicon Sequence Variant (ASV)-based surveys and the restricted range of silicified strain isolations:
populations in warmer waters are likely represented primarily by the flagellated form.

The restriction of the S-type phase appears to be governed by environmental parameters. Our
scaling analysis revealed two distinct regimes of S-type gene expression (**Fig. 5**), which, together with
the environmental modeling (**Fig. 6**), suggests that conditions favorable for biomass increase trigger
active silicification. In short, S-type cells appear to increase only under conditions favorable for growth,
while F-type cells appear to tolerate various conditions but their response to growth-favoring condition
is not as sharp as S-type. This interpretation aligns with culture experiments showing that the S-type is
cold-adapted with limited growth above 15°C (Ichinomiya et al., 2013), whereas the F-type exhibits
broader thermal tolerance (Stawiarski et al., 2016). Thus, the "ubiquitous" distribution of Clade IV in
ASV studies likely reflects the capacity of the F-type to inhabit warmer waters, while the S-type is
confined to niches satisfying specific growth conditions.

This niche separation is illustrated by the PON-MAG in tropical regions, where S-type
expression was substantially lower in surface waters than in the DCM, consistent with microscopic
observations of silicified parmaleans in deep photic layers (Fujita & Jordan, 2017). The historical reports
of parmalean "absence" in tropical surface waters likely reflect suppressed silicification below detection
limits. Conversely, F-type markers were robustly detected in these surface layers, suggesting that the
flagellated form inhabits this warm niche, remaining cryptic to morphological observation. The presence
of the F-type in these typically oligotrophic surface waters may reflect differences in trophic strategies.
Ban et al. (2023) hypothesized dimorphic trophic modes: primarily autotrophic for the S-type and
phago-mixotrophic for the F-type (Ban et al., 2023). Our data are compatible with this hypothesis, as S-
type markers prevailed in the nutrient-rich DCM whereas F-type markers dominated in nutrient-depleted

surface waters where bacterial grazing could sustain a mixotroph. Nevertheless, since the flagellated form is hypothesized to function as a gamete, its surface presence could partly reflect life-cycle transitions.

Our GAM analysis provides statistical support for these divergent strategies. S-type models demonstrated substantially higher explanatory power than F-type models, indicating that S-type expression is strongly coupled to abiotic variables while F-type expression is largely decoupled from measured physicochemical parameters. This contrast reinforces the hypothesis of distinct trophic modes: strict autotrophy would be directly regulated by inorganic nutrients, whereas a mixotrophic F-type driven by prey availability would not be captured by standard environmental metadata. Among the specific drivers, S-type expression showed consistent positive relationships with PO_4 in both MAGs. For the ARC-MAG, the additional association with lower salinity suggests an affinity for river-influenced, nutrient-rich Arctic waters, while the negative relationship with Si may reflect rapid drawdown by actively silicifying populations.

Building on these contrasting environmental dependencies, we propose a threshold-based model for the morphological switch in parmaleans. Our scaling analysis revealed that S-type accumulation consistently occurred above a minimum abundance threshold, suggesting that the transition from flagellated to silicified phases is not gradual but requires conditions sufficient to sustain stable growth. Below this threshold—as observed in oligotrophic surface waters—the community appears to persist primarily in the F-type form, whose mixotrophic capacity enables survival in nutrient-poor environments where strict autotrophy would be untenable. Once nutrient and other conditions reach a certain level that sustain S-type growth, the morphological switch is triggered and both phases accumulate within the community. This model reconciles the cryptic, F-type-dominated populations in warm oligotrophic waters with the conspicuous blooms of silicified cells reported in nutrient-rich regions (11).

These insights must be interpreted with caution. The low abundance of Parmales [<0.2% of the community (Ichinomiya et al., 2016; Kuwata et al., 2018)] limited the number of mapped reads, and missing environmental metadata necessitated sample exclusions that may have introduced bias. To validate the life cycle model, it is necessary to move beyond static distributional snapshots. While culture-based direct observation of S–F transitions remain the ultimate goal, field-based temporal dynamics data would offer an important alternative. Priority should be given to high-abundance regions absent from our dataset, such as the Western North Pacific (Oyashio) and the east coast of Australia, where Clade II blooms are known to occur (Ban et al., 2024). Integrating targeted environmental profiling will be an effective approach to elucidate the environmental drivers of morphological switching in this unique lineage.

Acknowledgement

We thank the *Tara* Oceans consortium and the people and sponsors who supported the *Tara* Oceans Expedition (http://www.embl.de/tara-oceans/) for making the data accessible. This is contribution number XXX of the *Tara* Oceans Expedition 2009–2013. Computational time was provided by the Supercomputer System, Institute for Chemical Research, Kyoto University.

Funding

This work was supported by the Collaborative Research Program of the Institute for Chemical Research, Kyoto University (2016-30), JSPS KAKENHI Grant Numbers JP26291085, JP15K14784, and

JP17H03724. The sequencing of *Triparma laevis* f. *inornata* strain NIES-2565 (BioProject PRJDB4267) was supported by JSPS KAKENHI Grant Number JP221S0002 (Grant-in-Aid for Scientific Research on Innovative Areas 'Genome Science').

Conflicts of Interest

The authors declare no conflicts of interest.

Reference

Armbrust, E. V. (2009). The life of diatoms in the world's oceans. *Nature*, *459*(7244), 185–192. https://doi.org/10.1038/nature08057

Ban, H., Endo, H., Kuwata, A., & Ogata, H. (2024). Global Distribution and Diversity of Marine Parmales. *Microbes and Environments*, *39*(1), ME23093. https://doi.org/10.1264/jsme2.ME23093

Ban, H., Sato, S., Yoshikawa, S., Yamada, K., Nakamura, Y., Ichinomiya, M., Sato, N., Blanc-Mathieu, R., Endo, H., Kuwata, A., & Ogata, H. (2023). Genome analysis of Parmales, the sister group of diatoms, reveals the evolutionary specialization of diatoms from phago-mixotrophs to photoautotrophs. *Communications Biology*, *6*(1), Article 1. https://doi.org/10.1038/s42003-023-05002-x

Booth, B. C., Lewin, J., & Norris, R. E. (1980). Siliceous nanoplankton I. Newly discovered cysts from the Gulf of Alaska. *Marine Biology*, *58*(3), 205–209. https://doi.org/10.1007/BF00391877

Booth, B. C., & Marchant, H. J. (1987). Parmales, a New Order of Marine Chrysophytes, with Descriptions of Three New Genera and Seven New Species1,2. *Journal of Phycology*, *23*(s2), 245–260. https://doi.org/10.1111/j.1529-8817.1987.tb04132.x

Carradec, Q., Pelletier, E., Da Silva, C., Alberti, A., Seeleuthner, Y., Blanc-Mathieu, R., Lima-Mendez, G., Rocha, F., Tirichine, L., Labadie, K., Kirilovsky, A., Bertrand, A., Engelen, S., Madoui, M.-A., Méheust, R., Poulain, J., Romac, S., Richter, D. J., Yoshikawa, G., … Wincker, P. (2018). A global ocean atlas of eukaryotic genes. *Nature Communications*, *9*(1), Article 1. https://doi.org/10.1038/s41467-017-02342-1

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, *34*(17), i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools and BCFtools. *GigaScience*, *10*(2), giab008. https://doi.org/10.1093/gigascience/giab008

Delmont, T. O., Gaia, M., Hinsinger, D. D., Frémont, P., Vanni, C., Fernandez-Guerra, A., Eren, A. M., Kourlaiev, A., d'Agata, L., Clayssen, Q., Villar, E., Labadie, K., Cruaud, C., Poulain, J., Da Silva, C., Wessner, M., Noel, B., Aury, J.-M., Sunagawa, S., … Jaillon, O. (2022). Functional repertoire convergence of distantly related eukaryotic plankton lineages abundant in the sunlit ocean. *Cell Genomics*, *2*(5), 100123. https://doi.org/10.1016/j.xgen.2022.100123

Emms, D. M., & Kelly, S. (2019). OrthoFinder: Phylogenetic orthology inference for comparative genomics. *Genome Biology*, *20*(1), 238. https://doi.org/10.1186/s13059-019-1832-y

Fujita, R., & Jordan, R. W. (2017). Tropical Parmales (Bolidophyceae) assemblages from the Sulu Sea and South China Sea, including the description of five new taxa. *Phycologia*, *56*(5), 499–509. https://doi.org/10.2216/16-128.1

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., … Regev, A. (2011). Full-length

transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology*, *29*(7), 644–652. https://doi.org/10.1038/nbt.1883

Green, J. C., Course, P. A., & Tarran, G. A. (1996). The life-cycle of Emiliania huxleyi: A brief review and a study of relative ploidy levels analysed by flow cytometry. *Journal of Marine Systems, EHUX (Emiliania Huxleyi)*, *9*(1), 33–44. https://doi.org/10.1016/0924-7963(96)00014-0

Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., Boutte, C., Burgaud, G., de Vargas, C., Decelle, J., del Campo, J., Dolan, J. R., Dunthorn, M., Edvardsen, B., Holzmann, M., Kooistra, W. H. C. F., Lara, E., Le Bescot, N., Logares, R., … Christen, R. (2013). The Protist Ribosomal Reference database (PR2): A catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids Research*, *41*(D1), D597–D604. https://doi.org/10.1093/nar/gks1160

Guillou, L., Chrétiennot-Dinet, M.-J., Medlin, L. K., Claustre, H., Goër, S. L., & Vaulot, D. (1999). Bolidomonas: A New Genus with Two Species Belonging to a New Algal Class, the Bolidophyceae (heterokonta). *Journal of Phycology*, *35*(2), 368–381. https://doi.org/10.1046/j.1529-8817.1999.3520368.x

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2018). UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution*, *35*(2), 518–522. https://doi.org/10.1093/molbev/msx281

Ichinomiya, M., dos Santos, A. L., Gourvil, P., Yoshikawa, S., Kamiya, M., Ohki, K., Audic, S., de Vargas, C., Noël, M.-H., Vaulot, D., & Kuwata, A. (2016). Diversity and oceanic distribution of the Parmales (Bolidophyceae), a picoplanktonic group closely related to diatoms. *The ISME Journal*, *10*(10), Article 10. https://doi.org/10.1038/ismej.2016.38

Ichinomiya, M., & Kuwata, A. (2015). Seasonal variation in abundance and species composition of the Parmales community in the Oyashio region, western North Pacific. *Aquatic Microbial Ecology*, *75*(3), 207–223. https://doi.org/10.3354/ame01756

Ichinomiya, M., Nakamachi, M., Shimizu, Y., & Kuwata, A. (2013). Growth characteristics and vertical distribution of Triparma laevis (Parmales) during summer in the Oyashio region, western North Pacific. *Aquatic Microbial Ecology*, *68*, 107–116. https://doi.org/10.3354/ame01606

Ichinomiya, M., Yoshikawa, S., Kamiya, M., Ohki, K., Takaichi, S., & Kuwata, A. (2011). Isolation and Characterization of Parmales (heterokonta/Heterokontophyta/Stramenopiles) from the Oyashio Region, Western North Pacific1. *Journal of Phycology*, *47*(1), 144–151. https://doi.org/10.1111/j.1529-8817.2010.00926.x

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., & Jermiin, L. S. (2017). ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods*, *14*(6), 587–589. https://doi.org/10.1038/nmeth.4285

Kamakura, S., Yamada, K., Arguelles, E. D. L. R., Kuwata, A., Nagumo, T., Takahashi, M., Sawada, K., Ichinomiya, M., & Sato, S. (2025). Reminiscent of the pre-diatom? A hitherto undescribed scaly bolidophyte Lepidoparma frigida gen. et sp. nov. in a new order Lepidoparmales based on morphology, phylogeny, and ecology. *Journal of Phycology*, *61*(4), 757–776. https://doi.org/10.1111/jpy.70043

Keeling, P. J., Burki, F., Wilcox, H. M., Allam, B., Allen, E. E., Amaral-Zettler, L. A., Armbrust, E. V., Archibald, J. M., Bharti, A. K., Bell, C. J., Beszteri, B., Bidle, K. D., Cameron, C. T., Campbell, L., Caron, D. A., Cattolico, R. A., Collier, J. L., Coyne, K., Davy, S. K., … Worden, A. Z. (2014). The Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): Illuminating the Functional Diversity of Eukaryotic Life in the Oceans through Transcriptome Sequencing. *PLOS Biology*, *12*(6), e1001889. https://doi.org/10.1371/journal.pbio.1001889

Kessenich, C. R., Ruck, E. C., Schurko, A. M., Wickett, N. J., & Alverson, A. J. (2014). Transcriptomic

Insights into the Life History of Bolidophytes, the Sister Lineage to Diatoms. *Journal of Phycology*, *50*(6), 977–983. https://doi.org/10.1111/jpy.12222

Kopylova, E., Noé, L., & Touzet, H. (2012). SortMeRNA: Fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics*, *28*(24), 3211–3217. https://doi.org/10.1093/bioinformatics/bts611

Kuwata, A., Yamada, K., Ichinomiya, M., Yoshikawa, S., Tragin, M., Vaulot, D., & Lopes dos Santos, A. (2018). Bolidophyceae, a Sister Picoplanktonic Group of Diatoms – A Review. *Frontiers in Marine Science*, *5*. https://www.frontiersin.org/articles/10.3389/fmars.2018.00370

Mann, D., & Marchant, H. J. (1989). The origin of the diatom and its life cycle. In *The Chromophyte Algae: Problems and Perspectives* (Vol. 38, pp. 305–321).

Manni, M., Berkeley, M. R., Seppey, M., & Zdobnov, E. M. (2021). BUSCO: Assessing Genomic Data Quality and Beyond. *Current Protocols*, *1*(12), e323. https://doi.org/10.1002/cpz1.323

Silver, M. W., Mitchell, J. G., & Ringo, D. L. (1980). Siliceous nanoplankton. II. Newly discovered cysts and abundant choanoflagellates from the Weddell Sea, Antarctica. *Marine Biology*, *58*(3), 211–217. https://doi.org/10.1007/BF00391878

Stawiarski, B., Buitenhuis, E. T., & Le Quéré, C. (2016). The Physiological Response of Picophytoplankton to Temperature and Its Model Representation. *Frontiers in Marine Science*, *3*. https://www.frontiersin.org/articles/10.3389/fmars.2016.00164

Sunagawa, S., Acinas, S. G., Bork, P., Bowler, C., Eveillard, D., Gorsky, G., Guidi, L., Iudicone, D., Karsenti, E., Lombard, F., Ogata, H., Pesant, S., Sullivan, M. B., Wincker, P., & de Vargas, C. (2020). Tara Oceans: Towards global ocean ecosystems biology. *Nature Reviews Microbiology*, *18*(8), Article 8. https://doi.org/10.1038/s41579-020-0364-5

Vasimuddin, Md., Misra, S., Li, H., & Aluru, S. (2019). Efficient Architecture-Aware Acceleration of BWA-MEM for Multicore Systems. *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS)*, 314–324. https://doi.org/10.1109/IPDPS.2019.00041

Wong, T. K. F., Ly-Trong, N., Ren, H., Baños, H., Roger, A. J., Susko, E., Bielow, C., Maio, N. D., Goldman, N., Hahn, M. W., Huttley, G., Lanfear, R., & Minh, B. Q. (2025). *IQ-TREE 3: Phylogenomic Inference Software using Complex Evolutionary Models*. https://ecoevorxiv.org/repository/view/8916/

Yamada, K., Yoshikawa, S., Ichinomiya, M., Kuwata, A., Kamiya, M., & Ohki, K. (2014). Effects of Silicon-Limitation on Growth and Morphology of Triparma laevis NIES-2565 (Parmales, Heterokontophyta). *PLoS ONE*, *9*(7), e103289. https://doi.org/10.1371/journal.pone.0103289

Figure 1. Genes specific to Parmales morphotypes. (a, b) UpSet plots illustrating the distribution of orthologs among
different Parmales strains with transcriptome data. The numbers above the vertical bars (Intersection Size) indicate the
number of orthologs shared by a specific combination of strains. The horizontal bars on the left (Set Size) show the
total number of orthologs within each strain. The red bar corresponds to the orthologs unique to the S-type
transcriptome, while the teal bar represents the orthologs identified in all F-type transcriptomes but not in the S-type.
Panel (a) includes all orthologs that were found to be universally conserved in eight Parmales strain with genomic
sequence data. In panel (b), only orthologs with an expression level of logCPM ≥ 5.0 are shown. (c) MA plot showing
genes specifically expressed in individual morphotypes. Red points indicate orthologs selected as S-type specific and
blue ones indicate orthologs selected as F-type specific.

Figure 2. Phylogenetic relationships inferred from concatenated single-copy genes. Maximum likelihood phylogenetic tree of concatenated alignments of single-copy orthologous genes. IDs shown in blue represent the phylogenetic placement of assemblies derived from RNA-seq data. IDs shown in red indicate the positions of parmalean metagenome-assembled genomes (MAGs) from the *Tara* Oceans dataset. Nodal support is provided by 1,000 ultrafast bootstrap replicates (UFB, first value) and SH-aLRT (SH, second value).

Figure 3. Expression profiles of S-marker and F-marker genes in two parmalean MAGs. Heatmaps displaying the
similarity of expression patterns among S- and F-markers. The Bray-Curtis dissimilarity among marker genes was
calculated based on the read counts of genes among samples. The color scale indicates the dissimilarity of expression
patterns across orthologs. The annotation bars on the top and left indicate the morphological types: green for F-type and
red for S-type. (a, b) show the clustering of all marker genes for the Arctic MAG (a) and Pacific Ocean North MAG
(b). (c, d) show the clustering of the 50 most highly expressed F-markers and all S-markers for the Arctic MAG (c) and
Pacific Ocean North MAG (d).

Figure 4. Global distribution and expression patterns of S- and F-type marker genes for two parmalean MAGs.
The maps show read counts from metagenomes (left column: a, c, e, g) and metatranscriptomes (right column: b, d, f, h). Bubble size is proportional to the number of reads, and color indicates the marker type (blue: F-type; red: S-type). Sites with read counts < 10 are shown as dots. Samples are organized by MAG and depth layer: (a, b) ARC-MAG in surface waters; (c, d) ARC-MAG in the Deep Chlorophyll Maximum (DCM) layer; (e, f) PON-MAG in surface waters; (g, h) PON-MAG in the DCM layer.

Figure 5. Scaling relationship between total gene abundance and morphotype-specific transcript abundance in ARC-MAGs and PON-MAGs. Log-log scatter plots showing the relationship between log10-transformed total gene abundance (metaG) and marker transcript abundance (metaT). (a, b) S-type markers for ARC-MAG and PON-MAG, respectively. Data points are divided into two groups: a manually defined subset showing a proportional response (Group 1, circles) and the remaining population (Group 2, triangles). (c, d) F-type markers for ARC-MAG and PON-MAG, respectively, showing uniform scaling without distinct regimes. Blue lines indicate linear regression fits with 95% confidence intervals (grey shading).

Figure 6. Generalized Additive Model (GAM) regression analysis of S-type specific expression variations based on environmental factors. Panels show partial effects of environmental variables on expression indices for the ARC-MAG dataset (n=146; a) and the PON-MAG dataset (n=142; b). The y-axis represents the partial effect (centered at zero) of each environmental variable on the expression index. Solid lines indicate the fitted smoothing curves, dashed lines represent 95% confidence intervals, and open circles show the partial residuals.

Table 1. Comparison of assembly statistics and BUSCO completeness between the de novo transcriptomes and reference Parmales genomes.

Species and strain	Accession	Number of sequences	Busco Completeness (percentage)
NIES_2565	PRJDB4267	29,568	589/697 (84.5 %)
CCMP1866	SAMN02740122	19,596	538/697 (77.2 %)
RCC1657	SAMN02740258	21,571	464/697 (66.6 %)
RCC208	SAMN02740123	23,225	473/697 (67.9 %)
RCC2347	SAMN02740277	19,213	425/697 (61.0 %)
Triparma laevis f.inornata	GCA_012489335.1	13,395	459/697 (65.9 %)
Tetraparma gracilis	GCA_030267505.1	15,987	383/697 (54.9%)
Triparma retinervis	GCA_030267625.1	13,713	456/697 (65.4 %)
Triparma laevis f.longispina	GCA_030267805.1	17,316	675/697 (96.8 %)
Triparma verrucosa	GCA_030267905.1	14,735	680/697 (97.6 %)
Triparma strigata	GCA_030267945.1	15,257	680/697 (97.6 %)
Triparma Columncea	GCA_030267965.1	14,430	667/697 (95.7 %)
Lepidoparma frigida	GCA_030267985.1	12,273	662/697 (95.0%)

Table 2. Coefficients of determination (R^2) between S-type and F-type marker abundances in metagenomic and metatranscriptomic datasets.

MAG	Dataset	Correlation (R^2)
ARC-MAG	Metagenome	0.5875
ARC-MAG	Metatranscriptome	0.3721
PON-MAG	Metagenome	0.8143
PON-MAG	Metatranscriptome	0.5834

Spatially, metagenomic distributions were synchronized between markers, whereas transcriptomic patterns diverged (Fig. 4). ARC-MAG transcription was restricted to polar regions, with F-type markers notably upregulated in the Atlantic inflow at the DCM layer (Fig. 4d). PON-MAG transcripts were concentrated in low-to-middle latitude DCM layers (Fig. 4h), contrasting with F-type dominance in surface waters (Fig. 4f) and localized S-type expression in the North Atlantic.