PARP1 regulates the genomic ribonucleotide processing activity of TOP1 to prevent the formation of toxic TOP1-DNA adducts and the associated mutations.

Edward James Sarrain[1#], Quan Wang[1#], Angel Caitlin Bondoy[1,2], Feng Guo[3], Qi Cao[4], Hengyao Niu[1]

[1] Department of Molecular and Cellular Biochemistry, Indiana University Bloomington, Bloomington, IN, 47405

[2] Present address: Department of Biochemistry and Cell Biology, Stony Brook University, Stony Brook, NY, 11794

[3] Department of Intelligent Systems Engineering, Indiana University Bloomington, IN, 47405

[4] Department of Urology, Feinberg School of Medicine, Northwestern University, Chicago, IL 60611

[#] Contributed equally.

To whom the correspondence should be addressed:

Hengyao Niu, Department of Molecular and Cellular Biochemistry, Indiana University Bloomington, Bloomington, IN 47405, USA
Email: hniu@iu.edu

Abstract

Ribonucleotides are frequently incorporated into the genome during DNA replication and are normally removed by RNase H2. Alternatively, DNA topoisomerase 1 (TOP1) can process genomic ribonucleotides through two sequential cleavage events, generating a 2′,3′-cyclic phosphate-terminated nick and, subsequently, a TOP1 cleavage complex (TOP1cc). In addition, TOP1cc formation can lead to 2- to 5-bp slippage deletions, recently identified as the ID4 cancer signature. Here, we show that human TOP1 is intrinsically mutagenic, with a tendency to undergo secondary cleavage and form TOP1 cleavage complexes. We further demonstrate that PARP1, a central single-strand break repair factor, suppresses TOP1 secondary cleavage, TOP1cc formation, and the resulting slippage mutations. This regulation requires PARP1 to interact with both TOP1 and the 2′,3′-cyclic phosphate-terminated nick produced by the initial TOP1 cleavage. Notably, a PARP1 mutant defective in TOP1 interaction is partially impaired in restoring PARP inhibitor sensitivity in PARP1-knockout cells, suggesting that PARP inhibitor cytotoxicity may partly result from PARP1 trapping at 2′,3′-cyclic phosphate-terminated nicks and blocking their processing by TOP1 and other factors. Together, these findings reveal a novel role for PARP1 in regulating TOP1-dependent ribonucleotide processing, thereby suppressing cancer-signature mutations and contributing to the mechanism of PARP inhibitor cytotoxicity.

Introduction

Ribonucleotides (rNMPs) are frequently incorporated into the genome by the major replicative polymerases. In mice, about one million ribonucleotides are incorporated into the genome per cell cycle, with human cells estimated to have an even higher accumulation (around 3 million ribonucleotides per genome).[1,2] The removal of single mis-incorporated ribonucleotides occurs mainly through the error-free ribonucleotide excision repair pathway (RER), which involves cleavage at the 5' of the embedded ribonucleotide by RNase H2 followed by strand displacement and DNA flap removal.[3,4] In humans, partial loss-of-function mutations in RNase H2 lead to a 10 to 25-fold increase in the abundance of genomic ribonucleotides.[5] Mutations in RNase H2-associated genes are linked to two types of cancers, chronic lymphocytic leukemia and metastatic castration-resistant prostate cancer, as well as autoimmune diseases, including Aicardi-Goutières syndrome and systemic lupus erythematosus.[6] Although the cancer pathology in RNase H2 mutants is understudied, the autoimmune pathology is believed to be the result of genomic instability in the absence of functional RNase H2 that subsequently leads to the activation of an innate immune response.[1,7] One of the potential sources for such genomic instability is the removal of genomic ribonucleotides by topoisomerase 1 (TOP1).

Biochemically, it has been long known that eukaryotic TOP1, including human TOP1, can cleave at the site of a single mis-incorporated ribonucleotide and generate a nick terminated by a 5'-OH and a 2', 3'-cyclic phosphate (CP).[8] Next, TOP1 can act at a CP nick in two ways. Besides a direct ligation that restores the embedded ribonucleotide, genetic studies from budding yeast indicate that TOP1 can initiate an alternative pathway to remove the ribonucleotide.[9-11] In this pathway, TOP1 removes the terminal CP via a secondary cleavage ~2-5 bp upstream of the ribonucleotide, which releases the CP-containing segment, but generates a trapped TOP1 cleavage complex (TOP1cc) due to the absence of a proximal end to ligate with. In DNA regions harboring short

91 tandem repeats, the TOP1cc from the secondary cleavage can catalyze ligation across the gap
92 following DNA misalignment, which leads to the formation of slippage mutations.[9] Notably, such
93 slippage mutations have been recently identified as the ID4 cancer signature, with a frequency
94 increase in human RNASEH2A knockout cells, RNase H2A being the catalytic subunit of the
95 RNase H2 trimeric complex (together with RNase H2B and H2C).[12,13]
96
While the TOP1-derived CP nick repair is well characterized in yeast, a major difference between
single-strand break (SSB) repair in yeast and humans is that human cells express poly(ADP-ribose)
polymerase 1 (PARP1), a universal SSB repair factor known for its ability to sense and signal
breaks in DNA. PARP1 is a nuclear enzyme that is activated by DNA breaks and catalyzes the
poly(ADP-ribosyl)ation (PARylation) of itself and other substrates to recruit and regulate different
repair proteins.[14-16] PARP inhibitors have become key agents in cancer therapy, where their
efficacy arises not only from blocking PARP1's catalytic use of NAD$^+$ ligand but also from
stabilizing PARP1-DNA complexes.[17,18] This process, known as PARP trapping, enhances
cytotoxicity in homologous recombination (HR)-deficient tumors, such as BRCA-deficient breast
and ovarian cancers.[19-21] Notably, RER-deficient cells are also hypersensitive to PARP inhibitor
treatments, linking the processing of genomic ribonucleotides by TOP1 to PARP-trapping
lesions.[22] This association prompted us to investigate genomic ribonucleotide excision by human
TOP1 and its regulation by PARP1. Our findings reveal the mutagenic nature of human TOP1 and
how PARP1 safeguards against the deleterious consequences of the ribonucleotide excision by
TOP1, including preventing the formation of TOP1-DNA adducts and associated cancer signature
mutations. Moreover, our work also sheds light on the mechanism of PARP inhibitor cytotoxicity.
113
114 **Results**
115
116 **Human TOP1 is prone to secondary cleavage and slippage mutations at genomic rNMP sites.**
117
TOP1 cleaves at a genomic rNMP site and generates a 2', 3'-cyclic phosphate (CP) terminated
nick.[8,23] First shown in yeast Top1, a second cleavage by TOP1 a few nucleotides (2-5 bp)
upstream from the CP nick may occur, which leads to the formation of TOP1cc and the release of
the CP-containing segment (**Fig 1a**).[24] When a duplex DNA substrate with a single embedded
ribonucleotide was treated with human TOP1 in a 20-min end point analysis, interestingly, the
predominant product type (as much as 60% of the total resulting species) was the TOP1cc product,
with a lesser amount of the CP nick-product (**Fig 1b)**. To determine whether the TOP1cc was
formed at the initial or the secondary cleavage site, the TOP1cc product was treated with proteinase
K, followed by yeast tyrosyl-DNA phosphodiesterase 1 (Tdp1) and DNA 3'-phophatase (Tpp1),
which together remove the TOP1 adduct and its bridging 3'-phosphate.[25,26] These treatments
generated a 25-nt product, corresponding to a secondary cleavage at 2-bp upstream of the CP nick
(**Fig 1c**). In a time-course analysis, the amount of TOP1cc product increased over time, while the
amount of CP nick product reduced slowly (**Supplementary Fig 2a**). We surmise that TOP1 is
prone to secondary cleavage rather than ligation following the CP nick formation. Indeed, when
TOP1 was titrated directly against a CP nick substrate, a clear preference for secondary cleavage
was revealed, with TOP1cc representing up to 67% of the resulting species and a lesser amount of
ligation product (up to 25% of the total products) (**Fig. 1d**). To further determine whether the
ligation product was derived from direct CP nick ligation or slippage ligation, we tested human

TOP1, with the yeast Top1 (yTop1) as a control, on a similar but shorter 45 bp CP nick substrate for a better separation of these two ligation products (**Fig 1e**). Interestingly, while a nearly equal amount of CP nick ligation and slippage ligation were observed for the human TOP1, yTop1 showcased a clear preference for CP nick ligation (over 3-fold more than the deletion product). Given the difference between human and yeast TOP1, we further compared their slippage mutation potential in the duplex substrate with an embedded ribonucleotide. In a time course analysis, human TOP1 again showcased an increased ability to generate deletion mutations compared to yTop1 (**Supplementary Fig 2b**). Notably, more substrate remained with yTop1 than with human TOP1 (~69% versus 29% at 20 nM, respectively). This difference in remaining substrate is not the result of a reduced ability of yTop1 to cleave rNMP, since both proteins displayed comparable activities on a suicide fork substrate that does not allow for ligation once the initial cut is made (**Fig. 1f**). Thus, the drastic difference between human and yeast TOP1 activity on embedded rNMP is likely because yTop1 prefers ligation at the CP nick site, while human TOP1 prefers the secondary cleavage.

PARP1 regulates TOP1 ribonucleotide processing by suppressing its secondary cleavage.

The secondary cleavage by TOP1 is directly linked to slippage mutations. The mutagenic nature of human TOP1 prompted us to probe its regulation for mutation suppression. Absent in yeasts, PARP1 is a first responder in single strand break repair in higher eukaryotes and has been reported to physically interact with TOP1, a known target for PARylation.[27-29] More recently, TOP1 cleavage of genomic ribonucleotides was shown to generate PARP-trapping lesions,[22] which motivated us to ask whether PARP1 is playing a role in regulating ribonucleotide excision by TOP1. PARP1 senses and is activated by DNA nicks. We first tested PARP1 in an electromobility shift assay (EMSA) using a mixture of equal amount of clean nick and CP nick substrates labeled with Cy5 and Cy3 respectively (**Supplementary Fig. 3a**). These results suggest that PARP1 recognizes both substrates equally well.

Interestingly, the addition of PARP1 to the TOP1 rNMP cleavage reaction drastically increased the amount of CP nick product (~3-fold), while reducing the amount of TOP1cc (~ 7-fold) (**Fig 2a**). Addition of NAD$^+$, the cofactor necessary for PARP1-catalyzed PARylation, exacerbated this change, biasing nearly 98% of the products towards CP nick. Surprisingly, the presence of NAD$^+$ in the reaction significantly increased the amount of remaining substrate (~80%) likely due to the inhibition of the initial rNMP cleavage, and not the promotion of CP nick ligation, since PARP1 inhibits both TOP1-catalyzed secondary cleavage and ligation when testing the CP nick substrate (**Fig 2b**). Importantly, substitution of PARP1 with PARP1-E988K, a catalytic dead mutant of PARP1, showed a similar effect as PARP1 without NAD$^+$ (**Supplementary Fig. 3b, c**), confirming PARP1 suppresses TOP1 secondary cleavage in both a catalytic-dependent and -independent manner, while it down-regulates TOP1-catalyzed initial rNMP cleavage only in a catalytic-dependent manner. To summarize, we surmise that upon the initial rNMP cleavage, PARP1 appears to protect the CP nick product from any further action by TOP1. The activated PARP1 also down-regulates TOP1 from further rNMP cleavage, but only in a catalytic-dependent manner.

PARP1 knockout increases the slippage mutation rate due to increased TOP1 secondary cleavage.

181
Our findings of PARP1 inhibition of TOP1 secondary cleavage strongly predict a role of PARP1
in suppressing slippage mutations. To test this hypothesis and ascertain TOP1 regulation by
PARP1 in cells, we adopted a slippage mutation reporter HeLa cell line that a 2-bp slippage
deletion will correct a frameshift mutation and restore puromycin resistance of the cells (kind gift
from Dr. Andrew P. Jackson, **Fig 3a**). We knocked out PARP1 in both the parental reporter cell
line and its derived RNASEH2A knockout cell lines (**Supplementary Fig 4a**). As expected from
previous studies, RNASEH2A knockout significantly increased the slippage mutation rate (~14-
fold) compared to the wild-type cell line (Fig 3a-i). Consistent with our hypothesis, PARP1
knockout alone increased the slippage mutation rate by over 7-fold when compared to the wild-
type. DNA sequencing of puromycin resistant clones from the PARP1 knockout cell line
confirmed all 10 clones picked were 2-bp slippage events at STRs (**Fig 3a-ii, Supplementary
Table 4**). Being previously reported as having G2/M arrest,[22] RNASEH2A PARP1 double
knockout cell line displayed a substantial slow growth phenotype, rendering the reporter assay
technically challenging (**Fig 3b**). In addition to the homozygous RNASEH2A knockout, a partial
deletion of RNASEH2A, namely RNASEH2A$^+$, was also reported with around 50% loss of RNase
H2 activity and causes a minor increase of slippage mutation rate.[12] The knocking out of PARP1
in RNASEH2A$^+$ led to an over 12-fold increase in the slippage mutation rate over the wild-type,
above 5-fold over RNASEH2A$^+$ and almost 2-fold over PARP1-KO (**Fig. 3a-i**). Hence, the
impacts of RNASEH2A deficiency and PARP1-KO on slippage mutation rate are additive.
Interestingly, the inhibition of PARP1 through PARP inhibitor talazoparib did not significantly
change the mutation rate observed in the wild-type reporter cell line (**Supplementary Fig 4b**).
This inhibition of deletions even in the absence of the catalytic activity of PARP1 indicates that
PARP-trapping might compensate by physically impeding the TOP1 secondary cleavage.
205
206 **Slow growth of RNASEH2A PARP1 double knockout cell line due to TOP1cc accumulation
207 and its suppression by TDP1 overexpression.**
208
Based on our biochemically derived model, we suspected that the markedly reduced proliferation
and G2/M accumulation in RNASEH2A-KO PARP1-KO cells is the result of excessive TOP1
secondary cleavage and thus the accumulation of stable TOP1cc. Navitoclax, a BCL2 inhibitor
(BCL2i) that blocks the apoptosis suppressor BCL2, has been shown to enhance the cytotoxicity
of the TOP1-trapping agent camptothecin in tumors, likely due to the pro-senescent role of
BCL2.[30-34] Consistent with our hypothesis of TOP1cc accumulation, the RNASEH2A PARP1
double knockout cells demonstrated hypersensitivity to navitoclax (**Fig. 4a**). To confirm the
TOP1cc accumulation in the double knockout cells, we performed a RADAR (rapid approach to
DNA adduct recovery) assay designed to detect the amount of TOP1cc.[35,36] Our results showed
that the double knockout cell line has a 3-fold increase in the formation of TOP1cc when compared
to wild-type and significantly higher than the single knockouts of either RNASEH2A or PARP1,
with a difference of ~1.5 and 2-fold respectively (**Fig 4b**). To test if the buildup of TOP1cc was
the cause for the slow growth, we overexpressed TDP1, a protein that is crucial for TOP1cc
removal in cells,[25] which almost completely rescued the growth defect of our double knockout
cells (**Fig 4c**). Moreover, the treatment of the double knockout cell lines with a TDP1 inhibitor
(TDP1i), CD00509,[37] led to an over 60% reduction in viability (**Fig 4c**). This TDP1i sensitivity
was further aggravated by BCL2 inhibition, affirming TOP1cc as the toxic lesion slowing the

proliferation of the RNASEH2A-KO PARP1-KO cells (**Fig. 4d**). These results confirmed the role of PARP1 in the suppression of the detrimental TOP1 secondary cleavage at genomic rNMP sites in cells.

A feedback loop of CP nick formation, PARP1 activation and TOP1 inhibition.

In the reconstituted system, catalytically, PARP1 inhibits TOP1-catalyzed secondary cleavage and CP nick ligation, as well as the initial rNMP cleavage. Analysis of the PARylation status of the proteins in the reaction containing the embedded ribonucleotide substrates confirmed that TOP1 was PARylated in the presence of NAD^+ (**Fig 5a, 5b**), which is consistent with previous proteomic studies.[28,29] Notably, the presence of TOP1 stimulated PARP1 auto-PARylation in the presence of rNMP embedded duplex DNA. Treating the reactions with Poly (ADP-ribose) glycohydrolase (PARG), an enzyme that removes the ribose chains from proteins and leaves a single ADP-ribose in the modified residues,[38] further confirmed TOP1 PARylation and revealed an over 5-fold increase in PARP1 auto-PARylation when TOP1 was present (**Fig 5b**). This stimulation in auto-PARylation is likely the result of the formation of CP nick that then activates PARP1, as this stimulation was not observed with the catalytically inactive TOP1-Y723F (**Fig 5c**).[39,40] Therefore, the CP nick formed after initial rNMP cleavage by TOP1 activates PARP1, which then PARylates TOP1 and inhibits TOP1-catalyzed cleavage of additional rNMP sites (diagram shown in **Fig 5d**).

Both DNA binding and TOP1 interaction are essential for PARP1 regulation of TOP1 ribonucleotide processing.

In addition to catalytical regulation of TOP1 by PARP1, we also found that PARP1 inhibits CP nick cleavage and ligation by TOP1 in a non-catalytic manner. However, the DNA binding domain (DBD) harboring zinc finger motif 1, 2 and 3 was unable to suppress the secondary cleavage by TOP1, nor does the N-terminal truncation that removes the DNA binding domain (DBD) of PARP1 (**Fig 6a, b**). Hence, the binding of PARP1 to CP nick is necessary, but not sufficient, for TOP1 secondary cleavage inhibition. This led us to ask if the reported physical interaction between PARP1 and TOP1 is also essential for TOP1 regulation.[27] To map the interaction between PARP1 and TOP1, we generated a series of PARP1 truncations, starting from the C-terminus, and tested them for TOP1 interaction using *in vitro* pull-down assays (**Fig 6c**). Interestingly, the removal of its adjoining helical domain (HD), an autoinhibitory motif that locks PARP1 activity before binding to its substrate DNA, completely disrupted PARP1 and TOP1 interaction (**Fig. 6d**).[41,42] These results suggest the HD domain as a key interface between PARP1 and TOP1.

PARP1-R735E mutation partially disrupts its interaction with TOP1 and fails to fully restore PARP inhibitor sensitivity in PARP1-KO cells.

It has been reported that portions of the core domain and C-terminus region TOP1 participate in PARP1 interaction.[27] Using Alphafold 3, we modeled the complex between TOP1-ΔN200, lacking the N-terminal disordered region of TOP1, and the PARP1 HD domain. Notably, among the top four out of the five models generated, the TOP1-HD interface consistently involved the same surface of the HD domain, which contains a cluster of conserved basic residues (K695, K700, R704, and R735) (**Supplementary Fig 5a, b**). Inspired by this finding, we generated three mutants that had their positive residue substituted with the negative charged glutamic acid to assess their

binding to TOP1. Our pull-down results showed that both the R735E and the concurrent K700E/R705E mutant largely abolished binding with TOP1 (**Fig 7a**). We chose to focus on the PARP1-R735E mutant, as this purified mutant is comparable to the wild-type PARP1 in nick binding and auto-PARylation, and sensitive to PARP inhibitor (PARPi), talazoparib, *in vitro* (**Supplementary Fig 5c and d**). Moreover, when expressed in PARP1-KO U2OS cell lines, PARP1-R735E retained its ability to rescue the PARP1-KO from camptothecin (CPT) sensitivity,[43,44] while displaying reduced interaction with TOP1 (**Supplementary Fig 5e, f and Fig 7b**). This deficiency in interaction of the R735E mutant with TOP1 appears to affect the inhibition of TOP1 secondary cleavage as TOP1cc formation was reduced nearly 3-fold with the R735E mutant compared to the wild-type PARP1 (at 50 nM PARP1 concentration) (**Fig 7c**). The R735E mutant also showcased a significant decrease in PARylation-dependent inhibition of TOP1 secondary cleavage and initial rNMP cleavage (**Fig 7d**). Importantly, the R735E mutant desensitized cells to PARP inhibitor talazoparib when compared to the cells expressing the wild-type PARP1, with a difference in half-maximal effective concentration (EC$_{50}$) of almost 4-fold (**Fig 7e**). This result suggests that the interaction with TOP1 might be crucial for the engagement of PARP1 at sites of ribonucleotide cleavage by TOP1. Overall, our findings highlight the relevance of the physical interaction between TOP1 and PARP1 for the regulation of ribonucleotide processing and PARP trapping.

Discussion

The processing of genomic ribonucleotides by TOP1 results in three major products that can compromise genome integrity: the CP nick, the TOP1cc, and the deletions at STRs. The human genome is densely populated with tandem repeats, which together account for approximately 3% of the genome, with dinucleotide repeats alone accounting for approximately 0.5% of the total genomic sequence.[45] This abundance of STRs, as well as the mutagenic nature of human TOP1, requires a mechanism to safeguard the genome in cells. Our findings reveal that PARP1 plays a crucial role in preventing these deletions by inhibiting TOP1 secondary cleavage at sites of genomic rNMPs. It does this through both catalytic-independent and -dependent mechanisms. Non-catalytically, PARP1 physically impedes the TOP1 secondary cleavage, which requires both nick binding and TOP1 interaction. Catalytically, following its activation upon binding to CP nick, PARP1 is capable of PARylating TOP1, which likely inhibits several activities of TOP1, including the rNMP cleavage, the secondary CP nick cleavage and CP nick ligation.

The concurrent deletion of both PARP1 and RNASEH2A led to a pronounced accumulation of TOP1cc. This is likely because failure of genomic rNMP removal by ribonucleotide excision repair shifts processing to TOP1-mediated cleavage, while the absence of PARP1 increases susceptibility of TOP1 to secondary cleavage. The resulting accumulation of trapped TOP1 proved to be deleterious to the growth of PARP1 and RNASHE2A double knockout cells, as it can be rescued by TDP1 overexpression. Notably, these cells were hypersensitive to BCL2 inhibitors, which induce apoptosis in senescent cells by blocking BCL2-mediated suppression of programmed cell death.[32,33] Consistent with our findings, TOP1cc accumulation has been linked to cellular senescence through the cGAS-mediated inflammatory senescence pathway.[31,34]

An important aspect of genomic rNMP removal by TOP1 is its link to inhibitor-induced PARP trapping.[22] *In vitro*, purified PARP1 binds CP nicks with an affinity comparable to that for clean nicks, supporting CP nicks as a relevant PARP1 trapping lesion. Notably, whereas PARP1 binding to clean nicks has little effect on ligation by ligase III,[46,47] binding to CP nicks blocked both TOP1-mediated CP nick ligation and CP nick cleavage. Thus, PARP1 trapping at CP nicks may impede their repair and cause replication fork collapse. Importantly, treatment with a PARP inhibitor had little effect on mutation rates, suggesting that trapped PARP1 efficiently blocks TOP1 secondary cleavage in a catalysis-independent manner. In contrast, PARP1-R735E, which retains DNA binding and catalytic activity but is defective in TOP1 interaction and suppression, partially rescued PARP inhibitor-induced cytotoxicity. This means that the physical interaction with TOP1 is relevant for engaging PARP1 at sites of ribonucleotide cleavage, such as due to recruitment of PARP1 by TOP1 or a handoff of DNA at the cleavage site between TOP1 and PARP1. Together, these findings establish a potential mechanism underlying PARP inhibitor cytotoxicity and its connection to TOP1 regulation during genomic rNMP removal.

Aicardi-Goutières syndrome (AGS), a neurodevelopmental disorder, and systemic lupus erythematosus (SLE) are both RNase H2-associated autoimmune disorders characterized by hyperactivation of the innate immune response in cells.[48] In both cases, this immune response is thought to arise from the genomic instability that occurs in the presence of a dysfunctional RNase H2 complex.[6,48-50] An improved understanding of genomic rNMP processing in the absence of RNase H2 in human cells will provide a framework to identify the source of the genomic instability that leads to this innate immune response in AGS and SLE. Given our findings that PARP1 limits the accumulation of TOP1cc in the absence of functional RNase H2, it would be interesting to assess which of these rNMP-associated lesions (CP nick and TOP1cc), if any, participate in the emergence of innate immune response in these disorders. TOP1cc, for example, has already been linked to cGAS activation.[31] Although most cells in the nervous system are post-mitotic, proliferative immune cells such as microglia can drive neuronopathic immune responses even when they are the only cells harboring AGS genetic mutations.[51] Furthermore, mice models of AGS mutations showcased increased phagocytosis activity in microglia, as well as elevated levels of type I interferon mainly within microglia.[52] These observations are particularly relevant because replicating cells are expected to freshly incorporate ribonucleotide and may therefore be more susceptible to genome instability arising from TOP1-mediated rNMP processing.

Methods

Plasmids and oligonucleotides list
The description of all the plasmids utilized has been provided in the **Supplementary Table 1**. The oligonucleotides used to generate the substrates for biochemical assays are provided in the **Supplementary Table 2**.

Expression and purification TOP1, yTop1 and TOP1-Y723F from insect cells

The coding sequence for TOP1 was amplified from TOP1 cDNA (Dharmacon Reagents) and cloned into a modified pFASTBac-HTB vector and fused in frame to the FLAG and His$_6$ tags within the vector.[53] The construct was transformed into DH10Bac *E. coli* strain (Invitrogen) to generate the bacmid, which was used to infect SF9 insect cells for TOP1 expression. To purify

TOP1, the insect cell pellet (~15g from 1 L culture) was resuspended in 50 mL K500 buffer (500 mM KCl, 20 mM KH$_2$PO$_4$, pH 7.4, 10% glycerol, 0.5 mM EDTA, 0.01% NP-40, 1mM β-mercaptoethanol) with a cocktail of protease inhibitors (aprotinin, chymostatin, leupeptin, and pepstatin A at 5 μg/ml each, and 1 mM phenyl-methylsulfonyl fluoride). The resuspended cells were disrupted by sonication for 1 minute, and the lysate was clarified by centrifugation (20,000 g for 20 minutes). The supernatant was then loaded onto a 3 mL Ni-NTA column (Thermo Fisher) using a BioRad Econo pump. The column was washed with 50 mL K500 buffer with 10 mM imidazole. TOP1 was eluted using 20 mL K500 buffer with 250 mM imidazole. The eluate was then incubated with 0.5-2 mL anti-FLAG M2 agarose resin (Sigma) overnight. The following day, the beads were spun down (1000 g for 5 minutes), and the supernatant was removed. The matrix was washed with K500 buffer with an adjusted 0.1% NP-40. TOP1 was finally eluted by incubating 1.2 mL of K500 containing 200 μg/mL FLAG peptide for 1 hour. The eluate was filter-dialyzed to remove excess FLAG peptide. The purified Top1 protein (~0.5 mg) was frozen in liquid nitrogen and stored at −80°C as small aliquots. To express TOP1-Y723F, the mutation was introduced via site-directed mutagenesis, and the protein was expressed similarly as WT, with a yield ~3 mg. The coding sequence of yTop1 was amplified from yeast genomic DNA and cloned into the modified pFASTBac-HTB with a Flag tag and a His$_6$ tag engineered at the amino-terminus. The same procedure was followed for the purification of yTop1 as reported.[54]

Expression and purification of recombinant PARP1 and its mutants from *E. coli*

The pET28 plasmid expressing full-length PARP1 (Addgene #192635, a kind gift from John Pascal)[55] was transformed into Rosetta 2 pLysS *E. coli* strain (Novagen) for protein expression. The cells were grown in 8 L of LB broth supplemented with 10 mM benzamide, and protein expression was induced at OD$_{600}$=0.6 with 0.5 mM IPTG and 100 μM ZnSO$_4$ was supplemented at the same time. After IPTG addition, protein expression was allowed to occur overnight at 18 °C. To purify PARP1, the bacterial pellet (~15g) was resuspended in 50 mL N500 buffer (500 mM NaCl, 20 mM Tris, pH 7.0, 10% glycerol, 0.5 mM EDTA, 0.01% NP-40, 1mM β-mercaptoethanol) with a cocktail of protease inhibitors (aprotinin, chymostatin, leupeptin, and pepstatin A at 5 μg/ml each, and 1 mM phenyl-methylsulfonyl fluoride). The resuspended cells were disrupted by sonication for 10 minutes, and the lysate was clarified by centrifugation (20,000 g for 20 minutes). The supernatant was then split onto two gravity columns with 2 mL HisPur Ni-NTA resin each (Thermo Fisher). Subsequently, the column was washed with 50 mL K500 buffer with 20 mM imidazole. PARP1 was eluted using 20 mL N500 buffer with 250 mM imidazole. The eluate was then diluted to 150mM salt concentration and loaded into two gravity columns with 2 mL Heparin Sepharose 6 Fast Flow resin (Cytiva). Each column was then washed with 25 mL N150 (150 mM NaCl, 20 mM Tris, pH 7.0, 10% glycerol, 0.5 mM EDTA, 0.01% NP-40, 1mM β-mercaptoethanol). The protein in each column was then eluted with 10 mL N500, and the combined protein elution was concentrated in an Ultracel-30K concentrator (Amicon). The eluted protein was finally loaded into a Superdex 200 increase gel filtration column (Cytiva) connected to an AKTA pure chromatography system (Cytiva) and eluted with K500 buffer. The eluted PARP1 (~1 mg) is free of DNA contamination and thus show minimal PARylation activity in the absence of DNA substrate. To generate PARP1 mutants, mutations were introduced via site-directed mutagenesis,

and the proteins were expressed similarly as WT. The C-terminal truncations of PARP1 (ΔC785
and ΔC660) were purified in the same way as WT. For these truncations, we used two pET28
constructs from the Conaway lab (Addgene #174804 and # 173942).[56] The ΔC785 truncation,
specifically, requires 12 L of cells rather than 8 L. The construct for the PARP1 DBD-only
truncation (N-366) was made by amplifying and cloning the DBD region of PARP1 into the
pETDuet-1 plasmid. This truncation was purified in a similar fashion to the previous PARP1
proteins. The N-terminal truncation of the DBD (ΔDBD) was also cloned in a pETDuet-1 with
His$_6$ and FLAG tags and purified with Ni and anti-FLAG affinity chromatography (similarly as
described with TOP1).
412
413 **Generating 2', 3'-cyclic phosphate terminated (CP) nick substrates**
414
To generate the 2', 3'-cyclic phosphate terminated nicked duplex, the H4 oligonucleotide
containing two consecutive ribonucleotides was treated with RNase I$_f$ (New England Biolabs) for
5 minutes at 37°C and subsequently heat inactivated. This reaction yields ~ 40-60% of 2', 3'-
cyclic phosphate terminated ends that are resistant to alkaline phosphatase (New England Biolabs).
The remaining RNase I$_f$ product corresponds to the 3'-phosphate-terminated oligonucleotide. To
obtain a pure cyclic phosphate product, the reactions were treated with calf intestinal alkaline
phosphatase and thermolabile exonuclease I (New England Biolabs) for 20 mins to convert the 3'-
phosphate ends into hydroxyl terminated oligonucleotides and subsequently digest them. The
reactions were then heat inactivated for 3 mins at 80°C, and the proteins were removed with oligo
cleaning columns (Zymogen). The purity of our cyclic phosphate terminated product was assessed
with alkaline phosphatase and T4 PNK treatment (New England Biolabs), in which the first
enzyme does not process the cyclic phosphate while the latter can remove it.[57] The product was
then 5'-P^{32} labelled using minus 3-phosphatase T4-PNK (New England Biolabs) and cleaned using
a Spin6 column (BIO-RAD). To make the final cyclic phosphate terminated nicked duplex, the
product was annealed with H3 and H5 oligonucleotides via slow cooling from 75 °C to avoid the
spontaneous hydrolysis of the cyclic phosphates. To make the shorter 45 bp CP nick substrate, the
treated DNA was H6 and it was annealed to H7 and H5 oligonucleotides. Streptavidin in our
reaction buffers will bind to the two biotinylated ends of the H3 or H7 oligonucleotide to prevent
the engagement of PARP1 at the blunt ends of the substrate. To make the CP nick containing
dumbbell substrate, the H12 oligonucleotide was treated with RNase I$_f$ as described in this section
and then annealed by immediate cooling on ice following heat denaturing at 95°C for 5 minutes to
minimize inter-molecular annealing. The same annealing procedure was followed for the regular
nick dumbbell substrate.
438
439 **TOP1 ribonuclease assay**
440
For TOP1 ribonuclease assays, the indicated amount of TOP1 in the reaction buffer (20 mM Tris-
HCl, pH 8.0, 2 mM MgCl$_2$, 100 μM TCEP, 100 μg/mL BSA, 150 mM KCl, 20 μg/mL streptavidin)
was incubated with 10 nM DNA substrate for 20 min at 37°C. The reaction was stopped by
treatment with SDS (0.2% final). When determining the identity of the TOP1 adduct, the reaction
was treated with proteinase K (0.5 mg/ml) before column purification (oligo clean & concentrator,
Cat. No: D4060, Zymo Research), followed by treatment with both yeast tyrosyl-DNA
phosphodiesterase 1 (Tdp1) and DNA 3'-phophatase (Tpp1). Equal volume of loading dye
containing 85% formamide with 25 mM EDTA, and 1 μM unlabeled H1 DNA strand (same

sequence as rNMP-containing oligonucleotide) was then added to reaction mixtures to prevent re-
annealing of the radiolabeled strand. The mixture was heated at 75 °C for 5 min and cooled on ice
for another 5 mins. The reactions were run in 12% denaturing polyacrylamide gel in TBE (Tris-
Boric Acid-EDTA) buffer, followed by phosphor imaging analysis.
453
454 **PARylation assay**
455
For *in vitro* PARylation assay, we followed a similar procedure as previously published, with slight
modifications for the experiments that include the rU-containing substrate.[58] Specifically, the auto-
PARylation assays with the dumbbell substrates (1µM) were incubated for 10 mins in a buffer
consisting of 20 mM Tris-HCl, pH 8.0, 2 mM $MgCl_2$, 100 µM TCEP, and 150 mM KCl. For the
PARylation assay using the rNMP containing substrate (10 nM), the reaction buffer was
supplemented with 20 µg/mL streptavidin to ensure the occlusion of the two blunt ends of our
substrate and thus preventing unwanted activation of PARP1 by double strand DNA ends. Other
than the negative controls and NAD^+ titrations, all the other reactions contain 1 mM NAD^+. Before
the addition of TOP1 or PARP1, the reactions were pre-incubated for 10 mins at 37°C to ensure
streptavidin is bound to the substrate ends. Once the proteins were added, the reactions were
incubated for 10 mins at 25°C. The reactions were stopped with 3x Laemmli SDS buffer and heated
for 5 minutes at 95°C. The reactions were run in a Mini Protean TGX Stain-Free 4-15% SDS gel
(BIO-RAD). The gels were first scanned with stain-free imaging in a BIO-RAD Gel Doc EZ
imager and subsequently transferred onto a nitrocellulose membrane for western blot analysis with
the Anti-Poly/Mono ADP-ribose rabbit monoclonal antibody (Cell Signaling, D9P7Z, 1:1,000
dilution). For detection, we used IRDye 800CW goat anti-rabbit antibody (LICOR, 926-32211,
1:10,000 dilution).
473
474 **DNA binding assay and binding competition assay**
475
Dumbbell substrates used in the DNA binding assays were generated as described in the
"Generating 2', 3'-cyclic phosphate terminated substrates" subsection of the Methods. Either the
CP nick dumbbell substrate, labeled internally with a Cy3 dye, or the clean nick dumbbell substrate,
labeled internally with a Cy5 dye (or both substrates in the case of the competition assay) was
incubated with indicated concentrations of PARP1 for 20 mins in a buffer containing 20 mM Tris-
HCl, pH 8.0, 2 mM $MgCl_2$, 100 µM TCEP, 100 µg/ml BSA, and 150 mM KCl. The reactions were
resolved using a 6% native gel and imaged using a ChemiDoc MP imaging system (BIO-RAD).
483
484 **Cell Lines**
485
486 All the human cell lines used are depicted in the **Supplementary Table 3**.
487
488 ***In vitro* affinity pulldown assay**
489
To test for TOP1-PARP1 interaction, (His)₆-tagged PARP1 (350 ng) was mixed with FLAG-
(His)₆-tagged TOP1 (750 ng) in 25 µL buffer (20 mM Tris-HCl, pH 8.0, 2 mM $MgCl_2$, 100 µM
TCEP, 100 µg/ml BSA, 150 mM KCl). The mixture was incubated with 25 µL anti-FLAG M2
agarose resin (Sigma) for 1 hour shaking at 25°C. Following incubation, the resin was washed
twice with 400 µl of wash buffer (150 mM KCl, 20 mM KH_2PO_4, pH 7.4, 10% glycerol, 0.5 mM

EDTA, 0.1% NP-40, 1mM β-mercaptoethanol), and 20 µl of 2% SDS was used to the elute proteins.
The supernatant (S) and SDS eluate (E) fractions, 10 µl each, were resolved by 8% SDS-PAGE
and immunoblotted with either HRP-conjugated anti-His antibodies or with their respective protein
specific antibodies.
499
500   **Fluctuation assay**
501
The engineered HeLa reporter cell lines including the WT, RNase H2A+, and RNase H2A-KO
cell lines were kindly provided by Dr. Andrew P. Jackson's lab to detect slippage mutation rate at
STRs. We further knocked out the PARP1 gene in these three cell lines (see "Cell Lines" in this
methods section). The fluctuation assays were performed as described by Dr. Andrew P. Jackson's
team.[12] For the PARP inhibitor pre-treatment in the fluctuation assay, we treated our cells in a
flask with either 1 µM talazoparib or a similar volume of DMSO (for the negative control) for 3
days before plating them for the fluctuation assay as outlined in the paper by Reijns, *et al*.
509
510   **Cell proliferation analysis**
511
To assess the proliferation of the different strains, 20,000 cells were seeded in multiple 12-well
plates (with triplicates in each plate, and each plate representing a different day to be read, from
day 0 to 5). Six hours after plating the day 0 plate, the plate was washed with 1x PBS. After
washing, the plates were fixed with a buffer consisting of 1% formaldehyde, 0.05% Crystal violet
(w/v), 1x PBS, and 1% methanol for 20 mins. The wells were washed three times with water and
allowed to air dry overnight. Once dried, 2 mL of 10% acetic acid was added to each well and
incubated for 20 mins with moderate shaking. The wells were mixed with a pipette, and the plates
were measured for 590 nm absorbance. Each day at the same time, a plate was fixed and allowed
to air dry. Once the data points for all the days were obtained, the growth curve was plotted by
normalizing the value of the signals using day 0 as the frame of reference.
522
523   **MTT viability assay**
524
The MTT assay to assess the viability of cells after treatments has been described previously.[59] To
assess viability after treatments with different inhibitors, we plated the different strains each with
8 repeats (one full column) in a 96-well plate (1000-2000 cells per well). The following day, we
treated them with the appropriate drug to be tested (untreated controls were also maintained for
each strain). After the treatment period (6 days), each well was supplemented with 20 µL 5mg/mL
thiazolyl blue tetrazolium bromide (MTT) solution (in PBS). The cells were then incubated for 3.5
hours in the 37 °C tissue culture incubator. The media was then removed and replaced with 150
µL MTT solvent (4 mM HCl and 0.1% NP-40 diluted in isopropanol), covered with aluminum foil,
and incubated in a shaker for 15 minutes. The absorbance at 590 nM with a 620 nM reference for
all the wells was measured using an Epoch plate reader (Biotek). The results were normalized to
the appropriate untreated controls for each strain.
536
537   **TOP1cc quantification RADAR assay**
538
539   The genomic DNA purification was conducted using the TOPOGEN ICE Assay Kit
540   (TG10201),[35,36] except that the cells were treated with 10 µM MG-132 protease inhibitor for 2

hours before DNA extraction to protect the TOP1-DNA adducts from degradation. Cells pre-treated with 20 μM camptothecin for 2-hour pre-DNA extraction were included as a positive control. The concentration of genomic DNA was quantified by nanodrop before and after dilution to 30 ng/μL using the sodium phosphate buffer. Each sample, 200 μL, was vacuum-blotted into both a nitrocellulose membrane, for protein detection, and an Amersham Hybond -N$^+$ membrane (GE Healthcare), for DNA detection (which also serves as a loading control). After blotting, both membranes were irradiated with 120 mJ/cm^2 using a UV Stratalinker 1800 (Stratagene). The nitrocellulose membrane was washed with TBST and immuno-stained with TOP1 antibody (provided in the TOPOGEN ICE kit) in 5% skim milk (w/v), TBS with 0.1% Tween-20 (v/v), while the Hybond membrane was washed with TBS and stained with Sybr Gold (Invitrogen) diluted in TBS. For detection of the TOP1 immuno-stained membrane, we used IRDye 800CW goat anti-rabbit antibody (LICOR). They were both imaged using a ChemiDoc MP imaging system (BIO-RAD).

Co-immunoprecipitation assay

U2OS cells were generated expressing either the wild-type or the R735E mutant of a 3xFLAG-tagged PARP1 in the PARP1-KO U2OS cells (a kind gift from Dr. Shan Zha).[60] The pCMV-PARP1-3xFlag-WT plasmid was a gift from Thomas Muir's lab (Addgene #111575).[61] To assess TOP1-PARP1 interaction in cells, 1 million cells were pelleted and frozen at -80 °C. The pellets were then resuspended in 400 μL PBS buffer supplemented with 0.01% NP-40, 1mM β-mercaptoethanol, and a protease inhibitors cocktail (aprotinin, chymostatin, leupeptin, and pepstatin A at 5 μg/ml each, and 1 mM phenyl-methylsulfonyl fluoride). The samples were sonicated at 40% amplification in a Fisher Scientific Model 120 sonicator for 5 secs to achieve proper extraction of the nuclear proteins. The lysate mixture was incubated with 25 μL anti-FLAG M2 agarose resin (Sigma) overnight at 4 °C. Following incubation, the resin was washed two times with 400 μl of wash buffer (1x PBS, 0.1% NP-40, 1mM β-mercaptoethanol) and eluted with 20 μl of 2x Laemmli sample buffer (BIO-RAD). The input and SDS eluate fractions were resolved by 10% SDS-PAGE and immunoblotted with either anti-TOP1 (Cell signaling # 79971S), anti-PARP1 (Millipore Sigma #SAB5701291), or anti-GAPDH (BioLegend #649203) antibody (the latter as a loading control). IRDye 800CW goat anti-rabbit antibody (LICOR # 926-32211) was used as secondary antibody for protein detection.

Acknowledgements

We are grateful to Shan Zha, Andrew P. Jackson, John Pascal, Thomas Muir, and Joan Conaway for providing plasmids and human cell lines. Research reported in this publication was supported by the National Cancer Institute of the National Institutes of Health (NIH) under Award Numbers F31CA268751 (E.S.) and R01CA285684 (H.N. and Q.C.). Part of effort for H.N was also supported by R35GM152207 and the American Cancer Society Research Scholar Award RSG-21-013-01-DMC. Part of effort for Q.C. was also supported by R01CA256741, R01CA278832 and R01CA300246, Prostate SPORE P50CA180995 Development Research Program and Polsky Urologic Cancer Institute of the Robert H. Lurie Comprehensive Cancer Center of Northwestern University at Northwestern Memorial Hospital.

References

Reijns, M. *et al.* Enzymatic removal of ribonucleotides from DNA is essential for mammalian genome integrity and development. *Cell* **149**, 1008–1022 (2012).

Clausen, A. R., Zhang, S., Burgers, P. M., Lee, M. Y. & Kunkel, T. A. Ribonucleotide incorporation, proofreading and bypass by human DNA polymerase δ. *DNA Repair (Amst)*, 121–127 (2013).

Sparks, J. L. *et al.* RNase H2-initiated ribonucleotide excision repair. *Molecular cell* **47**, 980–986 (2012). https://doi.org/10.1016/j.molcel.2012.06.035

Rydberg B, G. J. Excision of misincorporated ribonucleotides in DNA by RNase H (type 2) and FEN-1 in cell-free extracts. . *Proc Natl Acad Sci U S A.* , 16654–16659 (2002).

Lim, Y. W., Sanz, L. A., Xu, X., Hartono, S. R. & Chedin, F. Genome-wide DNA hypomethylation and RNA:DNA hybrid accumulation in Aicardi-Goutieres syndrome. *eLife* **4** (2015). https://doi.org/10.7554/eLife.08007

Ramantani, G. *et al.* Expanding the phenotypic spectrum of lupus erythematosus in Aicardi-Goutieres syndrome. *Arthritis Rheum* **62**, 1469–1477 (2010). https://doi.org/10.1002/art.27367

Gunther, C. *et al.* Defective removal of ribonucleotides from DNA promotes systemic autoimmunity. *J Clin Invest* **125**, 413–424 (2015). https://doi.org/10.1172/JCI78001

Sekiguchi, J. & Shuman, S. Site-specific ribonuclease activity of eukaryotic DNA topoisomerase I. *Molecular cell* **1**, 89–97 (1997).

Kim, N. *et al.* Mutagenic processing of ribonucleotides in DNA by yeast topoisomerase I. *Science* **332**, 1561–1564 (2011). https://doi.org/10.1126/science.1205016

Huang, S. Y., Ghosh, S. & Pommier, Y. Topoisomerase I alone is sufficient to produce short DNA deletions and can also reverse nicks at ribonucleotide sites. *J Biol Chem* **290**, 14068–14076 (2015). https://doi.org/10.1074/jbc.M115.653345

Sparks, J. L. & Burgers, P. M. Error-free and mutagenic processing of topoisomerase 1-provoked damage at genomic ribonucleotides. *EMBO J* **34**, 1259–1269 (2015). https://doi.org/10.15252/embj.201490868

Reijns, M. A. M., Parry, D.A., Williams, T.C. et al. Signatures of TOP1 transcription-associated mutagenesis in cancer and germline. . *Nature*, 623–631 (2022).

Tate, J. G. *et al.* COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res* **47**, D941–D947 (2019). https://doi.org/10.1093/nar/gky1015

D'Amours, D., Desnoyers, S., D'Silva, I. & Poirier, G. G. Poly(ADP-ribosyl)ation reactions in the regulation of nuclear functions. *Biochem J* **342 (Pt 2)**, 249–268 (1999).

Lindahl, T., Satoh, M. S., Poirier, G. G. & Klungland, A. Post-translational modification of poly(ADP-ribose) polymerase induced by DNA strand breaks. *Trends Biochem Sci* **20**, 405–411 (1995). https://doi.org/10.1016/s0968-0004(00)89089-1

Schreiber, V., Dantzer, F., Ame, J. & al., e. Poly(ADP-ribose): novel functions for an old molecule. *Nat Rev Mol Cell Biol* **7**, 517–528 (2006).

Murai, J. *et al.* Trapping of PARP1 and PARP2 by Clinical PARP Inhibitors. *Cancer Res* **72**, 5588–5599 (2012). https://doi.org/10.1158/0008-5472.CAN-12-2753

Lord, C. J. & Ashworth, A. PARP inhibitors: Synthetic lethality in the clinic. *Science* **355**, 1152–1158 (2017). https://doi.org/10.1126/science.aam7344

Farmer, H. *et al.* Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature* **434**, 917–921 (2005). https://doi.org/10.1038/nature03445

20   Bryant, H. E. *et al.* Specific killing of BRCA2-deficient tumours with inhibitors of
poly(ADP-ribose) polymerase. *Nature* **434**, 913–917 (2005).
https://doi.org/10.1038/nature03443
21   Ragupathi, A., Singh, M., Perez, A. M. & Zhang, D. Targeting the BRCA1/2 deficient
cancer with PARP inhibitors: Clinical outcomes and mechanistic insights. *Frontiers in Cell*
and Developmental Biology **Volume 11 - 2023** (2023).
https://doi.org/10.3389/fcell.2023.1133472
22   Zimmermann, M. *et al.* CRISPR screens identify genomic ribonucleotides as a source of
PARP-trapping lesions. *Nature* **559**, 285–289 (2018). https://doi.org/10.1038/s41586-018-
0291-z
23   Sekiguchi, J. & Shuman, S. Site-specific ribonuclease activity of eukaryotic DNA
topoisomerase 1. . *Mol. Cell* **1**, 89–97 (1997).
24   Cho, J. E., Kim, N., Li, Y. C. & Jinks-Robertson, S. Two distinct mechanisms of
Topoisomerase 1-dependent mutagenesis in yeast. *DNA Repair (Amst)* **12**, 205–211 (2013).
https://doi.org/10.1016/j.dnarep.2012.12.004
25   Pouliot, J. J., Yao, K. C., Robertson, C. A. & Nash, H. A. Yeast gene for a Tyr-DNA
phosphodiesterase that repairs topoisomerase I complexes. *Science* **286**, 552–555 (1999).
https://doi.org/10.1126/science.286.5439.552
26   Karumbati, A. S. *et al.* The role of yeast DNA 3'-phosphatase Tpp1 and rad1/Rad10
endonuclease in processing spontaneous and induced base lesions. *J Biol Chem* **278**,
31434–31443 (2003). https://doi.org/10.1074/jbc.M304586200
27   Bauer, P. I. *et al.* Molecular interactions between poly(ADP-ribose) polymerase (PARP I)
and topoisomerase I (Topo I): identification of topology of binding. *FEBS Lett* **506**, 239–
242 (2001). https://doi.org/10.1016/s0014-5793(01)02919-2
28   Zhang, Y., Wang, J., Ding, M. & Yu, Y. Site-specific characterization of the Asp- and Glu-
ADP-ribosylated proteome. *Nat Methods* **10**, 981–984 (2013).
https://doi.org/10.1038/nmeth.2603
29   Buch-Larsen, S. C. *et al.* Mapping Physiological ADP-Ribosylation Using Activated Ion
Electron Transfer Dissociation. *Cell Rep* **32**, 108176 (2020).
https://doi.org/10.1016/j.celrep.2020.108176
30   Chen, J. *et al.* The Bcl-2/Bcl-X(L)/Bcl-w inhibitor, navitoclax, enhances the activity of
chemotherapeutic agents in vitro and in vivo. *Mol Cancer Ther* **10**, 2340–2349 (2011).
https://doi.org/10.1158/1535-7163.MCT-11-0415
31   Zhao, B. *et al.* Topoisomerase 1 cleavage complex enables pattern recognition and
inflammation during senescence. *Nat Commun* **11**, 908 (2020).
https://doi.org/10.1038/s41467-020-14652-y
32   Yosef, R. *et al.* Directed elimination of senescent cells by inhibition of BCL-W and BCL-
XL. *Nat Commun* **7**, 11190 (2016). https://doi.org/10.1038/ncomms11190
33   Rysanek, D. *et al.* Synergism of BCL-2 family inhibitors facilitates selective elimination
of senescent cells. *Aging (Albany NY)* **14**, 6381–6414 (2022).
https://doi.org/10.18632/aging.204207
34   Zhang, J. W. *et al.* Autophagy inhibition switches low-dose camptothecin-induced
premature senescence to apoptosis in human colorectal cancer cells. *Biochem Pharmacol*
90, 265–275 (2014). https://doi.org/10.1016/j.bcp.2014.05.009

35   Meroni, A. & Vindigni, A. A RADAR method to measure DNA topoisomerase covalent
complexes. *Methods Enzymol* **672**, 369–381 (2022).
https://doi.org/10.1016/bs.mie.2022.03.036
36   Subramanian, D., Furbee, C. S. & Muller, M. T. ICE bioassay. Isolating in vivo complexes
of enzyme to DNA. *Methods Mol Biol* **95**, 137–147 (2001). https://doi.org/10.1385/1-
59259-057-8:137
37   Dean, R. A. *et al.* Identification of a putative Tdp1 inhibitor (CD00509) by in vitro and
cell-based assays. *J Biomol Screen* **19**, 1372–1382 (2014).
https://doi.org/10.1177/1087057114546551
38   Miwa, M., Tanaka, M., Matsushima, T. & Sugimura, T. Purification and properties of
glycohydrolase from calf thymus splitting ribose-ribose linkages of poly(adenosine
diphosphate ribose). *J Biol Chem* **249**, 3475–3482 (1974).
39   Stewart, L., Ireton, G. C. & Champoux, J. J. The domain organization of human
topoisomerase I. *J Biol Chem* **271**, 7602–7608 (1996).
https://doi.org/10.1074/jbc.271.13.7602
40   Stewart, L., Redinbo, M. R., Qiu, X., Hol, W. G. & Champoux, J. J. A model for the
mechanism of human topoisomerase I. *Science* **279**, 1534–1541 (1998).
https://doi.org/10.1126/science.279.5356.1534
41   Dawicki-McKenna, J. M. *et al.* PARP-1 Activation Requires Local Unfolding of an
Autoinhibitory Domain. *Molecular cell* **60**, 755–768 (2015).
https://doi.org/10.1016/j.molcel.2015.10.013
42   Langelier, M. F., Planck, J. L., Roy, S. & Pascal, J. M. Structural basis for DNA damage-
dependent poly(ADP-ribosyl)ation by human PARP-1. *Science* **336**, 728–732 (2012).
https://doi.org/10.1126/science.1216338
43   Ooka, M. *et al.* Chromatin remodeler ALC1 prevents replication-fork collapse by slowing
fork progression. *PLoS One* **13**, e0192421 (2018).
https://doi.org/10.1371/journal.pone.0192421
44   Das, B. B. *et al.* PARP1-TDP1 coupling for the repair of topoisomerase I-induced DNA
damage. *Nucleic Acids Res* **42**, 4435–4449 (2014). https://doi.org/10.1093/nar/gku088
45   Lander, E. S. *et al.* Initial sequencing and analysis of the human genome. *Nature* **409**, 860–
921 (2001). https://doi.org/10.1038/35057062
46   Kumamoto, S. *et al.* HPF1-dependent PARP activation promotes LIG3-XRCC1-mediated
backup pathway of Okazaki fragment ligation. *Nucleic Acids Res* **49**, 5003–5016 (2021).
https://doi.org/10.1093/nar/gkab269
47   Chin Sang, C. *et al.* PARP1 condensates differentially partition DNA repair proteins and
enhance DNA ligation. *EMBO Rep* **25**, 5635–5666 (2024). https://doi.org/10.1038/s44319-
024-00285-5
48   Pendergraft, W. F., 3rd & Means, T. K. AGS, SLE, and RNASEH2 mutations: translating
insights into therapeutic advances. *J Clin Invest* **125**, 102–104 (2015).
https://doi.org/10.1172/JCI78533
49   Aditi *et al.* Genome instability independent of type I interferon signaling drives
neuropathology caused by impaired ribonucleotide excision repair. *Neuron* **109**, 3962–
3979 e3966 (2021). https://doi.org/10.1016/j.neuron.2021.09.040
50   Crow, Y. J. *et al.* Mutations in genes encoding ribonuclease H2 subunits cause Aicardi-
Goutieres syndrome and mimic congenital viral brain infection. *Nat Genet* **38**, 910–916
(2006). https://doi.org/10.1038/ng1842

51 Nemec, K. M. *et al.* Microglia replacement by ER-Hoxb8 conditionally immortalized
macrophages provides insight into Aicardi-Goutieres Syndrome neuropathology. *bioRxiv*
(2025). https://doi.org/10.1101/2024.09.18.613629
52 Onizawa, H. *et al.* Aicardi-Goutieres syndrome-like encephalitis in mutant mice with
constitutively   active   MDA5.   *Int   Immunol*   **33**,   225–240   (2021).
https://doi.org/10.1093/intimm/dxaa073
53 Niu, H. *et al.* Mechanism of the ATP-dependent DNA End Resection Machinery from S.
cerevisiae. *Nature* **467**, 108–111 (2010).
54 Li, F. *et al.* Apn2 resolves blocked 3' ends and suppresses Top1-induced mutagenesis at
genomic   rNMP   sites.   *Nat   Struct   Mol   Biol*   **26**,   155–163   (2019).
https://doi.org/10.1038/s41594-019-0186-1
55 Langelier, M. F., Servent, K. M., Rogers, E. E. & Pascal, J. M. A third zinc-binding domain
of human poly(ADP-ribose) polymerase-1 coordinates DNA-dependent enzyme activation.
J Biol Chem **283**, 4105–4114 (2008). https://doi.org/10.1074/jbc.M708558200
56 Ooi, S. *et al.* Multiple roles for PARP1 in ALC1-dependent nucleosome remodeling. *Proc.*
Natl. Acad. Sci. **118 (36)**, epub (2021).
57 Honda, S., Morichika, K. & Kirino, Y. Selective amplification and sequencing of cyclic
phosphate-containing RNAs by the cP-RNA-seq method. *Nat Protoc* **11**, 476–489 (2016).
https://doi.org/10.1038/nprot.2016.025
58 Meng, Q. *et al.* EZH2 directly methylates PARP1 and regulates its activity in cancer. *Sci*
Adv **10**, eadl2804 (2024). https://doi.org/10.1126/sciadv.adl2804
59 Alley, M. C. *et al.* Feasibility of drug screening with panels of human tumor cell lines using
a microculture tetrazolium assay. *Cancer Res* **48**, 589–601 (1988).
60 Shao, Z. *et al.* Clinical PARP inhibitors do not abrogate PARP1 exchange at DNA damage
sites   in   vivo.   *Nucleic   Acids   Res*   **48**,   9694–9709   (2020).
https://doi.org/10.1093/nar/gkaa718
61 Liszczak, G., Diehl, K. L., Dann, G. P. & Muir, T. W. Acetylation blocks DNA damage-
induced chromatin ADP-ribosylation. *Nature chemical biology* **14**, 837–840 (2018).
https://doi.org/10.1038/s41589-018-0097-1

Figure Legends

Figure 1: Human TOP1 is prone to secondary cleavage and TOP1cc formation at genomic rNMP sites.

a) Diagram showcasing how TOP1 processes genomic ribonucleotides (rNMPs), and how this processing can lead to deletions in the context of short tandem repeats. b) (i) The titration of TOP1 against a 5'-^{32}P-labeled duplex DNA containing a single embedded rNMP was analyzed in a denaturing gel, (ii) with the quantification of the resulting species. c) TOP1cc products were processed with Proteinase K, Tdp1, and Tpp1 to reveal the TOP1 cleavage site. d) (i) TOP1 was titrated against a 5'-^{32}P-labeled CP nick substrate, (ii) with the quantification of the resulting species. e) (i) human TOP1 and yeast TOP1 (yTop1) were titrated against a CP nick substrate to examine their preference between CP nick ligation and slippage ligation. (ii) Quantification of the different products of TOP1 and yTop1. f) (i) 3'-Cy5-labeled duplex substrate and cleavage suicide substrate (fork) were tested with either human TOP1 or yTop1. (ii) The quantification of cleavage products.

Figure 2: PARP1 suppresses the secondary cleavage by TOP1 at genomic ribonucleotides.

a) (i) TOP1 ribonucleotide cleavage assay in the presence of PARP1 with or without NAD$^+$. To prevent PARP1 from binding, both ends of the substrate were occluded by biotin-bound streptavidin as diagramed with blue ovals. (ii) The quantification of the CP nick and TOP1cc products at 50 nM PARP1 concentration and (iii) of the remaining substrate are shown. b) (i) The processing of the CP nick substrate by TOP1 was assessed in the absence or presence of PARP1 with or without NAD+, with both sides of the substrate occluded with streptavidin. (ii-iii) The quantification of the TOP1cc product and of the ligation products are shown.

Figure 3: PARP1 prevents deletions at short tandem repeats, and its simultaneous deletion with RNASEH2A generates a slow growth phenotype.

a) (i) Fluctuation assay reporting on the mutation rate of the different HeLa reporter strains as indicated. (ii) Quantification of the mutation types observed in sequenced clones of the WT and PARP1-KO strain showcases the prevalence of 2 bp deletions in the absence of PARP1. The n reflects the total number of indels identified in the total sequenced clones. b) Growth curve showcasing the delayed growth of the RNASEH2A-KO PARP1-KO cell line.

Figure 4: Concurrent deletion of RNASEH2A and PARP1 generates a detrimental accumulation of TOP1cc.

a) Sensitivity of WT, RNASEH2A-KO, PARP1-KO and RNASEH2A-KO PARP1-KO cell lines to the BCL2 inhibitor (BCL2i) navitoclax was determined by the MTT viability assay. b) Slot blot of the RADAR assay showcasing the elevated accumulation of TOP1cc in the RNASEH2A-KO PARP1-KO cell line. Samples from treating WT cells with 10 μM of the TOP1-trapping agent camptothecin (CPT) were included as a positive control. The DNA loading control was stained with Sybr Gold. c) Growth curve showing that the overexpression of TDP1 (TDP1-OE) rescues the delayed growth in the RNASEH2A-KO PARP1-KO cell line. d) Viability measured by MTT assay after TDP1 inhibitor, CD00509, (TDPi) treatment demonstrates hypersensitivity of the RNASEH2A-KO PARP1-KO cell line to TDP1 inhibitor. e) MTT viability assay showcasing the

814 synthetic effect of TDP1i and BCL2i treatments to the viability of the RNASEH2A-KO PARP1-
815 KO cell line.
816
Figure 5: TOP1 initial cleavage at genomic ribonucleotides stimulates PARylation by PARP1.
a) (i) PARP1-catalyzed auto-PARylation and PARylation of TOP1 in the presence of the duplex
substrate with a single embedded ribonucleotide *in vitro*. Both DNA ends were occluded by biotin-
bound streptavidin. (ii) Quantification of the amount of unPARylated PARP1 in the absence or
presence of TOP1. b) (i) The *in vitro* PARylation assay using the ribonucleotide-containing
substrate where the products were further treated with PARG, leaving single ADP-ribosyl
modification that allows for precise determination of protein PARylation levels (ii) Quantification
of PARylation levels of PARP1 in the absence or presence of TOP1. c) (i) *in vitro* PARylation
assay with the ribonucleotide-containing substrate using the catalytic dead mutant TOP1-Y723F.
(ii) Quantification of the amount of unPARylated PARP1 in the absence or presence of TOP1-
Y723F. d) Diagram showcasing model on how PARP1 prevents the secondary cleavage by TOP1
at ribonucleotide sites both in a catalytically-independent (by physically restricting TOP1) and -
dependent manner (through the inhibitory PARylation of both TOP1 and itself).
830
Figure 6: The DNA-binding activity of PARP1 and its physical interaction with TOP1 are
important for the prevention of the TOP1 secondary cleavage.
a) Diagram showing the different domains of the full-length (FL) PARP1 and the two PARP1
truncations (PARP1-ΔDBD and PARP1-DBD) tested in the TOP1 ribonucleotide cleavage assay
b) (i) The TOP1 ribonucleotide cleavage assay in the absence or presence of the full-length
PARP1-FL, PARP1-ΔDBD or PARP1-DBD. (ii) Quantification of the cyclic phosphate (CP)
product and the TOP1cc product for the reactions with PARP1-FL, PARP1-ΔDBD and PARP1-
DBD at 50 nM concentration. c) Domain diagram of N-terminal truncations of PARP1 (PARP1-
ΔART and PARP1-ΔCAT) tested for TOP1-binding. d) The *in vitro* pulldown of FLAG-tagged
TOP1 against the full-length PARP1 and PARP1 N-terminal truncations (ΔART and ΔCAT).
Reactions without TOP1 were included as controls to assess non-specific binding of PARP1 to
the anti-FLAG M2 resin.
843
Figure 7: Characterization of PARP1-R735E mutant in TOP1 interaction, the regulation of
TOP1 ribonucleotide cleavage and sensitivity to PARP inhibitor, talazoparib.
a) The *in vitro* anti-FLAG-pull-down assay to test interaction of TOP1 with PARP1 mutants. The
membrane was blotted with anti-His antibody for the detection of TOP1 and PARP1, both His-
tagged. The supernatant (S) and elution (E) of the pull-down assay are shown. b) co-
immunoprecipitation analysis of interaction between TOP1 and PARP1 in cells, where anti-FLAG
M2 resin was used to immunoprecipitate either FLAG-tagged PARP1 or PARP1-R735E mutant
stably expressed in U2OS PARP1-KO cell line to test their interaction with TOP1. c) (i) The TOP1
ribonucleotide cleavage assay in the absence or presence of PARP1 or PARP1-R735E. (ii)
Quantification of the TOP1cc product from the reaction with either WT or the R735E mutant
PARP1. d) (i) TOP1 ribonuclease assay in the presence of NAD^+ comparing TOP1 inhibition by
the WT PARP1 and the R735E mutant. e) (i) Clonogenic survival curve testing the sensitivity to
talazoparib (TLZ) treatment of PARP1-KO U2OS cell lines expressing either no PARP1, WT
PARP1, or PARP1-R735E (normalized to untreated cells and fitted using a nonlinear least-squares

fit to a three-parameter dose–response model). (ii) A representative plate for each strain at the
different TLZ concentrations as indicated.
860

861 **Supplemental Figure 1: SDS-PAGE analysis of purified proteins.**
862 Purified proteins were fractionated in 4-15% SDS-PAGE or 10% SDS-PAGE and stained by
863 Coomassie blue G-250.
864

Supplemental Figure 2: Human TOP1 rNMP cleavage analysis and comparison with yeast
Top1 (yTop1).
a) (i) Time course analysis of ribonucleotide processing by human TOP1. (ii) Quantification of
different product species and the remaining substrate. b) (i) Comparison of ribonucleotide
processing between human TOP1 and yTop1. (ii)-(iii) Quantification of the remaining substrate
and the TOP1cc product for each TOP1 variant.

Supplemental Figure 3. CP nick activates PARP1 for TOP1 regulation.

a) (i) PARP1 binding competition assay using a Cy5 labeled dumbbell nick substrate and a Cy3
labeled dumbbell CP-nick substrate. Single channel and merged images are shown. (ii)
Quantification of the remaining unbound substrate for each condition. b) (i) TOP1 ribonucleotide
cleavage assay performed in the absence and presence of NAD^+ to examine the effect of NAD^+ on
TOP1. (ii) quantification of TOP1cc and CP nick products. c) (i) TOP1 ribonucleotide cleavage
assay performed in the absence or presence of indicated amount of PARP1 or the catalytically dead
PARP1-E988K mutant, with or without NAD^+. (ii) Quantification of TOP1cc and CP nick
products from each reaction. (iii) Quantification of the remaining substrate in reactions containing
PARP1 or PARP1-E988K in the presence of NAD^+.

Supplemental Figure 4: Confirmation of PARP1 knockout and TDP1 overexpression in the
reporter cell lines and examination of the effect of PARP inhibitor on the mutation rate in
the wild-type reporter cells.

a) PARP1 was knocked out in the WT, RNase H2A-KO, and RNase H2A+ reporter HeLa cells.
TDP1 was stably expressed in the RNase H2A-KO PARP1-KO cell line under the control of cmv
promoter. The protein samples were fractionated in 4-15% SDS-PAGE and transferred into a
nitrocellulose membrane for western blots. b) Fluctuation assay reporting on the mutation rate of
the WT HeLa reporter strain pre-treated with either PARP inhibitor talazoparib (TLZ) or DMSO,
showcasing that inhibition of PARP1 does not significantly affect the number of deletions at STRs.

Supplementary Figure 5: Though deficient in TOP1-interaction, PARP1-R735E is
comparable to the wild-type in DNA binding, autoPARylation and response to PARP
inhibitor *in vitro*, as well as complementing PARP1-KO for the sensitivity to camptothecin
in cells.

a) A model of the interface between TOP1 and the HD domain of PARP1 from AlphaFold
prediction. The targeted residues within the the HD domain (K695, K700, R704, and R735) are
labeled with the numbers 1-4, respectively. A close-up of the PARP1-TOP1 interface and the
electrostatic surface map of this interface. b) Alignment of the PARP1 HD domain region
containing the targeted residues across higher eukaryotes, including human, chicken, Xenopus frog,

zebrafish, and Arabidopsis thaliana. Arrows mark the sites tested for TOP1 interactions. Green arrows indicate sites whose mutation reduced TOP1 binding, whereas the red arrow indicates a site that retained substantial TOP1 binding after mutation. c) (i) DNA binding assay comparing WT PARP1 and the R735E mutant ability to bind to the CP-nicked dumbbell substrate. (ii) Quantification of the unbound substrate for both PARP1 variants. d) Coomassie stained 10% SDS-PAGE images showcasing the auto-PARylation activity and talazoparib sensitivity of both PARP1 and PARP1-R735E. e) Western blot showcasing the stable expression of PARP1-FLAG and PARP1-R735E-FLAG in the PARP1-KO U2OS cell lines. f) (i) Clonogenic assay assessing the sensitivity to camptothecin (CPT) of either the PARP1-KO U2OS cell line or the PARP1-KO cell line expressing either WT PARP1 or PARP1-R735E. (ii) Quantification of clonogenic survival for each strain.

Figure 1: Human TOP1 is prone to secondary cleavage and TOP1cc formation at genomic rNMP sites.
a) Diagram showcasing how TOP1 processes genomic ribonucleotides (rNMPs), and how this processing can lead to deletions in the context of short tandem repeats. b) (i) The titration of TOP1 against a 5'-^{32}P-labeled duplex DNA containing a single embedded rNMP was analyzed in a denaturing gel, (ii) with the quantification of the resulting species. c) TOP1cc products were processed with Proteinase K, Tdp1, and Tpp1 to reveal the TOP1 cleavage site. d) (i) TOP1 was titrated against a 5'-^{32}P-labeled CP nick substrate, (ii) with the quantification of the resulting species. e) (i) human TOP1 and yeast TOP1 (yTop1) were titrated against a CP nick substrate to examine their preference between CP nick ligation and slippage ligation. (ii) Quantification of the different products of TOP1 and yTop1. f) (i) 3'-Cy5-labeled duplex substrate and cleavage suicide substrate (fork) were tested with either human TOP1 or yTop1. (ii) The quantification of cleavage products.

Figure 2: PARP1 suppresses the secondary cleavage by TOP1 at genomic ribonucleotides.
a) (i) TOP1 ribonucleotide cleavage assay in the presence of PARP1 with or without NAD⁺. To prevent PARP1 from binding, both ends of the substrate were occluded by biotin-bound streptavidin as diagramed with blue ovals. (ii) The quantification of the CP nick and TOP1cc products at 50 nM PARP1 concentration and (iii) of the remaining substrate are shown. b) (i) The processing of the CP nick substrate by TOP1 was assessed in the absence or presence of PARP1 with or without NAD⁺, with both sides of the substrate occluded with streptavidin. (ii-iii) The quantification of the TOP1cc product and of the ligation products are shown.

Figure 3: PARP1 prevents deletions at short tandem repeats, and its simultaneous deletion with RNASEH2A generates a slow growth phenotype.

a) (i) Fluctuation assay reporting on the mutation rate of the different HeLa reporter strains as indicated. (ii) Quantification of the mutation types observed in sequenced clones of the WT and PARP1-KO strain showcases the prevalence of 2 bp deletions in the absence of PARP1. The n reflects the total number of indels identified in the total sequenced clones. b) Growth curve showcasing the delayed growth of the RNASEH2A-KO PARP1-KO cell line.

Figure 4: Concurrent deletion of RNASEH2A and PARP1 generates a detrimental accumulation of TOP1cc.
a) Sensitivity of WT, RNASEH2A-KO, PARP1-KO and RNASEH2A-KO PARP1-KO cell lines to the BCL2 inhibitor (BCL2i) navitoclax was determined by the MTT viability assay. b) Slot blot of the RADAR assay showcasing the elevated accumulation of TOP1cc in the RNASEH2A-KO PARP1-KO cell line. Samples from treating WT cells with 10 μM of the TOP1-trapping agent camptothecin (CPT) were included as a positive control. The DNA loading control was stained with Sybr Gold. c) Growth curve showing that the overexpression of TDP1 (TDP1-OE) rescues the delayed growth in the RNASEH2A-KO PARP1-KO cell line. d) Viability measured by MTT assay after TDP1 inhibitor, CD00509, (TDPi) treatment demonstrates hypersensitivity of the RNASEH2A-KO PARP1-KO cell line to TDP1 inhibitor. e) MTT viability assay showcasing the synthetic effect of TDP1i and BCL2i treatments to the viability of the RNASEH2A-KO PARP1-KO cell line.

Figure 5: TOP1 initial cleavage at genomic ribonucleotides stimulates PARylation by PARP1.
a) (i) PARP1-catalyzed auto-PARylation and PARylation of TOP1 in the presence of the duplex substrate with a single embedded ribonucleotide *in vitro*. Both DNA ends were occluded by biotin-bound streptavidin. (ii) Quantification of the amount of unPARylated PARP1 in the absence or presence of TOP1. b) (i) The *in vitro* PARylation assay using the ribonucleotide-containing substrate where the products were further treated with PARG, leaving single ADP-ribosyl modification that allows for precise determination of protein PARylation levels (ii) Quantification of PARylation levels of PARP1 in the absence or presence of TOP1. c) (i) *in vitro* PARylation assay with the ribonucleotide-containing substrate using the catalytic dead mutant TOP1-Y723F. (ii) Quantification of the amount of unPARy-lated PARP1 in the absence or presence of TOP1-Y723F. d) Diagram showcasing model on how PARP1 prevents the secondary cleavage by TOP1 at ribonucleotide sites both in a catalytically-independent (by physically restricting TOP1) and -dependent manner (through the inhibitory PARylation of both TOP1 and itself).

Figure 6: The DNA-binding activity of PARP1 and its physical interaction with TOP1 are important for the prevention of the TOP1 secondary cleavage.

a) Diagram showing the different domains of the full-length (FL) PARP1 and the two PARP1 truncations (PARP1-ΔDBD and PARP1-DBD) tested in the TOP1 ribonucleotide cleavage assay b) (i) The TOP1 ribonucleotide cleavage assay in the absence or presence of the full-length PARP1-FL, PARP1-ΔDBD or PARP1-DBD. (ii) Quantification of the 2', 3'-cyclic phosphate (CP) product and the TOP1cc product for the reactions with PARP1-FL, PARP1-ΔDBD and PARP1-DBD at 50 nM concentration. c) Domain diagram of N-terminal truncations of PARP1 (PARP1-ΔART and PARP1-ΔCAT) tested for TOP1-binding. d) The *in vitro* pulldown of FLAG-tagged TOP1 against the full-length PARP1 and PARP1 N-terminal truncations (ΔART and ΔCAT). Reactions without TOP1 were included as controls to assess non-specific binding of PARP1 to the anti-FLAG M2 resin.

Figure 7: Characterization of PARP1-R735E mutant in TOP1 interaction, the regulation of TOP1 ribonucleotide cleavage and sensitivity to PARP inhibitor, talazoparib.

a) The *in vitro* anti-FLAG-pull-down assay to test interaction of TOP1 with PARP1 mutants. The membrane was blotted with anti-His antibody for the detection of TOP1 and PARP1, both His-tagged. The supernatant (S) and elution (E) of the pull-down assay are shown. b) co-immunoprecipitation analysis of interaction between TOP1 and PARP1 in cells, where anti-FLAG M2 resin was used to immuno-precipitate either FLAG-tagged PARP1 or PARP1-R735E mutant stably expressed in U2OS PARP1-KO cell line to test their interaction with TOP1. c) (i) The TOP1 ribonucleotide cleavage assay in the absence or presence of PARP1 or PARP1-R735E. (ii) Quantification of the TOP1cc product from the reaction with either WT or the R735E mutant PARP1. d) (i) TOP1 ribonuclease assay in the presence of NAD+ comparing TOP1 inhibition by the WT PARP1 and the R735E mutant. e) (i) Clonogenic survival curve testing the sensitivity to talazoparib (TLZ) treatment of PARP1-KO U2OS cell lines expressing either no PARP1, WT PARP1, or PARP1-R735E (normalized to untreated cells and fitted using a nonlinear least-squares fit to a three-parameter dose–response model). (ii) A representative plate for each strain at the different TLZ concentrations as indicated.

Supplementary Table 1: List of plasmids used.

Construct	Description	reference
pFASTBac-HTB- FLAG-His$_6$-TOP1	Plasmid used to produce the Bacmid containing the 6xHis/FLAG-tagged TOP1 gene for expression in insect cells.	This work
pFASTBac-HTB- FLAG-His$_6$-TOP1-Y723F	Plasmid used to produce the Bacmid containing the His6/FLAG-tagged TOP1-Y723F gene for expression in insect cells.	This work
pET28a-His$_6$-PARP1	Plasmid for expressing full length PARP1 in *E. coli*, a kind gift from John Pascal's lab.	[1]
pET28a-His$_6$-PARP1-K695E	Plasmid for expressing PARP1-K695E mutant in *E. coli*.	This work
pET28a-His$_6$-PARP1-K700E+R704E	Plasmid for expressing PARP1-K700E+R704E mutant in *E. coli*.	This work
pET28a-His$_6$-PARP1-R735E	Plasmid for expressing full length PARP1-R735E in *E. coli*.	This work
pETDuet-His$_6$-PARP1-ΔDBD-FLAG	Bacterial expression of PARP1 lacking the DBD domain (379-1014).	This work
pETDuet- His$_6$-PARP1-N-366	Bacterial expression of DBD domain of PARP1.	This work
pET28_His$_6$-PARP1-ΔCAT (Addgene# 173942)	Bacterial expression of PARP1 lacking the CAT domain.	[2]
pET28_ His$_6$-PARP1-ΔART (Addgene #174804)	Bacterial expression of an inactive PARP1 mutant lacking the ART subdomain.	[2]
pcDNA3.1/Zeo(+)-TDP1	Plasmid for mammalian overexpression of TDP1.	This work
pCMV-PARP1-3xFlag (Addgene # 111575)	Plasmid for mammalian overexpression of 3xFLAG-PARP1	[3]
pCMV-PARP1-R735E-3xFLAG	Plasmid for mammalian overexpression of 3xFLAG-PARP1-R735E mutant	This work

Supplementary Table 2: List of oligonucleotides used to produce the substrates for biochemical assays.

Oligo name	Sequence (IDT standard)	Description
H1	CGGTTCGGACGTTCGAGCGTTGAAGATATGTGGCAAAACCTTTGTCTGGCTTCTGCTTGC	cold strand used for maintaining denatured labelled strand
H2	CGGTTCGGACGTTCGAGCGTTGAAGArUATGTGGCAAAACCTTTGTCTGGCTTCTGCTTGC	rU containing top strand for 60 bp rU containing substrate
H3	/5Biosg/GCAAGCAGAAGCCAGACAAAGGTTTTGCCACATATCTTCAACGCTCGAACGTCCGAACCG/3Bio/	biotinylated bottom strand for 60 bp substrate
H4	CGGTTCGGACGTTCGAGCGTTGAAGArUrAT	oligonucleotide used to make the CP-nick containing strand in 60 bp CP-nick substrate
H5	ATGTGGCAAAACCTTTGTCTGGCTTCTGCTTGC	strand that makes up the 5'-OH side of the nick in the 60 bp CP-nick substrate
H6	AGCGTTGAAGArUrAT	oligonucleotide used to make the CP-nick containing strand for 45 bp CP-nick substrate
H7	/5Biosg/GCAAGCAGAAGCCAGACAAAGGTTTTGCCACATATCTTCAACGCT/3Bio/	biotinylated bottom strand for 45 bp substrate
H8	AGCGTTGAAGArUATGTGGCAAAACCTTTGTCTGGCTTCTGCTTGC	rU containing top strand for 45 bp rU containing substrate
H9	GTCATTTGAATTCAAGCTrUTAGAGCTTAATTGCTGAATCT/3Cy5Sp/	3'-Cy5-labeled rU containing top strand of fork or duplex 41 bp substrate
H10	GACACTGCTTCCATTAGAGTAAAGCTTGAATTCAAATGAC	Bottom strand to produce 41 bp rU containing fork substrate
H11	*AGATTCAGCAATTAAGCTC*TAAAGCTTGAATTCAAATGAC	Bottom duplex to produce 41 bp rU containing duplex substrate
H12	GCTGAGCTTCTGGTGAAGCTCAGCACGCGGCAGCT/iCy3/GGTGCTGCCGCGrUrAT	Cy3-labeled oligonucleotide used to make the CP-nick containing dumbbell substrate
H13	GCTGAGCTTCTGGTGAAGCTCAGCACGCGGCAGCT/iCy5/GGTGCTGCCGCGT	Cy5-labeled nick dumbbell substrate

Supplementary Table 3: List of cell lines used.

Cell line	Description	reference
HeLa WT reporter cell	Previously published HeLa cell clone with 2-bp deletion reporter (from pTCW15) integrated at a single AAVS1 allele. Kind gift from Andrew P. Jackson.	[4]
HeLa RNASEH2A-KO reporter cell	Previously published RNase H2 null clone of the HeLa reporter cells with CRISPR/Cas9-mediated RNASEH2A knockout. Kind gift from Andrew P. Jackson.	[4]
HeLa RNASEH2A$^+$ reporter cell	Previously published RNase H2 partially proficient clone, resulting from CRISPR/Cas9 genome editing of HeLa parental reporter cells. Kind gift from Andrew P. Jackson.	[4]
HeLa PARP1-KO reporter cell	PARP1 null clone derived from the HeLa WT reporter cell. CRISPR/Cas9-mediated PARP1 knockout resulting from loss of function deletions in exon 2.	This work
HeLa RNASE H2A-KO PARP1-KO reporter cell	PARP1 null clone derived from the HeLa RNASEH2A-KO reporter cells. CRISPR/Cas9-mediated PARP1 knockout resulting from loss of function deletions in exon 2.	This work
HeLa RNASE H2A$^+$ PARP1-KO reporter cell	PARP1 null clone derived from the HeLa RNASEH2A$^+$ reporter cells. CRISPR/Cas9-mediated PARP1 knockout resulting from loss of function deletions in exon 2.	This work
U2OS parental cells	Cell line kindly provided by Shan Zha.	[5]
PARP1-KO U2OS cells	PARP1 null clone derived from the U2OS parental cells with CRISPR/Cas9-mediated PARP1 knockout. Kindly provided by Shan Zha.	[5]
PARP1-FLAG U2OS cells	PARP1-KO U2OS cells stably expressing PARP1-FLAG	This work
PARP1-R735E-FLAG U2OS cells	PARP1-KO U2OS cells stably expressing PARP1-R735E-FLAG.	This work

Supplementary Table 4: List of mutations found in the hygromycin-puromycin fusion reporter gene of isolated clones of either PARP1-KO or WT strains.

strain, clone #	original sequence (red for deletions and - for insertion location)	mutant	site, **type of indel**
PARP1-KO, clone 1	GAATCTCTCTCAAACCTC	GAA-TCTCTCAAACCTC	d378-379,**2nt deletion**
PARP1-KO, clone 2	TTTATATGCGC	TTT-ATGCGC	d451-452,**2nt deletion**
PARP1-KO, clone 3	AAACTGTGATGGA	AAAC-TGATGGA	d492-493,**2nt deletion**
PARP1-KO, clone 4	CTCAGAGAGCCT	CTC-AGAGCCT	d252-253, **2nt deletion**
PARP1-KO, clone 5	TGCGCGCATATATGC	TGCGCGC-ATATGC	d792-793, **2nt deletion**
PARP1-KO, clone 6	AAAGACAGATATGTATACAGACA	AAAGACAG-ATGTATACAGACA	d184-185, **2nt deletion**
PARP1-KO, clone 7	GCGCAGCGCTACCTA	GCGCA-GCTACCTA	d211-212, **2nt deletion**
PARP1-KO, clone 8	AAAGACAGATATGTATACAGACA	AAAGACAG-ATGTATACAGACA	d184-185, **2nt deletion**
PARP1-KO, clone 9	ATCTCTCTCAAACCT	ATC-TCTCAAACCT	d378-379, **2nt deletion**
PARP1-KO, clone 10	CTTCGCGAGCGCAGC	CTT-CGAGCGCAGC	d201-202, **2nt deletion**
WT, clone1	CCTCACATATTGTATATCCAGGCGCGCGCAAGGTGTAACACTACAAGATCTCC	CCTCACA-AGATCTCC	d265-302, **38bp deletion**
WT, clone2	AAGA-GTCGCGCGCATT	AAGAAGTCGCGCGCATT	between 98-99, **1nt insertion**
WT, clone3	CCTCACATATTGTATATCCAGGCGCGCGCAAGGTGTAACACTACAAGATCTCC	CCTCACA-AGATCTCC	d265-302, **38bp deletion**

WT, clone 4	GA**G**GGCGAA	GA-GGCGAA	d90, **1nt deletion (3 homopolymer)**
	T**G**GGAGGGAA	T-GGAGGGAA	d124, **1nt deletion**
WT, clone 5	CTCA**T**AGAG	CTCA-AGAG	d48, **1nt deletion**
	GTG**T**GTCAGA	GTG-GTCAGA	d66, **1nt deletion**
WT, clone 6	T**A**AACTCTTG	T-AACTCTTG	d150, **1nt deletion (3 homopolymer)**
	GA**G**GGCGAAG	GA-GGCGAAG	d90, **1nt deletion (3 homopolymer)**
WT, clone 7	CAT**AGAG**AAGTTC	CATAAGTTC	d48-51, **4nt deletion**
	AAAGGAA**TA**G	AAGGAA-G	d1024-1025, **2nt deletion**
	CATAGAGAAG-TTC	CATAGAGAAG**A**TTC	between 54-55, **1bp insertion**

Supplementary Table references

1   Langelier, M. F., Servent, K. M., Rogers, E. E. & Pascal, J. M. A third zinc-binding domain of human poly(ADP-ribose) polymerase-1 coordinates DNA-dependent enzyme activation. *J Biol Chem* **283**, 4105–4114 (2008). https://doi.org/10.1074/jbc.M708558200

2   Ooi, S. *et al.* Multiple roles for PARP1 in ALC1-dependent nucleosome remodeling. *Proc. Natl. Acad. Sci.* **118 (36)**, epub (2021).

3   Liszczak, G., Diehl, K. L., Dann, G. P. & Muir, T. W. Acetylation blocks DNA damage-induced chromatin ADP-ribosylation. *Nature chemical biology* **14**, 837–840 (2018). https://doi.org/10.1038/s41589-018-0097-1

4   Reijns, M. A. M., Parry, D.A., Williams, T.C. et al. Signatures of TOP1 transcription-associated mutagenesis in cancer and germline. . *Nature*, 623–631 (2022).

5   Shao, Z. *et al.* Clinical PARP inhibitors do not abrogate PARP1 exchange at DNA damage sites in vivo. *Nucleic Acids Res* **48**, 9694–9709 (2020). https://doi.org/10.1093/nar/gkaa718

Supplemental Figure 1: SDS-PAGE analysis of all purified proteins.
All proteins were fractionated in 4-15% SDS-PAGE or 10% SDS-PAGE and stained by Coomassie blue G-250.

Supplemental Figure 2: Human TOP1 cleavage analysis and comparison of rNMP processing with yeast Top1 (yTop1).
a) (i) Time course analysis of ribonucleotide processing by human TOP1. (ii) Quantification of the different product species and the remaining substrate. b) (i) Comparison of ribonucleotide processing between human TOP1 and yTop1. (ii)-(iii) Quantification of the remaining substrate and the TOP1cc product for each TOP1 variant.

Supplemental Figure 3: CP nick activates PARP1 for TOP1 regulation.
a) (i) PARP1 binding competition assay using a Cy5 labeled dumbbell nick substrate and a Cy3 labeled dumbbell CP-nick substrate. Single channel and merged images are shown. (ii) Quantification of the remaining unbound substrate for each condition. b) (i) TOP1 ribonucleotide cleavage assay performed in the absence and presence of NAD+ to examine the effect of NAD+ on TOP1. (ii) quantification of TOP1cc and CP nick products. c) (i) TOP1 ribonucleotide cleavage assay performed in the absence or presence of indicated amount of PARP1 or the catalytically dead PARP1-E988K mutant, with or without NAD+. (ii) Quantification of TOP1cc and CP nick products from each reaction. (iii) Quantification of the remaining substrate in reactions containing PARP1 or PARP1-E988K in the presence of NAD+.

Supplemental Figure 4: Confirmation of PARP1 knockout and TDP1 overexpression in the reporter cell lines and examination of the effect of PARP inhibitor on the mutation rate in the wild-type reporter cells.
a) PARP1 was knocked out in the WT, RNase H2A-KO, and RNase H2A+ reporter HeLa cells. TDP1 was stably expressed in the RNase H2A-KO PARP1-KO cell line under the control of cmv promoter. The protein samples were fractionated in 4-15% SDS-PAGE and transferred into a nitrocellulose membrane for western blots. b) Fluctuation assay reporting on the mutation rate of the WT HeLa reporter strain pre-treated with either PARP inhibitor talazoparib (TLZ) or DMSO, showcasing that inhibition of PARP1 does not significantly affect the number of deletions at STRs.

Supplementary Figure 5: Though deficient in TOP1-interaction, PARP1-R735E is comparable to the wild-type in DNA binding, autoPARylation and response to PARP inhibitor *in vitro,* as well as complementing PARP1-KO for the sensitivity to camptothecin in cells.

a) A model of the interface between TOP1 and the HD domain of PARP1 from AlphaFold prediction. The targeted residues within the the HD domain (K695, K700, R704, and R735) are labeled with the numbers 1-4, respectively. A close-up of the PARP1-TOP1 interface and the electrostatic surface map of this interface. b) Alignment of the PARP1 HD domain region containing the targeted residues across higher eukaryotes, including human, chicken, Xenopus frog, zebrafish, and Arabidopsis thaliana. Arrows mark the sites tested for TOP1 interactions. Green arrows indicate sites whose mutation reduced TOP1 binding, whereas the red arrow indicates a site that retained substantial TOP1 binding after mutation. c) (i) DNA binding assay comparing WT PARP1 and the R735E mutant ability to bind to the CP-nicked dumbbell substrate. (ii) Quantification of the unbound substrate for both PARP1 variants. d) Coomassie stained 10% SDS-PAGE images showcasing the auto-PARylation activity and talazoparib sensitivity of both PARP1 and PARP1-R735E. e) Western blot showcasing the stable expression of PARP1-FLAG and PARP1-R735E-FLAG in the PARP1-KO U2OS cell lines. f) (i) Clonogenic assay assessing the sensitivity to camptothecin (CPT) of either the PARP1-KO U2OS cell line or the PARP1-KO cell line expressing either WT PARP1 or PARP1-R735E. (ii) Quantification of the clonogenic survival for each strain.