Energetic misfires: Hybridization drives transgressive expression

in metabolic pathways in thermally divergent Icelandic stickleback

Matthew K Brachmann[1*], Bethany Smith[1*], Bjarni Kristjánsson[2], Colin Selman[1],

Kevin Parsons[1]

[1] School of Biodiversity, One Health, and Veterinary Medicine, University of Glasgow,
Glasgow, UK.

[2] Department of Aquaculture and Fish Biology, Hólar University, Saudárkrókur,
Iceland.

*Co-first authorship

Abstract

Climate change is causing rapid changes to freshwater environments, driving

selection for phenotypes that can cope with altered thermal conditions, while also

changing developmental environments. This may promote local adaptation,

migration to new habitats, and/or phenotypic plasticity. Climate change may also

increase hybridization rates between locally adapted phenotypes, as populations

migrate and spatially mix in new ways. Consequently, these conditions may facilitate

the production of variation, including both adaptive and maladaptive outcomes. To

examine this, we leveraged threespine stickleback (*Gasterosteus aculeatus*) that

have locally adapted to either geothermally warmed or ambient environments in

Iceland. We tested the effects of hybridization between thermal ecotypes on gene

expression using a common garden experiment, where pure-strain ecotypes and

their hybrids were raised under 12°C and 18°C. We performed RNA-seq on brain

and liver tissues to assess 1) ecotype divergence, 2) plasticity, and 3) the effects of

hybridization and inheritance patterns on gene expression. We identified a low

degree of expression divergence between locally adapted ecotypes, despite a high

degree of plasticity across rearing environments. Hybrid ecotypes were highly

divergent from both geothermal and ambient ecotypes and exhibited transgressive

expression under both rearing temperatures. Transgressive expression disrupted

gene networks extensively, with broader effects at 18°C than 12°C, that were

associated with metabolism and mitochondrial function. Hybridization between

locally adapted thermal ecotypes appears to largely disrupt genes associated with

energy balance and metabolic function. While demonstrating mechanisms underlying

the rapid evolution of reproductive isolation, these findings also provide insights for

how populations may cope or fail within a warming world.

Introduction

Climate change presents an ever-worsening threat to biodiversity (IPCC,

2021) as continued warming pushes species to thermal limits (Nunez et al., 2019).

Ectotherms are expected to be particularly vulnerable due to thermoregulation being

mediated by the environment (Fry, 1967; Paaijmans et al., 2013), and where

increasing temperatures destabilize thermoregulatory capacity (Haesemeyer, 2020;

Duffy et al., 2022). Climate change will require ectotherms to either migrate to more

favorable habitats or rapidly adapt to warming environments. Within ectotherms,

climate change has already been implicated as a cause for adaptive phenotypic

shifts across fishes (Crozier & Hutchings, 2014). However, the contribution of

phenotypic plasticity underlying such changes is unclear with a heritable basis being

untested in most cases (Crozier & Hutchings, 2014). Thus, there is a pressing need

to understand how climate change can induce adaptation through plasticity, genetic

evolution, and exactly how these factors interact. Determining the underlying

mechanisms is an area of growing interest as it can inform conservation strategies

(Reed et al., 2011; Campbell et al., 2017; Greenspoon & Spencer, 2021; Parsons,

2021; She et al., 2023; Sheldon et al., 2025; Wighard et al., 2025).

The capacity for adaptation to climate change is dependent on the amount of

phenotypic variation generated through genomic and developmental processes

(Pigliucci, 2008; Campbell et al., 2017; Riederer et al., 2022). The production of

adaptive phenotypic variation through development involves a combination of

environmental, genomic, and regulatory changes (Hansen, 2006; Riederer et al.,

2022). The inheritance of adaptive phenotypic variation can be both additive and

non-additive (Falconer & Mackay, 1996), with phenotypic expression being further

modified by cis- vs trans-regulatory elements (Wittkopp et al., 2004). While cis-

regulatory elements act on specific alleles and trans-regulatory elements influence

both alleles equally, allele specific expression allows for the disentanglement of the

inheritance of regulatory variation and differentiate between cis- and trans-regulatory

effects (Fan et al., 2020; Pierre et al., 2022). Transgressive expression, due to non-

additive inheritance, involves gene expression in hybrids that is greater or less than

parental phenotypes (Go & Civetta, 2020; Wang et al., 2022; Yazdi et al., 2022) and

can arise when cis- and trans-regulatory elements are decoupled, revealing cryptic

regulatory variation (Lemmon & Juenger, 2017; Jacobs et al., 2024). The relative

role of cis- vs trans-regulatory elements, and their interaction with both additive and

non-additive inheritance mechanisms, in response to changing environments has

been debated (Crispo et la., 2008; Lemos et al., 2008; Osada et al., 2017; McGirr &

Martin, 2021).

Hybridization between locally adapted species or ecotypes may promote

genomic and phenotypic variation within novel environments (Abbott et al., 2013;

Brice et al., 2021; Jacobs et al., 2024). Secondary contact and hybridization may

occur between locally adapted populations across both small and large scales

(Chunco, 2014). Hybrid offspring at zones of secondary contact are unlikely to have

phenotypes that match either parental phenotype (Gow et al., 2007; Rundle & Nosil,

2005) and these novel phenotypes may be selected against (Servedio, 2004;

Lemmon & Kirkpatrick, 2006; Lenormand, 2012; Butlin & Smadja, 2018). Hybrids are

generally predicted to be maladaptive in either environment due to a disruption of

adaptive gene by environment interactions (Carroll et al., 2003; Cenzer, 2017) and

can reinforce reproductive isolation (Lemmon & Juenger, 2017). In a warming world,

populations that are locally adapted to different thermal conditions may come into

contact at secondary contact zones as they begin to migrate (Chunco, 2014).

Therefore, addressing how parental and hybrid offspring of thermally adapted

populations respond to temperature variation could be insightful for understanding

how genomic and regulatory mechanisms are impacted by climate change.

Icelandic threespine stickleback (*Gasterosteus aculeatus*) provide a powerful

system to untangle the effects of hybridization on the molecular architecture of

adaptation to warming environments. While understanding adaptation to warming

environments often involves comparing populations from widely varying latitudes or

altitudes as a proxy, this confounds temperature variation with other environmental

factors (Daco et al., 2021; Pilakouta et al., 2023). However, populations of three-

spined stickleback (*Gasterosteus aculeatus*) avoid such confounding effects by

inhabiting geothermal and ambient habitats in close geographic proximity across

Iceland (Millet et al., 2013; Pilakouta et al., 2020; Pilakouta, et al., 2023). Geothermal

and ambient habitats exhibit a strong thermal gradient that typically differ by ~10°C

throughout the year, but with otherwise similar environments (Pilakouta et al., 2020).

Stickleback living across geothermal-ambient gradients demonstrate heritable

adaptive divergence in as little as 70 years, including large fitness tradeoffs

(Pilakouta, et al., 2023; Smith et al., 2024).

To understand the genomic and developmental mechanisms underlying

phenotypic divergence between thermal ecotypes we tested the influence of

plasticity and hybridization on gene expression. We tested three hypotheses: 1)

Geothermal and ambient stickleback ecotypes will show divergent and plastic gene

expression across temperature environments, reflecting local adaptation. 2) Hybrid

expression will differ from pure-strain ecotypes due to non-additive, transgressive

expression. 3) Hybridization between locally adapted ecotypes will disrupt co-

adapted cis-trans regulatory networks, leading to misexpression of specific alleles of

locally adapted genes and maladaptive phenotypes.

Methods

Creation of F1 hybrid and pure strain families.

Sexually mature sticklebacks were captured using un-baited minnow traps

(mesh size, 6.4mm) from an allopatric geothermal-ambient habitat pair in Northern

Iceland, near Sauðárkrókur (geothermal habitat: 65.732260^{o}, -19.618937^{o}, ambient

habitat 65.732191^{o}, -19.618574^{o}) in 2017 and brought to the University of Glasgow.

Ambient and geothermal summer temperatures were simulated in the aquaria using

12°C (±1°C) and 18°C (±1°C), respectively. Following two months of a simulated

winter light cycle (4hrs) fish were primed for reproduction with nutrient rich food

(bloodworms, mysis) and an extended light period (20-24hrs). *In vitro* fertilization

was performed to create four ambient families, four geothermal families, and two

hybrid families using two geothermal females and two ambient males (Figure 1). Egg

clutches were left to fertilize for fifteen minutes before being split evenly to 12°C

(±1°C) and 18°C (±1°C).

Fertilized clutches were kept in mesh baskets submerged within larger basins

of water fed by a recirculated continuous water drip system. Water was treated with

methylene blue (2.5 µg/ml) until hatching to reduce the risk of fungal infection (Ansari

et al., 2016). Containers were inspected daily, and dead or infected eggs were

removed immediately. After hatching, larvae and juveniles were fed live food (newly

hatched brine shrimp (*Artemia* salina), and microworms (*Panagrellus redivivus*) as
well as size-appropriate ZM powdered food (ZM100 and ZM200) (ZMsystems,
Twyford, UK). Upon reaching approximately 2cm, juveniles were moved, in their
family groups, to 10L tanks at standardized densities of 15-20 individuals. From 3cm
to maturation fish were fed trout pellets (Microstart, EWOS Ltd, Surrey, UK). The
photoperiod of the aquarium was kept at a 12:12 light cycle.

Tissue collection, RNA extraction and sequencing

Brain and liver samples were collected from stickleback for each treatment
condition. Sampled fish were one year old (±1 month), and that provide a size match
across treatments. Individuals were sexed by the identification of testes and ovaries.
Brain and liver samples were collected from 8 individuals per ecotype for each
temperature treatment with sampling from 3-4 separate families for the geothermal
and ambient ecotypes and 2 families for the hybrids, resulting in 48 total samples
(Figure 1). Each tissue was cut into 0.5cm^3 pieces and stored in RNAlater (Invitrogen
AM7020) at room temperature overnight before freezing at -80°C.

To obtain RNA, extractions were performed using a Qiagen RNeasy standard
kit (Qiagen Cat ID: 74104) for liver tissue (with 50% ethanol as recommended for
fatty tissue such as liver) and the Qiagen RNeasy lipid tissue kit (Qiagen Cat ID:
74804) for brain tissue. RNA was extracted from ~20mg of tissue and purity was
determined using a NanoDrop spectrophotometer (Thermo Fisher Scientific,
Massachusetts, US), where all samples had 260/230 or 260/280 ratios > 1.8.
Concentration was determined using a RNA Broad Range kit (Thermo Fisher
Scientific, Massachusetts, US) on a Qubit 4 Fluorometer using a Qubit RNA Broad
Range kit (Thermo Fisher Scientific, Massachusetts, US). RNA integrity was
assessed for all samples, using an Agilent 4200 Tapestation (Agilent Technologies,
California, US), to calculate RNA integrity number (RIN values). RIN values are a
numerical score ranging from 1-10 where a higher value indicates a higher degree of
RNA integrity (Sheng et al., 2017) and all samples has a RIN value >7.

The preparation of cDNA libraries and whole transcriptome sequencing were
performed by the Center for Genomic Research (CGR) a the University of Liverpool
using the NEBNext® Poly(A) mRNA Magnetic Isolation Module (NEB E7490) and
NEBNext® Ultra RNA Library Prep Kit for Illumina® (E7530). Sequencing was

performed on the Novaseq 6000 platform (Illumina, California, US) aiming for an

average read length of 150 base pairs and 20M sequenced reads per sample.

RNA-seq data analysis

Initial data processing and quality control

Read quality was investigated using *FastQC* (Andrews, 2010) and *MultiQC*

(Ewels et al., 2016). Fastq files were trimmed for the presence of Illumina adapter

sequences using *Cutadapt* version 1.2.1 (Martin, 2011) and further trimmed using

Sickle version 1.200 (Joshi & Fass, 2011) before alignment to version five of the

three-spined stickleback reference genome (Genbank GCA_016920845.1) (Peichel

et al., 2020) using the *Hisat2* aligner version 2.2.0 (Kim et al., 2019). The resulting

SAM files were then sorted and converted to BAM format using *Samtools sort* and

indexed using *Samtools index* (Danecek et al., 2021). Read counts were obtained

using *Htsqcount2* (Anders et al., 2015).

Differential expression between ecotypes and temperatures

Data analysis was performed in R (v4.5.1) (R Core Team, 2022). The

packages *EdgeR* (Robinson et al., 2009) and *LIMMA* (Ritchie et al., 2015) were used

in parallel to provide higher confidence through corresponding results. The read

count data was filtered for low-expressed genes using the *filterbyExpr* function in the

EdgeR package. Genes were filtered out if they had fewer than ten counts per million

(CPM) in more than 8 samples for each tissue type (Chen et al., 2016). Read counts

were normalized by converting observed library sizes into effective library sizes

using scaling factors calculated by *calcNormFactors* using the TMM method

(Robinson & Oshlack, 2010).

To address our first hypothesis, we tested for differences in gene expression

between ecotypes, hybrids, rearing temperatures, and their interaction using a single

contrast matrix. For all analyses the false discovery rate (FDR) was controlled using

the Benjamini-Hochberg method (Benjamini & Hochberg, 1995). Genes were

considered differentially expressed when the resulting adjusted p value was below

0.05. In our contrast matrix, we tested for divergent expression by subtracting

expression values for geothermal fish at a given rearing temperature from ambient

expression at the same temperature, resulting in two contrasts ("*ambient at 12°C –*

geothermal at 12°C" and "*ambient at 18°C – geothermal at 18°C"*). To examine

plasticity we compared expression at 18°C versus 12°C within each fish type
(geothermal, ambient, and hybrid). To address our second hypothesis, we then
tested for mismatches between hybrids and pure ecotypes by subtracting hybrid
expression at each temperature from a given pure ecotype expression at the same
temperature, resulting in four contrasts ("*ambient at 12°C – hybrid at 12°C", "ambient*
at 18°C – hybrid at 18°C", "geothermal at 18°C – hybrid at 18°C", and *"geothermal at*
12°C – hybrid at 12°C"). In *EdgeR* a negative binomial generalized linear model was
fitted to each gene. This analysis assumed that the variance of gene counts was
dependent on the negative binomial dispersion and the quasi-likelihood dispersion
(Ren & Kuan, 2020). Here, the negative binomial dispersion represented the
variability of the biological system, while the quasi-likelihood dispersion represented
gene-specific variability greater or lower than the overall level, capturing both
biological and technical sources of variability (Lun et al., 2016). The mean-dispersion
across all genes was used to estimate the negative binomial dispersion using the
estimateDisp function (Lun et al., 2016; Ren & Kuan, 2020). An empirical Bayes
approach was then used to estimate the quasi-likelihood dispersion and a
generalized linear model accommodating the design model matrix was fitted using
glmQLFit (Lun et al., 2016; Ren & Kuan, 2020). The top-ranked genes for each
contrast were then extracted using *TopTags*.

We then performed a complementary analysis using *LIMMA* to test
differences in gene expression within our four contrast matrices. Normalized read
counts were prepared for linear modelling in *LIMMA* by performing a voom
transformation, which incorporated the mean-variance relationship of the log-counts
into the precision weights for each observation (Law et al., 2014, 2018). Next, a
linear model was fitted to each gene using *lmFit* (Law et al., 2018). A corollary test
was then done using *eBayes* to perform a moderated F-statistic test and rank genes
by evidence of DE for each contrast (Law et al., 2018). The top-ranked genes for
each contrast were then extracted using *topTable*.

We then compared the results from *EdgeR* and *LIMMA* and selected genes
that showed differential expression in both analyses. For both approaches the fitted
models used contrast matrices designed to address our questions around
divergence, plasticity, and their interaction. These contrast matrices described the
linear combinations of parameters used to calculate the differences between groups

of interest (Law et al., 2020). We then tested for differences between fish types and

rearing temperatures using a principal component analysis (PCA) on all differentially

expressed genes for brain and liver tissues using the *prcomp* function.

Hybrid gene expression and inheritance mode categorization

To further address our second hypothesis, we assessed inheritance mode as

it was necessary to categorize patterns of gene expression. Thus, in each rearing

temperature hybrid expression was compared to pure strain fish. Genes with DE in

at least one of the previous hybrid to pure-strain contrasts were selected for

comparison, and inheritance modes were classified separately for each rearing

temperature. Specifically, genes were categorized based on the log fold changes

(LFC) of DE between hybrids and each pure-strain type with a threshold of 0.32

(1.25-fold difference, shown to be a useable cut-off by (Wang et al., 2022; Yazdi et

al., 2022)). DE genes were categorized as either additive, dominant, or transgressive

on the basis of their respective LFC threshold (Figure 2) Dominant gene expression

was assessed for each of the ecotypes where the hybrids had a LFC between -0.32

and 0.32, we could then assess dominance due to either the geothermal or ambient

ecotype. Additive gene expression was classified as being >0.32 LFC in one ecotype

and <0.32 in the other ecotype. Lastly, transgressive gene expression was classified

as the LFC being > or < 0.32 in both ecotypes.

Testing for divergence and disruption in gene co-expression

To assess our third hypothesis, we assessed the effects of hybridization on

gene co-expression networks linked up developmental pathways (Oldham et al.,

2006). Changes in gene networks between ecotypes can be used to discover key

drivers of evolutionary divergence (Oldham et al., 2006), while hybrids can display

disrupted co-expression networks that may alter phenotypic development and

decrease fitness (Filteau et al., 2013). Thus, gene networks associated with

divergence and plasticity were identified using the *GeneCoEx* workflow in R (Li et al.,

2023) whereby differentially expressed genes for each contrast and tissue type were

used to produce a correlation matrix and detect co-expression modules. First,

significant DEGs from each contrast were filtered to remove genes with invariable

expression profiles. A correlation matrix was then made for all genes and

correlations greater than 0.7 were retained and relationships between highly

correlated genes were calculated. For each contrast, modules of gene expression
were calculated using the *cluster_leiden* and *optimize_resolution* functions. The
cluster_leiden function performs Leiden clustering using the Leiden algorithm (Traag
et al., 2019) to find modularity in gene expression. To control false discovery and
spurious results, each module from the *optimize_resolution* function had to contain 5
or more genes with gene expression being calculated for each module. The gene
network was calculated using the *graph_from_data_frame* function which uses the
edges and nodes calculated with correlation coefficients greater than 0.7. Gene
networks for each contrast were then imported into *Cytoscape* (Ono et al., 2025) for
visualization and gene ontology analysis. Gene ontology (GO) analysis was then
performed for each gene network using *Enrichment Map* within *Cytoscape* (Merico et
al., 2010).

Comparing ecotype regulatory mechanisms through allele specific expression
analysis

To further address our third hypothesis, we assessed divergence in regulatory
mechanisms through investigation of allele-specific expression (ASE), defined as the
imbalanced expression of maternal or paternal alleles of a gene in a diploid
individual. ASE can influence evolution by enabling natural selection to act on
individual gene copies, allowing organisms to mask deleterious mutations or
adaptively overexpress beneficial ones (Wittkopp et al., 2004, 2008; Wittkopp &
Kalay, 2012). Thus, ASE can act as a regulatory mechanism that creates phenotypic
variation, driving rapid adaptation to environmental changes and influence complex
traits through tissue-specific and parent-of-origin effects. The premise for the
identification of ASE is that one copy of each pure-strain chromosome should be
present in each cell of an F1 hybrid, meaning that while the trans-regulatory
environment for each allele is the same, the cis-regulatory environment may differ.
Differences in cis-regulatory elements can therefore result in the differential
expression of each allele in hybrids.

To assess the presence of allele specific expression, aligned Bam files were
prepared for variant calling using the *GATK* best practices RNAseq short variant
discovery workflow (Brouard et al., 2019; Caetano-Anolles, 2023b). Post-alignment
quality was checked using *Picard ValidateSamFile* while mate-pair information was
verified and fixed, and duplicate reads tagged using *Picard's FixMateInformation* and

MarkDuplicates (Institute, 2023). *GATK's SplitNCigarReads* was used to reformat

alignments spanning introns for compatibility with *HaplotypeCaller*. An initial round of

variant calling was performed in order to bootstrap a set of known variants, which

were filtered (*GATK: VariantFiltration, SelectVariants*) using hard filters (Quality by

Depth (QD) <2.0, Fisher Strand (FS) > 60, RMS Mapping Quality (MQ) < 40, Strand

Odds Ratio (SOR) >4, as recommended by *GATK* best practices (Caetano-Anolles,

2023a)), for input into *GATK's Base Quality Score Recalibration*. Base quality

recalibration was performed using machine learning applied to each sample to detect

and correct patterns of systematic errors in the base quality scores using

BaseRecalibrator. A second round of variant calling using *HaplotypeCaller* provided

variants, which were also filtered (*VariantFiltration, SelectVariants*) using hard filters

(QD <2.0, FS > 60, MQ < 40, SOR >4) and SNPs were called for all fish

(geothermal, ambient, and hybrid). Pure-strain SNPs were filtered for homozygosity

across each ecotype to extract those unique to each.

SNPs identified in hybrid fish were filtered to include only those that were

heterozygous to target those that may be habitat divergent. Allele specific reads

were then output from the VCF files using GATK's *ASEReadCounter*. SNPs were

mapped to genes using *SNPeff* and filtered for a total read count of >20 with a

minimum read count for the reference and alternatives of >5. Allele specific

expression was then assessed per SNP using Allele-Specific Expression analysis in

a Population (ASEP) (Fan et al., 2020). ASEP is a method for gene-level detection of

allele-specific expression at the population level without individual genomic data,

using a pseudo-phasing procedure that uses allele-specific read counts to infer

haplotype phases through a majority voting procedure based on allele specific read

counts. A generalized linear mixed-effects model is then used to test for allele-

specific expression. Because differing levels of allele-specific expression between

rearing temperatures could indicate environmental effects on expression regulation

ASEP analysis was performed separately for hybrids reared at 12°C and 18°C. To

assess biological relevance SNPs identified by ASEP analysis were cross-

referenced to SNPs present in pure-strain fish that were unique to either ecotype.

Genes were annotated using Ensembl three-spine stickleback gene stable

IDs. To conduct ontology analysis Ensembl Stickleback gene stable IDs were

converted to Ensembl Zebrafish gene stable IDs using *biomaRt* (Durinck et al., 2009;

Kinsella et al., 2011). The Metascape package (Zhou et al., 2019) was then used on
converted data to interpret the list of DE genes produced from each analysis
(minimum overlap 3, p value cut-off 0.01 and minimum enrichment 1.5). Gene
ontology (GO) enrichment analysis relies on an accurate assessment of whether a
particular gene ontology is overrepresented in a dataset. Therefore, in order to avoid
sampling bias, background gene sets of only the expressed genes (non-zero total
read), rather than the whole zebrafish genome, were used for each tissue (Timmons
et al., 2015). As the recommended maximum for Metascape gene ontology analysis
is 3000 genes, gene lists that exceeded this threshold were filtered to remove genes
with an absolute LFC lower than 0.5. This analysis was performed only on modules
found to be significantly divergent, disrupted or plastic. The interpretation of module
gene ontology was further simplified by selecting the top-level gene ontology terms
containing the most significant lower level GO term (based on adjusted p values (log
q)).

Results

Divergence and plasticity of gene expression

Under common temperatures, heritable divergence in gene expression
occurred between geothermal and ambient ecotypes (Figure 2A-B). In the brain
there were a greater number of divergently expressed genes between geothermal
and ambient ecotypes at 12°C than at 18 °C (Figure 2; Table S1). Two genes were
divergent between geothermal and ambient ecotypes regardless of temperature,
loc120815948 and *trpc4a*. In the liver, there were no divergently expressed genes at
either temperature (Figure S1; Table S1).

We detected plastic responses across both 12°C and 18°C in ecotypes and
hybrids in the brain (Figure 3A-C; Table S2). The geothermal ecotype and hybrids
showed a greater number of plastic genes than the ambient ecotype (Table S2).
There were 47 shared plastic genes overlapping between the geothermal and
ambient ecotypes (Table S2), which makes up 77% of the plasticity genes exhibited
by the ambient ecotype. A similar proportion of plastic genes overlapped between
the ambient ecotype and the hybrids. While the geothermal ecotype and hybrids
shared 85 genes, the majority of their plastic responses were in genes unique to
each ecotype. Across ecotypes and hybrids there were 38 shared plastic responses

showing evidence for enrichment of RNA processing (6 genes) and splicing (5

genes). Comparatively, there was a lower degree of plasticity in gene expression

across 12°C and 18°C for ecotypes and hybrids in the liver relative to the brain

(Figure 2SA-C; Table S2), with only a single down-regulated gene in the liver, *ptpn6*.

Effects of hybridization on divergence and plasticity of gene expression

Divergence in gene expression for both ambient and geothermal ecotypes

relative to hybrids was extensive in the brain (Figure 4A-D; Table S3). In total, 2045

genes were differentially expressed between the ambient ecotype and hybrids

across rearing temperatures, with 553 and 531 genes being uniquely divergent

respectively at 12°C and 18°C. A large proportion (78% at 12°C and 79% at 18°C) of

differentially expressed genes were divergent at both temperatures between the

ambient ecotype and hybrids indicating a strong heritable component. A similar

pattern of divergence was observed between the geothermal ecotype and hybrids,

with 1559 divergent genes across both rearing temperatures (78% at 12°C and 75%

at 18°C), again indicating a strong heritable component of variation. There was a

high degree of divergence in gene expression between both the ambient and

geothermal ecotypes at 12°C and 18°C (Table S3). a total of 1395 genes were

divergent between hybrids and both ecotypes regardless of rearing temperature. In

contrast to the brain, there were only a small number of genes involved in expression

divergence between hybrids and both ecotypes in the liver (Figure S4A-D; Table S3).

There were only 2 divergent genes at 12°C and no divergent genes at 18°C between

ecotypes relative to hybrids, with no overlap of these genes between comparisons.

Regardless of temperature, there was substantial divergence of

transgressively expressed genes between the hybrids and both ecotypes in the brain

(Figure 4; Table S4). Hybrids were also clearly divergent from both ecotypes within

the PCA (Figure 4A) with 87% of differentially expressed genes at 12 and 18°C were

classified as transgressively expressed (Figure 5B-D; Table S4). Divergence

between hybrids and ecotypes displayed 2405 transgressively expressed genes

irrespective of rearing temperature. Of these, 354 were unique to 12°C, while 377

were unique to 18°C. In the liver, there were 15 transgressively expressed genes at

12°C and 4 at 18°C, with only 2 genes overlapping between temperatures.

Gene co-expression networks

In the brain, whole gene networks were associated with plastic responses for

each of the ambient and geothermal ecotypes and their hybrids (Figure S5A-C).

There were 49 genes connected by 140 edges making up the gene network of

plastic responses for the ambient ecotype (Figure S5A) with gene ontologies (GO)

associated with DNA binding and mitochondrial ABC transporters (Table S5).

Geothermal plastic response gene networks had 181 genes connected by 507 edges

(Figure S5B) with GO terms associated with mRNA processing, spliceosome,

defense responses, hydrolase activity, and cytoplasm (Table S5). Lastly, the gene

network of hybrid plastic responses had 160 genes connected by 461 edges (Figure

S5C) associated with the cytoplasm (Table S5). These three gene networks

overlapped at 23 genes, connected by 8 edges, corresponding to lysozyme activity,

CD4 receptor binding, and peptidoglycan muralytic activity (Table S5). All

overlapping plastic gene networks appear to function as part of the immune system

responses to increased temperature.

Divergence between ecotypes was dependent on temperature with

expression at 12°C associated with a whole network of genes, but this was not the

case at 18°C. The gene network associated with divergence at 12°C had 13 genes

and 20 edges connecting the genes (Figure S6). Gene ontology analysis of the

network identified *yrdc* as being associated with RNA binding and translational

fidelity and *mgst3a* being involved in glutathione-dependent detoxification of reactive

toxins. At 18°C, no genes had an association above 0.7 and a network could not be

built.

The gene networks associated with divergence between ecotypes and hybrids

contained more genes than any other network (Figure S7A-D), regardless of rearing

temperature. These gene networks were extensive and contained 2470 genes with

4092 edges at 12°C and 2468 genes with 4147 edges at 18°C (Figure S7A-B) with a

considerable number of overlapping genes (1945) and edges (1332) across rearing

temperatures (78% overlap in genes at 12°C and 18°C). The gene networks

associated with divergence between the geothermal ecotype and hybrids were

slightly less extensive containing 1895 genes and 3113 edges at 12°C and 1986

genes and 3327 edges at 18°C. Overlap was similar as to ambient ecotypes with

1480 genes and 973 edges that were concordant between rearing temperatures

(78% overlap at 12°C and 74% overlap at 18°C) and associated with divergence

between geothermal ecotypes and hybrids. Across all divergence networks there

were 1292 genes and 497 edges in common, but a relatively higher proportion of

overlapping gene networks associated with geothermal-hybrid divergence (68.2% at

12°C and 65.1% at 18°C) than ambient-hybrid divergence (52.3% at 12°C and

50.3% at 18°C). This overlapping network was overall less connected, measured by

overlapping edges, than the networks associated with ambient-hybrid divergence

(12.1% overlap at 12°C and 12.0% overlap at 18°C) and geothermal-hybrid

divergence (16.% overlap at 12°C and 14.9% overlap at 18°C).

We further identified gene networks associated with transgressive inheritance

patterns and their biological function, specifically involving genes that were over- or

under-expressed in hybrids relative to ecotypes. The gene network associated with

transgressive expression in hybrids relative to the ambient ecotype had 1142 genes

with 3242 edges at 12°C but was plastic by increasing to 1683 genes with 4762

edges when raised under 18°C. The geothermal ecotype showed a less extensive

network associated with transgressive expression in hybrids with 765 genes with

2157 edges at 12°C but this increased substantially to 1320 genes with 3757 edges

at 18°C. For both ecotypes there was overlap with 12°C conditions providing a gene

network associated with transgressive expression having 612 genes with 674 edges

and GO terms relating to the electron transport chain, oxidative phosphorylation, the

cytoskeleton, the proteasome, and the spliceosome (Figure 6A; Table S7). The

overlapping transgressive gene network at 18°C contained 1180 genes with 1481

edges (Figure 6B) with 42 significant GO terms primarily corresponding to

mitochondrial function, energy production, and use (Table S7). Across both

temperatures, the transgressive expression network contained 331 genes, 136

edges, and 7 GO terms related to energy production (Figure 6C; Table S7).

Proportion of inheritance pattern of transgressive loci.

Allele-specific expression (ASE) was detected in a small number of SNPs,

indicating cis-regulatory mechanisms for four specific genes that overlapped across

comparisons. Overall, SNPs within mfn1b, emsy, eef1a1, and fahd1 were identified

as being under cis-regulatory control and associated with temperature-dependent

expression differences between ecotypes. Three SNPs located in mfn1b, emsy, and

eef1a1 were inherited from the ambient ecotype, whereas seven SNPs distributed

across all four genes were inherited from the geothermal ecotype. ASE identified differential regulation among these loci where SNPs associated with eef1a1 and fahd1 exhibited reduced expression, while alleles at the remaining loci showed increased expression. The magnitude of these expression changes was temperature dependent, with larger differences observed at 12°C compared to 18°C, indicating that cid-regulatory variants at multiple loci contribute to temperature-associated divergence.

Discussion

We investigated the genomic consequences of hybridization between locally adapted thermal ecotypes of Icelandic stickleback using a two temperature common garden experiment. First, we tested for evidence of expression divergence (including plasticity) by comparing geothermal and ambient ecotypes across temperatures. While we identified a low degree of gene expression divergence between ecotypes, many genes were plastic to thermal variation in ways unique to each ecotype. Second, we tested whether hybrid gene expression differed from ecotypes. Supporting this, hybrid gene expression was highly divergent from both ecotypes regardless of developmental temperature indicating strong heritable effects. The majority of divergent genes exhibited transgressive expression and suggested substantial disruptions occurring within gene networks, with increasing effects under the warmer condition. This suggested that hybridization between ecotypes incurs asymmetric effects across temperatures, with gene networks of transgressive expression primarily associated with energy use and recruitment (Barreto et al., 2015a; Moran et al., 2021; Schwartz et al., 2024). Further, warmer environmental conditions disrupted a greater number of functional pathways suggesting maladaptive phenotypes may be more likely to arise. Third, we tested if hybridization between locally adapted ecotypes disrupted co-adapted cis-trans regulatory networks, leading to misexpression of specific alleles of locally adapted genes. Counter to our hypothesis, we found a surprisingly low degree of transgressively expressed SNPs within genes which suggests a greater degree of trans- vs cis-regulatory changes in gene expression due to hybridization. However, genes associated with cis-regulatory changes were associated with appear to have effects on DNA damage (*emsy*), metabolism (*eef1a1*), and mitochondrial dysfunction (*mfn1b* and *fahd1*). Thus, even though a low degree of genomic divergence occurred

between ecotypes (Brachmann et al., 2025), the impacts of hybridization could

involve metabolic dysfunction that may incur severe environmentally dependent

fitness consequences.

Divergence and plasticity of thermal ecotypes

The low degree of gene expression divergence between ecotypes could be

due to the rapid pace of evolution found in this system. The population pair used

here was established only ~70 years ago by the output of geothermal heating

systems in nearby residential buildings. Despite the geothermal environment being

colonized recently, and low levels of genome-wide allelic differentiation from ambient

counterparts (Brachmann et al., 2025), the gene expression patterns disrupted by

hybridization were key for energy use and metabolism. Such principles could

underlie other examples of rapid evolutionary divergence (Hairston et al., 2005; Kopp

& Matuszewski, 2014; Messer et al., 2016), including other cases of fish adapting to

warm habitats (Dayan et al., 2019). Our findings suggest that despite low levels of

expression divergence reproductive isolation could be enhanced by plasticity,

facilitating and maintaining phenotypic divergence between thermal habitats.

While there were low levels of divergence in gene expression, plasticity was

divergent. Ecotypes, as well as hybrids, showed unique plastic responses to

temperature, with geothermal ecotypes and hybrids showing a greater number of

plastic genes relative to the ambient ecotype. As a whole the plastic responses of

the ambient ecotype to warmer conditions were associated with carbohydrate

metabolism, indicating plastic shifts in the energy supply needed in warming

environments (Madeira et al., 2017; Middleton et al., 2024; Moore et al., 2025). In

contrast, geothermal ecotypes appear to adapt to warming environments through the

tuning of alternative splicing regulatory mechanisms (Schlaen et al., 2015; Steward

et al., 2022; Haltenhof et al., 2024) and overall metabolic architecture to manage

chronic metabolic and oxidative challenges (Seebacher et al., 2010; Madeira et al.,

2016; Lemieux & Blier, 2022). Divergent plastic responses, indicative of adaptation

to warming, is thus facilitated through divergence in metabolic strategies. In contrast,

hybrids had relatively biologically broad plastic responses indicating a release of

unrefined cryptic genetic variation (Gibson & Dworkin, 2004; Vallejo-Marín &

Hiscock, 2016; Mikkelsen & Irwin, 2021) that may be maladaptive in both

environments. The divergence in the plasticity of metabolic processes corresponds

too our previous findings of ecotype differences in metabolic rate (Pilakouta et al.,

2020), suggesting a phenotypic outcome inn line with our molecular responses.

Thus, hybridization could lead to maladaptive plasticity in metabolic rate

(Ghalamboret al., 2007; Chevin & Hoffmann, 2017; Campbell-Staton et al., 2021;

Eriksson et al., 2023).

While there were ecotype specific plastic responses to thermal variation, we

also detected shared plastic responses across ecotypes. For example, in the liver

ptptn6 showed a shared plastic response to thermal variation. *Ptpn6* is a known

regulator of the immune system through the *shp1* protein (Kanwal et al., 2013) and

has been implicated in regulating insulin resistance and lipid metabolism (Marín-Juez

et al., 2014; Xu et al., 2014, 2014; Kumar et al., 2023). In zebrafish (*Dania rerio*),

ptpn6/shp1 have previously been shown to regulate insulin resistance (Marín-Juez et

al., 2014). While shared, the ambient ecotype showed greater down-regulation of

ptpn6 in response to warmer temperatures than the geothermal ecotype or hybrids,

suggesting that the ambient ecotype may exhibit reduced insulin resistance or

immune function under warmer developmental environments. In the brain, genes

showing common plastic responses (the same gene and direction of plastic

response) comprised 62% of the total amount of plastic genes in the ambient

ecotype compared to only 16% and 20% in the geothermal ecotype and hybrids. The

high proportion of common plastic responses in the ambient ecotype indicated

conserved and possible ancestral plastic responses (Wund et al., 2008; Wood et al.,

2023; Coates et al., 2025), while the low proportion of common plastic responses in

the geothermal ecotype suggested a more derived and ecotype-specific responses

due to warming environments. The hybrids did not appear to combine parental

plastic responses which may indicate potential regulatory incompatibilities between

ecotypes (Go & Civetta, 2020; Lemmon & Kirkpatrick, 2006; Vallejo-Marín &

Hiscock, 2016).

Effects of hybridization on gene expression

548 Selection against hybridization can contribute to the formation and

549 maintenance of reproductive isolation between diverging populations (Butlin et al.,

550 2014; Kulmuni et al., 2020). Divergent populations are expected to be locally

adapted and hybrids inheriting intermediate or transgressive phenotypes are likely to face fitness consequences (Chhina et al., 2022; Gow et al., 2007; Hartman et al., 2013; Rice & McQuillan, 2018; Thompson et al., 2021). As ecotype specific gene expression is expected to be adapted to their native habitats (Ayroles et al., 2009; Brauer et al., 2017), uncoordinated gene expression or misexpression in hybrids is likely to be maladaptive in the wild. Hybrid gene expression differed greatly from both geothermal and ambient ecotypes at both rearing temperatures and predominantly showed transgressive patterns of inheritance. Hybrid mis-expression has been found to accompany severe (Barreto et al., 2015b; Ellison & Burton, 2008; Ortíz-Barrientos et al., 2007) or unpredictable (Brice et al., 2021; Díaz et al., 2023) fitness consequences (Rogers & Bernatchez, 2006; Rogers et al., 2007). Here, the disruption of gene co-expression networks was enhanced by the warmer temperature, suggesting thermally-dependent regulatory incompatibilities between ecotypes. Similar findings for hybrids has been shown in Atlantic silversides (*Menidia menidia*) (Jacobs et al., 2024), where a high proportion of genes were mis-expressed in relative to parental ecotypes. In Icelandic stickleback, transgressive mis-expression appeared to disrupt metabolic processes for energy use and production, consistent with hybrid metabolic dysfunction (Barreto et al., 2015). Our data indicate that the breakdown of metabolic processes in hybrids could be underlain by mitochondrial dysfunction, representing a severe physiological consequence (Barreto et al., 2015). Finally, the consequences of hybridization were enhanced in the warmer condition, further supporting the role of thermal plasticity in promoting reproductive isolation, although asymmetrically between thermal habitats.

Regulatory mechanisms and modes of inheritance underlying divergence

We found little evidence that additive variation contributed to divergence between geothermal and ambient sticklebacks. While transcriptional variation has been shown to be largely additive (Gilad et al., 2008; Kim & Gibson, 2010), our findings indicated less than 1% of such variation was additive across temperatures. While this would classically be interpreted as suggesting little potential for evolution (Falconer & Mackay, 1996), widespread non-additive gene expression inheritance has previously been found in other vertebrates (Debes et al., 2012), including stickleback (Leder et al., 2014). Hybridization transmitted a high degree of non-additive transcriptional variation indicating complex interactions due to dominance,

epistasis, and transgressive variation (Yazdi et al., 2022; Jacobs et al., 2024; Tsouris
et al., 2024). High levels of non-additive transcriptional variation is known to occur in
complex phenotypic traits, such as metabolic phenotypes (Zhong et al., 2019; Zhang
et al., 2025). However, a limitation of this study is that hybrid crosses were
generated from a single population of divergent ecotypes with no reciprocal crosses.
Future work using a fully reciprocal cross with multiple divergent populations would
allow for a more revealing examination of additive and non-additive transcriptional
variation.

Transgressive expression of genes in hybrids, relative to either parental
ecotype, was the dominant pattern of gene expression at both rearing temperatures,
suggesting substantial disruption of regulatory mechanisms. Generally, cis-
regulatory mechanisms are expected to underlie evolutionary divergence in gene
expression more often than trans-regulatory changes (Signor & Nuzhdin, 2018; Verta
& Jones, 2019; Wittkopp et al., 2004; Zhong et al., 2019). However, we identified a
low degree of cis-regulation and a high degree of trans-regulation of transgressively
expressed genes associated with metabolic divergence. Trans-regulation has
previously been shown to govern non-additive transcriptional variation (Tsouris et al.,
2024) and may play a large role temperature dependent regulation of metabolic
phenotypes.

Hybridization between locally adapted and rapidly evolving thermal ecotypes
appears to drive genomic incompatibilities and maladaptive phenotypic variation,
particularly under elevated temperatures. These findings have important implications
for climate change biology. As global temperatures rise, thermal gradients are
expected to shift, potentially increasing contact between locally adapted populations
that were previously segregated by microhabitat differences. Hybridization is often
considered a mechanism for evolutionary rescue and increased adaptive potential
under environmental change (Kelly & Phillips, 2016; Pfennig et al., 2016; Kulmuni et
al., 2023; Sexton et al., 2024). In contrast, our results demonstrated that even small
amounts of genomic divergence can generate temperature-dependent
incompatibilities that reduce fitness. In this system, warmer environments intensified
regulatory disruption, suggesting that climate warming may not only impose direct
physiological stress but also destabilize gene regulatory networks shaped by local
adaptation. These findings illustrate that population mixing or assisted gene flow may

dampen resilience to warming environments. Instead, the outcome of hybridization may critically depend on the degree of regulatory co-adaptation and the environmental context in which hybrids develop. As anthropogenic warming continues to alter thermal landscapes, understanding when hybridization facilitates adaptation versus when it uncovers cryptic incompatibilities will be essential for predicting evolutionary responses and guiding conservation strategies.

References

Abbott, R., Albach, D., Ansell, S., Arntzen, J. W., Baird, S. J. E., Bierne, N., Boughman, J., Brelsford, A., Buerkle, C. A., Buggs, R., Butlin, R. K., Dieckmann, U., Eroukhmanoff, F., Grill, A., Cahan, S. H., Hermansen, J. S., Hewitt, G., Hudson, A. G., Jiggins, C., … Zinner, D. (2013). Hybridization and speciation. *Journal of Evolutionary Biology*, *26*(2). https://doi.org/10.1111/j.1420-9101.2012.02599.x

Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq-A Python framework to work with high-throughput sequencing data. *Bioinformatics*, *31*(2), 166–169. https://doi.org/10.1093/bioinformatics/btu638

Andrews, S. (2010). *FastQC: A Quality Control Tool for High Throughput Sequence Data.*

Ansari, M. A., Fatima, Z., & Hameed, S. (2016). Antifungal Action of Methylene Blue Involves Mitochondrial Dysfunction and Disruption of Redox and Membrane Homeostasis in C. albicans. *The Open Microbiology Journal*, *10*, 12–22. https://doi.org/10.2174/1874285801610010012

Ayroles, J. F., Carbone, M. A., Stone, E. A., Jordan, K. W., Lyman, R. F., Magwire, M. M., Rollmann, S. M., Duncan, L. H., Lawrence, F., Anholt, R. R. H., & Mackay, T. F. C. (2009). Systems genetics of complex traits in Drosophila

melanogaster. *Nature Genetics*, *41*(3), 299–307.

https://doi.org/10.1038/ng.332

Barreto, F. S., Pereira, R. J., & Burton, R. S. (2015a). Hybrid Dysfunction and Physiological Compensation in Gene Expression. *Molecular Biology and Evolution*, *32*(3), 613–622. https://doi.org/10.1093/molbev/msu321

Barreto, F. S., Pereira, R. J., & Burton, R. S. (2015b). Hybrid dysfunction and physiological compensation in gene expression. *Molecular Biology and Evolution*, *32*(3), 613–622. https://doi.org/10.1093/molbev/msu321

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. In *Journal of the Royal Statistical Society. Series B (Methodological)* (Vol. 57, Issue 1, pp. 289–300).

Brachmann, M. K., Costa, A. P. B., Robertson, S., Donoghue, K., Pilakouta, N., Whitehead, M., Liu, X., Kristjansson, B., Skulason, S., Selman, C., & Parsons, K. (2025). *Parallel adaptation to geothermally-warmed habitats due to common structural variation and functional developmental pathways* (p. 2025.02.17.638600). bioRxiv. https://doi.org/10.1101/2025.02.17.638600

Brauer, C. J., Unmack, P. J., & Beheregaray, L. B. (2017). Comparative ecological transcriptomics and the contribution of gene expression to the evolutionary potential of a threatened fish. *Molecular Ecology*, *26*(24), 6841–6856. https://doi.org/10.1111/mec.14432

Brice, C., Zhang, Z., Bendixsen, D., & Stelkens, R. (2021). Hybridization Outcomes Have Strong Genomic and Environmental Contingencies. *The American Naturalist*, *198*(3), E53–E67. https://doi.org/10.1086/715356

Brouard, J. S., Schenkel, F., Marete, A., & Bissonnette, N. (2019). The GATK joint
genotyping workflow is appropriate for calling variants in RNA-seq
experiments. *Journal of Animal Science and Biotechnology, 10*(1), 1-6.
https://doi.org/10.1186/s40104-019-0359-0
Butlin, R. K., Saura, M., Charrier, G., Jackson, B., André, C., Caballero, A., Coyne,
J. a., Galindo, J., Grahame, J. W., Hollander, J., Kemppainen, P., Martínez-
Fernández, M., Panova, M., Quesada, H., Johannesson, K., & Rolán-Alvarez,
E. (2014). Parallel evolution of local adaptation and reproductive isolation
in the face of gene flow. *Evolution, 68*(4), 935-949.
https://doi.org/10.1111/evo.12329
Butlin, R. K., & Smadja, C. M. (2018). Coupling, Reinforcement, and Speciation.
The American Naturalist, 191(2), 155-172. https://doi.org/10.1086/695136
Caetano-Anolles. (2023a). *Hard-filtering germline short variants*. GATK Broad
Institute. https://gatk.broadinstitute.org/hc/en-us/articles/360035890471-
Hard-filtering-germline-short-variants
Caetano-Anolles, D. (2023b). *RNAseq short variant discovery (SNPs + Indels)*. GATK
Broad Institute. https://gatk.broadinstitute.org/hc/en-
us/articles/360035531192-RNAseq-short-variant-discovery-SNPs-Indels-
Campbell, C. S., Adams, C. E. ., Bean, C. W., & Parsons, K. J. (2017). Conservation
Evo-Devo: Preserving Biodiversity by Understanding Its Origins. *Trends in
Ecology and Evolution, 32*(10), 746-759.
https://doi.org/10.1016/j.tree.2017.07.002
Campbell-Staton, S. C., Velotta, J. P., & Winchell, K. M. (2021). Selection on
adaptive and maladaptive gene expression plasticity during thermal

adaptation to urban heat islands. *Nature Communications, 12*(1), 6195. https://doi.org/10.1038/s41467-021-26334-4

Carroll, S. P., Marler, M., Winchell, R., & Dingle, H. (2003). Evolution of cryptic flight morph and life history differences during host race radiation in the soapberry bug, Jadera haematoloma Herrich-Schaeffer (Hemiptera: Rhopalidae). *Annals of the Entomological Society of America, 96*(2), 135–143. https://doi.org/10.1603/0013-8746(2003)096%255B0135:EOCFMA%255D2.0.CO;2

Cenzer, M. L. (2017). Maladaptive plasticity masks the effects of natural selection in the red-shouldered soapberry bug. *American Naturalist, 190*(4), 521–533. https://doi.org/10.1086/693456

Chen, Y., Lun, A. T. L., & Smyth, G. K. (2016). From reads to genes to pathways: Differential expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline. *F1000Research, 5*, 1–51. https://doi.org/10.12688/F1000RESEARCH.8987.2

Chevin, L.-M., & Hoffmann, A. A. (2017). Evolution of phenotypic plasticity in extreme environments. *Philosophical Transactions of the Royal Society B: Biological Sciences, 372*(1723), 20160138. https://doi.org/10.1098/rstb.2016.0138

Chhina, A. K., Thompson, K. A., & Schluter, D. (2022). Adaptive divergence and the evolution of hybrid trait mismatch in threespine stickleback. *Evolution Letters, 6*(1), 34–45. https://doi.org/10.1002/evl3.264

Chunco, A. J. (2014). Hybridization in a warmer world. *Ecology and Evolution, 4*(10), 2019–2031. https://doi.org/10.1002/ece3.1052

Coates, S. E. R., Comeault, A. A., Wood, D. P., Fay, M. F., Creer, S., Osborne, O.

G., Dunning, L. T., & Papadopulos, A. S. T. (2025). Plastic responses to past

environments shape adaptation to novel selection pressures. *Proceedings of*

the National Academy of Sciences, 122(5), e2409541122.

https://doi.org/10.1073/pnas.2409541122

Crozier, L. G., & Hutchings, J. A. (2014). Plastic and evolutionary responses to

climate change in fish. *Evolutionary Applications, 7*(1), 68–87.

https://doi.org/10.1111/eva.12135

Daco, L., Colling, G., & Matthies, D. (2021). Altitude and latitude have different

effects on population characteristics of the widespread plant Anthyllis

vulneraria. *Oecologia, 197*(2), 537–549. https://doi.org/10.1007/s00442-

021-05030-6

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O.,

Whitwham, A., Keane, T., McCarthy, S. A., & Davies, R. M. (2021). Twelve

years of SAMtools and BCFtools. *GigaScience, 10*(2), 1–4.

https://doi.org/10.1093/gigascience/giab008

Dayan, D. I., Du, X., Baris, T. Z., Wagner, D. N., Crawford, D. L., & Oleksiak, M. F.

(2019). Population genomics of rapid evolution in natural populations:

Polygenic selection in response to power station thermal effluents 06

Biological Sciences 0604 Genetics. *BMC Evolutionary Biology, 19*(1), 1–20.

https://doi.org/10.1186/s12862-019-1392-5

Debes, P. V., Normandeau, E., Fraser, D. J., Bernatchez, L., & Hutchings, J. A.

(2012). Differences in transcription levels among wild, domesticated, and

hybrid Atlantic salmon (Salmo salar) from two environments. *Molecular*

Ecology, *21*(11), 2574–2587. https://doi.org/10.1111/j.1365-

294X.2012.05567.x

Díaz, F., Wolf, J., & de Brito, R. A. (2023). Transcriptional Misexpression in Hybrids

between Species Linked by Gene Flow Is Associated With Patterns of

Sequence Divergence. *Genome Biology and Evolution*, *15*(5), evad071.

https://doi.org/10.1093/gbe/evad071

Duffy, K., Gouhier, T. C., & Ganguly, A. R. (2022). Climate-mediated shifts in

temperature fluctuations promote extinction risk. *Nature Climate Change*,

12(11), 1037–1044. https://doi.org/10.1038/s41558-022-01490-7

Durinck, S., Spellman, P. T., Birney, E., & Huber, W. (2009). Mapping identifiers

for the integration of genomic datasets with the R/ Bioconductor package

biomaRt. *Nature Protocols*, *4*(8), 1184–1191.

https://doi.org/10.1038/nprot.2009.97

Ellison, C. K., & Burton, R. S. (2008). Genotype-dependent variation of

mitochondrial transcriptional profiles in interpopulation hybrids.

Proceedings of the National Academy of Sciences of the United States of

America, *105*(41), 15831–15836. https://doi.org/10.1073/pnas.0804253105

Eriksson, M., Kinnby, A., De Wit, P., & Rafajlović, M. (2023). Adaptive,

maladaptive, neutral, or absent plasticity: Hidden caveats of reaction

norms. *Evolutionary Applications*, *16*(2), 486–503.

https://doi.org/10.1111/eva.13482

Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize

analysis results for multiple tools and samples in a single report.

Bioinformatics.

Falconer, D. S., & Mackay, T. F. C. (1996). *Introduction to Quantitative Genetics*

(A. W. Longman, Ed.; 4th ed.). Longman Scientific & Technical.

https://doi.org/10.1002/9781118313718.ch4

Fan, J., Hu, J., Xue, C., Zhang, H., Susztak, K., Reilly, M. P., Reilly, M. P., Xiao,

R., & Li, M. (2020). ASEP: Gene-based detection of allele-specific expression

across individuals in a population by RNA sequencing. *PLoS Genetics*, *16*(5),

1-23. https://doi.org/10.1371/journal.pgen.1008786

Filteau, M., Pavey, S. A., St-Cyr, J., & Bernatchez, L. (2013). Gene coexpression

networks reveal key drivers of phenotypic divergence in lake whitefish.

Molecular Biology and Evolution, *30*(6), 1384-1396.

https://doi.org/10.1093/molbev/mst053

Fry, F. E. J. (1967). Responses of vertebrate poikilotherms to temperature. *In:*

Rose AH, Editor. Thermobiology., 375-409.

Ghalambor, C. K., McKAY, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive

versus non-adaptive phenotypic plasticity and the potential for

contemporary adaptation in new environments. *Functional Ecology*, *21*(3),

394-407. https://doi.org/10.1111/j.1365-2435.2007.01283.x

Gibson, G., & Dworkin, I. (2004). Uncovering cryptic genetic variation. *Nature*

Reviews. Genetics, *5*(9), 681-690. https://doi.org/10.1038/nrg1426

Gilad, Y., Rifkin, S. A., & Pritchard, J. K. (2008). Revealing the architecture of

gene regulation: The promise of eQTL studies. *Trends in Genetics*, *24*(8),

408-415. https://doi.org/10.1016/j.tig.2008.06.001

Go, A. C., & Civetta, A. (2020). Hybrid Incompatibilities and Transgressive Gene

Expression Between Two Closely Related Subspecies of Drosophila. *Frontiers*

in Genetics, *11*. https://doi.org/10.3389/fgene.2020.599292

Gow, J. L., Peichel, C. L., & Taylor, E. B. (2007). Ecological selection against

hybrids in natural populations of sympatric threespine sticklebacks. *Journal*

of Evolutionary Biology, 20(6), 2173–2180. https://doi.org/10.1111/j.1420-

9101.2007.01427.x

Greenspoon, P. B., & Spencer, H. G. (2021). Avoiding extinction under nonlinear

environmental change: Models of evolutionary rescue with plasticity.

Biology Letters, 17(12), 20210459. https://doi.org/10.1098/rsbl.2021.0459

Haesemeyer, M. (2020). Thermoregulation in fish. *Molecular and Cellular*

Endocrinology, 518, 110986. https://doi.org/10.1016/j.mce.2020.110986

Hairston, N. G., Ellner, S. P., Geber, M. A., Yoshida, T., & Fox, J. A. (2005). Rapid

evolution and the convergence of ecological and evolutionary time. *Ecology*

Letters, 8(10), 1114–1127. https://doi.org/10.1111/j.1461-

0248.2005.00812.x

Haltenhof, T., Preußner, M., & Heyd, F. (2024). Thermoregulated transcriptomics:

The molecular basis and biological significance of temperature-dependent

alternative splicing. *Biochemical Journal, 481*(15), 999–1013.

https://doi.org/10.1042/BCJ20230410

Hartman, Y., Uwimana, B., Hooftman, D. A. P., Schranz, M. E., van de Wiel, C. C.

M., Smulders, M. J. M., Visser, R. G. F., & van Tienderen, P. H. (2013).

Genomic and environmental selection patterns in two distinct lettuce crop-

wild hybrid crosses. *Evolutionary Applications, 6*(4), 569–584.

https://doi.org/10.1111/eva.12043

Institute, G. B. (2023). *Picard.*

Jacobs, A., Velotta, J. P., Tigano, A., Wilder, A. P., Baumann, H., & Therkildsen,

N. O. (2024). Temperature-dependent gene regulatory divergence underlies

local adaptation with gene flow in the Atlantic silverside. *Evolution, 78*(6),

1133-1149. https://doi.org/10.1093/evolut/qpae049

Joshi, N., & Fass, J. (2011). *Sickle: A sliding-window, adaptive, quality-based*

trimming tool for FastQ files.

Kanwal, Z., Zakrzewska, A., den Hertog, J., Spaink, H. P., Schaaf, M. J. M., &

Meijer, A. H. (2013). Deficiency in hematopoietic phosphatase ptpn6/Shp1

hyperactivates the innate immune system and impairs control of bacterial

infections in zebrafish embryos. *Journal of Immunology (Baltimore, Md.:*

1950), 190(4), 1631-1645. https://doi.org/10.4049/jimmunol.1200551

Kelly, E., & Phillips, B. L. (2016). Targeted gene flow for conservation.

Conservation Biology, 30(2), 259-267. https://doi.org/10.1111/cobi.12623

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based

genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature*

Biotechnology, 37(8), 907-915. https://doi.org/10.1038/s41587-019-0201-4

Kim, J., & Gibson, G. (2010). Insights from GWAS into the quantitative genetics of

transcription in humans. *Genetics Research, 92*(5-6), 361-369.

https://doi.org/10.1017/S001667231000056X

Kinsella, R. J., Kähäri, A., Haider, S., Zamora, J., Proctor, G., Spudich, G.,

Almeida-King, J., Staines, D., Derwent, P., Kerhornou, A., Kersey, P., &

Flicek, P. (2011). Ensembl BioMarts: A hub for data retrieval across

taxonomic space. *Database, 2011*, 1-9.

https://doi.org/10.1093/database/bar030

Kopp, M., & Matuszewski, S. (2014). Rapid evolution of quantitative traits:

Theoretical perspectives. *Evolutionary Applications, 7*(1), 169-191.

https://doi.org/10.1111/eva.12127

Kulmuni, J., Butlin, R. K., Lucek, K., Savolainen, V., & Westram, A. M. (2020).

838    Towards the completion of speciation: The evolution of reproductive

839    isolation beyond the first barriers. *Philosophical Transactions of the Royal*

840    *Society of London. Series B, Biological Sciences, 375*(1806), 20190528.

841    https://doi.org/10.1098/rstb.2019.0528

842 Kulmuni, J., Wiley, B., & Otto, S. P. (2023). On the fast track: Hybrids adapt more

843    rapidly than parental populations in a novel environment. *Evolution Letters,*

844    *8*(1), 128–136. https://doi.org/10.1093/evlett/qrad002

845 Kumar, A., Schwab, M., Laborit Labrada, B., Silveira, M. A. D., Goudreault, M.,

846    Fournier, É., Bellmann, K., Beauchemin, N., Gingras, A.-C., Bilodeau, S.,

847    Laplante, M., & Marette, A. (2023). SHP-1 phosphatase acts as a coactivator

848    of *PCK1* transcription to control gluconeogenesis. *Journal of Biological*

849    *Chemistry, 299*(9), 105164. https://doi.org/10.1016/j.jbc.2023.105164

850 Law, C. W., Alhamdoosh, M., Su, S., Dong, X., Tian, L., Smyth, G. K., & Ritchie, M.

851    E. (2018). RNA-seq analysis is easy as 1-2-3 with limma, Glimma and edgeR.

852    *F1000Research, 5*(1), 1–29.

853 Law, C. W., Chen, Y., Shi, W., & Smyth, G. K. (2014). Voom: Precision weights

854    unlock linear model analysis tools for RNA-seq read counts. *Genome Biology,*

855    *15*(2), 1–17. https://doi.org/10.1186/gb-2014-15-2-r29

856 Law, C. W., Zeglinski, K., Dong, X., Alhamdoosh, M., Smyth, G. K., Ritchie, M. E.,

857    Marini, F., & Drnevich, J. (2020). *A guide to creating design matrices for*

858    *gene expression experiments*. 1–45.

859 Leder, E. H., McCairns, R. J. S. S., Leinonen, T., Cano, J. M., Viitaniemi, H. M.,

860    Nikinmaa, M., Primmer, C. R., Merilä, J., Cano, M., Viitaniemi, H. M.,

861    Leder, E. H., McCairns, R. J. S. S., Leinonen, T., Cano, J. M., Viitaniemi, H.

M., Nikinmaa, M., Primmer, C. R., Merilä, J., Merila, J., … Leinonen, T.

(2014). The evolution and adaptive potential of transcriptional variation in

sticklebacks—Signatures of selection and widespread heritability. *Molecular*

Biology and Evolution, 32(3), 674–689.

https://doi.org/10.1093/molbev/msu328

Lemieux, H., & Blier, P. U. (2022). Exploring Thermal Sensitivities and Adaptations

of Oxidative Phosphorylation Pathways. *Metabolites, 12*(4).

https://doi.org/10.3390/metabo12040360

Lemmon, A. R., & Kirkpatrick, M. (2006). Reinforcement and the Genetics of

Hybrid Incompatibilities. *Genetics, 173*(2), 1145–1155.

https://doi.org/10.1534/genetics.105.048199

Lemmon, E. M., & Juenger, T. E. (2017). Geographic variation in hybridization

across a reinforcement contact zone of chorus frogs (Pseudacris). *Ecology*

and Evolution, 7(22), 9485–9502. https://doi.org/10.1002/ece3.3443

Lemos, B., Araripe, L. O., Fontanillas, P., & Hartl, D. L. (2008). Dominance and the

evolutionary accumulation of cis- and trans-effects on gene expression.

Proceedings of the National Academy of Sciences of the United States of

America, 105(38), 14471–14476. https://doi.org/10.1073/pnas.0805160105

Lenormand, T. (2012). From Local Adaptation to Speciation: Specialization and

Reinforcement. *International Journal of Ecology, 2012*(1), 508458.

https://doi.org/10.1155/2012/508458

Li, C., Deans, N. C., & Buell, C. R. (2023). "Simple Tidy GeneCoEx": A gene co-

expression analysis workflow powered by tidyverse and graph-based

clustering in R. *The Plant Genome, 16*(2), e20323.

https://doi.org/10.1002/tpg2.20323

Lun, A. T. L., Chen, Y., & Smyth, G. K. (2016). It's DE-licious: A recipe for

differential expression analyses of RNA-seq experiments using quasi-

likelihood methods in edgeR. *Methods in Molecular Biology*, *1418*, 391–416.

https://doi.org/10.1007/978-1-4939-3578-9_19

Madeira, D., Araújo, J. E., Vitorino, R., Costa, P. M., Capelo, J. L., Vinagre, C., &

Diniz, M. S. (2017). Molecular Plasticity under Ocean Warming: Proteomics

and Fitness Data Provides Clues for a Better Understanding of the Thermal

Tolerance in Fish. *Frontiers in Physiology*, *8*.

https://doi.org/10.3389/fphys.2017.00825

Madeira, D., Vinagre, C., & Diniz, M. S. (2016). Are fish in hot water? Effects of

warming on oxidative stress metabolism in the commercial species *Sparus*

aurata. *Ecological Indicators*, *63*, 324–331.

https://doi.org/10.1016/j.ecolind.2015.12.008

Marín-Juez, R., Jong-Raadsen, S., Yang, S., & Spaink, H. P. (2014).

Hyperinsulinemia induces insulin resistance and immune suppression via

Ptpn6/Shp1 in zebrafish. *Journal of Endocrinology*, *222*(2), 229–241.

https://doi.org/10.1530/JOE-14-0178

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.Journal*, *17*, 10–12.

Matthew A. Wund, John A. Baker, Brendan Clancy, Justin L. Golub, & Susan A.

Foster. (2008). A Test of the "Flexible Stem" Model of Evolution: Ancestral

Plasticity, Genetic Accommodation, and Morphological Divergence in the

Threespine Stickleback Radiation. *The American Naturalist*, *172*(4), 449–

462. https://doi.org/10.1086/590966

McGirr, J. A., & Martin, C. H. (2021). Few Fixed Variants between Trophic

Specialist Pupfish Species Reveal Candidate Cis-Regulatory Alleles

Underlying Rapid Craniofacial Divergence. *Molecular Biology and Evolution*,

38(2), 405–423. https://doi.org/10.1093/molbev/msaa218

Merico, D., Isserlin, R., Stueker, O., Emili, A., & Bader, G. D. (2010). Enrichment

Map: A Network-Based Method for Gene-Set Enrichment Visualization and

Interpretation. *PLOS ONE*, *5*(11), e13984.

https://doi.org/10.1371/journal.pone.0013984

Messer, P. W., Ellner, S. P., & Hairston, N. G. (2016). Can Population Genetics

Adapt to Rapid Evolution? *Trends in Genetics*, *32*(7), 408–418.

https://doi.org/10.1016/j.tig.2016.04.005

Middleton, E. K., Gilbert, M. J. H., Landry, T., Lamarre, S. G., & Speers-Roesch, B.

(2024). Environmental variation associated with overwintering elicits

marked metabolic plasticity in a temperate salmonid, Salvelinus fontinalis.

Journal of Experimental Biology, *227*(3), jeb246743.

https://doi.org/10.1242/jeb.246743

Mikkelsen, E. K., & Irwin, D. (2021). Ongoing production of low-fitness hybrids

limits range overlap between divergent cryptic species. *Molecular Ecology*,

30(16), 4090–4102. https://doi.org/10.1111/mec.16015

Millet, A., Kristjánsson, B. K., Einarsson, Á., & Räsänen, K. (2013). Spatial

phenotypic and genetic structure of threespine stickleback (Gasterosteus

aculeatus) in a heterogeneous natural system, Lake Mỳvatn, Iceland. *Ecology*

and Evolution, *3*(10), 3219–3232. https://doi.org/10.1002/ece3.712

Moore, B., McMahon, S., Izumiyama, M., Ryu, T., & Ravasi, T. (2025). Ocean

warming drives tissue-wide metabolic reprogramming in a fish. *iScience*,

28(9). https://doi.org/10.1016/j.isci.2025.113395

Moran, B. M., Payne, C., Langdon, Q., Powell, D. L., Brandvain, Y., & Schumer, M.

(2021). The genomic consequences of hybridization. *eLife*, *10*, e69016.

https://doi.org/10.7554/eLife.69016

Nunez, S., Arets, E., Alkemade, R., Verwer, C., & Leemans, R. (2019). Assessing

the impacts of climate change on biodiversity: Is below 2 °C enough?

Climatic Change, *154*(3), 351–365. https://doi.org/10.1007/s10584-019-

02420-x

Oldham, M. C., Horvath, S., & Geschwind, D. H. (2006). Conservation and

evolution of gene coexpression networks in human and chimpanzee brains.

Proceedings of the National Academy of Sciences of the United States of

America, *103*(47), 17973–17978. https://doi.org/10.1073/pnas.0605938103

Ono, K., Fong, D., Gao, C., Churas, C., Pillich, R., Lenkiewicz, J., Pratt, D., Pico,

A. R., Hanspers, K., Xin, Y., Morris, J., Kucera, M., Franz, M., Lopes, C.,

Bader, G., Ideker, T., & Chen, J. (2025). Cytoscape Web: Bringing network

biology to the browser. *Nucleic Acids Research*, *53*(W1), W203–W212.

https://doi.org/10.1093/nar/gkaf365

Ortíz-Barrientos, D., Counterman, B. A., & Noor, M. A. F. (2007). Gene expression

divergence and the origin of hybrid dysfunctions. *Genetica*, *129*(1), 71–81.

https://doi.org/10.1007/s10709-006-0034-1

Osada, N., Miyagi, R., & Takahashi, A. (2017). Cis- and Trans-regulatory Effects on

Gene Expression in a Natural Population of Drosophila melanogaster.

Genetics, *206*(4), 2139–2148. https://doi.org/10.1534/genetics.117.201459

Paaijmans, K. P., Heinig, R. L., Seliga, R. A., Blanford, J. I., Blanford, S., Murdock,

C. C., & Thomas, M. B. (2013). Temperature variation makes ectotherms

more sensitive to climate change. *Global Change Biology, 19*(8), 2373-2380.

https://doi.org/10.1111/gcb.12240

Parsons, K. J. (2021). Conservation biology meets evo-devo: How understanding

the emergence of variation can inform its management. *Evolution and*

Development, 23(4), 269-272. https://doi.org/10.1111/ede.12389

Peichel, C. L., McCann, S. R., Ross, J. A., Naftaly, A. F. S., Urton, J. R., Cech, J.

N., Grimwood, J., Schmutz, J., Myers, R. M., Kingsley, D. M., & White, M. A.

(2020). Assembly of the threespine stickleback y chromosome reveals

convergent signatures of sex chromosome evolution. *Genome Biology, 21*(1),

1-31. https://doi.org/10.1186/s13059-020-02097-x

Pfennig, K. S., Kelly, A. L., & Pierce, A. A. (2016). Hybridization as a facilitator of

species range expansion. *Proceedings of the Royal Society B: Biological*

Sciences, 283(1839), 20161329. https://doi.org/10.1098/rspb.2016.1329

Pierre, C. L. S., Macias-Velasco, J. F., Wayhart, J. P., Yin, L., Semenkovich, C. F.,

& Lawson, H. A. (2022). Genetic, epigenetic, and environmental

mechanisms govern allele-specific gene expression. *Genome Research,*

32(6), 1042-1057. https://doi.org/10.1101/gr.276193.121

Pigliucci, M. (2008). Opinion—Is evolvability evolvable? *Nat. Rev. Genet., 9*(1), 75-

82. https://doi.org/10.1038/nrg2278

Pilakouta, N., Humble, J. L., Hill, I. D. C., Arthur, J., Costa, A. P. B., Smith, B. A.,

Kristjansson, B. K., Skulason, S., Killen, S. S., Lindstrom, J., Metcalfe, N. B.,

& Parsons, K. J. (2023). Testing the predictability of morphological

evolution in contrasting thermal environments. *Evolution, 77*(1), 239–253.

https://doi.org/10.1093/evolut/qpac018

Pilakouta, N., Killen, S. S., Kristjánsson, B. K., Skúlason, S., Lindström, J.,

Metcalfe, N. B., & Parsons, K. J. (2020). Multigenerational exposure to

elevated temperatures leads to a reduction in standard metabolic rate in

the wild. *Functional Ecology, 34*(6), 1205–1214.

https://doi.org/10.1111/1365-2435.13538

R Core Team. (2022). *R: A language and environment for statistical computing. R*

Foundation for Statistical Computing.

REED, T. E., SCHINDLER, D. E., & WAPLES, R. S. (2011). Interacting Effects of

Phenotypic Plasticity and Evolution on Population Persistence in a Changing

Climate. *Conservation Biology, 25*(1), 56–63.

https://doi.org/10.1111/j.1523-1739.2010.01552.x

Ren, X., & Kuan, P. F. (2020). Negative binomial additive model for RNA-Seq data

analysis. *BMC Bioinformatics, 21*(1), 1–15. https://doi.org/10.1186/s12859-

020-3506-x

Rice, A. M., & McQuillan, M. A. (2018). Maladaptive learning and memory in

hybrids as a reproductive isolating barrier. *Proceedings of the Royal Society*

B: Biological Sciences, 285(1879). https://doi.org/10.1098/rspb.2018.0542

Riederer, J. M., Tiso, S., van Eldijk, T. J. B., & Weissing, F. J. (2022). Capturing

the facets of evolvability in a mechanistic framework. *Trends in Ecology and*

Evolution, 37(5), 430–439. https://doi.org/10.1016/j.tree.2022.01.004

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K.

(2015). Limma powers differential expression analyses for RNA-sequencing

and microarray studies. *Nucleic Acids Research, 43*(7), e47.

https://doi.org/10.1093/nar/gkv007

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2009). edgeR: A Bioconductor

package for differential expression analysis of digital gene expression data.

Bioinformatics, 26(1), 139–140.

https://doi.org/10.1093/bioinformatics/btp616

Robinson, M. D., & Oshlack, A. (2010). A scaling normalization method for

differential expression analysis of RNA-seq data. *Genome Biology, 11*(3), 1–

9.

Rogers, S. M., & Bernatchez, L. (2006). The genetic basis of intrinsic and extrinsic

post-zygotic reproductive isolation jointly promoting speciation in the lake

whitefish species complex (Coregonus clupeaformis). *Journal of*

Evolutionary Biology, 19(6), 1979–1994. https://doi.org/10.1111/j.1420-

9101.2006.01150.x

Rogers, S. M., Isabel, N., & Bernatchez, L. (2007). Linkage Maps of the dwarf and

Normal Lake Whitefish (Coregonus clupeaformis) Species Complex and Their

Hybrids Reveal the Genetic Architecture of Population Divergence. *Genetics,*

175(1), 375–398. https://doi.org/10.1534/genetics.106.061457

Rundle, H. D., & Nosil, P. (2005). Ecological speciation. *Ecology Letters, 8*(3), 336–

352. https://doi.org/10.1111/j.1461-0248.2004.00715.x

Runemark, A., Moore, E. C., & Larson, E. L. (2025). Hybridization and gene

expression: Beyond differentially expressed genes. *Molecular Ecology,*

34(15), e17303. https://doi.org/10.1111/mec.17303

Schlaen, R. G., Mancini, E., Sanchez, S. E., Perez-Santángelo, S., Rugnone, M. L.,

Simpson, C. G., Brown, J. W. S., Zhang, X., Chernomoretz, A., & Yanovsky,

M. J. (2015). The spliceosome assembly factor GEMIN2 attenuates the effects of temperature on alternative splicing and circadian rhythms. *Proceedings of the National Academy of Sciences*, *112*(30), 9382–9387. https://doi.org/10.1073/pnas.1504541112

Schwartz, L. C., González, V. L., Strong, E. E., Truebano, M., & Hilbish, T. J. (2024). Transgressive gene expression and expression plasticity under thermal stress in a stable hybrid zone. *Molecular Ecology*, *33*(9), e17333. https://doi.org/10.1111/mec.17333

Seebacher, F., Brand, M. D., Else, P. L., Guderley, H., Hulbert, A. J., & Moyes, C. D. (2010). Plasticity of Oxidative Metabolism in Variable Climates: Molecular Mechanisms. *Physiological and Biochemical Zoology*, *83*(5), 721–732. https://doi.org/10.1086/649964

Servedio, M. R. (2004). THE EVOLUTION OF PREMATING ISOLATION: LOCAL ADAPTATION AND NATURAL AND SEXUAL SELECTION AGAINST HYBRIDS. *Evolution*, *58*(5), 913–924. https://doi.org/10.1111/j.0014-3820.2004.tb00425.x

Sexton, J. P., Clemens, M., Bell, N., Hall, J., Fyfe, V., & Hoffmann, A. A. (2024). Patterns and effects of gene flow on adaptation across spatial scales: Implications for management. *Journal of Evolutionary Biology*, *37*(6), 732–745. https://doi.org/10.1093/jeb/voae064

She, H., Hao, Y., Song, G., Luo, X., Lei, F., Zhai, W., & Qu, Y. (2023). Gene expression plasticity followed by genetic change during colonization a high-elevation environment. *eLife*, *12*. https://doi.org/10.7554/eLife.86687.2

Sheldon, E. L., Martin, L. B., & Schrey, A. W. (2025). Integrating plasticity into conservation practice: Harnessing genetic estimates of epigenetic potential

to study phenotypic plasticity in wild populations. *Journal of Applied*

Ecology, *62*(4), 783–789. https://doi.org/10.1111/1365-2664.70019

Sheng, Q., Vickers, K., Zhao, S., Wang, J., Samuels, D. C., Koues, O., Shyr, Y., &

Guo, Y. (2017). Multi-perspective quality control of Illumina RNA sequencing

data analysis. *Briefings in Functional Genomics*, *16*(4), 194–204.

https://doi.org/10.1093/bfgp/elw035

Signor, S. A., & Nuzhdin, S. V. (2018). The Evolution of Gene Expression in cis and

trans. *Trends in Genetics*, *34*(7), 532–544.

https://doi.org/10.1016/j.tig.2018.03.007

Smith, B. A., Costa, A. P. B., Kristjánsson, B. K., & Parsons, K. J. (2024).

Experimental evidence for adaptive divergence in response to a warmed

habitat reveals roles for morphology, allometry and parasite resistance.

Ecology and Evolution, *14*(2), e10907. https://doi.org/10.1002/ece3.10907

Steward, R. A., de Jong, M. A., Oostra, V., & Wheat, C. W. (2022). Alternative

splicing in seasonal plasticity and the potential for adaptation to

environmental change. *Nature Communications*, *13*(1), 755.

https://doi.org/10.1038/s41467-022-28306-8

Thompson, K. A., Urquhart-Cronish, M., Whitney, K. D., Rieseberg, L. H., &

Schluter, D. (2021). Patterns, predictors, and consequences of dominance in

hybrids. *American Naturalist*, *197*(3), E72–E88.

https://doi.org/10.1086/712603

Timmons, J. A., Szkop, K. J., & Gallagher, I. J. (2015). Multiple sources of bias

confound functional enrichment analysis of global -omics data. *Genome*

Biology, *16*(1), 15–17. https://doi.org/10.1186/s13059-015-0761-7

Traag, V. A., Waltman, L., & van Eck, N. J. (2019). From Louvain to Leiden:

Guaranteeing well-connected communities. *Scientific Reports*, *9*(1), 5233.

https://doi.org/10.1038/s41598-019-41695-z

Tsouris, A., Brach, G., Schacherer, J., & Hou, J. (2024). Non-additive genetic

components contribute significantly to population-wide gene expression

variation. *Cell Genomics*, *4*(1), 100459.

https://doi.org/10.1016/j.xgen.2023.100459

Vallejo-Marín, M., & Hiscock, S. J. (2016). Hybridization and hybrid speciation

under global change. *New Phytologist*, *211*(4), 1170–1187.

https://doi.org/10.1111/nph.14004

Verta, J. P., & Jones, F. C. (2019). Predominance of cis-regulatory changes in

parallel expression divergence of sticklebacks. *eLife*, *8*, 1–30.

https://doi.org/10.7554/eLife.43785

Wang, H., Bruce, T. J., Su, B., Li, S., Dunham, R. A., & Wang, X. (2022).

Environment-Dependent Heterosis and Transgressive Gene Expression in

Reciprocal Hybrids between the Channel Catfish Ictalurus punctatus and the

Blue Catfish Ictalurus furcatus. *Biology*, *11*(1).

https://doi.org/10.3390/biology11010117

Wighard, S., Witte, H., & Sommer, R. J. (2025). Functional divergence of

conserved developmental plasticity genes between two distantly related

nematodes. *Scientific Reports*, *15*(1), 28518.

https://doi.org/10.1038/s41598-025-14207-5

Wittkopp, P. J., Haerum, B. K., & Clark, A. G. (2004). Evolutionary changes in cis

and trans gene regulation. *Nature*, *430*(6995), 85–88.

https://doi.org/10.1038/nature02698

Wittkopp, P. J., Haerum, B. K., & Clark, A. G. (2008). Regulatory changes

underlying expression differences within and between Drosophila species.

Nature Genetics, 40(3), 346–350. https://doi.org/10.1038/ng.77

Wittkopp, P. J., & Kalay, G. (2012). Cis-regulatory elements: Molecular

mechanisms and evolutionary processes underlying divergence. *Nature*

Reviews Genetics, 13(1), 59–69. https://doi.org/10.1038/nrg3095

Wood, D. P., Holmberg, J. A., Osborne, O. G., Helmstetter, A. J., Dunning, L. T.,

Ellison, A. R., Smith, R. J., Lighten, J., & Papadopulos, A. S. T. (2023).

Genetic assimilation of ancestral plasticity during parallel adaptation to zinc

contamination in Silene uniflora. *Nature Ecology & Evolution, 7*(3), 414–423.

https://doi.org/10.1038/s41559-022-01975-w

Xu, E., Forest, M.-P., Schwab, M., Avramoglu, R. K., St-Amand, E., Caron, A. Z.,

Bellmann, K., Shum, M., Voisin, G., Paquet, M., Montoudis, A., Lévy, E.,

Siminovitch, K. A., Neel, B. G., Beauchemin, N., & Marette, A. (2014).

Hepatocyte-specific Ptpn6 deletion promotes hepatic lipid accretion, but

reduces NAFLD in diet-induced obesity: Potential role of PPARγ. *Hepatology,*

59(5), 1803–1815. https://doi.org/10.1002/hep.26957

Yazdi, H. P., Ravinet, M., Rowe, M., Sætre, G. P., Guldvog, C., Eroukhmanoff, F.,

Marzal, A., Magallanes, S., & Runemark, A. (2022). Extensive transgressive

gene expression in testis but not ovary in the homoploid hybrid Italian

sparrow. *Molecular Ecology, 31*(15), 4067–4077.

https://doi.org/10.1111/mec.16572

Zhang, J., Zhang, W., Ding, C., Zhao, J., Su, X., Yuan, Z., Chu, Y., Huang, Q., &

Su, X. (2025). Non-Additive Gene Expression in Carbon and Nitrogen

Metabolism Drives Growth Heterosis in Populus deltoides. *Plant, Cell &*

Environment, *48*(5), 3529–3543. https://doi.org/10.1111/pce.15371

Zhong, H., Zhang, X., Xu, Q., Yan, J., Han, Z., Zheng, H., Xiao, J., Tang, Z., Wang,

F., Luo, Y., & Zhou, Y. (2019). Nonadditive and Asymmetric Allelic

Expression of Growth Hormone in Hybrid Tilapia. *Frontiers in Genetics*,

10(October), 1–11. https://doi.org/10.3389/fgene.2019.00961

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O.,

Benner, C., & Chanda, S. K. (2019). Metascape provides a biologist-oriented

resource for the analysis of systems-level datasets. *Nature Communications*,

10(1). https://doi.org/10.1038/s41467-019-09234-6

Tables and figures

The following is the table embedded in panel 4 of the figure:

Type	Rearing temperature	Brain	Liver
Ambient	12˚C	4 (1,1,3,3)	4 (1,1,3,3)
	18°C	3 (3,3,2)	4 (2,2,2,2)
Geothermal	12˚C	4 (3,1,3,1)	4 (3,1,3,1)
	18°C	3 (2,3,3)	3 (2,3,3)
Hybrid	12˚C	2 (4,4)	2 (4,4)
	18°C	2 (4,4)	2 (4,4)

Figure 1. Methodological approach for the creation of experimental hybrid and pure-strained crosses of geothermal and ambient Icelandic threespine stickleback. 1. Wild geothermal and ambient stickleback were collected from allopatric Saudárkrókur (SKR) populations in Iceland. 2. Hybrid and pure-strained geothermal and ambient crosses were made via invitro insemination. Eggs were striped from females and fertilized with sperm from the corresponding pure-strained male. Hybrid crosses were made by crossing ambient males with geothermal females. 3. Hybrid and pure-strained crosses were divided in half and raised under 12˚C and 18˚C rearing conditions until maturity.

Figure 2. Volcano plots showing differentially expressed genes between the geothermal and ambient pure-strained ecotypes under 12˚C (A) and 18˚C (B). The red dashed line represents the p-value cut-off of 0.05. Genes in blue are down-regulated and genes in orange are up-regulated.

Figure 3. Volcano plots representing up- and down-regulated genes corresponding to plastic responses between rearing temperatures for ambient (A), geothermal (B), and hybrid (C) ecotypes of Icelandic stickleback. A-C) all up- and down-regulated genes are shown as orange and blue, respectively. The dashed red line highlights the p-value cut-off of 0.05. D-F) Gene ontology results highlighting the gene ontology terms from unique plastic responses to 12˚C and 18˚C rearing conditions for the ambient (D), geothermal (F), and hybrid (G) ecotype. G) gene ontology terms associated with the 38 genes showing common plastic responses across all three ecotypes. The different colours in each GO plot represent the database the term belongs to.

Figure 4. Inheritance patterns of divergent genes in hybrids compared to pure-strain geothermal and ambient ecotypes. A) Principal component analysis of all differentially expressed genes. Dark and light blue corresponds to the ambient ecotype at 12°C and 18°C. Dark and light red corresponds to the geothermal ecotype at 12°C and 18°C. Green and yellow corresponds to the hybrid ecotype at 12°C and 18°C. B) Patterns of inheritance based on log fold change between hybrids relative to each pure-strained ecotype. Dominant inheritance (light and dark blue) patterns are between +/- 0.32 for each comparison. Additive inheritance (yellow) is defined as >0.32 in geothermal but <0.32 in the ambient ecotype or <0.32 in the geothermal ecotype but >0.32 in the ambient ecotype. Transgressive inheritance (purple) is defined as either < or > 0.32 in each of the pure-strained ecotypes. C) Patterns of inheritance in all differentially expressed genes at 12°C. D) Patterns of inheritance in all differentially expressed genes at 18°C.

Figure 5. Gene network analyses for transgressively inherited genes in hybrids relative to pure-strained ecotypes. A) Gene network for transgressively inherited genes at 12˚C and the significant gene ontology terms for the network. B) Gene network for transgressively inherited genes at 18˚C and the corresponding gene ontology terms for the network. C) Overlap between the 12˚C and 18˚C network of transgressively inherited genes and the associated gene ontology terms for the overlapping network.