Discovery and characterisation of OMVs produced by the bee gut microbiota

Rodney P. Eyles[1], Waldan K. Kwong[1,2]

[1] Gulbenkian Institute for Molecular Medicine, Rua da Quinta Grande 6, 2780-156 Oeiras, Portugal

[2] Faculty of Medicine, University of Lisbon, Av. Prof. Egas Moniz, 1649-028 Lisbon, Portugal

Correspondence: waldan.kwong@gimm.pt

Keywords: outer membrane vesicles, gut symbionts, bacterial secretion, RNA, *Apis mellfiera*

Subjects: biology, molecular biology, cellular biology

Abstract

Bacteria use diverse mechanisms to interact with each other and with eukaryotic hosts, thereby shaping microbiome composition and influencing host health. One of these mechanisms is the production of outer membrane vesicles (OMVs), nanoscale structures that bud off from bacterial cells into the extracellular space. OMVs can deliver bioactive cargoes, including enzymes, RNA and DNA, enabling functions such as cell-to-cell communication, nutrient acquisition and immunomodulation. However, the role of OMVs in beneficial host-associated microbiomes remains unclear. Here, we investigated OMV production in the gut bacteria of the western honey bee (*Apis mellifera*), which forms a highly conserved and stable microbial community. Using electron microscopy, fluorescence labelling, and nanoparticle tracking analysis, we detected OMV production in every gram-negative species of the normal bee microbiota that we investigated. Vesicles were observed in gut contents of wild and laboratory-inoculated bees, but absent in bees lacking a microbiota. OMVs contained nucleic acids, with more RNA than DNA. Bacterial strains varied in OMV properties, including abundance, size, and zeta potential. These findings indicate that OMVs are likely significant mediators of interbacterial and host-microbe interactions in the bee gut.

1. Introduction

Gut microbiomes are largely comprised of a core set of resident bacteria which form stable, structured communities that are highly adapted to the physical and chemical environment of the host. Gut microbes form symbiotic relationships with their host and contribute to its wellbeing, by increasing nutrient acquisition via breakdown of otherwise indigestible food material, strengthening and protecting the host gut lining, and boosting colonisation resistance from foreign or pathogenic bacteria [1]. Conversely, disruption to the gut microbiome (dysbiosis), is associated with a broad range of negative host health effects beyond those associated with digestion [2-4]. To maintain gut microbiome homeostasis,

bacteria use a variety of signalling and secretion mechanisms to interact with each other as well as their hosts. One of the least well-understood modes of interaction are through outer membrane vesicles (OMVs). These are nanoscale extracellular structures derived through budding of the bacterial outer membrane. They can perform a variety of roles including iron acquisition [5], delivery of toxins [6], facilitation of horizontal gene transfer [7] and as antimicrobial agents through, for example, acting as decoys against phage predation [8, 9]. OMVs can deliver various functional cargoes such as DNA, RNA and proteins, including toxic proteases and lipases [2, 6, 10-12]. They may also directly affect gene expression within target cells through transfer of noncoding RNAs [13]. Research to date has focused on OMVs derived from pathogenic bacteria and their impact on the host, while less attention has been placed upon their role in mutualistic relationships or in maintaining homeostasis among the resident gut microbiota. Developing a detailed understanding of the signalling factors contributing to microbiome homeostasis is challenging as the composition of most gut microbiomes is complex and may contain thousands of bacterial species [14-16]. In contrast, the gut microbiome of the western honey bee, *Apis mellifera*, consists of a small number (8–10) of vertically transmitted [17] bacterial genera, making up ~95% of its total population [18]. This composition is largely stable and resistant to displacement by non-core species [19]. This relative simplicity makes it an ideal model for gut microbiome research.

The majority of the bee gut microbiota reside in the distal compartments of the gut. The pylorus and ileum sections are dominated by gram-negative bacteria, including *Frischella perrara*, *Gilliamella apicola, Gilliamella apis,* and *Snodgrassella alvi*; in contrast, the rectum is predominantly colonised by the gram-positive *Bifidobacterium* spp., *Bombilactobacillus* spp., and *Lactobacillus* spp. Twelve distinct bacterial secretion systems, including OMVs, have been classified to date [20] and several of these have been bioinformatically predicted to exist in the genomes of the *A. mellifera* gut microbiome. Only type 5 and type 6 secretion systems have been functionally characterized, in *S. alvi* [20, 21] and *F. perrara* [22]. Given the ubiquitous nature of gram-negative associated OMVs and their multifaceted functions, detection and characterisation of OMVs in the bee gut is an important step to revealing novel mechanisms contributing to microbiome colonisation and homeostasis. In this study we show that OMVs are abundantly produced by each gram-negative member of the resident bee microbiota, and that they are produced *in vivo*. We also establish baseline quantity, size and zeta potential for OMVs from multiple species, under differing growth conditions and show that they can carry nucleic acid cargo.

2. Materials and Methods

(a) Bacterial culture and OMV collection

We examined members of the core bee gut microbiota for the production of OMVs. A complete list of strains and culture conditions used in this study is found in Table S1. For *in vitro* assays, liquid cultures were filtered through 0.2 μm pore membranes and centrifuged at 110,000–120,000 × g. The pellet was resuspended in dH$_2$O, spun in a 50 kDa molecular weight cut-off filter at 3,000 × g to remove particles <50 kDa. Retentate containing the

OMVs were collected and stored at 4°C. For *in vivo* assays, ilea contents from multiple adult
worker bees colonised with *S. alvi* wkB2 or taken directly from the hive were collected and
pooled, resuspended in dH_2O and then treated as per the *in vitro* assays. Detailed methods are
provided in Supplementary Methods.

(b) Electron microscopy

Gut ileum tissue sections, bacterial cells, and purified OMVs (from bacterial cultures and gut
ilea contents) were visualised with transmission electron microscopy (TEM) using negative
staining. Three-dimensional (3D) tomography imaging was done using a tilt series collected
at 23,000× magnification at 1° increments between -55° and 55°. Unstained, vitrified OMVs
were used for cryo-TEM imaging. Samples were prepared using standard protocols as
detailed in Supplementary Methods.

(c) Confocal imaging of OMV-associated nucleic acids

OMVs from *G. apis* ESL0172 were treated with DNase I and RNase A to remove non-
luminal nucleic acids. DNA or RNA spike-ins were used to determine digestion effectiveness.
For OMV-DNA co-localisation, OMV samples were stained with 1 μg/mL CM-DiI
membrane lipid stain and 1 μg/μL DAPI. For OMV-RNA co-localisation, samples were
stained with 1 μg/mL DiD membrane lipid stain and thiazole red (1 μM). Slides and
coverslips were cleaned with 0.02% Hellmanex in an ultrasonic bath for 30 min, rinsed, and
dried under a nitrogen stream before being activated in a Tergeo Plus Plasma Cleaner. Images
were captured on an Andor Dragonfly 200 (Oxford Instruments, UK) confocal microscope.

(d) OMV concentration, zeta (ζ) potential, and size measurement

Nanoparticle tracking analysis (NTA) measurements of OMV concentration, ζ potential, and
size were performed using a ZetaView x30 Mono (Particle Metrix, Germany). OMV
concentrations were recorded as the average of measurements at 11 positions. To determine
the number of OMVs per cell, calibration curves were generated by measuring the OD_{600}
absorbance of bacterial cultures using spectrophotometry and the corresponding count of
cells using flow cytometry. Zeta potential was calculated as the average of three technical
replicates each measured at two stationary layers. Determination of OMV size based on
electron microscopy imaging was performed using Fiji (ImageJ v1.54p). Briefly, the area of
OMVs with a circularity > 0.7 was calculated. Each vesicle was converted into an equivalent
circle and the diameter calculated. Structures < 66 nm were excluded from statistical analysis
as they could not reliably be identified as OMVs.

(e) Statistical analyses

To evaluate co-localisation of OMVs and nucleic acids, the Manders co-localisation
coefficient for each image was calculated using the Fiji (ImageJ v1.54p) Coloc2 plug-in. The
Costes method was used to determine the likelihood the co-localisation coefficient was a
result of chance, by calculating the frequency that a coefficient generated from randomised
co-localisation would exceed that of the observed co-localisation [23]. To determine if OMV
concentration, ζ potential, or size was correlated with cell density, t-tests of Pearson

correlation coefficients were performed. Reported sample concentrations were corrected for
culture OD_{600}, resuspension volume and dilution factors. Statistical analyses were conducted
in RStudio 2024.12.1 Build 563 or with Microsoft Excel 365 v2510.

3. Results

(a) The bee gut microbiota produces OMVs

We examined the nanoscale fraction from liquid cultures of bacterial strains using TEM
imaging. All gram-negative members of the core bee gut microbiota (*S. alvi*, *G. apicola*, *G.*
apis, *F. perrara*, and *B. apihabitans)* produced near-spherical, membrane-bound structures
(Figure 1a, Figure S1). These were consistent, in size and shape, with OMVs described in
other bacterial species, including *Pseudomonas aeruginosa*, *Vibrio cholerae*, *Klebsiella*
pneumoniae and *Xanthomonas campestris* [24-26], as well as with OMVs collected from the
hyper-vesiculating *Escherichia coli* strain BW25113 Δ*nlpI* (Figure S1). We did not observe
these structures in fractions isolated from the gram-positive bee gut symbiont *L. apis*; gram-
positive bacteria can also produce extracellular vesicles, although less frequently and
generally through different mechanisms than that of gram-negative bacteria [27]. These
structures were also absent in fractions from the gram-negative opportunistic bee gut
pathogen *Serratia marcescens* kz11. OMV production has been shown to be temperature-
dependent in another, human-derived, *S. marcescens* strain [28], thus, the cultivation
temperature used in our study (designed to mimic *in vivo* conditions) may have been
insufficient to trigger OMV production. Additionally, we observed OMV-sized blebs
emerging from intact cells of *G. apicola* wkB1 (Figure 1b), with a size and shape similar to
the vesicles seen within the extracellular space. The membranes of these emerging blebs were
continuous with the outer membranes of the parent cell. Taken together, we conclude that the
OMV-like structures observed from the bee gut bacteria are bonafide OMVs.

(b) OMVs are produced *in vivo*

Ilea sections from 2-week-old hive bees showed pronounced bacterial colonisation and the
presence of putative OMVs identical in appearance to those purified from liquid cultures
(Figure 1c). Using TEM tilt tomography on sectioned hive bee ilea, we produced 3D
reconstructions showing these vesicles to have a spherical morphology (Electronic
Supplementary Material, Video S1). These included vesicles that possess two membranes
(Video S2), suggesting that outer-inner membrane vesicles (O-IMVs) [29] may also be
produced by the bee gut microbiota. To further verify the existence of OMVs within bee guts,
we purified and imaged the liquid contents of pooled ilea from hive bees (pool of 18) or bees
mono-colonised with *S. alvi* wkB2 (pool of 22). Vesicles were observed in both the hive bee
and *S. alvi*-inoculated ilea (Figure 1c). In contrast, we did not observe any OMV-like
structures from two pools of ilea contents (12 and 30 ilea) from non-inoculated bees. NTA
measurement showed OMV abundance from hive bee ilea (pool of 18) and *G. apicola* wkB1-
inoculated ilea (pool of 21) were 2.8× and 3.3× fold higher, respectively, than from non-
inoculated ilea (pool of 30).

(c) Bee gut bacteria OMVs contain nucleic acid

Purified OMVs were treated with a fluorescent membrane stain in combination with either a fluorescent RNA-specific or DNA-specific stain. To remove nucleic acid not contained within the OMV lumen, samples were treated with RNase or DNase prior to visualisation (Figure S2). Confocal microscopy revealed significant OMV and nucleic acid co-localisation, with Costes $p < 0.0001$ (Figure 2a, 2b). We found the proportion of OMVs containing RNA to be 0.594 (Manders co-localisation, SEM 0.040, n = 12) and DNA to be 0.355 (SEM 0.043, n = 10), with consistently more OMVs containing RNA than DNA (Figure 2c, Figure S2). Extraction of nucleic acid from OMVs collected from 50 mL cultures yielded an average of 146.20 ng (SEM 23.41, n = 32) of RNA and 8.55 ng (SEM 3.26, n = 21) of DNA (Figure 2d).

(d) OMV characteristics vary by strain and species

NTA was used to estimate OMV concentration, ζ potential, and size. Comparing between strains, we found significant differences in the number of OMVs produced per cell (Figure 3a). *S. alvi* wkB2 produced considerably more OMVs than other strains (mean = 1161.7, SEM = 526.5, median = 260). The maximum mean OMVs per cell for any *Gilliamella* strain was 2.71 (SEM = 1.31, median = 1.40) for ESL0405. This was similar to *B. apihabitans* K-FP28 (mean = 1.72, SEM = 0.30, median = 1.43). OMV concentration was negatively correlated with cell density for all strains (Table S2). Differences were also observed in ζ potential between strains (Figure 3b). *B. apihabitans* K-FP28 and *Gilliamella* sp. ESL0405 had more negative ζ potentials (means < -20 mV), compared to *S. alvi* wkB2, *G. apicola* wkB1, and *F. perrara* PEB0191 (means > -15 mV). For *S. alvi* wkB2 OMVs, size positively correlated with increasing cell density, while OMVs from *Gilliamella* spp. strains and *B. apihabitans* K-FP28 showed weak to strong negative correlations. NTA measures the hydrodynamic diameter of vesicles and can therefore overestimate their size, while conventional TEM involves a degree of dehydration leading to changes in morphology [30]. Therefore, we used cryo-TEM, a technique which preserves hydration, to produce images of *S. alvi* wkB2 and *G. apicola* wkB1 for size analysis (Figure S3). This approach revealed that *S. alvi* and *G. apicola* produces OMVs of different sizes, with median diameters of 76 nm and 95 nm respectively (Figure 3d). A comparison of size by measurement technique is shown in Figure S3.

4. Discussion

OMVs have been widely reported in bacteria, but their role in shaping microbial communities, particularly within symbiotic host-microbe associations, remain poorly understood. We found that the gram-negative core species of the bee gut microbiome produce abundant OMVs. These were generated without induction, in liquid culture and were also observable directly within the ilea in both mono-inoculated bees and in those collected from the wild. Although similar in general appearance, we found substantial differences in concentration and colloidal properties between the tested strains.

S. alvi wkB2 produced OMVs at far higher quantities than other strains measured. Several
causes of OMV hypervesiculation have been described. These predominantly involve a
disruption to the rigidity of the cell envelope; for instance, null mutation phenotypes of cell
envelop proteins TolR, DegS, NlpI produce hyper-vesiculating phenotypes in *Salmonella* and
E. coli [31, 32]. In wildtype strains of *Salmonella*, disruption to cell envelope integrity is
often associated to stresses such as antibiotic exposure or acidity [33, 34]. Stress is also
induced by increased cell density as nutrients become depleted and waste levels increase [35].
We investigated if this may be the case in our strains by examining if OMV concentration
(OMVs/cell) correlated with cell density (Table S2). All five strains we examined, including
S. alvi wkB2, showed negative correlations with cell density (OD_{600}), with three of five being
significant ($p < 0.05$). Since the optimal conditions for growth are at the early timepoints of
in vitro culture, it is unlikely the higher OMV concentrations seen at lower cell densities, or
the higher concentrations overall for *S. alvi* wkB2, are stress related.

We observed a positive correlation with cell size and density in *S. alvi* wkB2 and negative
correlations for *G. apis* ESL0172 and *Gilliamella* sp. ESL0405. Size variation across growth
phases is associated with OMV functional change. For example, in *Helicobacter pylori* it has
been shown that OMV size determines the mechanism of entry into host gut epithelia cells
[36]. Additionally, the composition of cell envelop proteins in *H. pylori* culture varies with
growth phase [37], thereby suggesting a mechanism whereby OMV protein cargo is
regulated.

There was wide variation in ζ potential between strains, but this was generally uniform across
growth phases (Table S2). Cell surface protein composition is the principal source of ζ
potential; therefore, surface charge can change substantially with alterations to cell state and
can be variable across growth phases [38]. OMV ζ potential can also vary from that of its
parent cell and can have a greater or lesser negative charge [39, 40]. More negative ζ
potential contributes to repulsive forces between particles, preventing aggregation and aiding
even dispersal through the medium, a process extensively studied in red blood cells [41]. Less
negative ζ potential results in lower repulsive force which can lead to increased cell to cell
adhesion [42], potentially enabling OMV fusion to target cells and facilitating delivery of
cargo.

Luminal RNA and DNA were found in a high proportion of OMVs (Figure 2c), although the
amounts of DNA were significantly less than that of RNA. Our preliminary sequencing of
OMV-associated DNA from *S. alvi* wkB2 found most of it to be of prophage origin. Co-
fractionating with OMVs were other structures of a similar size, including capsids and other
phage components, which were noted during TEM imaging (Figure S3b, S3c). Therefore, we
cannot discount the possibility that the DNA was derived from phage remnants and not OMV
cargo. It is also possible that interactions between OMVs and phages can lead to phage DNA
being associated with OMVs. Phage recognition sites have been identified on *Vibrio* and
Salmonella OMVs, and these have been shown to function in phage resistance via adsorption
[9] and DNA ejection [43].

The finding of substantial amounts of OMV associated RNA suggests a possible mechanism
for intercellular communication and transcription modulation in the bee gut [44, 45]. OMV-
mediated small RNA transport between bacteria and eukaryotic hosts have been described in
multiple symbiotic systems, including in the squid light organ [46], plant roots and the human
lung [47]. In honey bees, double-stranded RNA from genetically engineered *S. alvi* in the gut
is able to translocate into host tissue and trigger an RNAi response [48]. OMVs may be a
mechanism for the protection and transport of these molecules, and potentially other factors
between gut bacteria and the host.

While we could not directly compare the OMV concentrations between *in vitro* cultures and
in vivo conditions, qualitatively, there were fewer observable OMVs from our ilea tissue
sections than in cultures. It is possible that OMV production is moderated *in vivo* by
signalling crosstalk between the microbiota and the bee gut epithelium in response to OMV
contact. It would not be unexpected to find that bee microbiota OMVs are engaged in a
signalling process facilitating gut colonisation and persistence. OMVs carry microbially
derived components (e.g., LPS and peptidoglycan), which can illicit immune responses such
as secretion of antimicrobial peptides and reactive oxygen species into the gut lumen [49, 50].
Gut colonisation by symbiotic bacteria is known to be accompanied by modulation of this
immune response — a process which can be mediated by OMVs [2, 51]. This signalling
adapts the local environment to conditions favourable to colonisation and symbiotic host-
microbe interactions.

Altogether, our results identify OMVs as pervasive features of the bee gut microbiota, and
implicate them as underappreciated mediators of microbial and host interactions within the
gut. These findings provide a foundation for future studies of bee bacterial OMVs, with
consequences for understanding intercellular communication in beneficial microbiomes and
for exploring vesicle-mediated pathways as potential targets for microbiome manipulation.

Ethics

This work did not require ethical approval from a human subject or animal welfare
committee.

Data accessibility

Electronic supplementary material is available online, including supplementary methods,
Figures S1–S3, Tables S1–S3, and Videos S1–S2.

Declaration of AI use

We have not used AI-assisted technologies in creating this article.

Authors' contributions

R.E. and W.K.: conceptualisation, methodology; R.E.: investigation, validation, formal
analysis, software, writing—original draft; W.K.: funding acquisition, supervision, Writing –
review & editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration

We declare we have no competing interests.

Funding

This work was supported by the European Research Council Starting Grant 101042912-BEE_GEMS, European Molecular Biology Organization Installation Grant 5045–2022, Fundação para a Ciência e a Tecnologia grant CEECIND/01358/2021, and the Fundação Calouste Gulbenkian.

Acknowledgements

We thank Philipp Engel (Univ. Lausanne), Kasie Raymann (North Carolina State Univ.), and Karina Xavier (Gulbenkian Institute for Molecular Medicine [GIMM]) for providing bacterial strains. We thank the technical support of the Bioimaging Platform of GIMM, funded by PPBI-POCI-01-0145-FEDER-022122, the GIMM Electron Microscopy Facility, the GIMM Flow Cytometry Facility and the Electron Microscopy and X-Rays Facility of the International Iberian Nanotechnology Laboratory.

References

1. Ebenberger, S.P., et al., *The activity of the quorum sensing regulator HapR is modulated by the bacterial extracellular vesicle (BEV)-associated protein ObfA of Vibrio cholerae.* J Extracell Vesicles, 2024. **13**(9): p. e12507.
2. Allin, K.H., T. Nielsen, and O. Pedersen, *Mechanisms in endocrinology: Gut microbiota in patients with type 2 diabetes mellitus.* Eur J Endocrinol, 2015. **172**(4): p. R167–77.
3. Pellegrini, C., et al., *Microbiota-gut-brain axis in health and disease: Is NLRP3 inflammasome at the crossroads of microbiota-gut-brain communications?* Prog Neurobiol, 2020. **191**: p. 101806.
4. Park, A.M. and I. Tsunoda, *Helicobacter pylori infection in the stomach induces neuroinflammation: the potential roles of bacterial outer membrane vesicles in an animal model of Alzheimer's disease.* Inflamm Regen, 2022. **42**(1): p. 39.
5. Lin, J., et al., *A Pseudomonas T6SS effector recruits PQS-containing outer membrane vesicles for iron acquisition.* Nat Commun, 2017. **8**(1): p. 14888.
6. Gao, H., et al., *Outer membrane vesicles from a mosquito commensal mediate targeted killing of Plasmodium parasites via the phosphatidylcholine scavenging pathway.* Nat Commun, 2023. **14**(1): p. 5157.
7. Shen, Z., et al., *Outer membrane vesicles mediating horizontal transfer of the epidemic bla(OXA-232) carbapenemase gene among Enterobacterales.* Emerg Microbes Infect, 2024. **13**(1): p. 2290840.
8. Park, J., et al., *A novel decoy strategy for polymyxin resistance in Acinetobacter baumannii.* Elife, 2021. **10**: p. e66988.

9. Reyes-Robles, T., et al., *Vibrio cholerae outer membrane vesicles inhibit bacteriophage infection.* Journal of bacteriology, 2018. **200**(15): p. 10.1128/jb. 00792–17.

10. Koeppen, K., et al., *A Novel Mechanism of Host-Pathogen Interaction through sRNA in Bacterial Outer Membrane Vesicles.* PLoS Pathog, 2016. **12**(6): p. e1005672.

11. Mondal, A., et al., *Cytotoxic and inflammatory responses induced by outer membrane vesicle-associated biologically active proteases from Vibrio cholerae.* Infection and Immunity, 2016. **84**(5): p. 1478–1490.

12. Chen, F., et al., *Outer membrane vesicle-associated lipase FtlA enhances cellular invasion and virulence in Francisella tularensis LVS.* Emerging Microbes & Infections, 2017. **6**(1): p. 1–12.

13. Swain, M.T., et al., *The RNA cargo of Myxococcus outer membrane vesicles.* Molecular Omics, 2024. **20**(2): p. 138–145.

14. Rosenberg, E., *Diversity of bacteria within the human gut and its contribution to the functional unity of holobionts.* NPJ Biofilms Microbiomes, 2024. **10**(1): p. 134.

15. Handl, S., et al., *Massive parallel 16S rRNA gene pyrosequencing reveals highly diverse fecal bacterial and fungal communities in healthy dogs and cats.* FEMS microbiology ecology, 2011. **76**(2): p. 301–310.

16. Levin, D., et al., *Diversity and functional landscapes in the microbiota of animals in the wild. Science 372: eabb5352.* 2021.

17. Koch, H., et al., *Diversity and evolutionary patterns of bacterial gut associates of corbiculate bees.* Molecular ecology, 2013. **22**(7): p. 2028–2044.

18. Kwong, W.K., et al., *Dynamic microbiome evolution in social bees.* Science advances, 2017. **3**(3): p. e1600513.

19. Steele, M.I., et al., *The Gut Microbiota Protects Bees from Invasion by a Bacterial Pathogen.* Microbiol Spectr, 2021. **9**(2): p. e0039421.

20. Maphosa, S., L.N. Moleleki, and T.E. Motaung, *Bacterial secretion system functions: evidence of interactions and downstream implications.* Microbiology, 2023. **169**(4): p. 001326.

21. Lariviere, P.J., et al., *Virulence-linked adhesin drives mutualist colonization of the bee gut via biofilm formation.* bioRxiv, 2024: p. 2024.10.14.618124.

22. Schmidt, K., et al., *Integration host factor regulates colonization factors in the bee gut symbiont Frischella perrara.* Elife, 2023. **12**: p. e76182.

23. Costes, S.V., et al., *Automatic and quantitative measurement of protein-protein colocalization in live cells.* Biophysical journal, 2004. **86**(6): p. 3993–4003.

24. Chutkan, H., et al., *Quantitative and qualitative preparations of bacterial outer membrane vesicles.* Methods Mol Biol, 2013. **966**: p. 259–72.

25. Bitto, N.J., et al., *Bacterial membrane vesicles transport their DNA cargo into host cells.* Scientific Reports, 2017. **7**(1): p. 7072.

26. Tran, T.M., et al., *Potentiation of plant defense by bacterial outer membrane vesicles is mediated by membrane nanodomains.* Plant Cell, 2022. **34**(1): p. 395–417.

27. Toyofuku, M., N. Nomura, and L. Eberl, *Types and origins of bacterial membrane vesicles.* Nature Reviews Microbiology, 2019. **17**(1): p. 13–24.

28. McMahon, K.J., et al., *Biogenesis of outer membrane vesicles in Serratia marcescens is thermoregulated and can be induced by activation of the Rcs phosphorelay system.* J Bacteriol, 2012. **194**(12): p. 3241–9.

29. Pérez-Cruz, C., et al., *New type of outer membrane vesicle produced by the Gram-negative bacterium Shewanella vesiculosa M7T: implications for DNA content.* Appl Environ Microbiol, 2013. **79**(6): p. 1874–81.

30.   Serrano-Pertierra, E., et al., *Extracellular Vesicles: Current Analytical Techniques for*
Detection and Quantification. Biomolecules, 2020. **10**(6).
31.   Fuentes, I., et al., *Hypervesiculation Meets Sec-Targeting: Enhancing Heterologous*
Protein Loading in Typhi Outer Membrane Vesicles for Delivery and Immune
Response International Journal of Molecular Sciences, 2025. **26**(9).
32.   Ojima, Y., et al., *Aberrant Membrane Structures in Hypervesiculating Escherichia*
coli Strain ΔmlaEΔnlpI Visualized by Electron Microscopy. Frontiers in Microbiology,
2021. **12**.
33.   Dehinwal, R., et al., *A pH-sensitive motif in an outer membrane protein activates*
bacterial membrane vesicle production. Nature Communications, 2024. **15**(1).
34.   Marchant, P., et al., *β-lactam-induced OMV release promotes polymyxin tolerance in*
sv. Typhi. Frontiers in Microbiology, 2024. **15**.
35.   Jaishankar, J. and P. Srivastava, *Molecular Basis of Stationary Phase Survival and*
Applications. Frontiers in Microbiology, 2017. **Volume 8 - 2017**.
36.   Turner, L., et al., *Outer Membrane Vesicle size Determines Their Mechanisms of host*
cell entry and Protein content. Frontiers in Immunology, 2018. **9**.
37.   Zavan, L., et al., *Growth Stage Determines the Size, Protein Composition, and*
Preferential Cargo Packaging of Outer Membrane Vesicles. Proteomics, 2019. **19**(1-
2).
38.   Tashiro, Y., et al., *Variation of physiochemical properties and cell association activity*
of membrane vesicles with growth phase in Pseudomonas aeruginosa. Appl Environ
Microbiol, 2010. **76**(11): p. 3732–9.
39.   Bril'kov, M.S., et al., *Bacterial extracellular vesicles: towards realistic models for*
bacterial membranes in molecular interaction studies by surface plasmon resonance.
Frontiers in Molecular Biosciences, 2023. **10**.
40.   Rogers, N.M.K., et al., *Comparative electrokinetic properties of extracellular vesicles*
produced by yeast and bacteria. Colloids Surf B Biointerfaces, 2023. **225**: p. 113249.
41.   Fernandes, H.P., C.L. Cesar, and L. Barjas-Castro Mde, *Electrical properties of the*
red blood cell membrane and immunohematological investigation. Rev Bras Hematol
Hemoter, 2011. **33**(4): p. 297–301.
42.   Kłodzińska, E., et al., *Effect of zeta potential value on bacterial behavior during*
electrophoretic separation. Electrophoresis, 2010. **31**(9): p. 1590–6.
43.   Stephan, M.S., et al., *In vitro analysis of O-antigen-specific bacteriophage P22*
inactivation by Salmonella outer membrane vesicles. Frontiers in Microbiology, 2020.
11: p. 510638.
44.   Ghosal, A., et al., *The extracellular RNA complement of.* Microbiologyopen, 2015.
4(2): p. 252–266.
45.   Sjöström, A.E., et al., *Membrane vesicle-mediated release of bacterial RNA.* Scientific
Reports, 2015. **5**.
46.   Moriano-Gutierrez, S., et al., *The noncoding small RNA SsrA is released by Vibrio*
fischeri and modulates critical host responses. PLoS biology, 2020. **18**(11): p.
e3000934.
47.   Koeppen, K., et al., *A novel mechanism of host-pathogen interaction through sRNA in*
bacterial outer membrane vesicles. PLoS pathogens, 2016. **12**(6): p. e1005672.
48.   Leonard, S.P., et al., *Engineered symbionts activate honey bee immunity and limit*
pathogens. Science, 2020. **367**(6477): p. 573–576.
49.   Martora, F., et al., *Isolation, characterization and analysis of pro-inflammatory*
potential of outer membrane vesicles. Microbial Pathogenesis, 2019. **136**.

50.   Tleiss, F., et al., *Spatial and temporal coordination of Duox/TrpA1/Dh31 and IMD*
415   *pathways is required for the efficient elimination of pathogenic bacteria in the*
416   *intestine of larvae.* Elife, 2024. **13**.
417   51.   Fábrega, M.J., et al., *Intestinal Anti-inflammatory Effects of Outer Membrane Vesicles*
418   *from Escherichia coli Nissle 1917 in DSS-Experimental Colitis in Mice.* Frontiers in
419   Microbiology, 2017. **8**.

Figures

Figure 1. OMVs are produced by bee gut microbiota members *in vitro* and *in vivo*. (a) Representative TEM micrographs of OMV-purified fractions from liquid cultures of bee gut bacteria strains showing presence of OMVs from gram-negative members of the normal microbiota. (b) OMV budding. OMVs can be found attached to the bacterial cell surface (unfilled arrows) as well as detached, in extracellular space (solid arrows). (c) Imaging of bee ilea tissue sections show presence of OMVs in wild-caught adult hive bees as well as lab-reared bees mono-colonised with *S. alvi* wkB2. Scale bars = 200 nm unless otherwise indicated.

Figure 2. Bee microbiota OMVs contain nucleic acids. Confocal fluorescence imaging of OMVs from *G. apis* ESL0172 shows (a) co-localisation of membrane lipid stain (DiD, red) and RNA stain (SYTO RNASelect, green), and (b) co-localisation of membrane lipid stain (DiI, yellow) and DNA stain (DAPI, blue). Scale bars = 10 μm. (c) Proportion of OMVs with co-localised nucleic acid (Manders correlation coefficient). (d) Quantification of nucleic acids in OMVs from stationary phase cultures. Error bars = SEM.

Figure 3. OMV characteristics differ by species. (a) OMV abundance and (b) ζ potential of strains grown in liquid cultures. Significant differences, indicated by letters above boxplots, tested with Welch's ANOVA and Games-Howell post hoc tests. (c) OMV sizes of *S. alvi* wkB2 and *G. apicola* wkB1, as measured by cryo-TEM. Significance tested with Wilcoxon rank-sum test. Dotted line indicates the size threshold (66 nm) for OMV detection.