A novel polymerase III promoter for gene editing in the agricultural pest *Ceratitis capitata*

Amber S. K. Hall[†], Samuel Matthew Shackleton-Chavez[†], Tracey Chapman, Philip T. Leftwich[*]

[†] These authors contributed equally to the work

[*] Correspondence to p.leftwich@uea.ac.uk

School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, U.K.

Keywords: Cas9; Gene Drive; Medfly; Mediterranean Fruitfly; Polymerase III

Abstract

We report the identification and functional validation of a 7SK RNA polymerase III promoter in the Mediterranean fruit fly, *Ceratitis capitata*. CRISPR/Cas9-based genetic control strategies for this global agricultural pest, including gene drives and precision guided sterile insect approaches, require efficient guide RNA expression, yet only a single U6 Pol III promoter had previously been validated for this purpose in *C. capitata*, and no 7SK promoter had been characterised in any Tephritid species. Using comparative genomics with *Drosophila* orthologues, we identified a previously unannotated 7SK gene in the *C. capitata* genome, confirmed its transcriptional activity by RT-PCR, and demonstrated that the cloned promoter drives functional guide RNA expression in CRISPR/Cas9-mediated knockouts of the *white* gene. Comparative analysis identified putative 7SK orthologues across the Tephritid fruit flies. The availability of this additional new Pol III promoter will enable multiplexed guide RNA strategies using distinct promoters, supporting more robust genetic control designs.

27 Graphical Abstract

Introduction

The Mediterranean fruit fly, *Ceratitis capitata* (medfly), is a globally significant agricultural pest
within the family Tephritidae, causing damage to over 200 fruit crop species (Christenson &
Foote, 1960). CRISPR/Cas9-based genetic control strategies, including homing gene drives,
pgSIT, and sex conversion approaches, are under active development in this species. These
strategies require the endogenous expression of two components: the Cas9 endonuclease,
driven by RNA polymerase II (Pol II) promoters, and the single guide RNA (sgRNA), driven by RNA
polymerase III (Pol III) promoters. Whilst the range of endonuclease promoters used in CRISPR
applications is broad, gRNA promoter choice remains comparatively restricted (Verkuijl et al.,
2026).

Several Pol II promoters have been evaluated for Cas9 expression in *C. capitata*, including
nanos, *vasa,* and *zpg* regulatory elements. In contrast,only a single Pol III promoter, *CcU6*, has
been characterised for sgRNA expression (Meccariello et al. 2021). This limitation has practical
consequences. A single sgRNA target site for gene drive homing permits the accumulation of
resistance alleles through non-homologous end joining repair, eventually rendering the drive
ineffective at the population level (Meccariello et al. 2024, Verkuijl et al. 2022). Multiplexed
guide RNA designs, which target multiple sites within the same gene, represent a promising
strategy to mitigate resistance (Anderson et al. 2024, Xu et al. 2025). However, expressing
multiple sgRNAs from repeated copies of the same promoter increases the risk of internal
recombination and cassette instability (Verkuijl et al. 2022). The use of distinct Pol III promoters
for each sgRNA cassette avoids this problem and has been demonstrated successfully in
mosquito gene drives (Anderson et al. 2024, Gonzalez et al. 2025).

Pol III promoters in insects are characterised by two conserved upstream elements: a proximal
sequence element A (PSEA) and a TATA box (Kim et al. 2020, Hernandez et al. 2006). These
motifs are conserved across Pol III promoter families, including U6 and 7SK classes, enabling
identification of orthologous sequences through comparative genomics (Gruber et al. 2008).

The 7SK small nuclear RNA is a component of the 7SK ribonucleoprotein complex involved in transcriptional regulation, and its promoter has been co-opted for sgRNA expression in several insect species. In *Aedes aegypti, Anopheles stephensi,* and *Culex quinquefasciatus*, 7SK-driven sgRNA expression has proved functional for CRISPR/Cas9 editing, including in gene drive contexts (Anderson et al. 2020, Anderson et al. 2024, Gonzalez et al. 2025, Purusothaman et al. 2021).

Comparative assessments of Pol III promoters for sgRNA expression in mosquitoes support the functional utility of 7SK-class promoters. In culicine mosquito cell lines, 7SK sequences have been found to be transcriptionally active and comparable in efficiency to U6 paralogues in driving reporter expression (Anderson et al. 2020). When tested in gene drive constructs *In vivo*, 7SK promoters have yielded among the highest inheritance bias (Gonzalez et al. 2025). These findings indicate that 7SK promoters can match or exceed the activity of established U6 sequences, and that the choice of Pol III promoter can materially affect gene drive performance. No 7SK promoter has yet been identified or tested in any Tephritid species.

Here, we report the identification, validation, and functional characterisation of a 7SK Pol III promoter in *C. capitata*. We demonstrate that the *Cc7SK* promoter drives sgRNA expression sufficient to generate CRISPR/Cas9-mediated knockout of the *white* gene (chromosome 5, GeneID_101458180), establishing a second validated Pol III promoter for this species. Through comparative genomic analysis, we additionally identify putative 7SK orthologues across nine Tephritidae species, suggesting that this promoter class may be broadly available within this family of agricultural pests.

Results & Discussion

Identification and validation of a C. capitata 7SK RNA gene

Using the *Drosophila melanogaster* 7SK RNA sequence (FBgn0065099) as a query, we identified a putative 7SK orthologue in the *C. capitata* genome (Ccap_2.1 and EGII-3.2.1 assemblies) that had not been previously annotated. Examination of the upstream region revealed a recognisable PSEA motif and a modified TATA box bearing a cytosine substitution at the 5' position (Figure 1). We extended this analysis across the Tephritidae, identifying a putative 7SK gene with recognisable PSEA and TATA box motifs in all nine species examined. While gene sequences were well conserved, TATA box sequences showed occasional substitutions (Figure 1, Table S1-S2)

To confirm that the identified locus is transcriptionally active, we performed RT-PCR on RNA extracted from adult *C. capitata* (Benakeion strain) and detected a product of the expected size (Figure S1, Table S3). This confirmed endogenous transcription from the *Cc7SK* locus.

FIGURE 1. Conservation of 7SK promoter elements across dipteran species and phylogenetic
relationships of 7SK genes. **(A)** Alignment of 7SK promoter sequences from *Ceratitis*, other tephritids,
Drosophila, mosquitoes, and *Plutella*, with annotated transcription start site (TSS), proximal sequence
element A (PSEA), and TATA box. **(B)** Phylogenetic relationships of 7SK genes among dipteran species,
inferred by neighbour-joining under the HKY substitution model, with *Plutella xylostella* 7SK as the
outgroup. Scale bar represents nucleotide substitutions per site.

Generation of white knockouts using Cc7SK-driven sgRNA expression

To test whether the *Cc7SK* promoter can drive functional sgRNA expression, we cloned a 600 bp
region upstream of the predicted transcription start site and used it to express a previously
validated sgRNA targeting exon 3 of the *white* gene (Meccariello et al. 2017). As a positive
control, we generated a parallel construct using the established CcU6 promoter with the
identical sgRNA sequence. Both constructs additionally carried an AmCyan-NLS fluorescent
marker to allow identification of successfully injected individuals.

Embryos for injection were generated by crossing OAMS21-Ccpub-Cas9 females (Davydova et
al. 2025), which provide both integrated and maternally deposited Cas9, to homozygous white-
eye mutant males. The resulting embryos were therefore heterozygous at the *white* locus,
carrying one functional and one disrupted allele, such that monoallelic disruption was required
for a visible phenotype change.

We injected embryos with either Cc7SK-white-eye-sgRNA (n = 584) or CcU6-white-eye-sgRNA (n
= 664) plasmids. Larvae were screened at 24–72 hours post-injection for AmCyan fluorescence
and separated by fluorescence score. Among adults that eclosed, we identified 9/73 (12%)
mosaic white-eyed individuals from Cc7SK-injected embryos and 6 /48 (12.5%) from CcU6-
injected embryos (full data in Table S4). Mosaic individuals displayed clusters of depigmented
(white) ommatidia against the wild-type dark eye background, consistent with somatic
disruption of *white* in a subset of cells. The recovery of mosaic phenotypes from Cc7SK-injected
embryos provided direct evidence that the Cc7SK promoter drives functional sgRNA expression
sufficient for Cas9-mediated cleavage in vivo (Figure 2).

Molecular confirmation of editing at the white *locus*

To obtain molecular evidence of target site editing, we performed a separate injection
experiment using the same constructs. Embryos injected with Cc7SK-white-eye-sgRNA (n =
435) or CcU6-white-eye-sgRNA (n = 458) were screened as L1 hatchlings for AmCyan
fluorescence, reared to the L2 stage, and collected for genomic DNA extraction. Fluorescence-
positive larvae were pooled by construct (Cc7SK, n = 20; CcU6, n = 19) and a 258 bp amplicon
spanning the sgRNA target site was sequenced using Illumina MiSeq (Amplicon-EZ, Genewiz).

Analysis with CRISPResso2 (Clement et al. 2019) revealed indels at the predicted Cas9
cleavage site in both samples (Figure 2). The predominant editing outcome under both
promoters was a 2 bp deletion immediately 5' of the PAM sequence (7SK: 39.38% and U6
32.91%). The *white* mutant allele, which carries a deletion spanning the majority of the sgRNA
recognition sequence, showed minimal editing, consistent with the loss of the target site in that
allele. The concordance of the dominant indel signature between both promoter constructs
provides molecular confirmation that the Cc7SK promoter drives production of the same
functional sgRNA as CcU6.

FIGURE 2. CRISPR-mediated editing of the *white* locus and resulting adult phenotypes. **(A)** Alignment of
the engineered sgRNA to the *white* reference sequence (GeneID: 101458180); the grey bar indicates the
sgRNA target region. Values denote the percentage of sequencing reads assigned to each sequence
variant for 7SK-driven and U6-driven sgRNA constructs, respectively. The upper rows show confirmed
wildtype and white mutant allele sequences; the lower rows show indel sequences identified by
CRISPResso2 analysis. Source data are deposited under ENA project PRJEB111759. **(B)** Representative
images of post-microinjection adult flies: wildtype (left) and mosaic mutant (right).

Implications for genetic control construct design

These data establish Cc7SK as the second validated Pol III promoter in *C. capitata*. It is
important to note that this proof-of-concept experiment, using transient plasmid-based
expression, does not allow quantitative comparison of transcriptional activity between Cc7SK
and CcU6. Differences in plasmid uptake during microinjection preclude meaningful
assessment of relative expression levels. Previous work in *Anopheles stephensi* demonstrated

that 7SK promoters can equal or exceed U6 activity for sgRNA expression when integrated and
directly compared (Gonzalez et al. 2025). A comparable integrated comparison in *C. capitata*,
measuring cutting or homing rates from the same genomic locus under identical Cas9
regulation, would be needed to determine the relative activity of CcU6 and Cc7SK in the
germline.

The availability of a second Pol III promoter has immediate practical implications. Multiplexed
gene drive designs in *Aedes aegypti* have used distinct Pol III promoters (U6 and 7SK) for each
sgRNA cassette to avoid repetitive sequences that promote recombination (Anderson et al.
2024). A similar approach is now feasible in *C. capitata*, which could improve the durability of
homing-based gene drives by targeting multiple sites within a gene such as transformer
(Meccariello et al. 2024). Additionally, different Pol III promoters may have distinct
spatiotemporal expression profiles. Having multiple promoter options therefore provides
flexibility to optimise sgRNA expression timing relative to Cas9 activity in the germline.

The identification of putative 7SK orthologues in eight further Tephritidae species (Table S1)
suggests that this approach may be transferable to related pest species for which genetic
control tools are currently lacking.

Methods

A putative 7SK small nuclear RNA gene was identified in the *C. capitata* genome (Ccap_2.1 and
EGII-3.2.1 assemblies) by BLASTN using the *D. melanogaster* 7SK RNA sequence as a query.
Upstream promoter elements (PSEA, TATA box) were identified computationally, and
orthologous sequences were searched across nine Tephritidae species (Supporting Information
S1-S2). Endogenous transcription was confirmed by RT-PCR on cDNA synthesised from adult
Benakeion RNA (Supporting Information S3–S7).

A 600 bp region upstream of the Cc7SK transcription start site was amplified from Benakeion
genomic DNA and cloned into an sgRNA expression plasmid carrying an AmCyan marker. A
parallel construct using the previously characterised CcU6 promoter (Meccariello et al. 2021)
with the identical white-eye sgRNA (Meccariello et al. 2017) served as a positive control. Both
plasmids were verified by Nanopore sequencing (Supporting Information S8).

Embryos from crosses of OAMS21-Ccpub-Cas9 females (Davydova et al. 2025) to homozygous
white-eye mutant males were injected with either construct at 500 ng/µl within 1–2 hours of
oviposition. For phenotypic assessment, larvae were reared to adulthood and scored for eye
colour mosaicism. For molecular confirmation, fluorescence-positive L2 larvae were pooled by
construct and a 258 bp amplicon spanning the target site was sequenced on an Illumina MiSeq
platform and analysed with CRISPResso2 (Clement et al. 2019). Full protocols are provided in
Supporting Information S9–S16.

Data Availability

183 Raw sequencing data are available through the European Nucleotide Archive under project
184 accession PRJEB111759. Scripts and raw data can be found at (https://github.com/Philip-
185 [Leftwich/A-novel-polymerase-3-promoter-for-gene-editing-in-the-agricultural-pest-Ceratitis-](https://github.com/Philip-Leftwich/A-novel-polymerase-3-promoter-for-gene-editing-in-the-agricultural-pest-Ceratitis-capitata)
186 [capitata](https://github.com/Philip-Leftwich/A-novel-polymerase-3-promoter-for-gene-editing-in-the-agricultural-pest-Ceratitis-capitata)). Complete information on analyses can be found in Supporting Information.

Acknowledgements

188 ASKH and SMSC supported by a UKRI BBSRC Norwich Research Park Biosciences Doctoral
189 Training Partnership (Grant No. BB/T008717/1 awarded to PTL and TC). We thank Angela
190 Meccariello and Jean-Charles de Coriolis for their advice and comments.

191 References

Anderson, M.A.E., Purcell, J., Verkuijl, S.A.N., Norman, V.C., Leftwich, P.T., Harvey-Samuel, T.
and Alphey, L. (2020) 'Expanding the CRISPR toolbox in culicine mosquitoes: in vitro validation
of Pol III promoters', ACS Synthetic Biology, 9(3), pp. 678–681. doi: 10.1021/acssynbio.9b00436.

Anderson, M.A.E., Gonzalez, E., Edgington, M.P., Ang, J.X.D., Purusothaman, D.-K., Shackleford,
L., Nevard, K., Verkuijl, S.A.N., Harvey-Samuel, T., Leftwich, P.T., Esvelt, K. and Alphey, L. (2024)
'A multiplexed, confinable CRISPR/Cas9 gene drive can propagate in caged *Aedes aegypti*
populations', Nature Communications, 15(1), 729. doi: 10.1038/s41467-024-44956-2.

Christenson, L.D. and Foote, R.H. (1960) 'Biology of fruit flies', Annual Review of Entomology, 5,
pp. 171–192. doi: 10.1146/annurev.en.05.010160.001131.

Clement, K., Rees, H., Canver, M.C., Gehrke, J.M., Farouni, R., Hsu, J.Y., Cole, M.A., Liu, D.R.,
Joung, J.K., Bauer, D.E. and Pinello, L. (2019) 'CRISPResso2 provides accurate and rapid genome
editing sequence analysis', Nature Biotechnology, 37(3), pp. 224–226. doi: 10.1038/s41587-019-
0032-3.

Davydova, S., Liu, J., Liu, Y., Prince, K., Mann, J., Kandul, N.P., Braswell, W.E., Champer, J.,
Akbari, O.S. and Meccariello, A. (2025) 'A self-limiting sterile insect technique alternative for
Ceratitis capitata', BMC Biology, 23, 88. doi: 10.1186/s12915-025-02201-2.

Gonzalez, E., Anderson, M.A.E., Ang, J.X.D., Nevard, K., Shackleford, L., Larrosa-Godall, M.,
Leftwich, P.T. and Alphey, L. (2025) 'Optimization of sgRNA expression with RNA pol III regulatory
elements in *Anopheles stephensi*', Scientific Reports, 15(1), 13408. doi: 10.1038/s41598-025-
98557-0.

Gruber, A.R., Kilgus, C., Mosig, A., Hofacker, I.L., Hennig, W. and Stadler, P.F. (2008) 'Arthropod
7SK RNA', Molecular Biology and Evolution, 25(9), pp. 1923–1930. doi: 10.1093/molbev/msn140.

Hernandez, G., Jr., Valafar, F. and Stumph, W.E. (2007) 'Insect small nuclear RNA gene
promoters evolve rapidly yet retain conserved features involved in determining promoter activity
and RNA polymerase specificity', Nucleic Acids Research, 35(1), pp. 21–34. doi:
10.1093/nar/gkl982.

Kim, J.-L. and Bhargava, P. (2020) or Kim, J.-L. and Stumph, W.E. (2020) 'Assembly of SNAPc,
Bdp1, and TBP on the U6 snRNA gene promoter in *Drosophila melanogaster*', Molecular and
Cellular Biology, 40(12), e00641-19. doi: 10.1128/MCB.00641-19.

Meccariello, A., Monti, S.M., Romanelli, A., Colonna, R., Primo, P., Inghilterra, M.G., Del
Corsano, G., Ramaglia, A., Iazzetti, G., Chiarore, A., Patti, F., Heinze, S.D., Salvemini, M.,
Lindsay, H., Chiavacci, E., Burger, A., Robinson, M.D., Mosimann, C., Bopp, D. and Saccone, G.
(2017) 'Highly efficient DNA-free gene disruption in the agricultural pest *Ceratitis capitata* by
CRISPR-Cas9 ribonucleoprotein complexes', Scientific Reports, 7(1), 10061. doi:
10.1038/s41598-017-10347-5.

Meccariello, A., Krsticevic, F., Colonna, R., Del Corsano, G., Fasulo, B., Papathanos, P.A. and
Windbichler, N. (2021) 'Engineered sex ratio distortion by X-shredding in the global agricultural
pest *Ceratitis capitata*', BMC Biology, 19, 78. doi: 10.1186/s12915-021-01010-7.

Meccariello, A., Hou, S., Davydova, S., Fawcett, J.D., Siddall, A., Leftwich, P.T., Krsticevic, F.,
Papathanos, P.A. and Windbichler, N. (2024) 'Gene drive and genetic sex conversion in the
global agricultural pest *Ceratitis capitata*', Nature Communications, 15, 372. doi:
10.1038/s41467-023-44399-1.

Purusothaman, D.-K., Shackleford, L., Anderson, M.A.E., Harvey-Samuel, T. and Alphey, L.
(2021) 'CRISPR/Cas-9 mediated knock-in by homology dependent repair in the West Nile Virus
vector *Culex quinquefasciatus* Say', Scientific Reports, 11, 14964. doi: 10.1038/s41598-021-
94065-z.

Verkuijl, S.A.N., Ang, J.X.D., Alphey, L., Bonsall, M.B. and Anderson, M.A.E. (2022) 'The
challenges in developing efficient and robust synthetic homing endonuclease gene drives',
Frontiers in Bioengineering and Biotechnology, 10, 856981. doi: 10.3389/fbioe.2022.856981.

Verkuijl, S.A.N., Cook, E.I., Bairu, L., Bonsall, M.B., Leftwich, P.T. and Windbichler N. (2026)
Cross-species meta-analysis reveals determinants of homing gene drive performance, bioRxiv,
doi: 10.64898/2026.02.28.708699.

Xu, X., Chen, J., Wang, Y., Liu, Y., Zhang, Y., Jia, X., Yang, J., Yang, X., He, Z. and Champer, J.
(2025) 'Gene drive-based population suppression in the malaria vector *Anopheles stephensi*',
Nature Communications, 16, 1007. doi: 10.1038/s41467-025-56290-2.