Amyloid plaques drive state-dependent long-range circuit reorganization in the hippocampus

Zhao Z[1,2], Locascio JJ[1,2], Li H[1], Gowravaram N[1], Green RJ[1], Kastanenka K[1,2], Bacskai B[1,2], Hyman BT[1,2], Gomperts SN[1,2,*]

1.   Department of Neurology, Massachusetts General Hospital; Boston MA, 02114, USA

2.   Harvard Medical School, Boston MA, 02115

* Corresponding author. Email: Gomperts.stephen@mgh.harvard.edu

Amyloid plaques are a pathological hallmark of Alzheimer's disease, but how they drive

widespread neuronal dysfunction remains unclear. While studies in anesthetized animals show that

plaques drive local hyperactivity[1,2], it is unknown how this pathology shapes functional

hippocampal maps in freely behaving animals. We combined chronic 1-photon calcium imaging,

local field potential recordings, and post hoc 2-photon plaque imaging in freely behaving APP/PS1

mice across behavior and sleep to correlate real-time hippocampal activity and place coding with

precise plaque topography. Here we show that plaques exert nonlocal, long-range effects on

hippocampal activity that depend on plaque size, laminar position, and the animal's behavioral

state. Place cells, which encode spatial position and are normally uniformly distributed, are

preferentially enriched near plaques, revealing an aberrant reorganization of plaque-adjacent

neurons into the hippocampal map of space. In longitudinal experiments, pre-existing place cell

locations do not predict future plaque sites, whereas hyperactivity during slow-wave sleep weakly

predicts future amyloid deposition. These findings identify a mechanism by which amyloid

pathology reorganizes brain circuits, degrading the functional architecture of the hippocampus

and contributing to widespread dysfunction and cognitive impairment in Alzheimer's disease.

During exploratory behavior, many hippocampal neurons form stable spatial representations

known as place fields. Notably, spatial representations deteriorate in parallel with spatial memory as

plaques accumulate in murine amyloid-β_{42} (Aβ) models of Alzheimer's disease[3–5], mimicking early

navigational deficits in patients (e.g., [6]). Aberrantly hyperactive and silent neuronal subpopulations in

hippocampus and cortex of murine Alzheimer's disease models suggest that disrupted excitatory-

inhibitory balance underlies Aβ-associated circuit dysregulation[1,2,7]. Fibrillar and oligomeric

Aβ species are both neurotoxic and synaptotoxic [8–14] and are well-positioned to disrupt circuit function.

In anesthetized Alzheimer's disease model mice, aberrantly hyperactive cells have been found

to cluster exclusively near amyloid plaques, while silent cells have been found to be uniformly

distributed[1,2]. However, behavioral state critically influences hippocampal activity[15,16], and anesthetics

have complex effects on hemodynamics, neurophysiology, and calcium homeostasis[17–19]. Therefore,

whether plaque-associated hyperactivity occurs during normal behaviors and across the sleep-wake

cycle, and how it impacts place coding, remains unknown. Furthermore, the impact of aberrant

clustering of hyperactive cells proximal to plaques on place coding has not been established; this

clustering and hyperactivity is likely to undermine place coding and degrade pattern separation needed

for accurate storage of unique experiences.

To address these gaps, we quantified CA1 calcium dynamics and concomitant

electrophysiological activity in freely-behaving APP/PS1 mice, and then co-registered these data to

high-resolution plaque topography. We show that plaques induce nonlocal, long-range, state-dependent

effects on neuronal activity and place coding that depend on plaque size and laminar location within

CA1. These findings provide a new framework in which amyloid plaques act as foci of network-level

circuit reorganization, rather than as purely focal lesions.

Behavioral state transforms the spatial influence of amyloid plaques on hippocampal circuits.

To investigate how amyloid plaques impact hippocampal function across behavioral states, we acquired simultaneous chronic GCaMP6f calcium imaging and local field potential (LFP) recordings in the CA1 region of freely behaving APP/PS1 mice across the sleep-wake cycle, followed by high-resolution plaque imaging (**Fig. 1a,b** and **Supplementary Fig. 1;** see **Methods**). We stratified neurons into three activity classes, high activity, moderately active, and quiescent cells, based on state-specific calcium event rates (**Fig. 1b,c**).

In contrast to previous observations made in anesthetized Alzheimer's mice[1,2], high-activity cells were not restricted to the immediate vicinity of amyloid plaques, but rather were detected at all distances from the nearest plaque across all behavioral states (**Fig. 1d and Supplementary Fig. 2**). Although the proportion of high-activity cells visually decreased with distance from the nearest plaque during running and slow-wave sleep, Spearman correlations between distance and the fraction of cells in each activity class were not significant in any state after Bonferroni correction (**Fig. 1d**). By contrast, CA1 activity profiles varied significantly with behavioral state (**Fig. 2a**), indicating that behavioral state, rather than local plaque pathology, primarily determines the distribution of activity strata.

We next modeled calcium rate as a continuous function of distance to the nearest plaque. Proximity to plaques was associated with higher activity in high-activity cells across states, with effects extending over hundreds of microns, and with lower activity in quiescent cells during running (**Fig. 2b**). Moderately active cells showed a nuanced response, with slightly lower activity near plaques during running and slightly higher activity during QW, SWS, and REM sleep. Importantly, activity class was state-dependent rather than an intrinsic property of the cell, supporting plaque-induced dysfunction as a dynamic circuit-level phenomenon. The specific relationship between plaque proximity and neuronal activity varied markedly across the sleep-wake cycle (**Fig. 2b**). These results suggest that plaques induce differential effects on neuronal activity that depend on both activity type and behavioral state.

Plaque size and laminar position determine the magnitude of spatial circuit disruption

Amyloid plaques varied in size and laminar position, with some centered in the pyramidal cell

layer and others in dendritic layers of stratum radiatum and stratum oriens (**Fig. 3a**). Because amyloid

plaques disrupt both the neuronal processes that traverse them[20] and local inhibitory synapses[21], we

next determined whether plaque size or laminar position within CA1 correlated with specific activity

patterns.

Neuronal activity levels were strongly associated with both plaque size and laminar position

(**Fig. 3b**). Large pyramidal-layer plaques were associated with the highest plaque-adjacent activity and

the steepest decay with distance. Small pyramidal-layer plaques showed intermediate effects, and

dendritic-layer plaques had the smallest impact. Thus, the spatial extent and magnitude of amyloid-

driven neuronal dysfunction are shaped by specific plaque characteristics and anatomical location.

The spatial organization of hippocampal maps relates to amyloid plaque topography

Spatial memory impairment is a hallmark of Alzheimer's disease[22] that is well-recapitulated in

the APP/PS1 model[3–5]. Given that hippocampal place cell recruitment during navigation is critical for

spatial memory formation[23], we evaluated the impact of amyloid plaques on hippocampal spatial

representations. APP/PS1 mice exhibited a lower proportion of place cells than age-matched wild-type

controls (**Fig. 4a-c**). Although the percentage of time spent running was similar, APP/PS1 mice ran

more slowly, a potential confound that may contribute to reduced place cell yield. Thus, despite

reduced place cell yield and altered physiology[3–5,16], the core hippocampal network required for place

coding remained functional in APP/PS1 mice, although future work will be required to disentangle

disease effects from speed-related influences on place cell recruitment.

To test whether plaques shape place cell topography, we compared the distances between place

cells in APP/PS1 and WT mice and related place cell identity to plaque proximity. The distribution of

place cell distances in APP/PS1 mice over-represented short distances (**Fig. 4d,e**). In addition, the

probability of a neuron being a place cell increased significantly with decreasing distance to the nearest

plaque (**Fig. 4f**; see **Methods**). Notably, this effect was greatest near dendritic-layer plaques. In

contrast, plaque proximity did not affect place cell spatial information (**Supplementary Fig. 3**),

indicating that plaques bias place cell recruitment rather than the quality of their spatial tuning.

Together, these results show that place cells are clustered near amyloid plaques.

Place cell topography does not prefigure future amyloid deposition sites

Neuronal activity drives release of Aβ42[24] and contributes to its concentration in interstitial

fluid[25]. To determine whether place cells aggregate near amyloid plaques or whether pre-existing place

cell clusters drive future plaque deposition, we acquired dynamic calcium imaging in CA1 of young

APP/PS1 mice prior to plaque formation. We then related the location of these place cells to the

topography of future plaques imaged in the same animals 5 months later (**Fig. 5a; see Methods**).

Place cell identity before plaque formation did not predict future plaque locations (**Fig. 5b**). By

contrast, elevated neuronal activity during slow-wave sleep was weakly associated with future plaque

locations (**Fig. 5c**), suggesting that state-specific hyperactivity during sleep may promote local amyloid

deposition. These findings suggest that while the functional architecture of the hippocampal map is

reorganized by plaques, its initial layout does not dictate the spatial distribution of amyloid pathology.

Discussion

Alzheimer's disease is characterized by progressive hippocampal failure, yet local plaque density

correlates poorly with cognitive impairment. Our findings resolve this paradox by showing that amyloid

plaques act as drivers of long-range, state-dependent circuit dysfunction in both neuronal activity and the

hippocampal map of space. This framework implies that the cognitive burden of the disease reflects

network-level collapse due to nonlocal plaque-associated disturbances, rather than simply the volume or

density of local plaque deposition.

Place coding is a key function of the hippocampus, and its deterioration in Alzheimer's patients

and in Alzheimer's models is well established[22]. The biased participation of plaque-adjacent cells in the

representation of space reveals a novel mechanism whereby Aβ deposits reorganize existing functional

circuits to degrade the functional architecture of the hippocampus. With accumulation of amyloid plaques

in Alzheimer's disease, excessive reliance on a reduced pool of available neurons to represent space

within and across environments would be expected to progressively impair both hippocampal remapping

and pattern separation, fundamental features of hippocampal mnemonic function[26].

Our results also suggest that amyloid plaques drive the clustering of place cells rather than vice

versa, as our longitudinal studies did not detect a predictive effect of place cell locations on future

plaque positions. Although the mechanisms underlying the reorganization of place cells to cluster near

amyloid plaques are unknown, loss of inhibitory synapses in the region around amyloid plaques, as

recently reported[21], is well-positioned to contribute, as may the disruption of local processes[20].

Alternatively, as a smaller proportion of neurons was found to represent space in this Alzheimer's

model compared to WT mice, another possibility is that place coding was lost in cells far from plaques.

Unexpectedly, we detected long-range gradients of neuronal activity centered on amyloid plaques

rather than highly local plaque effects. Although the biological basis of these effects is not yet clear, and

our estimates are necessarily limited by the spatial extent of the imaging window, we anticipate that

spatial gradients of plaque-derived soluble Aβ oligomers may contribute, as these species may diffuse

through the interstitial space to disrupt synapses far beyond the physical boundaries of the plaque[10,12].

Prior reports of aberrantly high and low activity of hippocampal and cortical neurons in amyloid

plaque-bearing Alzheimer's models have supported the concept of perturbed excitatory-inhibitory

balance in Alzheimer's disease[1,2,7]. Behavioral state has a marked effect on such imbalance[16]. The present

results extend these findings, showing that the spatial effects of plaques on neuronal activity are also

markedly state-dependent. In Run behavior, aberrant hyper- and hypoactivity was greatest close to

plaques. However, the plaque-associated dysfunction declined faster with distance in non-Run states than

in Run, and hypoactivity was unaffected by proximity in non-Run states. Thus, the brain's functional

state dictates the spatial extent of the plaque-associated pathophysiology.

Consistent with the toxicity of Aβ and the local impact of amyloid plaques on the brain

parenchyma, large plaques were more disruptive to neuronal activity than small plaques. Interestingly,

large plaques deposited in the pyramidal cell layer were most disruptive, while plaques in the dendritic

layers were least disruptive. In contrast, we observed that place cells were clustered more around

dendritic layer plaques than cell layer plaques. These observations – which suggest a laminar-specific

vulnerability of the hippocampal map – may reflect differential effects on inhibitory versus excitatory

synapses, given their laminar distribution within CA1[27] as well as the possible contribution of loss of

local inhibitory synapses[21,28].

Intriguingly, we found that the locations of hyperactive cells in SWS predicted future plaque

locations, albeit weakly. As SWS is the one behavioral state associated with amyloid clearance[29,30], it is

possible that release of Aβ in SWS by such hyperactive cells overwhelms local clearance mechanisms,

enabling local Aβ concentrations to rise above the local critical threshold for oligomerization and

deposition. Future work is needed to evaluate this hypothesis, and larger longitudinal cohorts will be

required to determine the robustness of this association.

Key highlights of this study include linking spatial representations and neuronal activity across

behavioral states to the local influence of amyloid plaques in the hippocampus, a region critically

involved in early Alzheimer's pathology and memory loss. A further strength was the use of 2P

microscopy to image amyloid plaques, providing an accurate assessment of plaque burden across the

laminar extent of CA1. While dynamic calcium imaging provides a robust measure of large-scale

neuronal ensembles, future studies incorporating high-density electrophysiology may provide

additional temporal resolution to the relationship between place cells and amyloid plaques[15,16]. Co-

registration of 1P and 2P datasets using angiogram fiducials was associated with low error. Although

sample sizes of mice were modest, we were able to sample 85 plaques and 3,762 neurons across

behavioral states. Another limitation was the inability to assess the effects of amyloid plaques on

calcium in non-somatic cellular compartments. In addition, this work utilized an amyloid-focused

APP/PS1 model[31]; future studies in models with robust tau pathology will be important to determine

how these circuit-level mechanisms interact with intracellular neurofibrillary pathology.

Together, these findings demonstrate that amyloid plaques are associated with state-dependent,

long-range spatial gradients of neuronal activity and a fundamental reorganization of the hippocampal

map. Such network-level collapse provides a mechanistic link between focal amyloid deposits and

widespread cognitive impairment in Alzheimer's disease, and suggests general principles by which focal

pathology can remap brain circuits in neurodegenerative diseases.

References

1. Busche, M. A. *et al.* Clusters of hyperactive neurons near amyloid plaques in a mouse model of Alzheimer's disease. *Science* **321**, 1686–1689 (2008).

2. Busche, M. A. *et al.* Critical role of soluble amyloid-β for early hippocampal hyperactivity in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* **109**, 8740–8745 (2012).

3. Janus, C., Flores, A. Y., Xu, G. & Borchelt, D. R. Behavioral abnormalities in APPSwe/PS1dE9 mouse model of AD-like pathology: comparative analysis across multiple behavioral domains. *Neurobiol Aging* **36**, 2519–2532 (2015).

4. Kilgore, M. *et al.* Inhibitors of class 1 histone deacetylases reverse contextual memory deficits in a mouse model of Alzheimer's disease. *Neuropsychopharmacology* **35**, 870–880 (2010).

5. Lalonde, R., Kim, H. D., Maxwell, J. A. & Fukuchi, K. Exploratory activity and spatial learning in 12-month-old APP(695)SWE/co+PS1/DeltaE9 mice with amyloid plaques. *Neurosci Lett* **390**, 87–92 (2005).

6. Lester, A. W., Moffat, S. D., Wiener, J. M., Barnes, C. A. & Wolbers, T. The Aging Navigational System. *Neuron* **95**, 1019–1035 (2017).

7. Busche, M. A. & Konnerth, A. Neuronal hyperactivity--A key defect in Alzheimer's disease? *Bioessays* **37**, 624–632 (2015).

8. Arbel-Ornath, M. *et al.* Soluble oligomeric amyloid-β induces calcium dyshomeostasis that precedes synapse loss in the living mouse brain. *Mol Neurodegener* **12**, 27 (2017).

9. Cleary, J. P. *et al.* Natural oligomers of the amyloid-beta protein specifically disrupt cognitive function. *Nat Neurosci* **8**, 79–84 (2005).

10. Cline, E. N., Bicca, M. A., Viola, K. L. & Klein, W. L. The Amyloid-β Oligomer Hypothesis: Beginning of the Third Decade. *J Alzheimers Dis* **64**, S567–S610 (2018).

11. Faucher, P., Mons, N., Micheau, J., Louis, C. & Beracochea, D. J. Hippocampal Injections of

Oligomeric Amyloid β-peptide (1-42) Induce Selective Working Memory Deficits and Long-lasting

Alterations of ERK Signaling Pathway. *Front Aging Neurosci* **7**, 245 (2015).

12. Ferreira, S. T., Lourenco, M. V., Oliveira, M. M. & De Felice, F. G. Soluble amyloid-β oligomers as

synaptotoxins leading to cognitive impairment in Alzheimer's disease. *Front Cell Neurosci* **9**, 191

(2015).

13. Nicole, O. *et al.* Soluble amyloid beta oligomers block the learning-induced increase in

hippocampal sharp wave-ripple rate and impair spatial memory formation. *Sci Rep* **6**, 22728

(2016).

14. Shankar, G. M. *et al.* Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair

synaptic plasticity and memory. *Nat Med* **14**, 837–842 (2008).

15. Zhou, H. *et al.* Cholinergic modulation of hippocampal calcium activity across the sleep-wake

cycle. *Elife* **8**, e39777 (2019).

16. Zhou, H. *et al.* Disruption of hippocampal neuronal circuit function depends upon behavioral state

in the APP/PS1 mouse model of Alzheimer's disease. *Sci Rep* **12**, 21022 (2022).

17. Krnjević, K. Cellular and synaptic actions of general anaesthetics. *Gen Pharmacol* **23**, 965–975

(1992).

18. Munting, L. P. *et al.* Influence of different isoflurane anesthesia protocols on murine cerebral

hemodynamics measured with pseudo-continuous arterial spin labeling. *NMR Biomed* **32**, e4105

(2019).

19. Lustig, B., Wang, Y. & Pastalkova, E. Oscillatory patterns in hippocampus under light and deep

isoflurane anesthesia closely mirror prominent brain states in awake animals. *Hippocampus* **26**,

102–109 (2016).

20. Knowles, R. B. *et al.* Plaque-induced neurite abnormalities: implications for disruption of neural

networks in Alzheimer's disease. *Proc Natl Acad Sci U S A* **96**, 5274–5279 (1999).

21. Kurucu, H. *et al.* Inhibitory synapse loss and accumulation of amyloid beta in inhibitory

presynaptic terminals in Alzheimer's disease. *Eur J Neurol* **29**, 1311–1323 (2022).

22. Silva, A. & Martínez, M. C. Spatial memory deficits in Alzheimer's disease and their connection to

cognitive maps' formation by place cells and grid cells. *Front Behav Neurosci* **16**, 1082158 (2022).

23. Girardeau, G. & Lopes-Dos-Santos, V. Brain neural patterns and the memory function of sleep.

Science **374**, 560–564 (2021).

24. Cirrito, J. R. *et al.* Synaptic activity regulates interstitial fluid amyloid-beta levels in vivo. *Neuron*

48, 913–922 (2005).

25. Bero, A. W. *et al.* Neuronal activity regulates the regional vulnerability to amyloid-β deposition.

Nat Neurosci **14**, 750–756 (2011).

26. Marks, W. D., Yokose, J., Kitamura, T. & Ogawa, S. K. Neuronal Ensembles Organize Activity to

Generate Contextual Memory. *Front Behav Neurosci* **16**, 805132 (2022).

27. Megías, M., Emri, Z., Freund, T. F. & Gulyás, A. I. Total number and distribution of inhibitory and

excitatory synapses on hippocampal CA1 pyramidal cells. *Neuroscience* **102**, 527–540 (2001).

28. Montero-Crespo, M., Domínguez-Álvaro, M., Alonso-Nanclares, L., DeFelipe, J. & Blazquez-

Llorca, L. Three-dimensional analysis of synaptic organization in the hippocampal CA1 field in

Alzheimer's disease. *Brain* **144**, 553–573 (2021).

29. Xie, L. *et al.* Sleep drives metabolite clearance from the adult brain. *Science* **342**, 373–377 (2013).

30. Roh, J. H. *et al.* Disruption of the sleep-wake cycle and diurnal fluctuation of β-amyloid in mice

with Alzheimer's disease pathology. *Sci Transl Med* **4**, 150ra122 (2012).

31. Jankowsky, J. L. *et al.* Co-expression of multiple transgenes in mouse CNS: a comparison of

strategies. *Biomol Eng* **17**, 157–165 (2001).

32. Li, H. *et al.* Impaired hippocampal circuit function underlying memory encoding and consolidation

precede robust Aβ deposition in a mouse model of Alzheimer's disease. *bioRxiv* 2024.05.26.595168

(2025) doi:10.1101/2024.05.26.595168.

33. Klunk, W. E. *et al.* Imaging brain amyloid in Alzheimer's disease with Pittsburgh Compound-B.

Ann Neurol **55**, 306–319 (2004).

34. Lee, Y. F. *et al.* Optogenetic targeting of astrocytes restores slow brain rhythm function and slows

Alzheimer's disease pathology. *Sci Rep* **13**, 13075 (2023).

35. Zhou, P. *et al.* Efficient and accurate extraction of in vivo calcium signals from microendoscopic

video data. *Elife* **7**, e28728 (2018).

36. Markus, E. J., Barnes, C. A., McNaughton, B. L., Gladden, V. L. & Skaggs, W. E. Spatial

information content and reliability of hippocampal CA1 neurons: effects of visual input.

Hippocampus **4**, 410–421 (1994).

37. Myronenko, A. & Song, X. Point set registration: coherent point drift. *IEEE Trans Pattern Anal*

Mach Intell **32**, 2262–2275 (2010).

38. Gatti, A. A. & Khallaghi, S. PyCPD: Pure NumPy Implementation of the Coherent PointDrift

Algorithm. *JOSS* **7**, 4681 (2022).

     Figure Legends

Figure 1. Plaque topography modulates hippocampal ensemble activity across behavioral states.

a, Experimental workflow: simultaneous 1-photon (1P) miniscope hippocampal dynamic calcium

imaging and local field potential (LFP) recording in CA1 of freely behaving APP/PS1 mice, followed

by *in vivo* 2-photon (2P) imaging of amyloid plaques. **b**, Representative LFPs (top) and corresponding

calcium transients (bottom) during exploratory behavior (RUN), quiet wakefulness (QW), slow-wave

sleep (SWS), and rapid eye movement sleep (REM) sleep. Neurons are stratified by activity: high-

activity (red), moderate (green), and quiescent (blue). **c**, 1P-imaged GCaMP6f somatic activity (left,

white) co-registered with 2P-imaged Methoxy-X04 labeled plaques (middle, red). Right: Overlay of

somas (green) and plaques (red) with neurons from (**b**) indicated. **d**, Proportion of quiescent (blue),

moderate (green), and high-activity (red) neurons as a function of distance from the nearest plaque (n=7

mice) across behavioral states. No significant correlations were observed between distance and the

proportion of cells in any activity stratum (Spearman correlation, Bonferroni-adjusted). Data are mean

± s.e.m. (n=7 mice, 3762 cells).

Figure 2. Behavioral state dictates the spatial influence of amyloid plaques on neuronal activity.

a, Representative spatial maps of hippocampal calcium activity co-registered with amyloid plaques (white) across behavioral states. Color bars indicate the state-specific thresholds for quiescent and high-activity cells. Scale bar, 100 μm. **b,** Variations in calcium activity as a function of distance from the nearest plaque for high-activity, quiescent, and moderately active strata. Proximity to plaques is associated with distinct activity profiles across behavioral states, extending over hundreds of microns (n=7 mice, 3762 cells; mixed effects model, p<0.0001). Data represent the model's best fit $\pm$ 95% confidence intervals.

Figure 3. Plaque size and laminar position determine the spatial gradient of neuronal hyperactivity. a, Representative 2-photon images of amyloid plaques (red) relative to GCaMP6F-expressing neurons in the CA1 pyramidal cell layer (green). (**Top**) Laminar view distinguishing 'cell layer' versus 'dendritic' plaques. (**Bottom**) Superior (XY) view. Scale bar, 100μm. **b**, Distribution of plaque diameters for cell layer and dendritic plaque populations (n=7 mice). The solid gray line indicates the threshold determined by k-means clustering. **c**, Spatial decay of calcium activity as a function of plaque diameter and laminar location (n=7 mice, 3762 cells; mixed effects model, p=0.0039). Large plaques in the pyramidal cell layer are associated with the steepest activity gradients. Fits are computed for moderate activity cells during slow-wave sleep (SWS) at a representative edge distance of 136.6 μm. Data represent the model's best fit $\pm$ 95% confidence intervals.

Figure 4. Amyloid plaques are associated with the aberrant spatial clustering of hippocampal

place cells. a, Spatial tuning of a representative CA1 neuron: Calcium transients (red vertical bars) are

plotted against mouse position on a linear track. **b**, Heatmaps of place cell activity sorted by place field

location for inbound (left-right; LR) and outbound (right-left; RL) directions. Color bar indicates

probability. Yellow indicates the peak field location. **c**, Reduced proportion of place cells in APP/PS1

mice (n=7) compared to age-matched wild-type (WT) controls (n=4). Box plots indicate median (center

line), $25^{th}/75$th percentiles (bounds), and full range (whiskers). **d**, Representative spatial map showing

co-registered place cells (cyan), non-place cells (gray), and amyloid plaques (white). **e**, (**Top**)

Distribution of inter-cell distances for WT (blue) and APP/PS1 (red) place cells, showing an over-

representation of short distances in APP/PS1 mice (vertical line, 75 μm). (**Bottom**) Empirical

cumulative distribution function (eCDF) of inter-cell distances ($p < 10^{-14}$, Kolmogorov-Smirnov test)

(Inset: full eCDF). **f**, In APP/PS1 mice, the probability of a neuron being a place cell increases with

proximity to the nearest amyloid plaque ($p < 0.0001$, mixed effects model). This effect is modulated by

distance from the edge of the field of view (inset) (mixed effects model, $p < 0.0001$). Place cell

probability is higher near dendritic layer plaques than cell layer plaques ($p < 0.005$). * $p < 0.05$. WT: n=4

mice (559 place cells, 2468 total cells); APP/PS1: n=7 mice (410 place cells, 3762 total cells). Data are

best fit ± 95% confidence intervals.

Figure 5. Pre-existing place cell topography does not predict future amyloid deposition sites. a,

Experimental workflow for longitudinal co-registration. Angiogram fiducials obtained via 1-photon

(1P) imaging prior to plaque formation (left) were co-registered with 2-photon (2P) angiograms after

plaque deposition (middle) to map initial dynamic calcium activity to future plaque topography. **b**,

Proximity to nearest future plaque sites did not differ significantly between place cells and nonplace

cells ($p > 0.05$, n.s.). Box plots indicate median (center line), $25^{th}/75$th percentiles (bounds), and full

range (whiskers). **c**, Relationship between pre-plaque neuronal activity and future plaque location.

Elevated calcium activity during slow-wave sleep (SWS), but not other behavioral states, was

associated with proximity to future amyloid plaques ($p=0.038$, mixed effects model). Data represent

best fit $\pm$ 95% confidence intervals (n=3 mice, 1080 cells).

347   Figure 2

349   Figure 3

353   Figure 5

Methods

All procedures were approved by the Institutional Animal Care and Use Committees of the Massachusetts General Hospital. The study was conducted in accordance with the ethical guidelines of the US National Institutes of Health.

Surgical preparation

The APP/PS1 mouse strain used for this research project, B6C3-Tg(APPswe,PSEN1dE9)85Dbo/Mmjax, RRID:MMRRC_034829-JAX, was purchased from the Mutant Mouse Resource and Research Center (MMRRC) at The Jackson Laboratory, an NIH-funded strain repository, and was donated to the MMRRC by David Borchelt, Ph.D., McKnight Brain Institute, University of Florida[31]. Surgical preparation has been previously prescribed[32]. Briefly, APP/PS1 mice and their age/sex matched non-transgenic control littermates underwent injection of 1 μL AAV-hSyn-GCaMP6f (Addgene, titer > $7x10^{12}$ vg/ml) into the hippocampal CA1 region (AP -2.1 mm, ML -1.65 mm, DV 1.4 mm from bregma) under anesthesia (isoflurane 1.5-2%). A gradient index (GRIN) lens (1 mm diameter, AP -2.1 mm, ML -1.50 mm, DV -1.2 mm from bregma) and a custom drivable probe holding 7 stereotrodes (AP -2.8 mm, ML 2.6 mm, -36° relative to the AP axis, -39° relative to the inter-aural axis; DV 1.2 mm) were implanted ipsilaterally 2 weeks later with cerebellar ground. The microscope baseplate was attached three weeks later. Animals were housed in individual cages with a 12 hr light/12hr dark standard light cycle.

1P Dynamic calcium imaging and electrophysiology recordings

The Inscopix™ nVista3 mini-microscope and acquisition system (IDAS) were used for dynamic calcium imaging, sampling at 20 fps. Local field potential (LFP) (Neuralynx™) was sampled at 32 kHz followed by 0.5-900 Hz filtering and down sampling to 1000 Hz. Animal position was measured with overhead camera tracking of two LED diodes mounted on the headstage.

Electrophysiological recordings and position sampling were acquired continuously. Simultaneous

dynamic calcium imaging was acquired in 10-min ON and 5-min OFF blocks to avoid photobleaching

and was synchronized with electrophysiological data with TTL pulses.

Behavior and sleep staging

Data were acquired as animals explored a linear track without reward and were then moved to

their home cage within the same recording room to sample subsequent quiet wakefulness (QW), slow-

wave sleep (SWS), and REM sleep. Animals were exposed to the track for 1 to 3 days (2.5 ± 0.3 days)

before experiments. Cortical delta (1-4 Hz), hippocampal theta (4-10 Hz) and ripples (100-300 Hz),

and mouse position and posture were used to identify behavioral states: Run was defined as animals

explored the linear track with speeds exceeding 1 cm/sec and was characterized by a strong theta

oscillation; QW was identified when animals were stationary (< 0.5 cm/sec) for at least 5 seconds with

their eyes open; SWS was identified by the presence of sharp wave ripples (SWRs) and high delta/theta

LFP power ratio (>2) in animals in a sleep posture; REM sleep was identified as epochs of high

theta/delta LFP power ratio (>2) in animals in a sleep posture[32].

2P imaging of amyloid plaques and acquisition of angiograms

Following 1P imaging of dynamic calcium activity, Methoxy-X04 (10 mg/kg, i.p.) was injected

to label amyloid plaques[33]. For 1P vascular imaging at least one day later, animals were then lightly

anesthetized with isoflurane (~1.5%; with concentration sufficient to inhibit dynamic calcium activity),

fluorescein dextran (0.3 ml, i.v.) was injected into the tail vein, and imaging was acquired. Animals

were then transferred to a 2P microscope and head-fixed for 2P vascular and amyloid plaque imaging

on a commercial multiphoton system as previously described[34] (Olympus Fluoview 1000MPE)

mounted on an Olympus BX61WI upright microscope. A mode-locked titanium/sapphire laser (MaiTai;

Spectra-Physics, Fremont, CA) was used to generate two-photon fluorescence with 800 nm excitation.

Detectors consisting of photomutliplier tubes (Hamamatsu, Ichinocho, Japan) collected light in the

following ranges: 380-480 and 500-540 nm. 2P imaging with 800 nm excitation of hippocampal CA1

was acquired through the GRIN lens, acquiring a stack of images across 165 to 344 μm spanning the

CA1 cell layer.

Data analysis. All analyses were performed using MATLAB, Python, or SAS unless otherwise

specified[32].

Calcium imaging: Inscopix Data Processing Software (IDPS, Inscopix, Palo Alto, California) was used

for movie preprocessing, spatial filtering, motion correction, normalization ΔF/F, and constrained non-

negative matrix factorization (CNMFE) cell identification[35]. Each identified cell was manually

examined and accepted with oval cell shape and high signal-noise-ratio. Calcium events were then

detected at 2 Z scores above the mean, with onset defined at 30% of peak amplitude[32]. Low rate cells

were defined as < 0.25 transients/min[32]; high rate cells were defined using a calcium event rate

threshold set at the top 5% of calcium rates for each behavioral state of each animal.

Place cell detection and analysis: Spatial tuning curves (3 cm bins) were constructed for each cell for

each running direction (LR and RL) using all detected calcium events acquired during running (speed >

1 cm/s). Spatial information was measured for each cell as the amount of information (in bits) conveyed

per calcium event, and was compared to 1000 shuffled spatial information distributions to test

significance (Monte Carlo p-value < 0.05)[36], and taken as the larger of the LR and the RL trajectories.

Single timepoint 2P-1P image co-registration and plaque characterization: Custom Python code was

used to co-register 1P and 2P angiograms, implementing a coherent point drift algorithm and Python

package pycpd 2.0.0[37,38]. In brief, landmark points marked on 1P and 2P angiograms were used to

compute the parameters for a similarity transformation (that included translation, rotation, and scaling)

that maximized their overlap. These transformation parameters were then applied to the 1P angiograms

and dynamic calcium imaging data to register them in the 2P space. Co-registration accuracy was

quantified by calculating the nearest-neighbor distance between corresponding features of the

transformed 1P and 2P angiogram datasets. Across the datasets, the root mean square error (RMSE)

was 1.9 ± 0.08 µm (mean $\pm$ s.e.m.; **Supplementary Fig. 1**). We measured individual plaque size as the

cross-sectional area (in µm^2) from these images using ImageJ software. Plaques were subsequently

dichotomized into "small" (radius < 10.7 µm^2) and "big" (radius ≥ 10.7 µm^2) categories based on k-

means clustering and further dichotomized by their laminar position (pyramidal cell layer vs. dendritic

layers; **Fig. 3b**).

<u>Young 1P- old 2P image co-registration</u>: The spatial alignment of 1P young calcium imaging data and 2P

amyloid plaque image stacks was achieved through a two-stage registration procedure. *Stage 1:*

Longitudinal 1P image co-registration. First, a similarity transformation was computed to align the initial

angiogram (1P young) with the follow-up angiogram (1P old). Landmark Selection: Corresponding

anatomical landmarks were identified on both the 1P young and 1P old angiograms. These landmarks

were used to compute the parameters for a first similarity transformation, which included translation,

rotation, and uniform scaling. The transformation parameters were optimized to maximize the spatial

overlap of the landmark points. The calculated transformation was then applied to the 1P young

angiogram and its associated dynamic calcium imaging data, registering them into the coordinate space

of the 1P old angiogram. *Stage 2: 2P-1P image co-registration.* To integrate the 1P and 2P datasets, a

second registration was performed. A new set of landmarks was selected from major vessels that

maintained a consistent structural configuration across the registered 1P young and 2P old angiograms.

A second similarity transformation was computed using these stable landmarks to maximize the overlap

between the registered 1P young angiogram and the 2P old angiogram. This second set of transformation

parameters was then applied to the registered 1P young angiogram and dynamic calcium imaging data to

map them into the 2P coordinate space. To evaluate the accuracy of this co-registration method, we

computed the nearest-neighbor distance between corresponding features of the transformed 1P young

and transformed 1P old angiograms (**Supplementary Fig. 4a**). Across all datasets, the RMSE of this

distance was 13.4 ± 0.2 µm (mean $\pm$ s.e.m.; **Supplementary Fig. 4d**). The vascular landmarks used for

this transformation covered $71.4 \pm 3.1\%$ of the total area containing imaged neurons and amyloid plaques

(**Supplementary Fig. 4b,c**).

Statistical analyses of neuronal activity – plaque relationships

<u>Neuronal activity as a function of proximity to amyloid plaques</u> was assessed using a backward

eliminated Mixed Effects Model for the dependent variable "Calcium Rate", using the "State" (Run,

QW, SWS, and REM) as a fixed predictor along with Type of Activity (Quiescent, Moderate, and

High), the Distance to the nearest plaque, and Edge Distance, including all 2-way interactions. The

random effect was Mouse.

The model was significant, revealing several significant 2-way interactions, including of greatest

interest, an interaction of distance to the nearest plaque and type of activity ($p<0.0001$) (**Fig. 2b,**

Supplementary Fig. 5a), an interaction of behavioral state and activity type ($p<0.0001$), an interaction

of distance to the nearest plaque and behavioral state ($p<0.0001$) (**Fig. 2b**), and a shallow but significant

interaction of edge distance and activity type ($p=0.0065$), consistent with the contribution of plaques

outside the field of view, such that with greater distance from the edge, the activity of hyperactive cells

slightly diminished, the activity of hypoactive cells slightly increased, while the activity of normally

active cells remained largely unchanged (**Supplementary Fig. 5b**), indicating that edge distance serves

only as a proxy for unobserved plaques beyond the imaging window. The proportion of the variance

accounted for by fixed effects in this model was 51.7%.

481   For **Fig. 1d**, the relationship between distance to the nearest plaque and the proportion of cells

482 in each activity class (high activity, moderate, quiescent) was assessed using Spearman correlations

483 computed separately for each behavioral state. P-values were Bonferroni-adjusted for multiple

484 comparisons.

485 Effect of plaque size and hippocampal laminar position on neuronal activity was assessed by adding

486 plaque size (big vs. small) and hippocampal laminar position (pyramidal cell layer vs. dendritic layers),

487 their interaction, and the interactions of these two terms with Distance to the nearest plaque and Edge

488 Distance to the previous activity model (which included State, Activity Type, Distance to Plaque, and

489 Edge Distance predictors).

490   In addition to the same significant 2-way interactions described above, the model detected a

491 significant 3-way interaction of plaque laminar position, plaque size, and distance to the nearest plaque

492 (p=0.0039). The model also revealed a small but significant interaction of plaque laminar position and

493 edge distance (p<0.0001), as well as a modest interaction of activity type and edge distance (p<0.0046)

494 similar to that observed in the previous model (**Supplementary Fig. 6**). The model accounted for

495 51.8% of the variance.

497 Place cell identity and place cell spatial information as a function of proximity to amyloid plaques were

498 assessed with two backward elimination mixed effects models: For both, the fixed effect predictors

499 were: Distance to the nearest plaque, the nearest plaque laminar position, nearest plaque size, and Edge

500 Distance (distance to the edge of the field of view, included to account for potential effects of

501 unobserved plaques outside the imaging area), and all two-way interactions. The Random Term was

502 Mouse.

503   The dependent variable for the first analysis was the binary variable of Place Cell Identity

504 (Yes/No; analyzed via a Generalized Linear Mixed Model assuming a binomial distribution and a logit

link function. This analysis revealed a significant interaction between distance to the nearest plaque and edge distance ($p<0.0001$; **Fig. 4f**). There was also a main effect for plaque laminar position ($p=0.005$).

The dependent variable for the second analysis was Spatial Information, where we found no significant effect of proximity to amyloid plaques (**Supplementary Fig. 3**).

Place cell identity as a function of proximity to future amyloid plaques was assessed with a mixed effects GLM for the binary dependent variable of pre-plaque Place Cell Identity (Yes/No; measured at timepoint 1). The fixed effect predictor was Distance to Nearest Future Plaque at timepoint 2. The Random Term was Mouse. The GLM assumed a binary distribution and a logit link function for the dependent variable. This analysis showed that place cell identity before plaques formed was not a significant predictor of future plaque location (**Fig. 5b**).

Effect of neuronal activity on proximity to future amyloid plaques was assessed with a mixed effects model for the dependent variable of Distance to Nearest Plaque at timepoint 2. Run, QW, SWS, and REM Calcium Rates (in separate analyses) were used as fixed predictors. The Random Term was Mouse. This analysis showed that higher activity in SWS – but not in Run, QW, or REM sleep – was weakly predictive of future plaque locations ($p=0.038$; **Fig. 5c**).

Acknowledgements

Funding:

National Institutes of Health grant R01 AG054551 (SNG)

National Institutes of Health grant R01 AG077611 (SNG)

Author contributions:

Conceptualization, Z.Z. and S.N.G.

Methodology, Z.Z., R.J.G, K.K., B.B., and S.N.G.

Software, Z.Z., H.L. N.G., and S.N.G.

Validation, Z.Z., H.L., and J.J.L.

Formal Analysis, Z.Z., J.J.L., H.L.

Investigation, Z.Z., H.L., and S.N.G.

Resources, Z.Z., K.K., B.B., and S.N.G.

Data Curation, Z.Z., H.L., and J.J.L.

Writing – Original Draft, Z.Z., and S.N.G.

Writing – Review & Editing, Z.Z., H.L., J.L., N.G., R.J.G., K.K., B.B., and S.N.G.

Visualization, Z.Z., H.L., J.J.L.

Supervision, S.N.G.

Project Administration, S.N.G.

Funding Acquisition, S.N.G.

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: The datasets generated and analyzed in the current study will be

available in the Harvard Dataverse repository.

Extended Data Figures

548
549

Supplementary Figure 1. Spatial co-registration of 1-photon and 2-photon imaging volumes.

a, **Representative co-registration of 1-photon (**1P) neuronal maps and 2-photon (2P) plaque

topography. Volumes were aligned via rigid transformation using hippocampal angiogram fiducials.

b, Precision of co-registration as measured by the root mean square error (RMSE) of the distances

between transformed 1P and imaged 2P vasculature (n = 7 co-registrations). Box plot indicate median

(center line), 25th/75th percentiles (bounds), and full range (whiskers).

561 **Supplementary Figure 2. Distribution of activity strata relative to plaque proximity using a 1%**

562 **hyperactivity threshold.** Proportion of high-activity (red), moderate (green), and quiescent (blue)

563 neurons as a function of distance from the nearest plaque across behavioral states. In this analysis,

564 high-activity cells are defined for each state as those cells in the top 1% of calcium event rates. Data are

565 presented as mean ± s.e.m. (n=7 mice, 3,762 cells).

575 **Supplementary Figure 3. Place cell spatial information does not depend on proximity to amyloid**

576 **plaques.** Linear least squares regression (r = 0.086, p=0.08; red line) and linear mixed-effects modeling

577 (p>0.05) reveal no relationship between a neuron's spatial information content and its distance to the

578 nearest plaque. Data represent individual place cells (n=410 cells from 7 mice).

Supplementary Figure 4. Longitudinal co-registration of hippocampal activity and future plaque topography. a, Mapping of hippocampal vasculature across time-points to align pre-plaque calcium activity with future plaque locations. Left: Representative 1-photon (1P) angiograms from a single mouse at 6.2 months (top) and 10.5 months of age (bottom). Middle: Automated identification of fiducial vessels. (Right) Superimposition of identified vessels using rigid body transformation (top) and the resulting spatial registration error (bottom). **b**, Representative field showing co-registered neuronal somas (6.2 months) and amyloid plaques (10.5 months). **c**, Vascular coverage across the co-registered field of view, illustrating the extent of the common imaging area. **d**, Precision of the longitudinal alignment as measured by the root mean square error (RMSE) of the distances between transformed young and old 1P vessels. Box plots indicate median (center line), 25th/75th percentiles (bounds), and full range (whiskers).

Supplementary Figure 5. State-dependent spatial gradients and edge-effect modeling of calcium

activity. a, High activity, moderate, and quiescent neurons exhibit distinct activity profiles as a function

of distance from the nearest plaque across behavioral states (n=7 mice, 3,762 cells; linear mixed

effects model, p<0.0001). **b**, Calcium activity as a function of distance from the Edge of the field of

view for each activity stratum (linear mixed effects model, p=0.0065). Data represent the model's best

fit ± 95% confidence intervals.

Supplementary Figure 6. Interaction between plaque laminar position and field-of-view edge

distance. Calcium activity as a function of distance from the Edge of the field of view, stratified by the

laminar position (cell layer vs. dendritic) of the nearest amyloid plaque (p<0.0001, linear mixed effects

model). Data represent the model's best fit ± 95% confidence intervals.