Regulator of G Protein Signaling 6 Negatively Regulates Platelet Activation and Arterial Thrombosis in Mice

Renat Roytenberg, Ph.D.[1#], Boyd R. Rorabaugh, Ph.D.[2#], Hong Yue, M.D., Ph.D.[1], Robbie Jividen, M.S.[1], Scott J. Cameron, M.D., Ph.D.[3,4,5], and Wei Li, M.D., Ph.D.[1]

1. Department of Biomedical Sciences, Joan C. Edwards School of Medicine at Marshall University, Huntington, WV.
2. Department of Pharmaceutical Sciences, Marshall University School of Pharmacy, Huntington, WV.
3. Heart, Blood, and Kidney Research, Cleveland Clinic Lerner College of Medicine, Cleveland, OH.
4. Department of Cardiovascular Medicine, Section of Vascular Medicine, Cleveland Clinic, Cleveland, OH
5. Department of Hematology, Taussig Cancer Institute, Cleveland Clinic, Cleveland, OH.

# Contributed equally.

Running title: RGS6 negatively regulates platelet function

Corresponding author:

Wei Li, MD, PhD, FAHA
Department of Biomedical Sciences, BBSC 241G
Joan C. Edwards School of Medicine
Marshall University
One John Marshall Drive, Huntington,
West Virginia 25755-9310, USA.
Phone: 1-304-696-7342
E-mail: liwe@marshall.edu

Abstract

Background: Platelet activation via G protein-coupled receptors (GPCRs) is central to arterial thrombosis. P2Y12 is a canonical Gi-coupled receptor mediating ADP-dependent platelet activation, yet the role of Regulator of G protein Signaling 6 (RGS6), a modulator of Gi signaling, in platelet function and thrombosis remains unclear.

Objectives: To determine the role of RGS6 in platelet activation and arterial thrombosis and to define its impact on $P2Y_{12}$/Gi signaling.

Methods: Arterial thrombosis was assessed using a FeCl□-induced carotid artery injury model in wild-type (WT) and $Rgs6^{-/-}$ mice. Platelet aggregation was measured ex vivo. Signaling pathways were analyzed by Western blot in ADP-stimulated platelets. $P2Y_{12}$/Gi signaling was further evaluated using a cAMP-responsive luciferase reporter assay in HEK293 cells.

Results: Male $Rgs6^{-/-}$ mice exhibited significantly accelerated thrombosis compared with WT controls. $Rgs6^{-/-}$ platelets showed enhanced ADP-induced aggregation, whereas collagen-induced aggregation was unchanged. In ADP-stimulated platelets, RGS6 deficiency altered signaling kinetics, characterized by delayed Akt phosphorylation and reduced PKA and VASP phosphorylation. In a heterologous cAMP-luciferase assay, RGS6 attenuated $P2Y_{12}$/Gi-mediated suppression of cAMP. Two-way ANOVA demonstrated significant effects of ADP and RGS6 expression on luciferase activity, with no interaction, indicating that RGS6 modulates signaling magnitude rather than agonist sensitivity. Pharmacologic inhibition of $P2Y_{12}$ with clopidogrel abolished the genotype-dependent difference in thrombosis in vivo.

Conclusions: RGS6 acts as a negative regulator of platelet $P2Y_{12}$/Gi signaling and thrombus formation. Loss of RGS6 enhances ADP-dependent platelet activation and accelerates arterial thrombosis, establishing RGS6 as an endogenous brake on platelet activation.

Key words: Platelet, G protein-coupled receptors, Regulator of G protein signaling 6, Thrombosis, Gi alpha subunit.

Introduction

Thrombotic events are a major cause of morbidity and mortality in the US and worldwide [1-3]. Platelet activation and increased reactivity at the site of vascular injury are the primary pathogenic components of thrombosis, which lead to vessel occlusion and causing myocardial infarction and ischemic stroke. Platelets have many surface receptors and, when activated, promoting platelet adhesion and initial activation in response to exposed collagen at the site of vascular injury [4-6]. Adherent, activated platelets release soluble agonists (e.g., ADP, thrombin, and thromboxane A_2 [TXA2]), which locally activate additional platelets via surface G protein-coupled receptors (GPCRs). The major platelet GPCRs that mediate platelet activation include ADP receptors $P2Y_{12}$ and $P2Y_1$, thrombin receptors protease-activated receptor (PAR) 1 and PAR4, and TXA2 receptor (TP) [5, 7, 8]. Consequently, the FDA has approved aspirin (non-selective COX inhibitor, which inhibits TXA2 production and thus reduces TP agonism), clopidogrel and prasugrel ($P2Y_{12}$ inhibitors), and vorapaxar (PAR1 inhibitor) as anti-platelet/thrombotic drugs, which are clinically indicated for both primary or secondary prevention of platelet-mediated thrombotic events [9-12]. However, these drugs have major side effects, including injury to the gastrointestinal mucosa, thrombocytopenia, and systemic hemorrhage [3, 9-12]. In addition, some patients do not respond to these regimens and have a high incidence of recurrent thrombosis [3]. Therefore, defining the molecular mechanisms for platelet activation in thrombus formation remains a significant clinical unmet need.

ADP and thrombin receptors are coupled to three major G proteins: $G_{\alpha q}$, $G_{\alpha 12/13}$, and $G_{\alpha i}$ [13, 14]. ADP-induced platelet activation requires concomitant signaling from both $P2Y_1$ and $P2Y_{12}$ receptors that couple to $G_{\alpha q}$ and $G_{\alpha i}$, respectively. Regulators of G protein signaling (RGS) proteins enhance the rate of GTP hydrolysis by the Gα subunit of heterotrimeric G proteins. This provides a mechanism for cells to control the magnitude and duration of signaling through GPCRs [15, 16]. Twenty canonical RGS proteins have been identified [17]. Members of the RGS protein family exhibit unique tissue distributions and selectivity for different Gα isoforms, making them attractive therapeutic drug targets. Previous work has demonstrated that RGS10, RGS16, and RGS18 are expressed in platelets where they regulate platelet activation and thrombosis [18-22]. RGS6 has also been identified in platelets at both the mRNA [23, 24] and protein levels [25, 26]. Notably, platelet RGS6 expression was reduced by approximately 30% in patients with COVID-19, irrespective of ICU admission status [23]. COVID-19 patients are a high-risk cohort to develop thrombosis. Interestingly, a recent pharmacogenomic study in Caribbean Hispanic patients identified an intronic RGS6 variant that is associated with clopidogrel responsiveness, further supporting a role for RGS6 in platelet $P2Y_{12}$ signaling [27]. However, the function of RGS6 in platelets has not been explored. In this study, for the first time, we demonstrated that RGS6 regulates ADP-induced platelet aggregation and is an important regulator of thrombosis following vascular injury.

Methods

Animals
$Rgs6^{-/-}$ mice were a generous gift from Dr. Rory Fisher (University of Iowa School of Medicine, Iowa City, Iowa). The generation of these mice has been previously described [28]. The $Rgs6^{+/-}$

mouse strain has been backcrossed onto the wildtype (WT) C57BL6/J (Jackson Labs, Bar Harbor, Maine; stock #000664) background more than 15 generations. $Rgs6^{-/-}$ mice were maintained in homozygous inbreeding. WT ($Rgs6^{+/+}$) mice purchased from The Jackson Labs were bred and housed within the same room as the $Rgs6^{-/-}$ mice and used as controls. All procedures and manipulations of animals have been approved by the Institutional Animal Care and Use Committee of Marshall University (#: 1033528, PI: WL, and #: 1174948, PI: BR).

Materials

Platelet agonists including ADP (P/N 384) and collagen (P/N 385) were purchased from Chrono-log (Havertown, PA). Antibodies to phosphorylated AKT (4060S), pan-AKT (2920S), phosphorylated PKA (4781S), pan-PKA (5842S), HRP-conjugated Pan-actin (12748S), HRP Conjugated anti-Rabbit (7074S) or mouse (7076S) IgG secondary antibody were purchased from Cell Signaling Technology (Danvers, MA). Antibodies to $P2Y_{12}$ (ab184411, ab183066) were purchased from abcam (Waltham, MA). All other chemical reagents were purchased from Millipore Sigma (Burlington, MA) except where specifically indicated.

Murine FeCl₃-injury-induced carotid artery thrombosis model

The ferric chloride ($FeCl_3$)-injury-induced carotid artery thrombosis model has been described previously [29, 30]. Briefly, mice of both sexes, 8 to 16 weeks old, were anesthetized by a mixture of ketamine/xylazine (100/10 mg/kg) via intraperitoneal injection. Platelets were labeled through direct jugular vein injection of 100 μl of rhodamine 6G solution (Sigma 252433-1G, 0.5 mg/ml in saline, 0.2 μm filtered). The carotid artery was exposed, and injury was induced by topically applying a piece of filter paper (1 x 2 mm) saturated with 7.5% $FeCl_3$ solution for 1 minute. Thrombus formation was observed in real-time using intravital microscopy with a Leica DM6 FS fluorescent microscope (Deerfield, IL, USA) attached to a Gibraltar Platform. Video imaging was conducted using a QImaging Retiga R1: 1.4 Megapixel Color CCD camera system with mono color mode (Teledyne Photometrics, Tucson, AZ, USA) and StreamPix version 7.1 software (Norpix, Montreal, Canada). The endpoints were set as 1) blood flow has ceased for > 30 seconds, or 2) occlusion is not seen 30 minutes after $FeCl_3$ injury. In the second case, 30 minutes was assigned as the data value for statistical analysis.

A different cohort of mice were fed a clopidogrel solution (clopidogrel bisulfate in ethanol, serially diluted in saline) via oral gavage at 1 mg/kg (uncertain dose to produce a significant antithrombotic effect) [31], once per day for 72h before experimentation.

Tail bleeding assay

Tail bleeding assay was conducted as previously described [31, 32]. Briefly, 10-12-week-old male mice were anesthetized with ketamine/xylazine (100/10 mg/kg). One centimeter from the tail tip was transected using a sharp scalpel, and the tail was immediately immersed into 37 °C warm saline. Bleeding time was recorded from the moment of tail transection until bleeding completely ceased, which is named as 1st bleed. After the 1st bleeding, the wound was nicked with the scalpel, the tail was immediately immersed in a new tube containing warm saline, and the time to bleeding cessation was recorded, which is designated as the 2nd bleed.

If bleeding persisted for longer than 6 minutes (3 times the mean bleeding time in WT mice), a gentle pressure was applied to the tail for 2 minutes to achieve hemostasis. For statistical analysis, a bleeding time of 6 minutes was assigned to these mice.

Whole blood count and platelet isolation
Mice were anesthetized with ketamine/xylazine (100/10 mg/kg), and 0.9 - 1 mL whole blood was collected through inferior vena cava puncture using 0.109 M sodium citrate as an anticoagulant. Blood cells were counted immediately using Hemavet 950FS. Modified Tyrode's buffer (concentration of components in mM: 137 NaCl, 2.7 KCl, 12 $NaHCO_3$, 0.4 NaH2PO$_4$, 5 HEPES, 0.1% glucose, and 0.35% BSA, pH 7.2) was added to the collected whole blood at 0.7 volumes, mixed, and then platelet-rich plasma (PRP) was isolated by centrifugation at 100 g for 10 min. The sediment containing red blood cells and leukocytes was further centrifuged at 13,000 rpm for 1 min to isolate platelet-poor plasma (PPP) [32].

Platelet aggregation assay
The platelet concentration in PRP was counted with a hemocytometer and adjusted to 2.5E+08/mL with PPP, and 0.4 ml of this platelet suspension was used for the platelet aggregation assay using Chrono-log 700 with an agitation speed of 1,200 rpm. $CaCl_2$/$MgCl_2$ was repleted at a final concentration of 1 mM immediately before adding a platelet agonist [31-34].

Evaluation of P2Y$_{12}$-mediated platelet signaling activation
Platelets in PRP (2.5E+08/ml) were pooled from 6 mice, divided into 4 aliquots, and then stimulated with 2.5 μM ADP for 0, 1, 3, or 5 min [34]. Platelet activation was stopped by adding a final concentration of 1 mM EDTA and 0.5 µM PGE1 to the reaction mixture. The platelets were then pelleted by centrifugation at 13,000 RPM for 15 seconds, immediately lysed in radioimmunoprecipitation assay (RIPA) buffer containing 1 x Halt™ Protease and Phosphatase Inhibitor Single-Use Cocktail, EDTA-Free (100 X stock solution, ThermoFisher, Cat# 78443). Protein concentration was determined using a Bio-Rad Protein assay. AKT, PKA, and VASP phosphorylation were assessed by immunoblotting assays.

cAMP-luciferase reporter assay
The GloResponse™ CRE-luc2P HEK293 cell line was purchased from Promega (Cat# 8500, Madison, WI) and has been used in previous studies [35]. HEK293 cells were seeded at 10^5 cells/well in 24-well plates in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (Atlanta Biologicals Inc, Flowery Branch, GA, USA) and 1 X Antibiotic-Antimycotic (Gibco). Six hours after seeding, cells were transfected using Lipofectamine™ 3000 (Cat# L3000-015, Thermo Fisher Scientific, USA) according to the manufacturer's protocol. Cells were transfected with pcDNA3.1-P2Y$_{12}$, pcDNA3.1-*Rgs6* (cDNA Resource Center, Bloomsburg, PA), or their combination, along with Renilla plasmids. Empty pcDNA3.1 vector was used to equalize the total DNA amount (500 ng/well) transfected.

After 24h, the medium was replaced with serum-free DMEM. Following a 4h incubation in serum-free DMEM, cells were treated with varying concentrations of ADP or 0.9% saline for 1h at 37°C. Cells were then lysed in passive lysis buffer provided by the Dual-Luciferase® Reporter Assay System (Cat #: E1910, Promega, USA). Bioluminescence readings for firefly luciferase and Renilla were obtained using a SpectraMax L microplate reader operated with

SoftMax Pro7 software (Molecular Devices). Luciferase activity was normalized to the corresponding Renilla signal and further normalized with the average value of cells transfected with empty pCDNA3.1.

Statistics
Data are expressed as mean ± SEM. Results were analyzed by 2-tailed Student's *t* test, Mann Whitney test, or 2-way/1-way ANOVA with Bonferroni post-hoc test for multiple comparisons using GraphPad Prism (version 10.2.2). In some cases, data were analyzed by Log-rank test using the Kaplan-Meier survival curve. $P < 0.05$ was considered statistically significant.

Results

***Rgs6* gene deficiency significantly enhanced thrombosis in male mice.**
To evaluate whether RGS6 regulates thrombosis, we employed the 7.5% FeCl□-induced carotid artery injury model [29, 30] to assess in vivo thrombus formation in WT and *Rgs6*$^{-/-}$ mice. As shown in **Fig.1A**, loss of RGS6 did not appreciably alter the initial phase of platelet adhesion and aggregation at the site of vascular injury, as observed 1 minute after injury. In contrast, the subsequent wave of platelet accumulation was markedly accelerated in *Rgs6*$^{-/-}$ male mice, resulting in a significantly shorter time to occlusive thrombus formation compared with WT controls. Overall, RGS6 deficiency accelerated the time to form an occlusive thrombus in male mice (**Fig. 1B**). Interestingly, while it was not statistically different, *Rgs6* deficiency in females tended to prolong the time to form an occlusive thrombus (**Fig. 1C**). However, due to the absence of statistical significance in females, all subsequent experiments were conducted using only male mice.

RGS6 deficiency did not affect complete blood cell counts (**Table 1**), suggesting that the prothrombotic phenotype observed in male mice was not attributable to alterations in circulating blood cell components. RGS6 deficiency also did not affect tail bleeding time (**Supplemental Fig. I**).

RGS6 deficiency significantly enhanced ADP-induced platelet aggregation.
We next investigated whether RGS6 affects platelet aggregation in response to two conventional platelet agonists, collagen and ADP. PRP was prepared from WT and *Rgs6*$^{-/-}$ mouse blood by centrifugation, and platelet counts were normalized to 2.5×10^8/mL using PPP. For each aggregation assay, 400 µL of the adjusted platelet suspension was used. As shown in **Fig. 2A**, collagen (1 µg/ml)-induced platelet aggregation was similar between WT and *Rgs6*$^{-/-}$ platelets. In contrast, stimulation with ADP produced significantly greater platelet aggregation in *Rgs6*$^{-/-}$ platelets than in WT controls (**Fig. 2B**). These findings suggest that loss of RGS6 preferentially augments ADP-mediated platelet activation rather than broadly increasing platelet responsiveness.

RGS6 deficiency enhances ADP-induced suppression of the cAMP/PKA pathway in platelets.
Since RGS6 is a potent negative regulator of Gi signaling [36] and P2Y$_{12}$ is a well-defined Gi-coupled platelet GPCR [37], whereas coupling of PAR1 and PAR4 to Gi in platelets has not been clearly established [38], we hypothesized that RGS6 may serve as a key regulator of P2Y$_{12}$

signaling. Consistent with this, ADP-induced platelet aggregation was significantly enhanced in $Rgs6^{-/-}$ platelets (Fig. 2B). To further investigate this, we examined the effect of RGS6 on platelet signaling in response to ADP stimulation. Platelets pooled from 6 WT and 6 $Rgs6^{-/-}$ mice were aliquoted and stimulated with ADP over a time course and effectors of the $P2Y_{12}$/Gi signaling axis were analyzed by Western blot.

As shown in **Fig. 3A**, RGS6 deficiency did not affect $P2Y_{12}$ expression on platelets. Phosphorylation of AKT, a downstream effector of PI3K and a common marker of $P2Y_{12}$-dependent signaling [31, 39], increased rapidly in WT platelets but was delayed and remained at low levels in $Rgs6^{-/-}$ platelets (Fig. 3A & **3B**). In contrast, phosphorylation of PKA (**Fig 3C & 3D**) and its substrate vasodilator□stimulated phosphoprotein phosphorylation (p-VASP, Fig. 3C and **3E**), a well-established readout of cAMP/PKA signaling and an inverse marker of $P2Y_{12}$ activity [40], were markedly reduced in $Rgs6^{-/-}$ platelets following ADP stimulation, indicating enhanced suppression of the cAMP/PKA pathway. Together, these data indicate that RGS6 deficiency alters the profile of ADP-induced $P2Y_{12}$ signaling. Specifically, loss of RGS6 enhances Gi-mediated suppression of the inhibitory cAMP/PKA pathway.

RGS6 modulates $P2Y_{12}$/G_i signaling
To further demonstrate that RGS6 regulates platelet ADP/$P2Y_{12}$ signaling, we performed a cAMP reporter assay using HEK293 cells expressing a luciferase reporter under the control of a cAMP-responsive element (CRE). Plasmids encoding murine RGS6 and $P2Y_{12}$ (and Renilla luciferase) were transfected into HEK293 cells and luciferase/Renilla activity was measured following ADP stimulation. Data from all experimental groups were normalized to cells transfected with empty vector (pcDNA3.1) alone.

Forskolin treatment induced more than a 10 fold increase in luciferase activity (**Supplemental Figure II**), confirming that the assay system was functional. Although HEK293 cells have been reported to express P2Y receptors [41], ADP did not significantly affect luciferase activity in cells transfected with pCDNA3.1 (**Fig. 4**, blue bar, $p = 0.28$), suggesting limited functional endogenous P2Y signaling under these conditions. Overexpression of $P2Y_{12}$ resulted in a modest reduction in luciferase activity following ADP stimulation (light blue, $p = 0.089$), consistent with Gi-mediated inhibition of cAMP signaling. Expression of RGS6 alone significantly increased basal level luciferase activity, which is reduced by ADP treatment (brown bar, $p = 0.033$), supporting the presence of low-level endogenous Gi signaling. Co-expression of RGS6 with $P2Y_{12}$ significantly increased luciferase activity compared with $P2Y_{12}$ alone at all ADP concentrations tested (0, 10, and 20 μM; $p = 0.0009$, 0.008, and 0.02, respectively), indicating attenuation of $P2Y_{12}$ signaling by RGS6.

Two-way ANOVA demonstrated significant main effects of ADP ($p = 0.0038$) and gene expression ($p = 0.0017$), with no significant interaction ($p = 0.533$), suggesting that the effect of RGS6 on $P2Y_{12}$ signaling is consistent across agonist concentrations. Together, these data support a role for RGS6 as a negative regulator of $P2Y_{12}$-dependent Gi signaling.

RGS6 regulates in vivo thrombosis through a $P2Y_{12}$-dependent mechanism
To determine whether the prothrombotic phenotype observed in $Rgs6^{-/-}$ mice was functionally dependent on $P2Y_{12}$/G_i signaling, WT and $Rgs6^{-/-}$ mice were treated with clopidogrel (1 mg/kg)

once per day for 72h and then subjected to the FeCl□-induced carotid artery injury model [31]. As shown in **Fig. 5**, clopidogrel treatment markedly prolonged the time to form an occlusive thrombus in both strains and diminished the difference in thrombotic responses between male WT and $Rgs6^{-/-}$ mice. These findings demonstrate that the prothrombotic phenotype associated with RGS6 deficiency is largely dependent on $P2Y_{12}$ signaling, consistent with the role of RGS6 as a regulator of Gi. These findings support the interpretation that the prothrombotic phenotype associated with RGS6 deficiency is largely $P2Y_{12}$-dependent.

Discussion

RGS proteins regulate GPCR by accelerating the rate of GTP hydrolysis by Gα subunits [42]. RGS6 has not been extensively studied in platelets. In this study, we identified RGS6 as a negative regulator of platelet-intrinsic arterial thrombosis. Genetic deletion of $Rgs6$ accelerated thrombus formation in vivo and enhanced ADP-induced platelet aggregation ex vivo. Importantly, pharmacologic inhibition of $P2Y_{12}$ with clopidogrel abolished the genotype-dependent difference in thrombosis in $Rgs6^{-/-}$ and WT mice, indicating that the prothrombotic phenotype associated with RGS6 deficiency is $P2Y_{12}$-dependent.

Our findings are consistent with clinical observations in COVID-19, a condition associated with increased thrombotic risk and reduced platelet RGS6 expression (~30%) [23]. In addition, a prior study identified the rs9323567C>T SNP within an intronic region of RGS6. Although RGS6 expression was not examined in that study, such variants are often associated with reduced gene function. Our finding may therefore provide a mechanistic explanation for the reported association between the RGS6 rs9323567C>T variant and decreased $P2Y_{12}$ Reaction Units (PRU) in patients treated with clopidogrel [27]. Moreover, these results raise the possibility that reduced RGS6 activity may contribute to variability in $P2Y_{12}$ inhibitor responsiveness, which could partially explain the limited clinical benefit observed in studies of COVID-19-related mortality [43].

Importantly, the cAMP-responsive luciferase assays demonstrated an inverse, dose-dependent relationship between ADP and $P2Y_{12}$/Gi signal, increased luciferase activity in the presence of RGS6, and no significant interaction between RGS6 expression and the dose-response. These data support our in vitro signaling findings, indicating that RGS6 suppresses the $P2Y_{12}$/Gi pathway. The lack of a significant RGS6 x ADP interaction suggests that RGS6 does not alter agonist sensitivity but instead modulates the overall magnitude of $P2Y_{12}$/Gi signaling. This interpretation is consistent with our in vivo thrombosis studies: under basal conditions, loss of RGS6 increases the ADP/$P2Y_{12}$ signal output, thereby accelerating occlusive thrombus formation (**Fig. 6**), whereas pharmacologic $P2Y_{12}$ blockade using clopidogrel abolishes this difference.

In addition to RGS6, other RGS proteins, including RGS10 [44], RGS18 [19], and RGS16 [21, 45], have been reported to target Gi and regulate platelet activation (**Table 2**). Although ADP-induced aggregation was increased in $Rgs6^{-/-}$ platelets, AKT phosphorylation was delayed and reduced. However, PKA and VASP phosphorylation were reduced at all time points following ADP stimulation. These findings suggest that RGS6-mediated modulation of the

P2Y$_{12}$/Gi/cAMP/PKA pathway may play a more prominent role than Akt signaling in regulating platelet activation.

Our results also support a potential sex-dependent role for RGS6 in thrombosis. Female $Rgs6^{-/-}$ mice exhibited a trend opposite to that observed in males, with prolonged occlusion times compared with WT mice. In our recent study, we found that P2Y$_{12}$ forms complexes with the Na$^+$/K$^+$-ATPase α1 subunit, and α1 haplodeficiency significantly inhibits thrombosis in males but not in females [32]. Given that female platelets express higher levels of α1 than male platelets, and female platelet are more sensitive to ADP [43], P2Y$_{12}$ signaling may be regulated in a context-dependent manner. These findings raise the possibility that RGS6 contributes to sex-specific regulation of platelet function, which warrants further investigation.

Members of the RGS family exhibit differences in their tissue distribution and have different specificities for various Gα isoforms [16]. Consequently, RGS proteins have emerged as potential new therapeutic targets. RGS inhibitors may provide a way to enhance the effects of endogenous agonists or may be combined with exogenous agonists to produce responses with greater tissue selectivity than can be achieved with an agonist alone [46]. Prior work suggests that RGS6 may provide a therapeutic target for the treatment of alcohol use disorder [47, 48], Parkinson's disease [49], and cardiac ischemic injury [50]. However, the data presented here suggest that pharmacotherapies that suppress RGS6 function may be limited by an increased risk of thrombosis-related morbidities such as myocardial infarction or ischemic stroke.

This study has several limitations. First, the lack of a reliable antibody for RGS6 limited direct assessment of protein expression, and mechanistic conclusions relied on downstream signaling readouts and a heterologous reporter system. While informative, this system represents a reductionist model and does not fully recapitulate the complexity of platelet signaling. Second, $Rgs6$ deletion was global, and contributions from non-platelet cell types cannot be excluded. However, the enhanced ADP-induced aggregation observed ex vivo and the normalization of thrombosis by P2Y$_{12}$ inhibition strongly support a platelet-intrinsic mechanism.

Conclusion
We identify RGS6 as a previously unrecognized negative regulator of platelet-intrinsic arterial thrombosis. Loss of RGS6 accelerated thrombus formation in vivo, enhanced ADP-induced platelet aggregation ex vivo, and altered signaling within the platelet ADP/P2Y$_{12}$ axis. cAMP reporter assays and pharmacologic inhibition studies indicate that RGS6 modulates the magnitude of P2Y$_{12}$/Gi signaling rather than altering agonist sensitivity. Together, these findings establish RGS6 as an important regulator of platelet function and thrombosis.

Sources of support: This work was supported in part by the following sources: the National Institutes of Health R15HL145573 and R01HL177493 (to WL), R01HL158801 (to SJC) and the NASA West Virginia Space Grant Consortium, NASA Agreement #80NSSC20M0055 (to RJ). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Competing Interests Statement:
None

Acknowledgements

We thank Dr. Rory Fisher (University of Iowa School of Medicine) for providing the $Rgs6^{-/-}$ mice used in this study.

References

Day ISCfWT. Thrombosis: a major contributor to the global disease burden. *J Thromb Haemost*. 2014; **12**: 1580-90. 10.1111/jth.12698.

Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012; **380**: 2197-223. 10.1016/S0140-6736(12)61689-4.

Jackson SP. Arterial thrombosis--insidious, unpredictable and deadly. *Nat Med*. 2011; **17**: 1423-36. 10.1038/nm.2515.

Ezumi Y, Shindoh K, Tsuji M, Takayama H. Physical and functional association of the Src family kinases Fyn and Lyn with the collagen receptor glycoprotein VI-Fc receptor gamma chain complex on human platelets. *The Journal of experimental medicine*. 1998; **188**: 267-76. 10.1084/jem.188.2.267.

Furie BC, Furie B. Tissue factor pathway vs. collagen pathway for in vivo platelet activation. *Blood Cells Mol Dis*. 2006; **36**: 135-8. 10.1016/j.bcmd.2005.12.010.

Stegner D, Nieswandt B. Platelet receptor signaling in thrombus formation. *J Mol Med (Berl)*. 2011; **89**: 109-21. 10.1007/s00109-010-0691-5.

Smyth SS, Woulfe DS, Weitz JI, Gachet C, Conley PB, Goodman SG, et al. G-protein-coupled receptors as signaling targets for antiplatelet therapy. *Arteriosclerosis, thrombosis, and vascular biology*. 2009; **29**: 449-57. 10.1161/ATVBAHA.108.176388.

Li S, Tarlac V, Hamilton JR. Using PAR4 Inhibition as an Anti-Thrombotic Approach: Why, How, and When? *Int J Mol Sci*. 2019; **20**. 10.3390/ijms20225629.

Depta JP, Bhatt DL. New approaches to inhibiting platelets and coagulation. *Annu Rev Pharmacol Toxicol*. 2015; **55**: 373-97. 10.1146/annurev-pharmtox-010814-124438.

Franchi F, Angiolillo DJ. Novel antiplatelet agents in acute coronary syndrome. *Nat Rev Cardiol*. 2015; **12**: 30-47. 10.1038/nrcardio.2014.156.

Desai NR, Bhatt DL. The state of periprocedural antiplatelet therapy after recent trials. *JACC Cardiovasc Interv*. 2010; **3**: 571-83. 10.1016/j.jcin.2010.04.008.

de Souza Brito F, Tricoci P. Novel anti-platelet agents: focus on thrombin receptor antagonists. *J Cardiovasc Transl Res*. 2013; **6**: 415-24. 10.1007/s12265-013-9454-3.

Offermanns S. Activation of platelet function through G protein-coupled receptors. *Circulation research*. 2006; **99**: 1293-304. 10.1161/01.RES.0000251742.71301.16.

Angiolillo DJ, Capodanno D, Goto S. Platelet thrombin receptor antagonism and atherothrombosis. *Eur Heart J*. 2010; **31**: 17-28. 10.1093/eurheartj/ehp504.

Stewart A, Fisher RA. Introduction: G Protein-coupled Receptors and RGS Proteins. *Prog Mol Biol Transl Sci*. 2015; **133**: 1-11. 10.1016/bs.pmbts.2015.03.002.

Masuho I, Balaji S, Muntean BS, Skamangas NK, Chavali S, Tesmer JJG, et al. A Global Map of G Protein Signaling Regulation by RGS Proteins. *Cell*. 2020; **183**: 503-21.e19. 10.1016/j.cell.2020.08.052.

O'Brien JB, Wilkinson JC, Roman DL. Regulator of G-protein signaling (RGS) proteins as drug targets: Progress and future potentials. *The Journal of biological chemistry*. 2019; **294**: 18571-85. 10.1074/jbc.REV119.007060.

Alshbool FZ, Karim ZA, Vemana HP, Conlon C, Lin OA, Khasawneh FT. The regulator of G-protein signaling 18 regulates platelet aggregation, hemostasis and thrombosis. *Biochemical and biophysical research communications*. 2015; **462**: 378-82. 10.1016/j.bbrc.2015.04.143.

Ma P, Ou K, Sinnamon AJ, Jiang H, Siderovski DP, Brass LF. Modulating platelet reactivity through control of RGS18 availability. *Blood*. 2015; **126**: 2611-20. 10.1182/blood-2015-04-640037.

Hensch NR, Karim ZA, Druey KM, Tansey MG, Khasawneh FT. RGS10 Negatively Regulates Platelet Activation and Thrombogenesis. *PloS one*. 2016; **11**: e0165984. 10.1371/journal.pone.0165984.

Hernandez KR, Karim ZA, Qasim H, Druey KM, Alshbool FZ, Khasawneh FT. Regulator of G-Protein Signaling 16 Is a Negative Modulator of Platelet Function and Thrombosis. *J Am Heart Assoc*. 2019; **8**: e011273. 10.1161/jaha.118.011273.

Karim ZA, Alshbool FZ, Vemana HP, Conlon C, Druey KM, Khasawneh FT. CXCL12 regulates platelet activation via the regulator of G-protein signaling 16. *Biochimica et biophysica acta*. 2016; **1863**: 314-21. 10.1016/j.bbamcr.2015.11.028.

Manne BK, Denorme F, Middleton EA, Portier I, Rowley JW, Stubben C, et al. Platelet gene expression and function in patients with COVID-19. *Blood*. 2020; **136**: 1317-29. 10.1182/blood.2020007214.

Kim SD, Sung HJ, Park SK, Kim TW, Park SC, Kim SK, et al. The expression patterns of RGS transcripts in platelets. *Platelets*. 2006; **17**: 493-7. 10.1080/09537100600758123.

Wijten P, van Holten T, Woo LL, Bleijerveld OB, Roest M, Heck AJ, et al. High precision platelet releasate definition by quantitative reversed protein profiling--brief report. *Arteriosclerosis, thrombosis, and vascular biology*. 2013; **33**: 1635-8. 10.1161/ATVBAHA.113.301147.

Burkhart JM, Vaudel M, Gambaryan S, Radau S, Walter U, Martens L, et al. The first comprehensive and quantitative analysis of human platelet protein composition allows the comparative analysis of structural and functional pathways. *Blood*. 2012; **120**: e73-82. 10.1182/blood-2012-04-416594.

Yang G, González P, Moneró M, Carrasquillo K, Renta JY, Hernandez-Suarez DF, et al. Discovery of ancestry-specific variants associated with clopidogrel response among Caribbean Hispanics. *NPJ Genom Med*. 2025; **10**: 20. 10.1038/s41525-025-00479-3.

Yang J, Huang J, Maity B, Gao Z, Lorca RA, Gudmundsson H, et al. RGS6, a modulator of parasympathetic activation in heart. *Circulation research*. 2010; **107**: 1345-9. 10.1161/circresaha.110.224220.

Li W, Nieman, M., Sen Gupta, A. Ferric Chloride-induced Murine Thrombosis Models. *J Vis Exp*. 2016; **115**: e54479. doi:10.3791/54479 (2016).

Li W, McIntyre T, Silverstein R. Ferric chloride-induced murine carotid arterial injury: a model of redox pathology. *Redox Biology*. 2013; **1**: 50-5. http://dx.doi.org/10.1016/j.redox.2012.11.001.

Belcher A, Zulfiker AHM, Li OQ, Yue H, Gupta AS, Li W. Targeting Thymidine Phosphorylase With Tipiracil Hydrochloride Attenuates Thrombosis Without Increasing Risk of Bleeding in Mice. *Arteriosclerosis, thrombosis, and vascular biology*. 2021; **41**: 668-82. 10.1161/atvbaha.120.315109.

Li OQ, Yue H, DeHart AR, Roytenberg R, Li AY, Bai F, et al. The Na/K-ATPase α1 subunit fine-tunes platelet P2Y12 function and mediates sex dimorphism–associated thrombosis. *Blood Advances*. 2026; **10**: 973-86. 10.1182/bloodadvances.2025016605.

Roytenberg R, Yue H, DeHart A, Kim E, Bai F, Kim Y, et al. Thymidine phosphorylase mediates SARS-CoV-2 spike protein enhanced thrombosis in K18-hACE2TG mice. *Thrombosis Research*. 2024; **244**: 109195. https://doi.org/10.1016/j.thromres.2024.109195.

Li W, Gigante A, Perez-Perez MJ, Yue H, Hirano M, McIntyre TM, et al. Thymidine phosphorylase participates in platelet signaling and promotes thrombosis. *Circulation research*. 2014; **115**: 997-1006. 10.1161/CIRCRESAHA.115.304591.

Morrell CN, Mix D, Aggarwal A, Bhandari R, Godwin M, Owens P, 3rd, et al. Platelet olfactory receptor activation limits platelet reactivity and growth of aortic aneurysms. *The Journal of clinical investigation*. 2022; **132**. 10.1172/jci152373.

Luo Z, Ahlers-Dannen KE, Spicer MM, Yang J, Alberico S, Stevens HE, et al. Age-dependent nigral dopaminergic neurodegeneration and α-synuclein accumulation in RGS6-deficient mice. *JCI Insight*. 2019; **5**. 10.1172/jci.insight.126769.

Cattaneo M. The platelet P2Y□□ receptor for adenosine diphosphate: congenital and drug-induced defects. *Blood*. 2011; **117**: 2102-12. 10.1182/blood-2010-08-263111.

Yeung J, Li W, Holinstat M. Platelet Signaling and Disease: Targeted Therapy for Thrombosis and Other Related Diseases. *Pharmacological Reviews*. 2018; **70**: 526-48. https://doi.org/10.1124/pr.117.014530.

Gachet C. P2Y(12) receptors in platelets and other hematopoietic and non-hematopoietic cells. *Purinergic Signal*. 2012; **8**: 609-19. 10.1007/s11302-012-9303-x.

Kuliczkowski W, Rychlik B, Chiżyński K, Watała C, Golański J. Comparison of the VASP assay and platelet aggregometry in the evaluation of platelet P2Y12 receptor blockade. *Pol Arch Med Wewn*. 2011; **121**: 115-21.

Schachter JB, Sromek SM, Nicholas RA, Harden TK. HEK293 human embryonic kidney cells endogenously express the P2Y1 and P2Y2 receptors. *Neuropharmacology*. 1997; **36**: 1181-7. https://doi.org/10.1016/S0028-3908(97)00138-X.

Ross EM, Wilkie TM. GTPase-activating proteins for heterotrimeric G proteins: regulators of G protein signaling (RGS) and RGS-like proteins. *Annu Rev Biochem*. 2000; **69**: 795-827. 10.1146/annurev.biochem.69.1.795.

Berger JS, Kornblith LZ, Gong MN, Reynolds HR, Cushman M, Cheng Y, et al. Effect of P2Y12 Inhibitors on Survival Free of Organ Support Among Non-Critically Ill Hospitalized Patients With COVID-19: A Randomized Clinical Trial. *Jama*. 2022; **327**: 227-36. 10.1001/jama.2021.23605.

Ma P, Gupta S, Sampietro S, DeHelian D, Tutwiler V, Tang A, et al. RGS10 shapes the hemostatic response to injury through its differential effects on intracellular signaling by platelet agonists. *Blood Advances*. 2018; **2**: 2145-55. 10.1182/bloodadvances.2017008508.

Hiol A, Davey PC, Osterhout JL, Waheed AA, Fischer ER, Chen CK, et al. Palmitoylation regulates regulators of G-protein signaling (RGS) 16 function. I. Mutation of amino-terminal cysteine residues on RGS16 prevents its targeting to lipid rafts and palmitoylation of an internal cysteine residue. *The Journal of biological chemistry*. 2003; **278**: 19301-8. 10.1074/jbc.M210123200.

Neubig RR, Siderovski DP. Regulators of G-protein signalling as new central nervous system drug targets. *Nat Rev Drug Discov*. 2002; **1**: 187-97. 10.1038/nrd747.

Spicer MM, Weber MA, Luo Z, Yang J, Narayanan NS, Fisher RA. Regulator of G protein signaling 6 (RGS6) in dopamine neurons promotes EtOH seeking, behavioral reward, and susceptibility to relapse. *Psychopharmacology (Berl)*. 2024; **241**: 2255-69. 10.1007/s00213-024-06631-8.

DeBaker MC, Mitten EH, Rose TR, Marron Fernandez de Velasco E, Gao R, Lee AM, et al. RGS6 negatively regulates inhibitory G protein signaling in dopamine neurons and positively

regulates binge-like alcohol consumption in mice. *British journal of pharmacology*. 2023; **180**: 2140-55. 10.1111/bph.16071.

49   Ahlers-Dannen KE, Spicer MM, Fisher RA. RGS Proteins as Critical Regulators of Motor Function and Their Implications in Parkinson's Disease. *Molecular pharmacology*. 2020; **98**: 730-8. 10.1124/mol.119.118836.

50   Rorabaugh BR, Chakravarti B, Mabe NW, Seeley SL, Bui AD, Yang J, et al. Regulator of G Protein Signaling 6 Protects the Heart from Ischemic Injury. *The Journal of pharmacology and experimental therapeutics*. 2017; **360**: 409-16. 10.1124/jpet.116.238345.

Figure legends

Fig. 1. RGS6 deficiency enhances arterial thrombosis in male mice, but not in females. (**A**) Representative video images of thrombus formation in the carotid artery of male mice following 7.5% FeCl$_3$ treatment. Platelets were labeled via direct intravenous injection of Rhodamine 6G. (**B**) Kaplan-Meier curve showing the time to occlusive thrombus formation after injury in male mice. (**C**) Survival curve data showing the time to occlusive thrombus formation after injury in female mice.

Fig. 2. RGS6 deficiency enhances ADP-induced platelet aggregation in vitro. Whole blood was collected from WT and $Rgs6^{-/-}$ mice using 0.109 M sodium citrate as an anticoagulant, Tyrode's buffer (0.7 of blood volume) was added and PRP was prepared by centrifugation. Platelet concentration was adjusted to 2.5 x 10^8 cells/mL with platelet-poor plasma (PPP). PRP (400 µL) was supplemented with CaCl□/MgCl□ to a final concentration of 1 mM, immediately before aggregation was initiated by addition of collagen (**A**) or ADP (**B**).

Fig. 3. RGS6 deficiency enhances ADP-mediated suppression of cAMP/PKA pathway. Platelet-rich plasma (PRP) pooled from 6 mice per genotype was divided into 4 aliquots and stimulated with 2.5 µM ADP for 0, 1, 3, and 5 min. Reactions were terminated by addition of EDTA (1 mM final) and PGE□ (0.5 µM final). Platelets were immediately pelleted by centrifugation and lysed in RIPA buffer containing protease and phosphatase inhibitors. Equal amounts of protein (30 µg) were subjected to Western blot analysis using the indicated antibodies. Band intensities were quantified using ImageJ. Data are presented as the ratio of phosphorylated (p-) protein to total (T-) protein, further corrected for loading by dividing by the relative loading-control intensity normalized to WT at time 0. **A&B**: GβγPI3K/AKT signaling, **C, D, and E:** Gai/AC/cAMP/PKA signaling.

Fig. 4. RGS6 negatively regulates P2Y12/Gi signaling. GloResponse™ CRE-luc2P HEK293 cells were seeded in 24-well plates at a density of 1×10□ cells per well. Cells were transfected with the indicated plasmids 6 h after seeding and incubated for an additional 24 h. Cells were then serum-starved for 4 h prior to stimulation with ADP at the indicated concentrations. "+" indicates 0.25 µg plasmid.

Fig. 5. Inhibition of P2Y$_{12}$ with clopidogrel abolishes the enhanced arterial thrombosis observed in $Rgs6^{-/-}$ mice. WT and $Rgs6^{-/-}$ mice were administered clopidogrel (1 mg/kg/day) by oral gavage for 3 days prior to induction of thrombosis.

Fig. 6. Schematic of RGS6-mediated negative regulation of P2Y12/Gi signaling. Black arrows indicate activation, and blue blunt-ended lines indicate inhibition. AC, adenylyl cyclase.

Fig. 1. RGS6 deficiency enhances arterial thrombosis in male mice, but not in females. (**A**) Representative video images of thrombus formation in the carotid artery of male mice following 7.5% FeCl₃ treatment. Platelets were labeled via direct intravenous injection of Rhodamine 6G. (**B**) Survival curve data showing the time to occlusive thrombus formation after injury in male mice. (**C**) Survival curve data showing the time to occlusive thrombus formation after injury in female mice.

Fig. 2. RGS6 deficiency enhances ADP-induced platelet aggregation in vitro. Whole blood was collected from WT and *Rgs6*⁻/⁻ mice using 0.109 M sodium citrate as an anticoagulant, Tyrode's buffer (0.7 of blood volume) was added and PRP was prepared by centrifugation. Platelet concentration was adjusted to 2.5 x 10⁸ cells/mL with platelet-poor plasma (PPP). PRP (400 μL) was supplemented with $CaCl_2/MgCl_2$ to a final concentration of 1 mM, immediately before aggregation was initiated by addition of collagen (A) or ADP (B).

Fig. 3. RGS6 deficiency enhances ADP-mediated suppression of cAMP/PKA pathway. Platelet-rich plasma (PRP) pooled from 6 mice per genotype was divided into 4 aliquots and stimulated with 2.5 μM ADP for 0, 1, 3, and 5 min. Reactions were terminated by addition of EDTA (1 mM final) and PGE_1 (0.5 μM final). Platelets were immediately pelleted by centrifugation and lysed in RIPA buffer containing protease and phosphatase inhibitors. Equal amounts of protein (30 μg) were subjected to Western blot analysis using the indicated antibodies. Band intensities were quantified using ImageJ. Data are presented as the ratio of phosphorylated (p-) protein to total (T-) protein, further corrected for loading by dividing by the relative loading-control intensity normalized to WT at time 0. **A&B**: GβγPI3K/AKT signaling, **C, D, and E**: Gαi/AC/cAMP/PKA signaling.

Fig. 4. RGS6 negatively regulates P2Y12/Gi signaling. GloResponse™ CRE-luc2P HEK293 cells were seeded in 24-well plates at a density of 1×10^5 cells per well. Cells were transfected with the indicated plasmids 6 h after seeding and incubated for an additional 24 h. Cells were then serum-starved for 4 h prior to stimulation with ADP at the indicated concentrations. "+" indicates 0.25 μg plasmid.

Fig. 5. Inhibition of P2Y$_{12}$ with clopidogrel abolishes the enhanced arterial thrombosis observed in *Rgs6*$^{-/-}$ mice. WT and *Rgs6*$^{-/-}$ mice were administered clopidogrel (1 mg/kg) by oral gavage for 3 days prior to induction of thrombosis.

Fig. 6. Schematic of RGS6-mediated negative regulation of P2Y$_{12}$/Gi signaling. Black arrows indicate activation, and blue blunt-ended lines indicate inhibition. AC, adenylyl cyclase.

	Rgs6⁻/⁻			WT			
	Mean	SEM	Sample size	Mean	SEM	Sample size	*P* value
WBC (K/uL)	4.478	0.319	4	4.370	0.935	6	0.931
NE (K/uL)	0.863	0.050	4	1.495	0.475	6	0.319
LY (K/uL)	3.288	0.190	4	2.513	0.629	6	0.362
MO (K/uL)	0.303	0.093	4	0.290	0.068	6	0.914
EO (K/uL)	0.005	0.003	4	0.058	0.036	6	0.269
BA (K/uL)	0.000	0.000	4	0.013	0.008	6	0.221
NE (%)	19.533	1.390	4	30.468	8.183	6	0.319
LY (%)	73.888	1.311	4	58.798	6.984	6	0.124
MO (%)	6.448	1.477	4	9.557	3.969	6	0.559
EO (%)	0.103	0.029	4	0.933	0.475	6	0.200
BA (%)	0.030	0.010	4	0.243	0.124	6	0.205
RBC (M/uL)	7.748	0.103	4	7.303	1.018	6	0.737
Hb (g/dL)	8.475	0.063	4	7.650	1.162	6	0.586
HCT (%)	35.350	0.463	4	32.267	4.412	6	0.592
MCV (fL)	45.650	0.119	4	44.433	0.409	6	0.047
MCH (pg)	10.950	0.206	4	10.233	0.414	6	0.223
MCHC (g/dL)	24.000	0.449	4	23.050	1.014	6	0.491
RDW (%)	18.425	0.217	4	17.467	0.342	6	0.071
PLT (K/uL)	612.250	24.098	4	520.167	68.948	6	0.327
MPV (fL)	4.450	0.096	4	4.717	0.098	6	0.101

Table 1. Whole blood cell counts in male WT and *Rgs6⁻/⁻* mice.

WBC: white blood cells; NE: neutrophils; LY: lymphocytes; MO: monocytes; EO: eosinophils; BA: basophils; RBC: Red blood cells; Hb: Hemoglobin; HCT: Hematocrit; MCV: mean corpuscular volume; MCH: Mean corpuscular hemoglobin; MCHC: Mean corpuscular hemoglobin concentration; RDW: red cell distribution width; PLT: Platelet; MPV: Mean platelet volume.

Table 2: Summary of RGS proteins with evidence on platelet activation

RGS protein	Gi activity	Platelet evidence	Deficiency on Akt (PI3K/Gβγ)	Deficiency on PKA / VASP (cAMP)	Interpretation
RGS10	Strong	Strong	↓ Akt	↑ PKA / ↑ VASP	Classic Gi inhibition → less PI3K, more cAMP
RGS18	Moderate	Strong	↓ Akt (context-dependent)	↑ PKA / ↑ VASP	Broad regulator; modulates both Gi and Gq, net inhibitory
RGS6	Strong (biochemical)	Emerging	~ no consistent ↑ (context-dependent; may ↓ or unchanged)	↑ PKA / ↑ VASP (baseline effect)	Kinetic regulator of Gi; strongest effect on cAMP/PKA, variable effect on Akt
RGS16		limited	Likely ↓ Akt (limited data)	Likely ↑ PKA	Expected Gi GAP behavior, but not well defined in platelets