1 Cryptic haplotypes of “*Candidatus Liberibacter africanus*”

2

3 Warrick R Nelson, The New Zealand Institute for Plant & Food Research Limited, Private Bag 4704,
4 Christchurch 8140, New Zealand

5 Sandrine Eveillard, Marie-Pierre Dubrana and Joseph M Bové, Université Bordeaux and INRA, UMR
6 13 32 Biologie du fruit et Pathologie, 71, Ave. Edouard Bourlaux, 33883 Villenave d’Ornon, France

7 Dedicated to the memory of Colette Bové-Dumeau (1927-2014).

8

9 Abstract

10 “*Candidatus Liberibacter africanus*” (Laf) has long been recognised as a causal agent of the
11 devastating citrus disease huanglongbing (HLB) or citrus greening. This species is currently restricted
12 to Africa, the Arabian Peninsula and some Indian Ocean islands and vectored by the African citrus
13 psyllid, *Trioza erytreae*. Blotchy mottle on citrus leaves is characteristic of the disease.

14 Somewhat similar symptoms in the Rutaceous tree *Calodendrum capensis* (Cape Chestnut) resulted
15 in the discovery of Laf outside commercial citrus crops in South Africa. This was classed as a
16 subspecies of Laf (capensis, hence LafC). In subsequent surveys of both commercial citrus crops and
17 *Calodendrum*, both natural and ornamental specimens, LafC was not found in the citrus crop, nor
18 has Laf been found in *C. capensis*. HLB was reported from Madagascar in 1968 but no sequences
19 from this source have so far been published.

20 Until fairly recently, only the reference 16S rRNA gene sequences of Laf (L22533) and LafC
21 (AF137368) had been deposited in GenBank. Both of these reference sequences contain a number
22 of unresolved nucleotides. Resolving these nucleotide positions by aligning against more recently
23 available sequences, it becomes evident that these unresolved positions represent one percentage
24 point difference in similarity between Laf and LafC. The originally reported 97.4% similarity is
25 therefore incorrect based on this new information. Recalculating the similarity on the full length 16S
26 rDNA sequence results in 99.54% similarity, a value too high to justify a subspecies status. LafC
27 should therefore be reduced to that of a haplotype of Laf.

28 Further, the six 16S rRNA gene sequences currently available in GenBank identified as the species Laf
29 separate into 2 haplotype groups. The 3 haplotypes of Laf are therefore LafA designated as the first
30 accession sequenced (L22533), LafC for the former capensis subspecies and to recognise the prior
31 use of this term, and LafB for the third haplotype not previously recognised. Thus the cryptic
32 presence of 3 haplotypes is revealed by this review of the Laf 16S rDNA sequences.

33 **Keywords** “*Candidatus Liberibacter africanus*”; haplotype, citrus

34

35 Introduction

Huanglongbing (HLB), also known as citrus greening, is a serious disease threatening the economic existence of many commercial citrus orchards. It has recently spread to the major production regions of Brazil, Florida and California (Wang & Trivedi, 2013). Currently, 3 species of "*Candidatus Liberibacter*", named for the continent of first discovery, are associated with symptomatic commercial citrus plants (Bové, 2006), "*Candidatus Liberibacter asiaticus*" (Las) being the most geographically widespread (Bové, 2014).

"*Candidatus Liberibacter africanus*" (Laf) is associated with HLB throughout East Africa, the Arabian Peninsula and Indian Ocean islands (Bové, 2006; Bové, 2014), including Madagascar (Bové & Cassin, 1968). The Rutaceous tree, *Calodendrum capense* (Cape Chestnut) has also been discovered to be infected by "*Ca. Liberibacter africanus subsp. capensis*" (LafC)(Garnier *et al.*, 2000). This has subsequently been found to be widespread in South Africa where this tree is both native and grown ornamentally (Phahladira *et al.*, 2012). In spite of the subspecies being widespread, only Laf is present in symptomatic commercial citrus orchards in South Africa (Pietersen *et al.*, 2010). Other native Rutaceae species in South Africa also tested positive for Laf and LafC and possibly another variant of the species, but interpreted as a series of subspecies of Laf (Roberts *et al.*, 2014).

The original description of LafC was based on three gene fragments (Garnier *et al.*, 2000), of which the 16S rRNA gene is taxonomically the most important (Weisburg *et al.*, 1991; Woese & Fox, 1977). This 16S sequence (GenBank accession AF137368) was noted as having 97.4% similarity with Laf (L22533). Three percent divergence over the full gene sequence is a commonly recognised requirement for bacterial species designation (Stackebrandt & Goebel, 1994), thus the subspecies status for LafC. Closer inspection of these sequences revealed 6 unresolved nucleotides on LafC and 5 in the Laf reference sequences. There is thus potentially one percentage point of difference derived simply from these unresolved nucleotide positions. A review of this subspecies status is now feasible with the availability of more recent accession sequences.

60

61 **Methods and Materials**

A literature review revealed further examples of both Laf and LafC 16S rDNA sequences, as well as some sequences for 16S-23S ISR, 23S and 50S rRNA gene regions. The megablast protocol in GenBank was used to find further closely related sequences.

These sequences were downloaded from the NCBI database (GenBank) and aligned in ClustalX 2.1 (Larkin *et al.*, 2007). Sequences for 16S rDNA from the genomes of two other *Liberibacter* species ("*Ca. L. asiaticus*" and "*Ca. L. solanacearum*", Las and Lso respectively) were also downloaded for additional comparison. These were chosen as phylogenies generally place them closest to Laf (Nelson *et al.*, 2013).

DNA material stored from symptomatic leaves of sweet orange collected in Madagascar during 2011, previously reported to be Laf by PCR (Bové, 2014), was retested using the primer pair OA1/Oi2c (Jagoueix *et al.*, 1996) and the resultant 16S rDNA samples sequenced. The rplJ component of the 50S rRNA gene was also sequenced.

74

75 Results

76 On the 16S rRNA gene, 15 sequences identified as Laf or LafC were found. Three of these proved to
77 contain only a very short component of the 16S gene and were discarded. Aligning the remaining 12
78 sequences showed a number of apparently random SNPs scattered across the accessions,
79 presumably arising as sequencing errors, but the LafC sequences separated clearly.

80 In the sequence L22533 (Laf), there are 5 nucleotide positions labelled “N”. From within the aligned
81 set of sequences, these can be unambiguously resolved to the nucleotides T, C, G, A and G.
82 Similarly, in the sequence AF137368 (LafC), these also resolve unambiguously to A, A, A, G, C, C.

83 Laf sequences also aligned into 2 groups over 7 SNPs (Table 1), here designated as haplotypes LafA
84 and LafB. Excluding the occasional single sequence SNPs, the number of SNPs separating the 2
85 haplotypes and subspecies is outlined in Table 2. Although the sequence of each accession varied in
86 length, these SNPs occur over an approximately 1100 bp length. Aligning the Laf sequences against
87 the full 16S gene sequences for Las and Lso indicated that all the SNPs in Table 1 occur within the
88 overlapping regions. *Liberibacter* 16S genes are approximately 1500 bp (Nelson *et al.*, 2015). Until
89 full Laf sequences are available, the similarities reported here might prove to be high if further SNPs
90 exist in the currently unknown regions. There are 8 SNPs between Las and Lso in these end regions
91 of the gene, representing 0.53% difference. Assuming a similar number of SNPs between the Laf
92 sequences in these end regions, we can calculate a likely worst-case scenario for the number of SNPs
93 between the proposed haplotypes and the percentage similarity, assuming a 16S rRNA gene length
94 of 1500 bp.

95 Only 4 sequences across the 16S-23S ISR and 23S partial gene regions were found in GenBank
96 (LAU61360, EU754741, FJ914622, JF819884, JF819885) for Laf and none for LafC. The LafC sequence
97 has been mentioned as being closer to Las than Laf, because only one tRNA (alanine) is present in
98 the ISR of Laf, while two tRNAs (alanine and isoleucine) occur in the ISR of Las and LafC (Garnier *et*
99 *al.*, 2000).

100 Over a common segment of 479 bp on the 50S gene region (β -operon) there are 68 SNPs between
101 Laf and LafC (14 and 4 accessions respectively)(Phahladira *et al.*, 2012; Pietersen *et al.*, 2010). The
102 available sequence material in the 50S gene confirms the basic Laf/LafC separation, but we feel that
103 separating LafA and LafB is as yet inconclusive with only 2 SNPs across the 4 accessions.

104 The 16S sequences derived from Madagascar material aligned with LafA and the rplJ sequence
105 (GenBank) aligned with EF122255, a sequence with metadata indicating it is derived from
106 commercial citrus material in South Africa.

107

Table 1 Proposed haplotypes and SNP differences on the 16S rRNA gene within “*Candidatus Liberibacter africanus*” (Laf) sequences. The nucleotide numbers count from the beginning of the reference sequence L22533 although the alignment was done using a 1057 bp length common to all sequences. Shaded SNPs are positions of ambiguity between the sequences within a series where the SNP is in the same position as the nucleotide in one of the other haplotypes.

GenBank	Source	Author	Haplotype	SNP description													
				g.149G>	g.329T>C	g.355G>A	g.411G>A	g.413G>A	g.427insA	g.486insG	g.507C>T	g.562A>G	g.565G>A	g.908C>T	g.920T>G	g.921T>A	g.1124insG
L22533	Nelspruit	(Jagoueix <i>et al.</i> , 1994)	Laf A	G	G	G	G	G	-	-	C	A	G	C	T	T	-
EU921620	Mpumalang a-UPCRI-06-0026	(Lin <i>et al.</i> , 2009)	Laf A	G	T	G	G	G	-	-	C	A	G	C	T	T	-
EU921619	Mpumalang a-UPCRI-05-0252	(Lin <i>et al.</i> , 2009)	Laf A	G	T	G	G	G	-	-	C	A	G	C	T	T	-
KJ152137	UPCRI 12-0004 (Zanthoxylum capensis), Knysna	(Roberts <i>et al.</i> , 2014)	Laf A	-	T	G	G	G	-	-	C	G	G	C	G	A	-
LN795909	Madagascar ,	This study	Laf A	G	T	G	G	G	-	-	C	A	G	C	T	T	-
EU921621	Mpumalang a-UPCRI-06-0071	(Lin <i>et al.</i> , 2009)	Laf B	G	C	A	A	A	A	G	C	A	G	C	T	T	G
EU754741	Mpumalang a-UPCRI-06-0071	(Doddapane ni <i>et al.</i> , 2008)	Laf B	G	C	A	A	A	A	G	C	A	A	C	A	T	G
FJ914622	Mpumalang a-UPCRI-05-0232	(Lin <i>et al.</i> , 2009)	Laf B	G	C	A	A	A	A	G	C	A	G	C	T	T	G
AF137368	LafC (from Calodendrum, Western Cape province)	(Garnier <i>et al.</i> , 2000)	Laf C	-	T	G	G	G	-	-	T	G	A	T	G	A	-

KJ152127	UPCRI 10-2267 (Calodendrum capense), Good Hope	(Roberts <i>et al.</i> , 2014)	Laf C	-	T	G	G	G	-	-	T	G	A	T	G	A	-
KJ152131	UPCRI 11-4505 (Clauseniana anisata), Enseleni	(Roberts <i>et al.</i> , 2014)	Laf C	-	T	G	G	G	-	-	T	G	A	T	G	A	-
KJ197226	UPCRI 11-4270 (Zanthoxylum capense), St Francis Bay	(Roberts <i>et al.</i> , 2014)	Laf C	-	T	G	G	G	-	-	T	G	A	T	G	A	-
KJ152134	UPCRI 11-4044 (Vepris lanceolata), Knysna	(Roberts <i>et al.</i> , 2014)	Laf C	G	T	G	G	G	-	-	T	G	A	T	G	A	-

Table 2 Percent identity similarity between “*Candidatus Liberibacter africanus*” (Laf) haplotypes based on 16S rRNA gene sequences, % similarity calculated on a 1500 bp length. An alternate % similarity figure is indicated in parentheses assuming a further 8 SNPs exists in the currently unknown segments of the gene.

Comparison identity	SNPs	% similarity
LafA:LafB	7 (15)	99.53 (99.0)
LafA:LafC	7 (15)	99.53 (99.0)
LafB:LafC	14 (22)	99.07 (98.54)

Discussion

The unresolved nucleotides in the sequences for Laf and LafC (11 positions in total) are readily resolved to their correct nucleotide by comparison against the other 10 sequences now available in GenBank. These newly resolved positions represent one percentage point of similarity between Laf and LafC, thus the originally described 97.4% (Garnier *et al.*, 2000) similarity becomes 98.4%. However, this similarity calculation was conducted on the known sequence proportion of the gene since at the time the full length of the gene was not known. Although the full 16S gene sequence for Laf is still not available, it is likely to be approximately 1500 bp in length as found in 4 other species of *Liberibacter* (Nelson *et al.*, 2015). Across the sequences currently available, only approximately 40 bases at the beginning and 30 at the end of the gene sequence remain unknown, thus allowing a closer estimate of similarity between the species and the subspecies (Table 2).

Variation across the 16S rRNA gene appears to be quite large with 14 SNPs apparent across the Laf/LafC accessions available, 7 of them between LafA/LafB and a different 7 between LafA/LafC, with all 14 being different between LafB/LafC. For comparison, Las appears to be homogenous across accessions on the 16S rRNA gene (Moreno-Enríquez *et al.*, 2014; Nelson, 2012), while for Lso 5 haplotypes have been described, with only 7 SNPs defining them on the 16S rRNA gene (Nelson *et al.*, 2011; Nelson *et al.*, 2013; Teresani *et al.*, 2014).

The unexpected separation into 2 groups on the 16S rRNA gene within the currently known Laf species suggests 2 haplotypes, here designated LafA and LafB (Table 2). No prior study has indicated that there were genetic differences within Laf on commercial citrus. This suggests 3 haplotypes within the Laf species, 2 primarily indicated by sequences derived from commercial citrus crops, and the previous subspecies designation “capensis” can be resolved to a haplotype but retaining the short form of LafC. Caution is required in this conclusion since the LafB designation is based on only 3 available sequences and 2 of these, although sequenced by different laboratories, are clearly derived from the same field sample. The metadata associated with these sequences give little indication of geographical or plant/insect host separation. For LafA/LafB they are primarily from commercial citrus crops and LafC from native Rutaceous species. This provides a stark contrast to the analogous situation in Lso where the haplotypes also express partial plant/insect host and geographic separation (Nelson *et al.*, 2011; Nelson *et al.*, 2013; Teresani *et al.*, 2014). If the A/B haplotypes can be confirmed, they indicate 2 separate pathosystem events from the respective currently unknown African native plant hosts into commercial citrus, since *Citrus* is not native to Africa (Beattie *et al.*, 2008), and LafC has yet to make this jump as it is not yet known from citrus (Phahladira *et al.*, 2012).

The 50S rRNA gene sequences show clear separation between Laf and LafC as expected, with only a suggestion on 4 of 15 Laf sequences showing SNPs. Caution is suggested in considering this indicative of LafA/B haplotypes as these SNPs occur together and only 4 bases from the end of the sequence. Further, there is no indication from the metadata to tie these 4 sequences together with the 3 suggested as LafB from the 16S gene.

Although previously known that Laf was the species responsible for HLB on Madagascar by both epidemiological and PCR studies (Bové & Cassin, 1968; Bové, 2014), this study confirms not only the species as Laf via both 16S and rplJ genes, but that it is also haplotype LafA. This very strongly suggests an incursion event of LafA from Africa to Madagascar, rather than a Gondwanan origin, in spite of the position of Madagascar in Gondwana between East Africa and India.

The very recent suggestion of a further three subspecies of Laf (Roberts *et al.*, 2014) can be resolved within this proposal of haplotypes rather than subspecies within Laf by giving them a biotype designation, recognising the current host plant differences. Laf subspecies *vepridis* is a biotype of LafA, while Laf subspecies *zanthoxyli* and Laf subspecies *clausenae* are biotypes of LafC.

Conclusion

Instead of the species Laf and subspecies *capensis*, re-analysis of the phylogenetically important 16S rRNA gene suggests the subspecies is not that far removed from the species and should be revised to haplotype status. Further, existing sequences of Laf indicate that this species also comprises two haplotypes. Therefore two haplotypes (LafA and LafB) are known symptomatically in commercial citrus orchards while the third (LafC) is known only from very mild symptoms in native Rutaceous plants but not (yet) in citrus. Thus the cryptic presence of three haplotypes is revealed by this review of the Laf 16S rDNA sequences.

References

- Beattie, G., Holford, P., Mabblerley, D., Haigh, A. & Broadbent, P. (2008).** On the origins of *Citrus*, Huanglongbing, *Diaphorina citri* and *Trioza erytreae*. *International Research Conference on Huanglongbing, Orlando, Florida*, 25-57.
- Bové, J. M. (2006).** Huanglongbing: a destructive, newly-emerging, century-old disease of citrus. *Journal of Plant Pathology* **88**, 7-37.
- Bové, J. M. (2014).** Heat-tolerant Asian HLB meets Heat-sensitive African HLB on the Arabian Peninsula. Why?. *Journal of Citrus Pathology* **1**, 1-78.
- Bové, J. M. & Cassin, J. P. (1968).** Problèmes de l'agrumiculture malgache; comperendu de mission. *Document IRFA*: 52 p.
- Doddapaneni, H., Liao, H., Lin, H., Bai, X., Zhao, X., Civerolo, E., Irely, M., Coletta-Filho, H. & Pietersen, G. (2008).** Comparative phylogenomics and multi-gene cluster analyses of the Citrus Huanglongbing (HLB)-associated bacterium *Candidatus Liberibacter*. *BMC Res Notes* **1**, 72.
- Garnier, M., Jagoueix-Eveillard, S., Cronje, P., Le Roux, H. & Bové, J. (2000).** Genomic characterization of a liberibacter present in an ornamental rutaceous tree, *Calodendrum capense*, in the Western Cape province of South Africa. Proposal of "*Candidatus Liberibacter africanus* subsp. capensis". *Int J Syst Evol Microbiol* **50**, 2119-2125.
- Jagoueix, S., Bové, J. M. & Garnier, M. (1994).** The phloem-limited bacterium of greening disease of citrus is a member of the α -subdivision of the Proteobacteria. *Int J Syst Bacteriol* **44**, 379-386.
- Jagoueix, S., Bové, J. M. & Garnier, M. (1996).** PCR detection of the two "*Candidatus*" liberobacter species associated with greening disease of citrus. *Mol Cell Probes* **10**, 43-50.
- Larkin, M. A., Blackshields, G., Brown, N. P., Chenna, R., McGettigan, P. A., McWilliam, H., Valentin, F., Wallace, I. M., Wilm, A. & other authors (2007).** Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947-2948.
- Lin, H., Doddapaneni, H., Munyaneza, J. E., Civerolo, E. L., Sengoda, V. G., Buchman, J. L. & Stenger, D. C. (2009).** Molecular characterization and phylogenetic analysis of 16S rRNA from a new species of "*Candidatus Liberibacter*" associated with Zebra chip disease of potato (*Solanum tuberosum* L.) and the potato psyllid (*Bactericera cockerelli* Sulc). *Journal of Plant Pathology* **91**, 215-219.
- Moreno-Enríquez, A., Minero-García, Y., Ramírez-Prado, J. H., Loeza-Kuk, E., Uc-Varguez, A. & Moreno-Valenzuela, O. A. (2014).** Comparative analysis of 16S ribosomal RNA of '*Candidatus Liberibacter asiaticus*' associated with Huanglongbing disease of Persian lime and Mexican lime reveals a major haplotype with worldwide distribution. *African Journal of Microbiology Research* **8**, 2861-2873.
- Nelson, W. (2012).** First record of *Trioza vitreoradiata* (Maskell)(Hemiptera: Triozidae) in citrus. *Citrus Research & Technology* **33**, 35-38.

- 211 **Nelson, W. R., Fisher, T. W. & Munyaneza, J. E. (2011).** Haplotypes of "*Candidatus Liberibacter*
212 *solanacearum*" suggest long-standing separation. *European Journal of Plant Pathology* **130**, 5-12.
- 213 **Nelson, W. R., Munyaneza, J. E., McCue, K. F. & Bové, J. M. (2013).** The Pangaeian origin of
214 "*Candidatus Liberibacter*" species. *Journal of Plant Pathology* **95**, 455-461.
- 215 **Nelson, W. R., Wulff, N. A. & Bové, J. M. (2015).** Intragenomic homogeneity on *Liberibacter* 16S
216 rDNA confirms phylogeny and explains ecological strategy. *bioRxiv* doi:
217 <http://dx.doi.org/10.1101/016188>.
- 218 **Nelson, W., Sengoda, V., Alfaro-Fernandez, A., Font, M., Crosslin, J. & Munyaneza, J. (2013).** A new
219 haplotype of "*Candidatus Liberibacter solanacearum*" identified in the Mediterranean region.
220 *European Journal of Plant Pathology* **135**, 633-639.
- 221 **Phahladira, M., Viljoen, R. & Pietersen, G. (2012).** Widespread occurrence of "*Candidatus*
222 *liberibacter africanus* subspecies *capensis*" in *Calodendrum capense* in South Africa. *European*
223 *Journal of Plant Pathology* **134**, 39-47.
- 224 **Pietersen, G., Arrebola, E., Breytenbach, J., Korsten, L., le Roux, H., la Grange, H., Lopes, S., Meyer,**
225 **J., Pretorius, M. & other authors (2010).** A survey for '*Candidatus Liberibacter*' species in South
226 Africa confirms the presence of only '*Ca. L. africanus*' in commercial citrus. *Plant Dis* **94**, 244-249.
- 227 **Roberts, R., Steenkamp, E. T. & Pietersen, G. (2014).** Novel lineages of '*Candidatus Liberibacter*
228 *africanus*' associated with native Rutaceae hosts of *Trioza erythrae* in South Africa. *Int J Syst Evol*
229 *Microbiol* , doi:10.1099/ijs.0.069864-0.
- 230 **Stackebrandt, E. A. & Goebel, B. M. (1994).** Taxonomic note: a place for DNA-DNA reassociation
231 and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol*
232 **44**, 846-849.
- 233 **Teresani, G. R., Bertolini, E., Alfaro-Fernandez, A., Martínez, C., Tanaka, F. A. O., Kitajima, E.,**
234 **Rosello, M., Sanjuan, S., Ferrandiz, J. C. & other authors (2014).** Association of '*Candidatus*
235 *Liberibacter solanacearum*' with a vegetative disorder of celery in Spain and development of a real-
236 time PCR method for its detection. *Phytopathology* **104**, 804-811.
- 237 **Wang, N. & Trivedi, P. (2013).** Citrus Huanglongbing: a newly relevant disease presents
238 unprecedented challenges. *Phytopathology* **103**, 652-665.
- 239 **Weisburg, W., Barns, S., Pelletier, D. & Lane, D. (1991).** 16S ribosomal DNA amplification for
240 phylogenetic study. *J Bacteriol* **173**, 697-703.
- 241 **Woese, C. & Fox, G. (1977).** Phylogenetic structure of the prokaryotic domain: The primary
242 kingdoms. *Proceedings of the National Academy of Sciences* **74**, 5088-5090.
- 243