Beyond 2/3 and 1/3: the complex signatures of sex-biased admixture on the X chromosome

Amy Goldberg*
Noah A Rosenberg

Department of Biology, Stanford University, Stanford, CA, 94305-5020 USA

Abstract Sex-biased demography, in which parameters governing migration and population size differ between females and males, has been studied through comparisons of X chromosomes, which are inherited sex-specifically, and autosomes, which are not. A common form of sex bias in humans is sex-biased admixture, in which at least one of the source populations differs in its proportions of females and males contributing to an admixed population. Studies of sex-biased admixture often examine the mean ancestry for markers on the X chromosome in relation to the autosomes. A simple framework noting that in a population with equally many females and males, $\frac{2}{3}$ of X chromosomes appear in females, suggests that the mean X-chromosomal admixture fraction is a linear combination of female and male admixture parameters, with coefficients $\frac{2}{3}$ and $\frac{1}{3}$, respectively. Extending a mechanistic admixture model to accommodate the X chromosome, we demonstrate that this prediction is not generally true in admixture models, though it holds in the limit for an admixture process occurring as a single event. For a model with constant ongoing admixture, we determine the mean X-chromosomal admixture, comparing admixture on female and male X chromosomes to corresponding autosomal values. Surprisingly, in reanalyzing African-American genetic data to estimate sex-specific contributions from African and European sources, we find that the range of contributions compatible with the excess African ancestry on the X chromosome compared to autosomes has a wide spread, permitting scenarios either without male-biased contributions from Europe or without female-biased contributions from Africa.

*Corresponding author. Phone: 650-724-5122. Fax: 650-724-5114. Email: agoldb@stanford.edu

Introduction

Comparisons of the X chromosome and the autosomes provide a strategy for understanding the history of sex-biased demography [1–14]. Unlike the autosomes, the X chromosome follows a sex-specific inheritance pattern, with females inheriting two copies, one from the mother and one from the father, and males inheriting only a single copy from the mother. As a consequence, demographic differences between females and males—in such phenomena as the breeding population size, the variance of reproductive success, and migration rates—can be studied by examining differences in patterns of genetic variation between X chromosomes and autosomes.

Many of the best-known cases of sex-biased patterns in human demography involve recently admixed populations [15–31]. During the formation of such populations, sex-biased admixture occurs if one or more of the source populations contributes different fractions of the females and males to the admixed group. What patterns are expected for X chromosomes and autosomes in an admixed population formed through a sex-biased admixture process? An initial hypothesis, reflecting the fact that in a population with equally many females and males, $\frac{2}{3}$ of the X chromosomes are in females and $\frac{1}{3}$ are in males, is that if s_1^f is the fraction of females originating from population 1 and s_1^m is the fraction of males originating from population 1, then the fraction of ancestry from population 1 for a site on the X chromosome is [19]

$$H_1^X = \frac{2}{3}s_1^f + \frac{1}{3}s_1^m. \quad (1)$$

This simple linear combination has been used to estimate the sex-specific contributions from females and males to African-American and Latino populations [19, 31]. As we will show, however, it presumes a very specific history for the admixture process, a history often not viewed as reasonable for practical admixture scenarios.

Here, we extend a two-sex mechanistic admixture model [32] to incorporate the genetic signatures of sex-specific admixture patterns on the X chromosome. We derive a recursive expression for the expectation of the X-chromosomal admixture fraction as a function of sex-specific admixture parameters, demonstrating that the X-chromosomal admixture is obtained from a more complex formula than in the simple $\frac{2}{3}-\frac{1}{3}$ weighting (eq. (1)). The limiting mean X-chromosomal admixture is a predictable function of female and male contributions from the source populations, but among cases we consider, only has the $\frac{2}{3}-\frac{1}{3}$ weighting for a single admixture event that takes place at a single point in time. During the approach to this limit, the behavior of the mean X-chromosomal admixture is dependent on the time since admixture. For a single admixture event and for constant ongoing admixture, we characterize the difference between the limit and the mean X-chromosomal admixture under the model as a function of time. We reinterpret admixture patterns in recently admixed African Americans, demonstrating that consideration of both the admixture model and the time since admixture is important for estimating sex-specific admixture contributions.

Materials and Methods

A mechanistic, sex-specific model for admixture histories

We follow the discrete-time model and notation of Verdu & Rosenberg [33] and Goldberg et al. [32], in which two source populations, S_1 and S_2 , contribute to an admixed population, H (Fig. 1). For

each population, female and male contributions are considered separately. At generation g , the contribution of sex δ , with $\delta \in \{f, m\}$, from source S_α , with $\alpha \in \{1, 2\}$, is $s_{\alpha,g}^\delta$. Corresponding contributions from H are denoted h_g^δ . Thus, for example, $s_{\alpha,g}^\delta$ denotes the fraction in generation g of individuals of sex δ originating in the previous generation in population S_α . The female and male contributions from S_1 , H , and S_2 at generation g are $s_{1,g}^f$, $s_{1,g}^m$, h_g^f , h_g^m , $s_{2,g}^f$, and $s_{2,g}^m$.

We recall key relations among the sex-specific parameters [32, eqs. 1-6]. First, the total contribution from each population is the mean of female and male contributions: $s_{\alpha,g} = (s_{\alpha,g}^f + s_{\alpha,g}^m)/2$ and $h_g = (h_g^f + h_g^m)/2$. Also, as each parameter is a probability, the total female and male contributions separately sum to one: $s_{1,g}^f + h_g^f + s_{2,g}^f = 1$ and $s_{1,g}^m + h_g^m + s_{2,g}^m = 1$. For $g = 0$, the admixed population does not yet exist, so that $h_0 = h_0^f = h_0^m = 0$ and $s_{1,0}^f + s_{2,0}^f = s_{1,0}^m + s_{2,0}^m = 1$.

Under this general framework, we define sex bias as a difference in the contributions from one or more source populations to the admixed population. That is, an admixture process is considered sex-biased if $s_{1,g}^f \neq s_{1,g}^m$ or $s_{2,g}^f \neq s_{2,g}^m$, or both. Thus, sex bias involves differences between females and males entering from a source population, rather than a difference between parameters from the two source populations; after the founding, sex bias can occur in one source but not the other.

Considering the three populations (S_1, H, S_2) from which the parents of an individual from the admixed population can originate, we have nine possible ordered parental pairings (Table 1). As in Goldberg et al. [32], we study the random variable representing the admixture fraction of a random individual in the admixed population. Whereas Goldberg et al. [32] examined admixture only for the autosomes, here we also study the X chromosome. We define $H_{1,g,\delta}^\gamma$ as the admixture fraction of chromosomal type γ sampled in individuals of sex δ from the admixed population H at generation g . Thus, $H_{1,g,\delta}^\gamma$ is the probability that a random site on an autosome ($\gamma = A$) or the X chromosome ($\gamma = X$) in a random individual of sex δ in H in generation g ultimately traces to S_1 .

In contrast to autosomal DNA, for which both female and male offspring receive a single copy of each chromosome from each parent, the X chromosome is inherited sex-specifically. That is, female offspring inherit one copy of the X chromosome from the mother and one from the father, but males inherit only a maternal copy. Therefore, whereas the autosomal admixture fractions sampled in females and males from the admixed population, $H_{1,g,f}^A$ and $H_{1,g,m}^A$, are identically distributed, the female and male X-chromosomal admixture fractions, $H_{1,g,f}^X$ and $H_{1,g,m}^X$, have different distributions. As a result, whereas Goldberg et al. [32] could uncouple $H_{1,g,f}^A$ and $H_{1,g,m}^A$ and consider recursions for these two quantities separately (both were identical and only one needed to be studied), here we examine the distributions of $H_{1,g,f}^X$ and $H_{1,g,m}^X$ as a coupled pair of recursions.

Table 1 reports the probabilities that a random individual of a given sex from the admixed population has each possible set of parents, ℓ , as well as the admixture fraction of the individual for source population 1 conditional on the parental pairing. The table provides a basis for recursively computing the fraction of admixture from S_1 for a random individual of sex δ from the admixed population at generation g , $H_{1,g,\delta}^X$. For $g = 1$, the founding event of the admixed population, the admixed population does not yet exist, so only cases 1, 3, 7, and 9, the four cases in which both parents are from the source populations, are considered.

Conditional on the previous generation, $H_{1,g,f}^X$ and $H_{1,g,m}^X$ are independent random variables. The female X-chromosomal admixture fraction depends on both the female and male admixture fractions in the previous generation, $H_{1,g-1,f}^X$ and $H_{1,g-1,m}^X$, but the male X-chromosomal admixture fraction depends only on $H_{1,g-1,f}^X$. Therefore, the female X-chromosomal admixture fraction can

equivalently be written as a function of the female X-chromosomal admixture fractions in the previous generation, $H_{1,g-1,f}^X$, and from two generations ago, $H_{1,g-2,f}^X$.

Expectation of the X-chromosomal fraction of admixture

Following Verdu & Rosenberg [33] and Goldberg et al. [32], we can find the expectation of the X-chromosomal admixture fraction of an individual of specified sex randomly chosen from the admixed population. Using the law of total expectation to consider the random parental pairing L , we sum over the nine possible pairings ℓ (Table 1). For the mean X-chromosomal admixture fraction of a randomly chosen admixed individual of sex δ sampled at generation g , we have

$$\mathbb{E}[H_{1,g,\delta}^X] = \mathbb{E}_L [\mathbb{E}[H_{1,g,\delta}^X|L]] = \sum_{\ell \in \left\{ \begin{array}{l} S_1 S_1 \\ S_1 H \\ S_1 S_2 \\ H S_1 \\ H H \\ H S_2 \\ S_2 S_1 \\ S_2 H \\ S_2 S_2 \end{array} \right\}} P(L = \ell) \mathbb{E}[H_{1,g,\delta}^X|L = \ell]. \quad (2)$$

Applying eq. (2) using Table 1 to consider X chromosomes sampled in females and males from the admixed population, for the first generation, the initial condition is

$$\begin{bmatrix} \mathbb{E}[H_{1,1,f}^X] \\ \mathbb{E}[H_{1,1,m}^X] \end{bmatrix} = \begin{bmatrix} s_{1,0} \\ s_{1,0}^f \end{bmatrix}. \quad (3)$$

For $g \geq 2$, we have

$$\begin{bmatrix} \mathbb{E}[H_{1,g,f}^X] \\ \mathbb{E}[H_{1,g,m}^X] \end{bmatrix} = \begin{bmatrix} s_{1,g-1} \\ s_{1,g-1}^f \end{bmatrix} + \begin{bmatrix} \frac{h_{g-1}^f}{2} & \frac{h_{g-1}^m}{2} \\ h_{g-1}^f & 0 \end{bmatrix} \begin{bmatrix} \mathbb{E}[H_{1,g-1,f}^X] \\ \mathbb{E}[H_{1,g-1,m}^X] \end{bmatrix}. \quad (4)$$

For the autosomes, for each sex δ , the expectation of the admixture fraction depends only on the total (non-sex-specific) contributions from the source populations and the corresponding expectation in the previous generation [32, eq. 19],

$$\mathbb{E}[H_{1,g,\delta}^A] = \begin{cases} s_{1,0} & g = 1 \\ s_{1,g-1} + h_{g-1} \mathbb{E}[H_{1,g-1,\delta}^A] & g \geq 2. \end{cases} \quad (5)$$

By contrast, the expectations of the female and male X-chromosomal admixture fractions depend also on the sex-specific contributions in the previous generation (eq. (4)).

Because the inheritance pattern is identical for X chromosomes and autosomes in females, the mean X-chromosomal admixture in females in eq. (4) matches that of the autosomes before equality of the female and male autosomal means is applied [32, eq. 18]. Whereas female and male autosomal admixture random variables are identically distributed and their expectations can be written with a

generic δ , the corresponding random variables differ for X chromosomes, and the coupled recursion in eqs. (3) and (4) cannot be quickly reduced to one equation. The mean X-chromosomal admixture in males (eq. (4)), however, has a similar form to the autosomal mean (eq. (5)).

We use the general mechanistic model presented here to derive closed-form expressions for the expected X-chromosomal admixture under two specific models of admixture, a single admixture event and constant admixture over time (Results).

Data analysis: estimating admixture in African Americans

Many admixture studies have reported X-chromosomal and autosomal admixture estimates for African Americans (Table 2). For illustration, we focus on data from Cheng et al. [34], who provided one of the largest samples, reporting estimates of quantities close to those that appear in our model. Cheng et al. [34] estimated admixture fractions of the X chromosome and autosomes for 15,280 African-Americans from 14 studies.

The mean admixture on the X chromosome reported by Cheng et al. [34] combines females and males into a single estimate, whereas we consider separate quantities for X-chromosomal admixture in females and males. In our notation, the mean X-chromosomal admixture across all individuals, $\mathbb{E}[H_{1,g}^X]$, is a weighted average of admixture in females and males that takes into account the fraction of the sample that is female. Thus, for a sample divided into females and males in proportions p_f and $p_m = 1 - p_f$, respectively, we can write

$$\mathbb{E}[H_{1,g}^X] = p_f \mathbb{E}[H_{1,g,f}^X] + p_m \mathbb{E}[H_{1,g,m}^X]. \quad (6)$$

To compare admixture estimates from data to the mean admixture across individuals produced under mechanistic models, we compute the Euclidean distance between predictions for $\mathbb{E}[H_{1,g}^A]$, $\mathbb{E}[H_{1,g,f}^X]$, and $\mathbb{E}[H_{1,g,m}^X]$ and observed values in the data, weighting the predicted female and male X-chromosomal admixture by eq. (6). Denoting by q_A and q_X the observed mean admixture in a specific source group for autosomes and the X chromosome, we evaluate

$$D = \sqrt{\left[q_A - \mathbb{E}[H_{1,g}^A]\right]^2 + \left[q_X - \left(p_f \mathbb{E}[H_{1,g,f}^X] + p_m \mathbb{E}[H_{1,g,m}^X]\right)\right]^2}. \quad (7)$$

This quantity D measures the fit of a prediction under a model to the actual estimates of X-chromosomal and autosomal admixture for a data set that reports three quantities: estimates of admixture for a specific source population for autosomes and for the X chromosome, and the fraction of the admixed sample consisting of females.

Results

To analyze the properties of X-chromosomal admixture in relation to the female and male contributions of two source populations, we consider two specific versions of our general admixture model. We then reinterpret African-American admixture estimates in light of these scenarios.

A single admixture event

First, we study a special case in which no further contributions from the source population occur after the admixed population is founded in generation $g = 1$. In this scenario of a single admixture event [35], $s_{1,g}^f = s_{1,g}^m = s_{2,g}^f = s_{2,g}^m = 0$, and $h_g^f = h_g^m = h_g = 1$, for all $g \geq 1$.

Applying eqs. (3) and (4), we have for $g = 1$,

$$\begin{bmatrix} \mathbb{E}[H_{1,1,f}^X] \\ \mathbb{E}[H_{1,1,m}^X] \end{bmatrix} = \begin{bmatrix} s_{1,0} \\ s_{1,0}^f \end{bmatrix}. \quad (8)$$

For $g \geq 2$,

$$\begin{bmatrix} \mathbb{E}[H_{1,g,f}^X] \\ \mathbb{E}[H_{1,g,m}^X] \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \mathbb{E}[H_{1,g-1,f}^X] \\ \mathbb{E}[H_{1,g-1,m}^X] \end{bmatrix}. \quad (9)$$

For $g \geq 2$, the expected admixture fraction for a male X chromosome is simply the expected admixture fraction for a female X chromosome from the previous generation, or $\mathbb{E}[H_{1,g,m}^X] = \mathbb{E}[H_{1,g-1,f}^X]$. We can use this identity between the expected female admixture in generation $g - 1$ and male admixture in generation g to simplify the system of equations. For $g \geq 3$, we then have

$$\mathbb{E}[H_{1,g,f}^X] = \frac{\mathbb{E}[H_{1,g-1,f}^X] + \mathbb{E}[H_{1,g-2,f}^X]}{2}. \quad (10)$$

Denoting by y_g the quantity $2^{g-1}\mathbb{E}[H_{1,g,f}^X]$, using eq. (10), for $g \geq 3$, we have

$$\frac{y_g}{2^{g-1}} = \frac{\frac{y_{g-1}}{2^{g-2}} + \frac{y_{g-2}}{2^{g-3}}}{2}.$$

Multiplying both sides by 2^{g-1} , we have a recursion $y_g = y_{g-1} + 2y_{g-2}$, with $y_1 = s_{1,0}$ and $y_2 = s_{1,0} + s_{1,0}^f$. Then for $g \geq 3$, y_g can be written $A_g s_{1,0} + B_g s_{1,0}^f$, where A_g and B_g satisfy

$$A_g = A_{g-1} + 2A_{g-2} \quad (11)$$

$$B_g = B_{g-1} + 2B_{g-2}, \quad (12)$$

with $A_1 = 1$, $A_2 = 1$, $B_1 = 0$, and $B_2 = 1$. Noting that $B_3 = 1$, we immediately observe that for $g \geq 3$, $B_g = A_{g-1}$, so that $y_g = A_g s_{1,0} + A_{g-1} s_{1,0}^f$. Then

$$\mathbb{E}[H_{1,g,f}^X] = \frac{A_g s_{1,0} + A_{g-1} s_{1,0}^f}{2^{g-1}}. \quad (13)$$

The recursion for A_g in eq. (11) with the initial conditions $A_0 = 0$ and $A_1 = 1$ gives the recursion for the Jacobsthal numbers, for which the closed form is $A_g = [2^g - (-1)^g]/3$ (Online Encyclopedia of Integer Sequences A001045 website). We then have for $g \geq 1$,

$$\mathbb{E}[H_{1,g,f}^X] = \left[2 + \left(-\frac{1}{2}\right)^g \right] \frac{s_{1,0}^f}{3} + \left[1 - \left(-\frac{1}{2}\right)^g \right] \frac{s_{1,0}^m}{3}. \quad (14)$$

Using eq. (9), for $g \geq 2$,

$$\mathbb{E}[H_{1,g,m}^X] = \left[2 + \left(-\frac{1}{2}\right)^{g-1} \right] \frac{s_{1,0}^f}{3} + \left[1 - \left(-\frac{1}{2}\right)^{g-1} \right] \frac{s_{1,0}^m}{3}. \quad (15)$$

For the case of a single admixture event, with no further gene flow from the source populations to the admixed population, we use eqs. (14) and (15) to understand the behavior over time of the mean admixture on the X chromosome. Notably, unlike for the autosomes under a single-admixture scenario [32], the mean X-chromosomal admixture depends on the sex-specific contributions from the source populations, as well as on the time since admixture. With no sex bias ($s_{1,0}^f = s_{1,0}^m = s_{1,0}$), the mean X-chromosomal admixture is the same as the autosomal admixture, which is constant in time, equaling simply $s_{1,0}$, the total contribution from S_1 .

Eqs. (14) and (15) oscillate over time (Fig. 2), as they incorporate a negative fraction raised to a power. The long-term limit of the X-chromosomal admixture fraction is

$$\lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,f}^X] = \lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,m}^X] = \frac{2}{3}s_{1,0}^f + \frac{1}{3}s_{1,0}^m. \quad (16)$$

Because $\mathbb{E}[H_{1,g,m}^X] = \mathbb{E}[H_{1,g-1,f}^X]$ for all $g \geq 2$, the expected admixture fraction sampled in males approaches the same limit over time as that sampled in females. The limit of the mean X-chromosomal admixture fraction fits the 2:1 ratio by which the X chromosome is inherited, following eq. (1), with the variables in eq. (1) viewed as the initial admixture values $s_{1,0}^f$ and $s_{1,0}^m$. Thus, the limiting admixture, but not the transient admixture, matches the simple linear combination.

For recent admixture, we can calculate the difference between the expected admixture under the model and the limiting value. For $g \geq 1$, we have

$$\mathbb{E}[H_{1,g,f}^X] - \lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,f}^X] = \frac{s_{1,0}^f - s_{1,0}^m}{3} \left(-\frac{1}{2}\right)^g, \quad (17)$$

and for $g \geq 2$,

$$\mathbb{E}[H_{1,g,m}^X] - \lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,m}^X] = \frac{s_{1,0}^f - s_{1,0}^m}{3} \left(-\frac{1}{2}\right)^{g-1}. \quad (18)$$

The differences in eqs. (17) and (18) provide a measure of the difference of a transient single-admixture model from the simpler linear combination in eq. (1), which is constant in time and does not consider differences between the mean admixture in females and males from the admixed population. The $g \rightarrow \infty$ limit in eq. (16) agrees with eq. (1), but for small g , the differences in eqs. (17) and (18) can be as large as $\pm \frac{1}{3}$, decreasing by a factor of 2 each generation. For a fixed g , the maximal absolute difference occurs when the sex bias is largest, that is, when $s_{1,0}^f - s_{1,0}^m$ is 1 or -1 . At the other extreme, with no sex bias and $s_{1,0}^f = s_{1,0}^m$, the simple linear combination exactly describes the X-chromosomal admixture throughout the history of the admixed population.

For six values of g , Figure 3 plots eq. (17) as a function of the difference between female and male contributions from S_1 , $s_{1,0}^f - s_{1,0}^m$. With no sex bias, $s_{1,0}^f = s_{1,0}^m$, eq. (17) is zero, and our model follows eq. (1). Additionally, as g increases, the difference between our time-dependent model and eq. (1) becomes smaller, as eq. (1) gives the limiting behavior of the single-admixture model.

Constant, non-zero admixture over time

Next, we consider the special case of constant, non-zero contributions from the source populations to the admixed population over time. As in Goldberg et al. [32], we can rewrite the sex-specific parameters as constants, $(s_{1,g}^f, s_{1,g}^m, s_{2,g}^f, s_{2,g}^m) = (s_1^f, s_1^m, s_2^f, s_2^m)$, for all $g \geq 1$. We maintain separate

parameters for the founding contributions, $s_{1,0}^f$, $s_{1,0}^m$, $s_{2,0}^f$, and $s_{2,0}^m$. In this setting, h_f and h_m cannot both be 1, and at least one among s_1^f , s_1^m , s_2^f , and s_2^m must be nonzero.

Using a generating function approach, we derive a closed-form solution for the mean X-chromosomal admixture fractions, $\mathbb{E}[H_{1,g,f}^X]$ and $\mathbb{E}[H_{1,g,m}^X]$. The mean depends on the number of generations of constant admixture, g , the initial conditions $s_{1,0}^f$ and $s_{1,0}^m$, and the sex-specific contributions from the two source populations, s_1^f , s_1^m , s_2^f , and s_2^m . In the Appendix, we show that for each $g \geq 1$,

$$\mathbb{E}[H_{1,g,f}^X] = P_1 r_1^{g+1} + P_2 r_2^{g+1} + P_3, \quad (19)$$

where P_1 , P_2 , and P_3 are defined in eqs. (36)-(38), and r_1 and r_2 in eqs. (34) and (35). Using eqs. (27) and (19), we can write an expression for the male X-chromosomal admixture fraction,

$$\mathbb{E}[H_{1,g,m}^X] = \begin{cases} s_{1,0}^f & g = 1 \\ s_1^f + h^f (P_1 r_1^g + P_2 r_2^g + P_3) & g \geq 2. \end{cases} \quad (20)$$

The limits of the mean X-chromosomal admixture fractions in eqs. (19) and (20) are

$$\lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,f}^X] = \frac{s_1^f + s_1^m + s_1^f h^m}{s_1^f + s_1^m + s_2^f + s_2^m + h^m (s_1^f + s_2^f)} \quad (21)$$

$$\lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,m}^X] = \frac{2s_1^f + h^f s_1^m}{s_1^f + s_1^m + s_2^f + s_2^m + h^m (s_1^f + s_2^f)}. \quad (22)$$

Unlike in the case of a single admixture event, the limit over time of the mean fraction of admixture from S_1 for a constant admixture process is not a simple 2:1 weighting of the female and male contributions from S_1 . Notably, recalling $s_1^f + h^f + s_2^f = s_1^m + h^m + s_2^m = 1$, the limiting admixture fraction from S_1 depends on the contributions both from S_1 and from S_2 . The mean X-chromosomal admixture depends on the sex-specific contributions and cannot be reduced in terms of s_1 and s_2 only, as was possible for the autosomes [32, eq. 37].

For the autosomes, the limiting mean is the ratio of contributions from S_1 to the total contributions from S_1 and S_2 combined [32, eq. 37], $\mathbb{E}[H_{1,g,\delta}^A] = s_1/(s_1 + s_2)$. As was observed for the autosomes, the limiting mean X-chromosomal admixture in females can be viewed as the fractional contribution of X chromosomes from S_1 in relation to the total number of X chromosomes from both source populations. Therefore, in eq. (21), the numerator has the same form as the denominator, incorporating only contributions from S_1 . X chromosomes from S_1 present in a population of females from the admixed population come from one of three origins in the limit, identified by the three terms of the numerator. They can be directly contributed from S_1 females or males, giving the terms s_1^f and s_1^m . Alternatively, they can be contributed from admixed males who in turn received them from S_1 females, producing the term $s_1^f h^m$; viewed in the limit, contributions from admixed females are already subsumed in the s_1^f and s_1^m terms. Similar reasoning can be used to understand the limiting male admixture fraction, noting that the limiting mean for males is the sum of two quantities, the product of the limiting female mean and the fraction of females from the admixed population, h^f , and the new influx of female contributions from S_1 , or s_1^f (eq. (27)).

The limits over time of the means in eqs. (21) and (22) depend only on the continuing contributions, not on the founding parameters. With no sex bias, $s_1^f = s_1^m = s_1$ and $h^f = h^m = h$, and we can

simplify the limiting mean female and male X-chromosomal admixture fractions:

$$\lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,f}^X] = \lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,m}^X] = \frac{2s_1 + hs_1}{2 - h - h^2} = \frac{s_1}{1 - h} = \frac{s_1}{s_1 + s_2}. \quad (23)$$

This limit is equivalent to the limiting mean of autosomal admixture from Verdu & Rosenberg [33, eq. 31]. That is, with no sex bias, the limiting X-chromosomal mean matches the limiting autosomal mean, $\lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,\delta}^A] = \lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,f}^X] = \lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,m}^X]$.

Figure 4 plots the expectations of the X-chromosomal and autosomal admixture fractions over time for two scenarios with the same difference in female and male contributions, but different directions of sex bias. In Figure 4A, more males enter from S_1 and more females from S_2 , leading to a lower mean admixture on the X chromosome than on the autosomes, with the mean in males smaller than the mean in females. Conversely, in Figure 4B, more females enter from S_1 and more males from S_2 , leading to a larger mean for X chromosomes than for autosomes. The parameters are set so that for both panels, autosomal admixture is constant in time at $\frac{1}{2}$ [32, eq. 37]. In both cases, the limiting mean X-chromosomal admixture in females lies intermediate between the mean autosomal admixture and the mean X-chromosomal admixture in males.

Figure 5 plots the mean X-chromosomal admixture fraction for females and males (eqs. (19) and (20)) with the mean autosomal admixture fraction when both source populations have a female bias in contributions, $s_1^f > s_1^m$ and $s_2^f > s_2^m$. Though S_1 has an excess of females, the mean X-chromosomal admixture is less than the autosomal admixture. Whereas an excess of females from a given source population might be viewed as generating elevated X-chromosomal admixture, when both populations have a sex bias in the same direction, the source population with the larger sex bias dominates the signal. It is tempting to interpret lower X-chromosomal than autosomal admixture as a signal of male-biased admixture from S_1 and female-biased admixture from S_2 , but Figure 5 plots an example where admixture is female-biased in both source populations, and the simple interpretation of opposite biases in the two source populations is incorrect. A similar example could also have been produced with male bias in both source populations.

We note that in the limit as the parameters s_1^f , s_1^m , s_2^f , and s_2^m simultaneously approach zero, the constant-admixture model approaches the single-event model. Thus, we expect the limiting mean female and male X-chromosomal admixture fractions in the constant-admixture model to approach the corresponding limits in the single-event model. Indeed, by taking the limits of eqs. 19 and 20 as s_1^f , s_1^m , s_2^f , and s_2^m approach 0, we obtain eqs. 14 and 15, respectively. Interestingly, as we expect the trajectories of the X-chromosomal admixture fractions under constant admixture to approach corresponding single-event trajectories, we also expect the oscillatory pattern of the single-event model to occur in the constant admixture model for sufficiently small s_1^f , s_1^m , s_2^f , and s_2^m . Indeed, we can find instances of oscillation around the limit in the constant-admixture model as well as in the single-admixture model, for example, for $s_1^f = s_1^m = s_2^f = s_2^m = 0.05$ and initial condition $(s_{1,0}^f, s_{1,0}^m, s_{2,0}^f, s_{2,0}^m) = (0.25, 0.5, 0.75, 0.5)$ (Figure 6). As the continuing contributions in this case have no sex bias, the mean female and male X-chromosomal admixture and the mean autosomal admixture tend to the same limit.

Sex-biased admixture in African Americans

Studies of the genetic admixture history of African-American populations have consistently reported evidence for male-biased gene flow from Europe [15, 16, 19, 22, 23, 31]. Lind et al. [19] and Bryc et

al. [31] estimated female and male contributions from Africa and Europe using mean admixture estimates for the X chromosome and autosomes. Both analyses followed the linear combination method of eq. (1), in which the mean autosomal admixture averages the female and male contributions, while the mean X-chromosomal admixture weights the female and male contributions by $\frac{2}{3}$ and $\frac{1}{3}$, respectively. In our notation, the framework can be written

$$\mathbb{E}[H_{1,g,\delta}^A] = \frac{1}{2}s_1^f + \frac{1}{2}s_1^m \quad (24)$$

$$\mathbb{E}[H_{1,g,\delta}^X] = \frac{2}{3}s_1^f + \frac{1}{3}s_1^m, \quad (25)$$

with $\mathbb{E}[H_{1,g,\delta}^A]$ and $\mathbb{E}[H_{1,g,\delta}^X]$ estimated from data.

We have demonstrated, however, that in autosomal admixture models, eq. (24) suggests a single admixture event [32], and in X-chromosomal admixture models, eq. (25) suggests the limit of a single admixture process (replacing parameters for the continuing contributions in these equations by corresponding parameters for admixture in the initial generation, $s_{1,0}^f$ and $s_{1,0}^m$, as in eq. (16)). For recent admixture in the single-admixture model, or in a model with continuing admixture after the founding of the admixed population, the mean admixture fractions from S_1 for the X chromosome depend on the sex-specific contributions in a more complex way, incorporating the sex-specific contributions from S_2 (eqs. (14), (15), (19)-(22)). Use of eqs. (24) and (25) implies consideration of a single-admixture model in its temporal limit, or otherwise is unsuited to the single-admixture and continuing admixture scenarios.

We used our refined predictions about $\mathbb{E}[H_{1,g,\delta}^X]$ to estimate the sex-specific contributions for the African-American population from Cheng et al. [34] under two different models, a single admixture event with various times since admixture, and a constant admixture process with $g = 15$. Although both models underestimate the spatial and temporal complexity of the true admixture history of African Americans, these approximations enable us to illustrate the way in which estimates of the sex bias depend on assumptions about the admixture model.

The exact timing of the onset of significant admixture in the African-American admixture process is unknown, but assuming a generation time of 20-27 years, the Trans-Atlantic slave trade to North America can be regarded as having begun 14-20 generations ago [36]. As the mean X-chromosomal admixture fraction is near its limit for g in this range, both in the single-admixture model and in the constant-admixture model, the specific $g \in [14, 20]$ only minimally changes the results. For our constant-admixture analysis, we chose a single value $g = 15$ for consistency.

Single admixture event: Using the mean X-chromosomal and autosomal admixture reported by Cheng et al. [34] (Table 2), we calculated the sex-specific contributions $s_{1,0}^f$ and $s_{1,0}^m$ for specified values of $g \in [2, 5]$, and in the $g \rightarrow \infty$ limit. As Cheng et al. [34] reported admixture in a combined sample of females and males, we used eq. (6) to calculate the X-chromosomal admixture fraction $\mathbb{E}[H_{1,g}^X]$ as a function of p_f , p_m , $\mathbb{E}[H_{1,g,f}^X]$, and $\mathbb{E}[H_{1,g,m}^X]$. Writing $\mathbb{E}[H_{1,g,f}^X]$ and $\mathbb{E}[H_{1,g,m}^X]$ as functions of g , $s_{1,0}^f$, and $s_{1,0}^m$ (eqs. (14) and (15)) then generated an equation for $\mathbb{E}[H_{1,g}^X]$ in terms of p_f , p_m , g , $s_{1,0}^f$, and $s_{1,0}^m$, into which we inserted the reported values $q_X = 0.845$ for $\mathbb{E}[H_{1,g}^X]$ and $(p_f, p_m) = (0.557, 0.443)$. We obtained a second equation for the autosomal admixture fraction from Goldberg et al. [32, eq. 34], $\mathbb{E}[H_{1,g}^A] = s_{1,0} = (s_{1,0}^f + s_{1,0}^m)/2$, using $q_A = 0.807$ for the empirical observation of $\mathbb{E}[H_{1,g}^A]$. We then solved the pair of equations for $s_{1,0}^f$ and $s_{1,0}^m$, with g fixed.

For small g (2, 3, 4, 5) as well as in the $g \rightarrow \infty$ limit, $s_{1,0}^f$, representing the contribution of females from the African source population, exceeds 0.9, and $s_{2,0}^f$, representing the female European contribution, is below 0.1 (Table 3). For males, the African contribution $s_{1,0}^m$ is ~ 0.7 , whereas the European contribution $s_{2,0}^m$ is ~ 0.3 . Because of the oscillation of the mean X-chromosomal admixture (Fig. 2), the estimated ratio of male to female contributions oscillates around the limiting values. For contributions from Europe, in the $g \rightarrow \infty$ limit we estimate $s_{2,0}^m/s_{2,0}^f \approx 3.89$. That is, in the temporal limit of a model of a single admixture event, for each female from Europe, ~ 3.89 males contributed to the gene pool of African Americans. The direction of this ratio is reversed for African contributions, with ~ 1.33 females for every African male. If we assume that the admixture was recent, for example $g \leq 5$, then these values do vary substantially, with larger deviations from the limiting value of 3.89 occurring under more recent admixture.

Constant admixture over time: Assuming $g = 15$, we computed the mean female and male X-chromosomal admixture (eqs. (19) and (20)), and mean autosomal admixture given the four sex-specific contributions (eq. (5)), on a grid of possible parameter values, $s_1^f, s_1^m, s_2^f, s_2^m \in [0, 1]$ using 0.01 increments. We fixed the initial contributions $s_{1,0}^f = s_{1,0}^m = 0.5$, though by $g = 15$ the mean nears its limit, erasing the signal of the initial conditions (eqs. (21) and (22)). We calculated the distance D between the Cheng et al. [34] data and the computed values (eq. (7)).

The parameters s_1^f, s_1^m, s_2^f , and s_2^m are not uniquely identifiable from the data, as we continue to have two equations, one describing autosomal admixture and one for X-chromosomal admixture, but now we consider four unknowns. We therefore examined sets of parameter values that produced $D \leq 0.01$. Distance cutoffs of 0.001 and 0.1 gave rise to similar ranges for each parameter.

Figure 7 plots the set of parameter values that generate $D \leq 0.01$. Because the female and male contributions separately sum to one, the possible range of each set of contributions is represented by a unit simplex. Figure 7A plots the marginal distributions of the set of parameter values with $D \leq 0.01$. Contributions from S_1 (Africans) vary over most of the permissible parameter space, but the S_2 contributions (Europeans) take their values from a narrower range. Figure 7B plots D on the space of female contributions, (s_1^f, h^f, s_2^f) , for values of (s_1^m, h^m, s_2^m) fixed at the median of the set of parameter vectors that produce $D \leq 0.01$, $s_1^m = 0.47$, $h^m = 1 - s_1^m - s_2^m = 0.34$, and $s_2^m = 0.19$. Figure 7C plots D as a function of the male contributions, (s_1^m, h^m, s_2^m) , with female contributions fixed at the median from the distribution of parameter values for which $D \leq 0.01$, $(s_1^f, h^f, s_2^f) = (0.64, 0.29, 0.07)$. In both panels, the parameter sets closest to the data in terms of D permit a large range of possible contributions from S_1 , Africans, but only a smaller range of possible contributions, both female and male, from Europeans, S_2 .

Figure 8A plots the natural logarithm of the ratio of male to female contributions for parameter sets with $D \leq 0.01$. With no sex bias, this quantity is zero, with more males than females it is positive, and with more females than males it is negative. Most points plotted from S_1 , Africans, are female-biased, whereas contributions from S_2 , Europeans, are predominantly male-biased. The median ratio of males to females from Europe is 2.67 males per female, compared to a median 1.32 females per male from Africa. Note, however, that these patterns do not hold for all parameter sets. For 26.15% of values with $D \leq 0.01$, more male than female contributions occur from Africa. For 9.25% of values with $D \leq 0.01$, more female than male contributions occur from Europe.

Although the pattern is initially surprising, scenarios of female bias from Europe or male bias from Africa accord with the phenomenon depicted in Figure 5. In the former scenario, both populations

have an excess of females, and the strong female bias in the contributions from Africa can overwhelm the number of European X chromosomes in the gene pool of African Americans, producing greater African admixture on the X chromosome than on autosomes. This effect is demonstrated in Figure 8B, which plots the logarithm of the ratio of male to female contributions from S_2 against the corresponding quantity from S_1 . The ratios of male to female contributions in S_2 on the y-axis and S_1 on the x-axis are correlated. The values that are negative on the y-axis, indicating a female bias from S_2 , also show the strongest female bias from S_1 , that is, the most negative $\ln(s_1^m/s_1^f)$ on the x-axis (an additional 2.34% of parameter sets show no sex bias in the European population; the female and male contributions are equal and $\ln(s_1^m/s_1^f) = 0$). Analogously, the strongest male bias from Africa is also observed when the excess of males from Europe is largest (Fig. 7B).

Discussion

Under a two-sex mechanistic admixture model with sex-biased admixture, we have demonstrated that the relationship between X-chromosomal and autosomal admixture fractions depends both on the time since admixture and on the model of admixture, and does not simply follow a prediction from the fractions of X chromosomes and autosomes present in females. Using the mechanistic framework, we have reinterpreted African American admixture values computed from a non-mechanistic perspective, estimating sex-specific parameters and levels of sex bias in African and European source populations. This analysis uncovers a counterintuitive case in which female-biased or male-biased contributions in the same direction occur both from Africans and Europeans in a manner consistent with estimated mean X-chromosomal and autosomal admixture fractions in African Americans, rather than an African female bias and a European male bias.

Estimating sex bias using X-chromosomal and autosomal admixture

Differences between X-chromosomal and autosomal admixture estimates are sometimes used to demonstrate the occurrence of sex bias, even without estimating sex-specific contributions [21, 23, 24, 26, 27, 37]. Higher estimated admixture from a population S_1 for X chromosomes than for autosomes is taken as evidence of a female bias from S_1 and a male bias from a second population S_2 . We have found, however, that the pattern is in principle compatible with two additional possibilities (Fig. 7): a female bias from both populations, with a larger female bias from S_1 , or a male bias from both populations, with a larger male bias from S_2 .

We considered a general mechanistic model of admixture, focusing specifically on models of a single admixture event and constant admixture. For a single admixture event, mean autosomal admixture over time is constant, depending only on the total contributions from the source populations, and not on the sex-specific contributions or time [32]. Mean X-chromosomal admixture, on the other hand, is variable over time—in such a way that given the number of generations g and observed levels of X-chromosomal and autosomal admixture, the initial sex-specific contributions from the two source populations are identifiable. Because of the oscillation of the mean X-chromosomal admixture over time, depending on the time since admixture, the sex bias in admixture contributions can be either overestimated or underestimated using the $\frac{2}{3}-\frac{1}{3}$ linear combination (eq. (1)).

For constant admixture, estimated mean X-chromosomal and autosomal admixture values do not uniquely identify the female and male contributions over time. Therefore, instead of point estimates for the sex-specific contributions and their ratio, we have reported a measure of the compatibility

of parameter values and data over a range of values of the parameters (Figs. 6 and 7). It is possible, however, that by examining higher moments of the distribution of admixture estimates across individuals, the sex-specific contributions might become identifiable [32].

Theoretical population genetics of the X chromosome

The complex signature of admixture we have observed for the X chromosome is reminiscent of other X-chromosomal phenomena in theoretical population genetics, including results related to effective population size, allele-frequency dynamics, and numbers of ancestors. For effective population size, N_e , when female and male population sizes are equal, the ratio of X-chromosomal to autosomal values is $\frac{3}{4}$ [38–40]. In the same way that various forms of sex difference between females and males—in such parameters as the number of individuals, the variance of reproductive success, and migration rates—cause the basic X-to-autosomal N_e ratio to differ from $\frac{3}{4}$ [1, 3, 4, 6, 8, 12–14, 39, 41], transient dynamics and ongoing admixture cause the X-chromosomal admixture fraction to differ from a linear combination of female and male contributions with coefficients $\frac{2}{3}$ and $\frac{1}{3}$.

The X-chromosomal admixture fraction in our single-admixture model also has similarities to the allele-frequency trajectory in the approach to Hardy-Weinberg equilibrium in a one-locus X-chromosomal model. In that model [42–44], the frequency of an allele on female X chromosomes depends on the frequency in both females and males in the previous generation, whereas the frequency on male X chromosomes depends only on the frequency in females. If an allele frequency differs between females and males at the first generation, then during the approach to equilibrium, the frequency in males matches the corresponding frequency in females of the previous generation; both the female and male frequencies oscillate around the same limit. The equilibrium frequency in turn has a $\frac{2}{3}$ contribution from the initial female frequency and $\frac{1}{3}$ from the male frequency.

All these phenomena from the one-locus model—female values dependent on both female and male values from the previous generation, male values dependent only on the female value, males lagging one generation behind females, oscillations around the same limit, and a limiting linear combination of $\frac{2}{3}$ and $\frac{1}{3}$ —appear in our X-chromosomal admixture model with a single admixture event. An additional parallel lies in the difference between the X chromosome and the autosomes: in the one-locus model for the X chromosome, Hardy-Weinberg equilibrium is achieved in the limit, and for the autosomes, it is achieved in one generation. For admixture on the autosomes, the mean admixture is constant after the founding of the admixed population, and for the X chromosome, the mean X-chromosomal admixture approaches a limit rather than remaining constant in time.

Yet another connection to theoretical population genetics of the X chromosome comes from the form of the recursive sequence in eq. (13). The Jacobsthal sequence in the single-admixture model is obtained from a generalization of the Fibonacci sequence [45]. Both the Jacobsthal and Fibonacci sequences appeared in breeding system models as early as a century ago [42, 46–48], with the Jacobsthal sequence providing the numerators of allele frequencies at a sex-linked locus in a model that amounts mathematically to a special case of our single-event admixture model [42, 47]; the Fibonacci sequence is well-known as the number of genealogical ancestors of a haplodiploid system g generations ago [49]. Specifically, for the pair of X chromosomes in a female, the number of ancestors is entry $g + 2$ in the Fibonacci sequence F_n , or equivalently, the sum of the numbers of maternal and paternal ancestors, F_{g+1} and F_g , respectively. The number of genealogical ancestors for a male X chromosome in generation g is F_{g+1} . Our recursion $A_n = A_{n-1} + 2A_{n-2}$ (eq. (11)) generates a different sequence, but its form is similar to the Fibonacci recursion $F_n = F_{n-1} + F_{n-2}$.

Sex-biased admixture in African-Americans

Our analysis of sex-biased admixture in African Americans has similarities but a number of noteworthy differences from earlier nonmechanistic analyses. Under our single-admixture model, the estimate of ~ 4 European males for every European female accords with estimated ratios of 3 to 4 from previous studies [19, 31]. For even g , more African females and European males contribute than in the $g \rightarrow \infty$ limit; odd g values produce the opposite pattern.

For the constant admixture model, the median estimated ratios of male to female contributions from Europe and Africa are ~ 2.67 males per female from Europe and ~ 1.32 females per male from Africa (Figs. 6 and 7). The estimated male bias in contributions from Europe is lower for a constant admixture history than for a single admixture event.

For both a single admixture event and constant admixture over time, the point estimates of the ratio of males to females have a larger male contribution from Europe and female contribution from Africa. In the single-admixture model, the excess African ancestry on the X chromosome compared to the autosomes implies a female-biased contribution from Africa and a male-biased contribution from Europe. For constant admixture, surprisingly, a sex bias in one source alone can produce the observed pattern without a sex bias in the other (Fig. 7). In fact, $\sim 27\%$ of the parameter sets most similar to the data ($D \leq 0.01$) show no sex bias or have larger male than female contributions from Africa; $\sim 12\%$ show no sex bias or have larger female than male contributions from Europe. Note that in obtaining this result, we have given equal weight to all values of D below a cutoff, rather than giving more weight to parameter choices producing lower D ; however, owing to the existence of ridges in the parameter space that have similarly small values of D , for different choices of the cutoff, similar results are produced.

An excess of African mitochondrial and European Y-chromosomal haplotypes has been described in African Americans [15, 16, 19, 22, 50]. Similar phenomena to those we observed for X chromosomes and autosomes could affect the Y-mtDNA comparisons: in other words, a strong excess of males from Europe compared to European females, even if more males than females contribute from African populations, could give rise to a larger fraction of European Y chromosomes in the African-American gene-pool without a female bias for the African source population. Such a process could also explain the excess African ancestry of mtDNA.

Conclusions

We have presented a model to describe the effect of admixture on the X chromosome, deriving a theoretical framework that considers the impact of sex bias during the admixture process. The model can be used to understand the sex-specific contributions from source populations to an admixed population. We have found that because of model dependence, time dependence, and a lack of identifiability of admixture parameters from mean admixture alone, a variety of admixture processes and parameter values might be compatible with estimates of the mean admixture on X chromosomes and the autosomes. Factors we have not studied, including mutation, recombination, selection, and genetic drift can differentially affect the X chromosome and autosomes [10, 13, 41, 51–55], potentially further complicating the estimation of admixture parameters. Examination in the context of sex-biased models of more detailed summaries of admixture patterns, including higher moments of the admixture distributions [32, 33, 56] and distributions of the lengths of admixture tracts [56–58], will assist in refining the estimation of sex-biased admixture histories.

We demonstrated that female and male X-chromosomal admixture have different expectations under mechanistic admixture models. With the exception of studies restricted to males [19], however, past studies of sex-biased admixture have generally reported composite X-chromosomal admixture estimates only in pooled collections of females and males. We recommend that such studies report estimates separately in females and males. Even if the values are similar, the difference between X-chromosomal admixture estimates in females and males contains information about parameters of the admixture process. Autosomal admixture estimates obtained only in females and only in males are identically distributed [32], so that a pooled estimate is more sensible than in the X-chromosomal case; nevertheless, autosomal admixture estimates reported separately in females and males can enable an informative comparison with a corresponding pair of X-chromosomal values.

Curiously, we have found that African American admixture patterns do not necessarily imply a simultaneous African female bias and European male bias in the genetic ancestry of the population, and that in both source populations, it is in principle possible on the basis of genetic admixture patterns that both source populations had female bias or that both had male bias, albeit at different magnitudes. The latter interpretation, of male biases both in Europeans *and* in Africans is plausible in light of historical scholarship on demographic contributions to African Americans [36, 59–61], documenting both the well-known contributions of male European slave owners and asymmetric mating practices involving African females and European males, as well as an overrepresentation of males among African slaves arriving in North America. For less well-documented cases, our model highlights that if a difference in X-chromosomal and autosomal admixture is observed, it is important to consider the possibility that rather than opposite sex biases in the two populations, both populations might have the same type of sex bias.

Acknowledgments. We thank Carlos Bustamante, Filippo Disanto, Michael D. Edge, Marc Feldman, Ethan Jewett, Hua Tang, and Paul Verdu for useful discussions. We acknowledge support from NIH grant R01 HG005855 and a National Science Foundation Graduate Research Fellowship.

Web Resources

On-Line Encyclopedia of Integer Sequences A001045 website, <http://oeis.org/A001045>.

References

- [1] Ramachandran, S., Rosenberg, N. A., Zhivotovsky, L. A., and Feldman, M. W. (2004). Robustness of the inference of human population structure: a comparison of the X-chromosomal and autosomal microsatellites. *Human Genomics* 1, 87–97.
- [2] Wilkins, J. F. and Marlowe, F. W. (2006). Sex-biased migration in humans: what should we expect from genetic data? *Bioessays* 28, 290–300.
- [3] Hammer, M. F., Mendez, F. L., Cox, M. P., Woerner, A. E., and Wall, J. D. (2008). Sex-biased evolutionary forces shape genomic patterns of human diversity. *PLoS Genetics* 4, e1000202.
- [4] Ramachandran, S., Rosenberg, N. A., Feldman, M. W., and Wakeley, J. (2008). Population differentiation and migration: coalescence times in a two-sex island model for autosomal and X-linked loci. *Theoretical Population Biology* 74, 291–301.
- [5] Bustamante, C. D. and Ramachandran, S. (2009). Evaluating signatures of sex-specific processes in the human genome. *Nature Genetics* 41, 8–10.

- [6] Keinan, A., Mullikin, J. C., Patterson, N., and Reich, D. (2009). Accelerated genetic drift on chromosome X during the human dispersal out of Africa. *Nature Genetics* *41*, 66–70.
- [7] Casto, A. M., Li, J. Z., Absher, D., Myers, R., Ramachandran, S., and Feldman, M. W. (2010). Characterization of X-linked SNP genotypic variation in globally distributed human populations. *Genome Biol* *11*, R10.
- [8] Emery, L. S., Felsenstein, J., and Akey, J. M. (2010). Estimators of the human effective sex ratio detect sex biases on different timescales. *American Journal of Human Genetics* *87*, 848–856.
- [9] Hammer, M. F., Woerner, A. E., Mendez, F. L., Watkins, J. C., Cox, M. P., and Wall, J. D. (2010). The ratio of human X chromosome to autosome diversity is positively correlated with genetic distance from genes. *Nature Genetics* *42*, 830–831.
- [10] Labuda, D., Lefebvre, J.-F., Nadeau, P., and Roy-Gagnon, M.-H. (2010). Female-to-male breeding ratio in modern humans—an analysis based on historical recombinations. *American Journal of Human Genetics* *86*, 353–363.
- [11] Lambert, C. A., Connelly, C. F., Madeoy, J., Qiu, R., Olson, M. V., and Akey, J. M. (2010). Highly punctuated patterns of population structure on the X chromosome and implications for African evolutionary history. *American Journal of Human Genetics* *86*, 34–44.
- [12] Gottipati, S., Arbiza, L., Siepel, A., Clark, A. G., and Keinan, A. (2011). Analyses of X-linked and autosomal genetic variation in population-scale whole genome sequencing. *Nature Genetics* *43*, 741–743.
- [13] Heyer, E., Chaix, R., Pavard, S., and Austerlitz, F. (2012). Sex-specific demographic behaviours that shape human genomic variation. *Molecular Ecology* *21*, 597–612.
- [14] Arbiza, L., Gottipati, S., Siepel, A., and Keinan, A. (2014). Contrasting X-linked and autosomal diversity across 14 human populations. *American Journal of Human Genetics* *94*, 827–844.
- [15] Parra, E. J., Marcini, A., Akey, J., Martinson, J., Batzer, M. A., Cooper, R., Forrester, T., Allison, D. B., Deka, R., Ferrell, R. E., et al. (1998). Estimating African American admixture proportions by use of population-specific alleles. *American Journal of Human Genetics* *63*, 1839–1851.
- [16] Parra, E. J., Kittles, R. A., Argyropoulos, G., Pfaff, C. L., Hiester, K., Bonilla, C., Sylvester, N., Parrish-Gause, D., Garvey, W. T., Jin, L., et al. (2001). Ancestral proportions and admixture dynamics in geographically defined African Americans living in South Carolina. *American Journal of Physical Anthropology* *114*, 18–29.
- [17] Bedoya, G., Montoya, P., García, J., Soto, I., Bourgeois, S., Carvajal, L., Labuda, D., Alvarez, V., Ospina, J., Hedrick, P. W., et al. (2006). Admixture dynamics in Hispanics: a shift in the nuclear genetic ancestry of a South American population isolate. *Proceedings of the National Academy of Sciences* *103*, 7234–7239.
- [18] Wilkins, J. F. (2006). Unraveling male and female histories from human genetic data. *Current Opinion in Genetics and Development* *16*, 611–617.
- [19] Lind, J. M., Hutcheson-Dilks, H. B., Williams, S. M., Moore, J. H., Essex, M., Ruiz-Pesini, E., Wallace, D. C., Tishkoff, S. A., O’Brien, S. J., and Smith, M. W. (2007). Elevated male European and female African contributions to the genomes of African American individuals. *Human Genetics* *120*, 713–722.
- [20] Berniell-Lee, G., Plaza, S., Bosch, E., Calafell, F., Jourdan, E., Césari, M., Lefranc, G., and Comas, D. (2008). Admixture and sexual bias in the population settlement of La Réunion Island (Indian Ocean). *American Journal of Physical Anthropology* *136*, 100–107.
- [21] Wang, S., Ray, N., Rojas, W., Parra, M. V., Bedoya, G., Gallo, C., Poletti, G., Mazzotti, G., Hill, K., Hurtado, A. M., et al. (2008). Geographic patterns of genome admixture in Latin American Mestizos. *PLoS Genetics* *4*, e1000037.
- [22] Stefflova, K., Dulik, M. C., Pai, A. A., Walker, A. H., Zeigler-Johnson, C. M., Gueye, S. M., Schurr, T. G., and Rebbeck, T. R. (2009). Evaluation of group genetic ancestry of populations from Philadelphia and Dakar in the context of sex-biased admixture in the Americas. *PLoS One* *4*, e7842.

- [23] Bryc, K., Auton, A., Nelson, M. R., Oksenberg, J. R., Hauser, S. L., Williams, S., Froment, A., Bodo, J.-M., Wambebe, C., Tishkoff, S. A., et al. (2010). Genome-wide patterns of population structure and admixture in West Africans and African Americans. *Proceedings of the National Academy of Sciences USA* *107*, 786–791.
- [24] Bryc, K., Velez, C., Karafet, T., Moreno-Estrada, A., Reynolds, A., Auton, A., Hammer, M., Bustamante, C. D., and Ostrer, H. (2010). Genome-wide patterns of population structure and admixture among Hispanic/Latino populations. *Proceedings of the National Academy of Sciences USA* *107*, 8954–8961.
- [25] Quintana-Murci, L., Harmant, C., Quach, H., Balanovsky, O., Zaporozhchenko, V., Bormans, C., van Helden, P. D., Hoal, E. G., and Behar, D. M. (2010). Strong maternal Khoisan contribution to the South African coloured population: a case of gender-biased admixture. *American Journal of Human Genetics* *86*, 611–620.
- [26] Beleza, S., Campos, J., Lopes, J., Araújo, I. I., Almada, A. H., Correia e Silva, A., Parra, E. J., and Rocha, J. (2012). The admixture structure and genetic variation of the archipelago of Cape Verde and its implications for admixture mapping studies. *PLoS One* *7*, e51103.
- [27] Moreno-Estrada, A., Gravel, S., Zakharia, F., McCauley, J. L., Byrnes, J. K., Gignoux, C. R., Ortiz-Tello, P. A., Martínez, R. J., Hedges, D. J., Morris, R. W., et al. (2013). Reconstructing the population genetic history of the Caribbean. *PLoS Genetics* *9*, e1003925.
- [28] Verdu, P., Becker, N. S., Froment, A., Georges, M., Grugni, V., Quintana-Murci, L., Hombert, J.-M., Van der Veen, L., Le Bomin, S., Bahuchet, S., et al. (2013). Sociocultural behavior, sex-biased admixture, and effective population sizes in Central African Pygmies and non-Pygmies. *Molecular Biology and Evolution* *30*, 918–937.
- [29] Marcheco-Teruel, B., Parra, E. J., Fuentes-Smith, E., Salas, A., Buttenschön, H. N., Demontis, D., Torres-Español, M., Marín-Padrón, L. C., Gómez-Cabezas, E. J., Álvarez-Iglesias, V., et al. (2014). Cuba: exploring the history of admixture and the genetic basis of pigmentation using autosomal and uniparental markers. *PLoS Genetics* *10*, e1004488.
- [30] Salzano, F. M. and Sans, M. (2014). Interethnic admixture and the evolution of Latin American populations. *Genetics and Molecular Biology* *37*, 151–170.
- [31] Bryc, K., Durand, E. Y., Macpherson, J. M., Reich, D., and Mountain, J. L. (2015). The genetic ancestry of African Americans, Latinos, and European Americans across the United States. *American Journal of Human Genetics* *96*, 37–53.
- [32] Goldberg, A., Verdu, P., and Rosenberg, N. A. (2014). Autosomal admixture levels are informative about sex bias in admixed populations. *Genetics* *198*, 1209–1229.
- [33] Verdu, P. and Rosenberg, N. A. (2011). A general mechanistic model for admixture histories of hybrid populations. *Genetics* *189*, 1413–1426.
- [34] Cheng, C.-Y., Kao, W. H. L., Patterson, N., Tandon, A., Haiman, C. A., Harris, T. B., Xing, C., John, E. M., Ambrosone, C. B., Brancati, F. L., et al. (2009). Admixture mapping of 15,280 African Americans identifies obesity susceptibility loci on chromosomes 5 and X. *PLoS Genetics* *5*, e1000490.
- [35] Long, J. C. (1991). The genetic structure of admixed populations. *Genetics* *127*, 417–428.
- [36] Eltis, D. and Richardson, D. (2010). *Atlas of the Transatlantic Slave Trade*. (New Haven: Yale University Press).
- [37] Cox, M. P., Karafet, T. M., Lansing, J. S., Sudoyo, H., and Hammer, M. F. (2010). Autosomal and X-linked single nucleotide polymorphisms reveal a steep Asian–Melanesian ancestry cline in eastern Indonesia and a sex bias in admixture rates. *Proceedings of the Royal Society of London Series B: Biological Sciences* *277*, 1589–1596.
- [38] Wright, S. (1933). Inbreeding and homozygosis. *Proceedings of the National Academy of Sciences USA* *19*, 411–420.

- [39] Caballero, A. (1995). On the effective size of populations with separate sexes, with particular reference to sex-linked genes. *Genetics* *139*, 1007–1011.
- [40] Hartl, D. L. and Clark, A. G. (2007). *Principles of Population Genetics* 4th ed. (Sunderland: Sinauer).
- [41] Charlesworth, B. (2001). The effect of life-history and mode of inheritance on neutral genetic variability. *Genetical Research* *77*, 153–166.
- [42] Jennings, H. S. (1916). The numerical results of diverse systems of breeding. *Genetics* *1*, 53–89.
- [43] Wright, S. (1969). *Evolution and the Genetics of Populations: Vol. 2. The Theory of Gene Frequencies*. (Chicago: University of Chicago Press).
- [44] Crow, J. F. and Kimura, M. (1970). *An Introduction to Population Genetics Theory*. (Minneapolis: Burgess).
- [45] Koshy, T. (2001). *Fibonacci and Lucas numbers with applications*. (Hoboken: John Wiley & Sons).
- [46] Jennings, H. S. (1914). Formulae for the results of inbreeding. *American Naturalist* *48*, 693–696.
- [47] Jennings, H. S. (1916). The numerical results of diverse systems of breeding. *Proceedings of the National Academy of Sciences USA* *2*, 45–50.
- [48] Wright, S. (1921). Systems of mating II. The effects of inbreeding on the genetic composition of a population. *Genetics* *6*, 124–143.
- [49] Basin, S. (1963). The Fibonacci sequence as it appears in nature. *Fibonacci Quarterly* *1*, 53–56.
- [50] McLean Jr., D. C., Spruill, I., Argyropoulos, G., Page, G. P., Shriver, M. D., and Garvey, W. T. (2005). Mitochondrial DNA (mtDNA) haplotypes reveal maternal population genetic affinities of Sea Island Gullah-speaking African Americans. *American Journal of Physical Anthropology* *127*, 427–438.
- [51] Hedrick, P. W. (2007). Sex: differences in mutation, recombination, selection, gene flow, and genetic drift. *Evolution* *61*, 2750–2771.
- [52] Pool, J. E. and Nielsen, R. (2007). Population size changes reshape genomic patterns of diversity. *Evolution* *61*, 3001–3006.
- [53] Ellegren, H. (2009). The different levels of genetic diversity in sex chromosomes and autosomes. *Trends in Genetics* *25*, 278–284.
- [54] Veeramah, K. R., Gutenkunst, R. N., Woerner, A. E., Watkins, J. C., and Hammer, M. F. (2014). Evidence for increased levels of positive and negative selection on the X chromosome versus autosomes in humans. *Molecular Biology and Evolution* *31*, 2267–2282.
- [55] Ségurel, L., Wyman, M. J., and Przeworski, M. (2014). Determinants of mutation rate variation in the human germline. *Annual Review of Genomics and Human Genetics* *15*, 47–70.
- [56] Gravel, S. (2012). Population genetics models of local ancestry. *Genetics* *191*, 607–619.
- [57] Pool, J. E. and Nielsen, R. (2009). Inference of historical changes in migration rate from the lengths of migrant tracts. *Genetics* *181*, 711–719.
- [58] Liang, M. and Nielsen, R. (2014). The lengths of admixture tracts. *Genetics* *197*, 953–967.
- [59] Fredrickson, G. M. (1982). *White Supremacy: A Comparative Study of American and South African History*. (Oxford: Oxford University Press).
- [60] Painter, N. I. (2006). *Creating Black Americans: African-American History and its Meanings, 1619 to the Present*. (Oxford: Oxford University Press).
- [61] Davis, F. J. (2010). *Who is Black? One Nation’s Definition*. (University Park: Penn State Press).
- [62] Tian, C., Hinds, D. A., Shigeta, R., Kittles, R., Ballinger, D. G., and Seldin, M. F. (2006). A genomewide single-nucleotide-polymorphism panel with high ancestry information for African American admixture mapping. *American Journal of Human Genetics* *79*, 640–649.

Appendix: Solving for $\mathbb{E}[H_{1,g,\delta}^X]$ under constant admixture

Here, we obtain the closed form for the mean X-chromosomal admixture fraction in a random female and a random male from the admixed population in the special case of constant admixture over time. Using the constant sex-specific parameters, s_1^f , s_1^m , s_2^f , and s_2^m , we rewrite the expectations in eq. (4). For $g = 1$, $\mathbb{E}[H_{1,1,f}^X] = s_{1,0}$, and for $g \geq 2$, we have

$$\mathbb{E}[H_{1,g,f}^X] = s_1 + \frac{1}{2} \left(h^f \mathbb{E}[H_{1,g-1,f}^X] + h^m \mathbb{E}[H_{1,g-1,m}^X] \right). \quad (26)$$

Similarly, sampling males in the admixed population, for $g = 1$, $\mathbb{E}[H_{1,1,m}^X] = s_{1,0}^f$. For $g \geq 2$,

$$\mathbb{E}[H_{1,g,m}^X] = s_1^f + h^f \mathbb{E}[H_{1,g-1,f}^X]. \quad (27)$$

Using equations (26) and (27), we derive a generating function for the mean female X-chromosomal admixture fraction, which we then use to find closed-form expressions for $\mathbb{E}[H_{1,g,f}^X]$ and $\mathbb{E}[H_{1,g,m}^X]$. Because $\mathbb{E}[H_{1,g,m}^X]$ depends only on constants and $\mathbb{E}[H_{1,g-1,f}^X]$, we first find an expression for $\mathbb{E}[H_{1,g,f}^X]$, which we then use to report $\mathbb{E}[H_{1,g,m}^X]$.

First, using eq. (27), we can rewrite eq. (26) as a second-order recursion of a single variable,

$$\mathbb{E}[H_{1,g,f}^X] = \begin{cases} s_{1,0} & g = 1 \\ s_1 + \frac{1}{2}(s_{1,0}h^f + s_{1,0}^f h^m) & g = 2 \\ s_1 + \frac{h^f}{2} \mathbb{E}[H_{1,g-1,f}^X] + \frac{h^m}{2}(s_1^f + h^f \mathbb{E}[H_{1,g-2,f}^X]) & g \geq 3. \end{cases} \quad (28)$$

We simplify the notation by defining $z_g = \mathbb{E}[H_{1,g,f}^X]$. For $g \geq 3$, we have,

$$z_g = c_1 + c_2 z_{g-1} + c_3 z_{g-2}, \quad (29)$$

with $c_1 = s_1 + s_1^f h^m/2$, $c_2 = h^f/2$, and $c_3 = h^f h^m/2$. Eq. (28) gives $z_1 = s_{1,0}$ and $z_2 = s_1 + \frac{1}{2}(s_{1,0}h^f + s_{1,0}^f h^m)$.

Define a generating function $Z(x) = \sum_{g=1}^{\infty} z_g x^g$ whose coefficients z_g represent the values of $\mathbb{E}[H_{1,g,f}^X]$ in each generation. As the admixed population does not yet exist in generation 0, $\mathbb{E}[H_{1,g,f}^X]$ and $Z(x)$ are undefined for $g = 0$. For convenience, we therefore work with $W(x) = Z(x)/x$, setting $w_g = z_{g+1}$ for $g \geq 0$. We then have

$$W(x) = \sum_{g=0}^{\infty} z_{g+1} x^g = \sum_{g=0}^{\infty} w_g x^g, \quad (30)$$

and $w_g = c_1 + c_2 w_{g-1} + c_3 w_{g-2}$ for $g \geq 2$. Using eq. (30), it follows that

$$\begin{aligned} W(x) &= w_0 + w_1 x + \sum_{g=2}^{\infty} (c_1 + c_2 w_{g-1} + c_3 w_{g-2}) x^g \\ &= w_0 + w_1 x + c_1 \sum_{g=2}^{\infty} x^g + c_2 x \sum_{g=2}^{\infty} w_{g-1} x^{g-1} + c_3 x^2 \sum_{g=2}^{\infty} w_{g-2} x^{g-2} \\ &= w_0 + w_1 x + c_1 \left(\frac{x^2}{1-x} \right) + c_2 x [W(x) - w_0] + c_3 x^2 W(x). \end{aligned} \quad (31)$$

Solving for $W(x)$, we have

$$W(x) = \frac{(w_0 + w_1x - c_2w_0x)(1-x) + c_1x^2}{(1-x)(1-c_2x - c_3x^2)}. \quad (32)$$

We can decompose the expression in eq. (32), producing

$$W(x) = \frac{P_1}{1-r_1x} + \frac{P_2}{1-r_2x} + \frac{P_3}{1-x}, \quad (33)$$

where r_1 and r_2 are reciprocals of the two roots of $1 - c_2x - c_3x^2$,

$$r_1 = \frac{-c_2 + \sqrt{c_2^2 + 4c_3}}{2c_3} = \frac{h^f + \sqrt{(h^f)^2 + 8h^fh^m}}{4} \quad (34)$$

$$r_2 = \frac{-c_2 - \sqrt{c_2^2 + 4c_3}}{2c_3} = \frac{h^f - \sqrt{(h^f)^2 + 8h^fh^m}}{4}. \quad (35)$$

Setting eq. (32) equal to eq. (33), we have

$$P_1 = \frac{(w_0r_1 + w_1 - c_2w_0)(r_1 - 1) + c_1}{(r_1 - r_2)(r_1 - 1)} \quad (36)$$

$$P_2 = \frac{(w_0r_2 + w_1 - c_2w_0)(r_2 - 1) + c_1}{(r_2 - r_1)(r_2 - 1)} \quad (37)$$

$$P_3 = \frac{c_1}{(1-r_1)(1-r_2)}. \quad (38)$$

The Taylor expansion of eq. (33) around $x = 0$ then gives

$$\begin{aligned} W(x) &= P_1 \sum_{g=0}^{\infty} r_1^g x^g + P_2 \sum_{g=0}^{\infty} r_2^g x^g + P_3 \sum_{g=0}^{\infty} x^g, \\ &= \sum_{g=0}^{\infty} (P_1 r_1^g + P_2 r_2^g + P_3) x^g. \end{aligned}$$

Therefore, for $g \geq 0$, $w_g = P_1 r_1^g + P_2 r_2^g + P_3$, and the closed-form expression for the X-chromosomal female mean admixture fraction in generation $g \geq 1$, $\mathbb{E}[H_{1,g,f}^X]$, is $z_g = P_1 r_1^{g+1} + P_2 r_2^{g+1} + P_3$. We report this result in the main text as eq. (19), using it to obtain $\mathbb{E}[H_{1,g,m}^X]$ in eq. (20).

Because for $h^f, h^m \in [0, 1]$, r_1 monotonically increases in h^f and h^m and r_2 monotonically decreases, the maxima and minima of r_1 and r_2 occur at the boundaries of the closed interval $[0, 1]$. Using eqs. (34) and (35), we have $r_1 \in [0, 1)$ and $r_2 \in (-\frac{1}{2}, 0]$; we exclude $r_1 = 1$ and $r_2 = -\frac{1}{2}$, as h^f and h^m cannot both be 1. Because $|r_1|, |r_2| < 1$, the mean X-chromosomal admixture fractions in eqs. (19) and (20) approach limits as $g \rightarrow \infty$. Using eqs. (19) and (20), we can find expressions for the limits of the mean of the X-chromosomal admixture fractions,

$$\begin{aligned} \lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,f}^X] &= P_3 \\ \lim_{g \rightarrow \infty} \mathbb{E}[H_{1,g,m}^X] &= s_1^f + h^f P_3, \end{aligned}$$

which can be simplified to give eqs. (21) and (22).

Figures

Figure 1: Sex-specific mechanistic model of admixture. At each generation g , females and males from each of two source populations, S_1 and S_2 , contribute to the admixed population, H . Contributions can vary in time. The fraction of admixture from source population $\alpha \in \{1, 2\}$ for chromosomal type $\gamma \in \{A, X\}$ in an admixed individual of sex $\delta \in \{f, m\}$ is $H_{\alpha,g,\delta}^\gamma$. This model, by considering different chromosomal types, generalizes the model of Goldberg et al. [32].

Figure 2: The mean of the X-chromosomal admixture fraction over time in females (red) and males (blue) from the admixed population for a single admixture event (eqs. (14) and (15)). The mean X-chromosomal admixture oscillates in approaching its limit (eq. (16)). The limiting mean, shown in black, is the same for X chromosomes in both females and males. (A) $s_{1,0}^f = 0.25$, $s_{1,0}^m = 0.5$. (B) $s_{1,0}^f = 0.5$, $s_{1,0}^m = 0.25$.

Figure 3: The difference between the expectation of the female X-chromosomal admixture fraction and its limit for $g \in [1, 6]$ as a function of the difference between female and male contributions from S_1 for a single admixture event (eq. (17)). As $g \rightarrow \infty$, this quantity approaches zero. For small g , however, the limit $\frac{2}{3}s_{1,0}^f + \frac{1}{3}s_{1,0}^m$ (eq. (16)) is a poor approximation. The difference oscillates so that the slope of the line is negative when g is odd and positive when g is even.

Figure 4: The expectation of the mean X-chromosomal and autosomal admixture fractions over time, with their associated limits, for constant ongoing admixture. (A) Male-biased admixture from population S_1 . (B) Female-biased admixture from population S_1 . The initial condition is $(s_{1,0}^f, s_{1,0}^m, s_{2,0}^f, s_{2,0}^m) = (0.5, 0.5, 0.5, 0.5)$. The autosomal admixture is constant over time because $s_{1,0} = s_{2,0} = \frac{1}{2}$ and $s_1 = s_2$; it is the same in both panels because it does not depend on the sex-specific contributions. The X-chromosomal admixture is different in females and males; it is smaller than the autosomal mean for male biased-admixture from population 1, and larger for female-biased admixture. The X-chromosomal mean is plotted using eqs. (19)-(22). The autosomal mean uses eq. 37 from Goldberg et al. [32].

Figure 5: The expectation of the mean X-chromosomal and autosomal admixture fractions over time for constant admixture, with female-biased contributions from both source populations. The initial condition is $(s_{1,0}^f, s_{1,0}^m, s_{2,0}^f, s_{2,0}^m) = (0.5, 0.5, 0.5, 0.5)$. The mean X-chromosomal admixture from S_1 , $\mathbb{E}[H_{1,g,\delta}^X]$, is smaller than the mean autosomal admixture, $\mathbb{E}[H_{1,g,\delta}^A]$, even though S_1 has an excess of females, $s_1^f > s_1^m$. The expectation of X-chromosomal admixture is plotted using eqs. (19) and (20). The autosomal mean uses eq. 37 from Goldberg et al. [32].

Figure 6: The expectation of the mean X-chromosomal and autosomal admixture fractions over time for constant admixture, in a case with small continuing contributions. The initial condition is $(s_{1,0}^f, s_{1,0}^m, s_{2,0}^f, s_{2,0}^m) = (0.25, 0.5, 0.75, 0.5)$. With sex bias in the founding generation and small continuing contributions, the mean X-chromosomal admixture has a pattern resembling the oscillating behavior seen for a single admixture event. The expectation of X-chromosomal admixture is plotted using eqs. (19) and (20). The autosomal mean uses eq. 37 from Goldberg et al. [32].

Figure 7: Sex-specific contributions estimated from the data of Cheng et al. [34]. (A) The range, median, and 25th and 75th percentiles of the sets of sex-specific contributions for which the Euclidean distance D (eq. (7)) between model-predicted admixture and observed admixture from Cheng et al. [34], was at most 0.01. The range of values for s_2^f and s_2^m , the contributions representing Europeans (S_2), is smaller than those representing Africans (S_1). (B) Female contributions

as a function of D . (C) Male contributions as a function of D . For (B), the male contributions are fixed at their median values producing $D \leq 0.01$, $(s_1^m, h^m, s_2^m) = (0.47, 0.34, 0.19)$. For (C), the female contributions are fixed in a corresponding manner, at $(s_1^f, h^f, s_2^f) = (0.64, 0.29, 0.07)$. The space of possible values for (s_1^m, h^m, s_2^m) or (s_1^f, h^f, s_2^f) is the unit simplex. Parameter values were tested at increments of 0.01 for each quantity.

Figure 8: The natural logarithm of the ratio of male to female contributions in African Americans, as inferred from the data of Cheng et al. [34]. (A) The range (excluding infinity, produced when a parameter value is zero), median, and 25th and 75th percentiles of the natural logarithm of the ratio of male to female contributions from S_1 (Africans) and S_2 (Europeans) separately for the sex-specific contributions that produced $D \leq 0.01$ (eq. (7)). Values from Africans, S_1 , are largely below 0, or female-biased, whereas contributions from Europeans, S_2 , are mostly above 0 and male-biased. Approximately 26.15% of the contributions from Africans are male-biased and 9.25% from Europeans are female-biased. This pattern is typically observed when a still larger sex bias occurs in the other population. (B) The logarithm of the ratio of male and female contributions from S_2 on the y-axis and the corresponding ratio for S_1 on the x-axis, plotted by the density of points in 0.05 square bins. For the cases with male bias in Africa ($\ln(s_1^m/s_1^f) > 0$), the level of male bias in Europe is also positive; for the cases with female bias in Europe ($\ln(s_2^m/s_2^f) < 0$), the level of female bias from Africa is also negative. Parameter sets in which at least one parameter is 0, and therefore, we have values of $+\infty$ or $-\infty$ for $\ln(s_1^m/s_1^f)$ or $\ln(s_2^m/s_2^f)$, appear in bins on the edge of the plot for convenience. These bins contain a substantial number of parameter sets.

Table 1: Recursion for the X-chromosomal and autosomal admixture fractions of a randomly chosen female or male from an admixed population at generation g , given a set of parents. With two source populations and the contributions from the admixed population to itself, a random individual has nine possible parental pairings, for each of which the probability is listed.

Table 2: Studies that published estimates of the X-chromosomal and autosomal admixture fractions of an African American population. As the X chromosomes of females and males were considered jointly in the studies, we list the fraction of the sample that was female. In each study, the X chromosome shows elevated African ancestry compared to the autosomes. Values in the table were compiled from quantities reported in the studies, or by taking 1 minus an African estimate to obtain a European estimate, or vice versa, with the following exceptions. For Bryc et al. [31], the “autosome” estimates are genome-wide, including the X chromosome; the percentage of females was kindly provided by K. Bryc (pers. comm.). For Bryc et al. [23], we obtained an African X-chromosomal estimate by visual inspection of their Figure 4g, and we counted the number of females from their supplementary material. For Tian et al. [62], we computed the values from information reported in their Figure 5 caption.

Table 3: Estimates of the sex-specific contributions from Africa and Europe to African Americans, based on the Cheng et al. [34] data, inferred under our model for a single admixture event. The estimates depend on the time since admixture. All estimates show male-biased contributions from Europe and female-based contributions from Africa.

Figure 1

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Case	Female parent's population	Male parent's population	Probability	Admixture in males		Admixture in females	
				$H_{1,g,m}^A$	$H_{1,g,m}^X$	$H_{1,g,f}^A$	$H_{1,g,f}^X$
1	S_1	S_1	$s_{1,g-1}^f s_{1,g-1}^m$	1	1	1	1
2	S_1	H	$s_{1,g-1}^f h_{g-1}^m$	$\frac{1 + H_{1,g-1,m}^A}{2}$	1	$\frac{1 + H_{1,g-1,m}^A}{2}$	$\frac{1 + H_{1,g-1,m}^X}{2}$
3	S_1	S_2	$s_{1,g-1}^f s_{2,g-1}^m$	$\frac{1}{2}$	1	$\frac{1}{2}$	$\frac{1}{2}$
4	H	S_1	$h_{g-1}^f s_{1,g-1}^m$	$\frac{1 + H_{1,g-1,f}^A}{2}$	$H_{1,g-1,f}^X$	$\frac{1 + H_{1,g-1,f}^A}{2}$	$\frac{1 + H_{1,g-1,f}^X}{2}$
5	H	H	$h_{g-1}^f h_{g-1}^m$	$\frac{H_{1,g-1,f}^A + H_{1,g-1,m}^A}{2}$	$H_{1,g-1,f}^X$	$\frac{H_{1,g-1,f}^A + H_{1,g-1,m}^A}{2}$	$\frac{H_{1,g-1,f}^X + H_{1,g-1,m}^X}{2}$
6	H	S_2	$h_{g-1}^f s_{2,g-1}^m$	$\frac{H_{1,g-1,f}^A}{2}$	$H_{1,g-1,f}^X$	$\frac{H_{1,g-1,f}^A}{2}$	$\frac{H_{1,g-1,f}^X}{2}$
7	S_2	S_1	$s_{2,g-1}^f s_{1,g-1}^m$	$\frac{1}{2}$	0	$\frac{1}{2}$	$\frac{1}{2}$
8	S_2	H	$s_{2,g-1}^f h_{g-1}^m$	$\frac{H_{1,g-1,m}^A}{2}$	0	$\frac{H_{1,g-1,m}^A}{2}$	$\frac{H_{1,g-1,m}^X}{2}$
9	S_2	S_2	$s_{2,g-1}^f s_{2,g-1}^m$	0	0	0	0

Table 1

Reference	Mean ancestry estimates (%)				Percent female
	African autosome ($E[H_{1,g}^A]$)	European autosome ($E[H_{2,g}^A]$)	African X chromosome ($E[H_{1,g}^X]$)	European X chromosome ($E[H_{2,g}^X]$)	
Cheng et al. 2009 ³⁴	80.7	19.3	84.5	15.5	55.7
Bryc et al. 2015 ³¹	73.2	24.0	76.9	19.8	52.3
Bryc et al. 2010a ²³	77	23	81	19	66.3
Lind et al. 2007 ¹⁹	80	20	87.9	12.1	0
Tian et al. 2006 ⁶²	81.8	18.2	87.1	12.9	Not reported

Table 2

Generations since admixture	African contributions			European contributions		
	$s_{1,0}^f$	$s_{1,0}^m$	$s_{1,0}^f/s_{1,0}^m$	$s_{2,0}^f$	$s_{2,0}^m$	$s_{2,0}^m/s_{2,0}^f$
2	0.943	0.671	1.40	0.057	0.329	5.77
3	0.912	0.702	1.30	0.088	0.298	3.39
4	0.926	0.688	1.35	0.074	0.312	4.22
5	0.919	0.695	1.32	0.081	0.305	3.77
∞	0.921	0.693	1.33	0.079	0.307	3.89

Table 3