1 Title

2 Successful asexual lineages of the Irish potato Famine pathogen are triploid

3

4 Authors

5 Ying Li^{1*}, Qian Zhou^{1,2}, Kun Qian¹, Theo van der Lee³, Sanwen Huang^{1,2*}

6

7 **The oomycete *Phytophthora infestans* was the causal agent of the Irish Great Famine**
8 **and is a recurring threat to global food security¹. The pathogen can reproduce both**
9 **sexually and asexually and has a potential to adapt both abiotic and biotic environment².**
10 **Although in many regions the A1 and A2 mating types coexist, the far majority of**
11 **isolates belong to few clonal, asexual lineages³. As other oomycetes, *P. infestans* is**
12 **thought to be diploid during the vegetative phase of its life cycle³, but it was observed**
13 **that trisomy correlated with virulence and mating type locus⁴ and that polyploidy can**
14 **occur in some isolates^{5,6}. It remains unknown about the frequency of polyploidy**
15 **occurrence in nature and the relationship between ploidy level and sexuality. Here we**
16 **discovered that the sexuality of *P. infestans* isolates correlates with ploidy by comparison**
17 **of microsatellite fingerprinting, genome-wide polymorphism, DNA quantity, and**
18 **chromosome numbers. The sexual progeny of *P. infestans* in nature are diploid, whereas**
19 **the asexual lineages are mostly triploids, including successful clonal lineages US-1 and**
20 **13_A2. This study reveals polyploidization as an extra evolutionary risk to this notorious**
21 **plant destroyer.**

22

23

¹Department of Biotech, Institute of Vegetables and Flowers, Chinese Academy of Agricultural Sciences, Beijing 100081, China. ²Biointeractions & Plant Health, Wageningen University & Research Center, Plant Research International, Wageningen 6700AA, The Netherlands. ³Agricultural Genome Institute at Shenzhen, Chinese Academy of Agricultural Sciences, Shenzhen 518124, China.

24 Understanding the mechanisms of rapid adaption in devastating plant pathogens is of
 25 critical importance for disease management. Plant oomycete pathogens employed a flexible
 26 reproduction system to gain dual advantages in creating both genotype diversity via sexual
 27 crosses and in achieving large population expansion via asexual cycles². Compared to other
 28 *Phytophthora* sister species, *P. infestans* has a much larger genome that is enriched in
 29 repetitive DNA (~74%). The genome can be largely divided into gene-dense and gene-sparse
 30 compartments, with the latter containing fast-evolving effector repertoire that is required for
 31 virulence and host adaptation^{7,8}.

32 An increasing number of reports showed that most *P. infestans* field strains from wide
 33 geographical distribution were identified as belonging to a few asexual, clonal lineages⁹. The
 34 first recorded clonal lineage HERB-1 was dominant in both Europe and North America
 35 between 1845 and 1896, which triggered the Great Irish Potato Famine⁵. Its replacement, the
 36 US-1 clonal lineage of the A1 mating type, firstly emerged in North America in the 20th
 37 century and dominated the globe^{5,10}. Rather recently (1990's), US-1 was supplanted by more
 38 aggressive lineages of the A2 mating type, including 13_A2 in Europe and Asia¹¹ as well as
 39 US-22 in North America^{12,13}. As accumulation of deleterious mutations would lead to rapid
 40 extinction of asexual lineages, a phenomenon coined as the Muller's ratchet¹⁴, it is puzzling
 41 that a few clonal lineages could adapt to a wide geographical area for a considerably long
 42 period. What is their common denominator? Although *P. infestans* was known as diploid for
 43 decades³, several studies indicated trisomy and polyploidy^{5,6}. Polyploidy can enhance the
 44 vigour in plants and buffer mutational load in asexual reproduction by masking deleterious
 45 alleles as reported in both fungi, plants and animals^{9,15,16}. It remains unclear to what extent
 46 polyploidy contributed to evolutionary advantage of clonal lineages and whether there is link
 47 between sexuality and ploidy.

Previously we used multiplex microsatellites to fingerprint 520 *P. infestans* isolates (Supplementary Table 1), including 397 isolates from nine major clonal lineages in four continents and 123 isolates from three sexual populations from the Netherlands, Mexico and Tunisia^{9,17,18}. In previous studies, the analysis was only based on the allele size, although differences of allele dosage were observed by peak height. To survey the frequency of polyploidy occurrence and to gain insights into the relationship of reproduction systems and ploidy levels, we re-analyzed the fingerprinting dataset by including scoring of allele peak height, besides molecular size of alleles. The ratio of allele peak heights allows assessing allele dosage at each microsatellite locus (Supplementary Fig.). Surprisingly, we discovered that eight of nine asexual lineages showed high ratio of triallelic loci (0.73-0.99), whereas the value (0.14-0.35) in the three sexual populations is significantly lower ($p=0.008$). The only exception is the clonal lineage in Northern China (CN-Northern), which is similar to sexual populations in the percentage of triallelic loci. CN-Northern belongs to the A1 mating type and it is a rather old population, as sub-populations evolved in the same lineage¹⁷. In China, seed potatoes are produced in the North and shipped elsewhere. Strict quarantine is taken for seed tubers to enter the area domestically and abroad. Therefore, the *P. infestans* population there should be rather isolated and remains as diploid.

To further analyze the genome-wide ploidy level, we re-sequenced 11 representative isolates for >50X genome depth from a meta-population from the Netherlands, a major source of international seed potato trade. The meta-population includes three asexual lineages (NL 13_A2, NL 06_A1, and NL 08_A1) and two sexual populations (NL pop 2 and NL pop3, Table 1). Sequences were aligned to the T30-4 reference genome to determine single nucleotide polymorphism (SNP). Relative read depth at each heterozygous SNP was used to determine ploidy level. For diploid genome, the mean of read counts at heterozygous

positions should have a single mode at 0.5 (1/2) when the allele ratio is 1:1, while there should be two modes, 0.33 and 0.67 (1/3 and 2/3) when the allele ratio is 1:2 and 2:1 for triploid genomes (Supplementary Table 2). Consistent with the microsatellite fingerprinting analyses, all the nine isolates from the sexual populations displayed diploid mode, whereas the three isolates from the asexual lineages showed triploid mode (Table 2, Fig. 1 and Supplementary Table 2). We also re-analyzed sequences of the seven isolates from a previous report⁵ (Table 2 and Fig. 1), which confirmed that all the four isolates from asexual lineage US-1 and EC-1 showed also triploid mode. These analyses indicate that sexuality and ploidy are strongly associated.

83

In addition to deep re-sequencing, we used flow cytometry and microscope observation to directly evaluate the DNA content and count chromosome number, respectively (Fig. 2). The isolate NL07434 (A2) from the sexual population NL pop2 was used as control, since it had been investigated previously⁵ as well as in this study. DNA content of the isolate NL08797 (A2) from NL pop 2 is the same as NL07434. As expected, DNA content of the three asexual isolate NL08080 (A1), NL08452 (A1), and NL07041 (A2) is about 1.5 times of that of NL07434 and NL08797, confirming that they are triploid. The average of chromosome number for isolates from sexual populations is 13.6, whereas 20.6 for isolates from asexual lineages (Table 2, Supplementary Table 3). DNA content measurement and chromosome counting further support that these asexual isolates are triploid.

94

In summary, *P. infestans* can no longer be considered to be only diploid, which should be taken into account in future studies. The observed triploidy must play an imperative role in the successful global epidemic of modern asexual lineages such as US-1, NL 13_A2, TU-1, NL 06_A1 and NL 08_A1. Polyploidization in *P. infestans* is likely to enhance fitness, as

99 reported in yeast that tetraploid had a higher beneficial mutation rate than haploid and diploid
100 strains¹⁹. Ancient or old asexual lineages such as HERB1 and CN Northern are diploids,
101 therefore it is worth further investigation on when and how asexual lineages became triploid.
102 Was the transition of the ploidy level due to survival pressure from disease resistance genes
103 that were incorporated into potato breeding after the Great Famine or linked to the frequent
104 use of pesticide deployed in the last 50 years to control this disease? This study revealed a
105 new dimension of genomic feature of the great evolutionary risk of *P. infestans*, which should
106 also be considered in future agricultural management.

107

Methods

Definition of asexual and sexual populations

This study only focuses on field isolates. We adopted the definition of clonality widely accepted in papers dealing with the population structure of pathogens²⁰. The definition of asexuality (clonal lineage) and sexuality (sexual population) in this study do not refer to the cytological mechanism of reproduction, but rather to the population structure that results from an absence or restriction of genetic recombination²⁰. The asexuality obtains wherever isolates with the same mating type show multilocus genotypes (MLGs) that are identical or nearly identical. In contrast sexual progeny shows extensive recombination of alleles.

Isolate collection

In this study, isolates from China¹⁷, The Netherlands⁹, Ecuador¹⁸, Tunisia (manuscript prepared), and Mexico were investigated (Table 2). The clonal lineages and sexual populations have been defined by MLGs in those previous studies. The clonal lineages and sexual populations used in this study have been defined by MLGs in the previous studies.

Microsatellite analysis

Twelve microsatellite markers were used²¹⁻²⁴. Amplification of the SSR markers was carried out as described by Li et. al²⁴. The amplicon was capillary electrophoresed on an automated ABI 3730 according to the manufacturer's instructions. SSR allele sizing was performed using GeneMapper v3.7 software (Applied Biosystems, USA).

Re-sequencing data and SNP calling

The reference genome sequence of the artificial strain T30-4 and the re-sequencing data used in this study were published in previous studies^{5,7,10,11}. The method of read mapping was used as described⁵. The re-sequenced isolates selected from previous studies were analyzed again here. Since the reference genome has 4921 supercontigs, we only analyzed the first 100 longest supercontigs.

The isolates were sequenced using the Illumina Hi-Seq 2000 sequencer. The sequencing averagely generated 100 million 100-bp paired-end reads for each isolate. The sequencing reads of each isolate were mapped to the T30-4 reference genome using BWA²⁵ and SNP calling was conducted subsequently using Sequence Alignment/Maptools²⁶. Several criteria was considered in SNP filtering: (1) a SNP should be bi-allelic between the isolate and T30-4 genomes; (2) at the SNP locus, the phred quality score of base sequencing and score of read mapping should be both higher than 30; (3) each allele of a SNP should be supported by at least 4 reads; (4) the non-reference allele frequency should be between 0.2-0.8 at a heterozygous SNP locus.

Flow cytometry

The nuclei were collected by simply chopping hyphae with a scalpel blade in cold phosphate buffered saline (PBS)²⁷. Another method used was zoospore cultures. Zoospores were induced at 4°C for two hours and harvested in excess of 10⁴ spores ml⁻¹. The nuclei were stained by 10ug/ml propidium iodide (Sigma). The samples were delivered into a laser BD FACSCalibur Flow Cytometer for data analysis.

Microscopy observation of chromosomes

The young hyphae stained by 50ug/ml propidium iodide (Sigma) were transferred to a microscope slide for observation. The observation was preformed with Zeiss LSM 780 Laser Scanning Confocal Microscope.

References

- 1 Jones, N. Planetary disasters: It could happen one night. *Nature***493**, 154-156 (2013).
- 2 McDonald, B. A. & Linde, C. Pathogen population genetics, evolutionary potential, and durable resistance. *Annu Rev Phytopathol***40**, 349-379 (2002).
- 3 Grunwald, N. J. & Flier, W. G. The biology of *Phytophthora infestans* at its center of origin. *Annu Rev Phytopathol***43**, 171-190 (2005).
- 4 van der Lee, T., Testa, A., Robold, A., van 't Klooster, J. & Govers, F. High-density genetic linkage maps of *Phytophthora infestans* reveal trisomic progeny and chromosomal rearrangements. *Genetics***167**, 1643-1661 (2004).
- 5 Yoshida, K. *et al.* The rise and fall of the *Phytophthora infestans* lineage that triggered the Irish potato famine. *Elife***2**, e01108 (2013).
- 6 Hamed, B. H. & Gisi, U. Generation of pathogenic F1 progeny from crosses of *Phytophthora infestans* isolates differing in ploidy. *Plant Pathology***62**, 708-718 (2013).
- 7 Haas, B. J. *et al.* Genome sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*. *Nature***461**, 393-398 (2009).
- 8 Raffaele, S. *et al.* Genome evolution following host jumps in the Irish potato famine pathogen lineage. *Science***330**, 1540-1543 (2010).
- 9 Li, Y. *et al.* Population dynamics of *Phytophthora infestans* in the Netherlands reveals expansion and spread of dominant clonal lineages and virulence in sexual offspring. *G3 (Bethesda)***2**, 1529-1540 (2012).
- 10 Martin, M. D. *et al.* Reconstructing genome evolution in historic samples of the Irish potato famine pathogen. *Nature Communication***4**, 2172 (2013).
- 11 Cooke, D. E. *et al.* Genome analyses of an aggressive and invasive lineage of the Irish potato famine pathogen. *PLoS Pathogen***8**, e1002940 (2012).
- 12 Hu, C.-H. *et al.* Recent genotypes of *Phytophthora infestans* in the Eastern United States reveal clonal populations and reappearance of mefenoxam sensitivity. *Plant Disease***96**, 1323-1330 (2012).
- 13 Danies, G. *et al.* An ephemeral sexual population of *Phytophthora infestans* in the Northeastern United States and Canada. *PLoS One***9**, e116354 (2014).
- 14 Muller, H. J. The relation of recombination to mutational advance. *Mutat Res***106**, 2-9 (1964).
- 15 Simon, J. C., Delmotte, F., Rispe, C. & Crease, T. Phylogenetic relationships between parthenogens and their sexual relatives: the possible routes to parthenogenesis in animals. *Biological Journal of the Linnean Society***79**, 151-163 (2003).
- 16 Kearney, M. Hybridization, glaciation and geographical parthenogenesis. *Trends Ecol Evol***20**, 495-502 (2005).
- 17 Li, Y. *et al.* Population structure of *Phytophthora infestans* in China – geographic clusters and presence of the EU genotype Blue_13. *Plant Pathology***62**, 932-942 (2013).
- 18 Delgado, R. A. *et al.* Large subclonal variation in *Phytophthora infestans* populations associated with Ecuadorian potato landraces. *Plant Pathology***62**, 1081-1088 (2013).
- 19 Selmecki, A. M. *et al.* Polyploidy can drive rapid adaptation in yeast. *Nature***519**, 349-352 (2015).
- 20 Tibayrenc, M. & Ayala, F. J. Reproductive clonality of pathogens: a perspective on pathogenic viruses, bacteria, fungi, and parasitic protozoa. *Proc Natl Acad Sci U S A***109**, E3305-3313 (2012).
- 21 Knapova, G. & Gisi, U. Phenotypic and genotypic structure of *Phytophthora infestans* populations on potato and tomato in France and Switzerland. *Plant Pathology***51**, 641-653 (2002).
- 22 Lees, A. K. *et al.* Novel microsatellite markers for the analysis of *Phytophthora infestans* populations. *Plant Pathology***55**, 311-319 (2006).
- 23 Li, Y. *et al.* A new set of highly informative SSR markers for *Phytophthora infestans* population analysis assembled into an efficient multiplex. *Molecular Ecology Resources***10**, 1098-1105 (2010).
- 24 Li, Y., Cooke, D. E. L., Jacobsen, E. & van der Lee, T. Efficient multiplex simple sequence repeat genotyping of the oomycete plant pathogen *Phytophthora infestans*. *Journal of Microbiological Methods***92**, 316-322 (2013).
- 25 Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics***25**, 1754-1760 (2009).
- 26 Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics***25**, 2078-2079 (2009).
- 27 Catal, M. *et al.* Heterokaryotic nuclear conditions and a heterogeneous nuclear population are observed by flow cytometry in *Phytophthora infestans*. *Cytometry A***77**, 769-775 (2010).

Supplementary Information includes one figure and three tables and is available in attached files.

Acknowledgements We thank G. J. T. Kessel and G. B. M. van den Bosch (Wageningen University) for providing *P. infestans* isolates. We appreciate Z. Zhang, B. Xie and S. Dong for useful discussion and E. Jacobsen (Wageningen University) and T. Städler (ETH Zurich) for critical reading. This work was supported by Chinese Special Fund for Agro-scientific Research in the Public Interest (201303018), National High Technology Research and Development Program (2013AA102603), National Natural Science Foundation of China (31201257), Ministry of Finance in China (1251610601001), Science and Technology Innovation Program of Chinese Academy of Agricultural Sciences (CAAS-ASTIP-IVFCAAS).

Author Contributions Y. L. and S. H. designed the study and wrote the paper. Y. L. performed SSR analysis and chromosome observation. Q. Z. analysed re-sequencing data. Q. K. performed flow cytometry. T. L. provided the raw SSR data.

Author Information The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to Y. L. (liying05@caas.cn) and S. H. (huangsanwen@caas.cn).

231 **Table 1. The allele dosage of microsatellite loci in asexual and sexual populations**

Lineage / Population	Area	MT ^a	mtDNA ^b	No. isolate	No. locus	3-allele loci ^c	2-allele loci ^c	1-allele loci ^c	3-allele ratio ^d
<i>Asexual lineages</i>									
CN Northern	Northern China	A1	IIa	67	10	106	213	346	0.33
CN 13_A2	Southwest China	A2	Ia	64	10	375	25	240	0.94
CN Fujian	Eastern China	A1	IIb	36	10	153	17	190	0.90
NL 13_A2	Netherlands	A2	Ia	73	12	591	5	280	0.99
NL 06_A1	Netherlands	A1	Ib	7	12	52	5	17	0.91
NL 08_A1	Netherlands	A1	Ia	16	12	91	15	47	0.86
EC-1	Ecuador	A1	IIa	73	12	422	124	330	0.77
TU-1	Tunisia	A1	Ia	57	12	312	114	251	0.73
US-1	US	A1	Ib	4	12	25	9	14	0.74
<i>Sexual population</i>									
NL pop2	Netherlands	A1, A2	Ia, IIa	37	10	45	83	241	0.35
Mex	Mexico	A1, A2	/	44	10	25	154	253	0.12
TU-2	Tunisia	A1, A2	Ia	42	10	56	105	238	0.35

232 ^aMating type

233 ^bMitochondria haplotype

234 ^c3-allele loci means triallelic loci, 2-allele loci means biallelic loci, and 1-allele loci means monoallelic loci

235 ^d=triallelic loci/(triallelic loci + biallelic loci)

236

237

Table 2. Ploidy analysis of isolates by genome re-sequencing, flow cytometry and microscope observation

Isolate	MT ^a	Lineage / Population	Seq. Depth	DNA Content (pg)	No. chromosome	Ploidy Mode	Reference
NL00150	A1	NL pop3	53	nd ^b	13.5±1.0	Diploid	This study
NL04092	A1	NL pop3	54	nd	nd	Diploid	This study
NL04106	A2	NL pop3	39	nd	nd	Diploid	This study
NL05159	A1	NL pop3	51	nd	14.2±0.8	Diploid	This study
NL05385	A1	NL pop3	55	nd	nd	Diploid	This study
NL05387	A1	NL pop3	51	nd	13.2±0.9	Diploid	This study
NL05890	A2	NL pop3	40	nd	12.8±0.8	Diploid	This study
NL07434	A2	NL pop2	54	0.56 ±0.02	14.2±0.7	Diploid	Ref. 5
NL08797	A2	NL pop2	23	0.56 ±0.01	13.8±0.4	Diploid	This study
P17777	A1	nd	48	nd	nd	Diploid	Ref. 5
06_3928A ^c	A2	NL 13_A2	50	nd	nd	Triploid ^c	Ref. 5
DDR7602	A1	US-1	13	nd	nd	Triploid	Ref. 5
LBUS5	A1	US-1	9	nd	nd	Triploid	Ref. 5
NL07041	A2	NL 13_A2	106	0.81±0.03	20.7±0.4	Triploid	This study
NL08080	A1	NL 08_A1	43	0.82±0.03	20.3±0.9	Triploid	This study
NL08452	A1	NL 06_A1	41	0.83±0.02	20.8±0.4	Triploid	This study
P13527	A1	EC-1	20	nd	nd	Triploid	Ref. 5
P13626	A1	EC-1	34	nd	nd	Triploid	Ref. 5

^aMating type

^bNot determined

^c06_3928A appeared to be tetraploid in previous research⁵. However, our analysis showed it is triploid.

Figure 1. Sexuality and ploidy. The phylogeny tree generated in previous study was used for sampling strategy. Green dots show isolates with A2 mating type, while A1 mating type was marked by orange cycles. Three representative isolates (NL07041, NL08080 and NL08452) were selected from three major clonal lineages in the Netherlands, and nine sexual isolates were from genetic neighbor clades. The ploidy mode was determined with the mean

251 frequency of read counts at heterozygous positions, a single and diploid mode at 0.5, while
252 two modes, 0.33 and 0.67 were for triploid genomes.

Figure 2. DNA content and chromosome number. Two isolates from asexual lineage (NL08452 and NL07041) and two isolates from sexual populations (NL07434 and NL08797) were investigated. DNA content was determined by flow cytometry. The chromosome number was counted by adjusting the focal levels on Laser Scanning Confocal Microscope (LSM780, Zeiss).