

Two Similarity Metrics for Medical Subject Headings (MeSH):

An Aid to Biomedical Text Mining and Author Name

Disambiguation

Neil R. Smalheiser* and Gary Bonifield

Department of Psychiatry, University of Illinois at Chicago, 1601 W. Taylor Street MC912,

Chicago, IL 60612 USA

Correspondence: neils@uic.edu

Abstract

In the present paper, we have created and characterized several similarity metrics for relating any two Medical Subject Headings (MeSH terms) to each other. The article-based metric measures the tendency of two MeSH terms to appear in the MEDLINE record of the same article. The author-based metric measures the tendency of two MeSH terms to appear in the body of articles written by the same individual (using the 2009 Author-ity author name disambiguation dataset as a gold standard). The two metrics are only modestly correlated with each other ($r = 0.50$), indicating that they capture different aspects of term usage. The article-based metric provides a measure of semantic relatedness, and MeSH term pairs that co-occur more often than expected by chance may reflect relations between the two terms. In contrast, the author metric is indicative of how individuals practice science, and may have value for author name disambiguation and studies of scientific discovery. We have calculated article metrics for all MeSH terms appearing in at least 25 articles in MEDLINE (as of 2014) and author metrics for MeSH terms published as of 2009. The dataset is freely available for download and can be queried at http://arrowsmith.psych.uic.edu/arrowsmith_uic/mesh_pair_metrics.html.

Keywords

Scientometrics, authorship, scientific publication, MEDLINE, interdisciplinarity, text mining, author name disambiguation, scientific journals, bibliometrics, discovery, novelty.

Background

Text mining analyses often involve estimating the similarity of two terms or concepts. In the biomedical domain, MEDLINE records include manual indexing by experts of topics discussed in each article, using a standardized hierarchical terminology of Medical Subject Headings (MeSH terms) that is employed to assist in retrieval of articles on a given topic. Various schemes have been proposed for relating different MeSH terms to each other in terms of their similarity. In general, these schemes can be classified as a) semantic, e.g., the path distance separating the two MeSH terms on the hierarchical tree; b) contextual, e.g., to what extent the two MeSH terms co-occur within the same articles; and c) lexical, e.g., the edit distance involved in transforming one term into another (Zhou et al, 2015). Co-occurring MeSH terms have been studied as an indicator of relations discussed in articles (e.g., Burgun and Bodenreider, 2001; Srinivasan and Hristovski, 2004; Kastrin et al, 2014) and MeSH-based similarity metrics have been employed in clustering of topically related articles (e.g., Lee et al, 2006; Zhu et al, 2009; Boyack et al, 2011). Several text mining models devoted to literature-based discovery have utilized similarity of two MeSH terms, or of two UMLS concepts, as features (e.g., Cohen et al, 2010; Theodosiou et al., 2011; Workman et al., 2013, 2015).

In the present work, we have computed and characterized two different MeSH term pair similarity metrics. The first involves calculating how often two different MeSH terms co-occur in the same articles, relative to the expected chance level (i.e., due to the frequencies of each MeSH term considered independently). We confirm that this metric captures topical similarity as judged by human raters, and point out some potential new uses for the metric in text mining. The second metric is novel: how often two different MeSH terms co-occur in the body of articles written by the same individual, relative to the expected chance level. As we will show, this

author-based metric has potential value for author name disambiguation modeling. Both person-centered and article-centered metrics are being released openly as comprehensive datasets and can be viewed via public web interfaces at http://arrowsmith.psych.uic.edu/mesh_pair_metrics.html.

Methods

Article-based metric. For each article included in the 2014 baseline version of MEDLINE, we extracted the Medical Subject Headings (MeSH) associated with the article, and calculated the number of times that each pair of MeSH terms co-occurred within the same article, as well as the total number of articles in which each MeSH term occurred. A stoplist of the 20 most frequent MeSH terms (D'Souza and Smalheiser, 2014) was employed to remove them from consideration, since highly frequent terms would appear to be similar to all other MeSH terms. Only those MeSH terms appearing in at least 25 articles were considered in calculating term similarity measures, since lower values would be highly subject to noise. The final number of included MeSH terms is 25,548.

Author-based metric. The 2009 Author-ity dataset (Torvik et al, 2005; Torvik and Smalheiser, 2009) is based on a snapshot of PubMed (which includes both MEDLINE and PubMed-not-MEDLINE records) taken in July 2009, including a total of 19,011,985 Article records, 61,658,514 author name instances and 20,074 unique journal names. Each instance of an author name is uniquely represented by the PMID and the position on the paper (e.g., 10786286_3 is the third author name on PMID 10786286). Thus, each predicted author-individual cluster is associated with a list of predicted PMIDs written by that individual. For each author-individual

cluster included in 2009 version of Author-ity, we extracted the MeSH terms found in each cluster (each term is counted once in each cluster, regardless of how many articles it appeared in within that cluster). We then calculated the number of times that each pair of MeSH terms co-occurred within the same author-individual cluster, across all clusters in the dataset. Only MeSH terms that were included for calculating article-based similarity (see above) were considered for calculating author-based similarity; a total of 25,007 MeSH terms were included in the author-based metric.

There are 37,385,852 pairs of MeSH terms included in the article similarity metric. 201,136,960 pairs of MeSH terms were included in the author similarity metric. The number of pairs calculated for author metric is greater than included in the article metric, since MeSH terms were counted as co-occurring if they were mentioned in ANY articles written by a given individual, even if they never co-occurred in the same article. Conversely, the article metric contains 729,894 pairs that are not included in the author-based metric (i.e., involving MeSH terms which were added to MEDLINE after 2009). Finally, 36,655,958 pairs of MeSH terms were included in both Author and Article similarity metrics, and could be directly compared to see how the two metrics capture different aspects of similarity.

Calculation of odds ratios. For any pair of MeSH terms, the number of co-occurrences needs to be normalized by the total number of occurrences of each MeSH term, in order to assess properly how meaningful it is to find two terms co-occurring (in the same article, or in the set of articles published by a given author). Two very common MeSH terms might be expected to co-occur often just by chance, whereas it will be highly significant if one observes any co-occurrence of

two very rarely occurring MeSH terms. We computed the co-occurrence score that would be expected simply by chance (for two MeSH terms of their size), separately for the article-based and author-based metrics. This was done by ranking all MeSH pairs by the geometric mean of their individual document occurrences, dividing into bins of 5,000 pairs (i.e., each having roughly the same size), and calculating the average co-occurrence score across all MeSH pairs in the same bin. Finally, we calculated the MeSH odds ratio for each pair of MeSH terms present in that bin, by taking the observed co-occurrence score divided by the average co-occurrence score for that bin. This is similar to the manner in which odds ratios were computed for journal similarity metrics in D'Souza and Smalheiser (2014).

Statistics

We employed correlation measures to characterize the relationship between two metrics, which allowed us to estimate the similarity of the metrics. In general, the nonparametric Spearman rho rank correlation coefficient is more appropriate for these comparisons, because the metrics are generally not linear. However, we also present the parametric Pearson r correlation coefficient as well, since there is some value in comparing the Pearson and Spearman values (e.g., if both are high, the relationships are likely to be linear, whereas if Pearson is very low and Spearman is very high, the relationships are likely to be nonlinear). Because each correlation was computed across millions of data points, statistical significance is generally extremely high and p-values are not displayed.

Results

As one might expect, the article-based and author-based MeSH odds ratios were significantly correlated, but perhaps surprisingly, the correlations were only about 0.5 (Pearson $r = 0.501$, Spearman $\rho = 0.558$). In other words, the two metrics do not simply measure the same thing. Rather, the tendency of two MeSH terms to co-occur in the same article reflects somewhat different aspects of similarity than the tendency of the same MeSH terms to co-occur within the body of work published by the same author.

The article-based metric, which counts co-occurrence of two MeSH terms in the same article, is subject to some limitations and constraints since a single article tends to have only 8-20 MeSH terms, and since MEDLINE indexers follow complex rules by which they decide to pick a given MeSH term (e.g., if more than one term is applicable but they lie vertically within the hierarchical tree, they are instructed to choose only the most specific term). Nevertheless, the article-based metric corresponded well to human ratings of semantic relatedness. Pedersen et al (2007) compiled a list of 29 UMLS concept (CUI) pairs annotated by physicians on a 1 to 4 scale of semantic similarity (Table 1). We mapped these to the corresponding MeSH term pairs as far as possible, and found that physician ratings correlated very well with the article-based metric ($r = 0.67$). A similar finding was observed with ratings by medical coders (Table 1).

In contrast, these ratings showed a much lower correlation with the author-based metric ($r = 0.38$). Note that one of the test pairs (Cholangiocarcinoma and Colonoscopy) co-occurred relatively infrequently within the same article (odds ratio = 0.44), but had a high author-based odds ratio (= 8.02), indicating that certain individuals, presumably GI specialists, tended to

publish on both topics. Seven of the 29 MeSH pairs had no co-occurrences at all within the same article (and hence have article-based similarity scores of 0), yet all of these had author-based co-occurrences such that the odds ratios were greater than zero. This may suggest that the author-based metric is more sensitive in detecting indirect similarities.

Another feature of the author-based metric is its “smoothing” effect relative to the article-based metric. If an author has published 7 articles, and each has 8 MeSH terms, potentially there is a pool of 56 MeSH terms to be considered pairwise, compared to only 8 MeSH terms for each article. This makes the author metric relatively robust and less influenced by fluctuations due to low sampling, particularly for MeSH terms that occur in relatively few articles. For any given MeSH term, its article-based odds ratio tended to achieve higher maximal values than did the author-based odds ratios (article-based maximal odds ratio = 73.525 mean \pm 52.16 SD vs. author-based maximal odds ratio = 49.213 mean \pm 36.11 SD, a difference that is highly significant ($p < 0.0001$, one-tailed unpaired t-test)).

One way to compare the article-based and author-based metrics is to examine the datasets as they can be queried on the Arrowsmith project MeSH Pair Metrics page http://arrowsmith.psych.uic.edu/arrowsmith_uic/mesh_pair_metrics.html. The user selects any MeSH term from a drop-down menu, and the site displays the top 20 most related MeSH terms ranked according to either the article-based or author-based metrics (Table 2). For each MeSH pair, the site also displays the number of articles in which each MeSH term occurs, the number of co-occurrences (in articles or author-individual clusters), the average number of co-occurrences expected for two MeSH terms by chance (based on their size), and the calculated

odds ratio. It is interesting to view how the article-based and author-based metrics sometimes emphasized different dimensions of similarity. For example, consider the top 20 terms related to the MeSH term “Tennis” (Table 3). The article-based metric lists 8 terms related to physical therapy and disorders that affect tennis players (vs. 4 terms listed under the author-based metric), whereas the author-based metric listed 10 other sports (vs. 5 sports listed under the article metric). Simply put, articles on tennis talked more about disorders afflicting tennis players, and did not generally include other sports in the same articles, whereas authors who wrote about tennis wrote more often about a variety of other sports. Another interesting example is “Abbreviations as Topic” (Table 4). The top 20 terms according to the article-based metric included 7 terms that were related to nursing and medications (vs. 2 listed under the author-based metric), whereas the author-based metric included 14 terms related to information science (vs. 8 under the article-based metric).

Discussion

The present paper describes and provides comprehensive article-based and author-based similarity metrics for pairs of Medical Subject Heading (MeSH) terms. We also present a web query interface that allows users to retrieve, for any specified MeSH term, the top 20 most related MeSH terms according to either the article-based or author-based metric.

As discussed in the introduction, article-based MeSH term pair similarity has been previously discussed by others and utilized in studies of information retrieval, document clustering, and literature-based discovery. Here, we have confirmed that the article-based metric does correspond to judgments of semantic relatedness made by human raters. We note that indexing an article with two MeSH terms suggests that the article may discuss a potential relation between the two topics – especially if the article-based odds ratio of that pair is greater than 1, i.e., if the two MeSH terms co-occur in the same articles more often than expected by chance. We deem a MeSH term pair as “important” if their article odds ratio >1 . This further suggests a new kind of similarity metric for relating different articles to each other. That is, for any pair of articles, one can score the number of “important” MeSH term pairs that they share, perhaps weighted so that MeSH term pairs with higher odds ratios count more. Counting MeSH term pairs will be more stringent and restrictive than counting individual shared MeSH terms.

The author-based MeSH term pair similarity measures the tendency of the same individual to discuss two different topics at some time during their career, i.e., in the body of articles that they have authored or co-authored. This says as much (or more) about the individual, and his or her range of interests, as it does any overt relation between the two topics. For example, the first author of this paper (NS) has written on a variety of subjects ranging from extracellular matrix

biochemistry to natural language processing to a biography of the early neuroscientist Walter Pitts. The two MeSH terms “Dystroglycans” and “History, 20th Century” are not obviously related to each other, and in fact, do not co-occur in any single article in PubMed. Yet one might hypothesize that an investigator who has worked on both topics might be psychologically or otherwise better poised to detect new knowledge that bridges these two fields, or that requires assembling different pieces of knowledge from each field, than someone who has only worked in one field. Although novel discoveries often involve combining topics together in new ways (e.g., Uzzi et al, 2013; Chen, 2014; Mishra and Torvik, 2014), one can hypothesize that MeSH term pairs which do not co-occur at all in the same articles, yet have high author-based odds ratios, may draw upon a pool of prepared minds (to quote Pasteur) and be particularly likely to be linked in new discoveries. Those which have very low author-based odds ratios might be less likely to be assembled into a new finding by an individual investigator. Multi-disciplinary teams may have a particular advantage for the investigation and discovery of findings that involve putting together topics that have very low author-based odds ratios. These hypotheses are testable, and may further our understanding of why and how particular novel combinations of topics lead to new discoveries.

Our original reason for studying author-based MeSH term similarity was to create an additional feature that can be used to disambiguate author names on PubMed articles comprehensively (Torvik et al., 2005; Torvik and Smalheiser, 2009). The most relevant measure of similarity for disambiguation is not topic centered, but rather author-centered. For example, two journals may cover the same topic (e.g., *Scandinavian Journal of Immunology* vs. *Iranian Journal of Immunology*), yet the same person may have very little likelihood of publishing articles in both

journals. Thus, the 2009 Author-ity disambiguation dataset was earlier mined to create a metric that comprehensively measures the tendency of individuals to publish articles in any two journals (D'Souza and Smalheiser, 2014). Here, the author metric measures the tendency of individuals to publish a body of articles that is indexed by any two different MeSH terms during their careers.

We are currently using the author-based metric in modeling author disambiguation, as follows:

Given two articles sharing the same author (lastname, firstinitial), we compute all MeSH terms for each article and consider the pairs that are formed across articles (i.e., one MeSH term in article 1 paired with another MeSH term in article 2). The three highest author-based odds ratios are averaged and used as the similarity score for the two articles. This feature is one of several new features that we are using to update and improve the performance of our author name disambiguation model.

Implementation

The datasets and readme.pdf are freely available for download from the Arrowsmith project website (http://arrowsmith.psych.uic.edu/arrowsmith_uic/mesh_pair_metrics.html) as well as from the UIC Institutional Repository, INDIGO, under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike CC BY-NC-SA license 4.0. The MeSH term pair data is contained in mesh_pair_metrics.txt (16 GB uncompressed, 5 GB compressed).

Acknowledgements

This research was supported by National Institutes of Health grants R01LM010817 and P01AG039347. We thank Vetle I. Torvik for advice regarding computing, and Arushi Chawla for assistance in data analysis.

Competing interests

The authors declare that no competing interests exist.

Authors' contributions

NRS designed the experiments. GB performed the data analyses and created the datasets. NRS wrote the paper.

References

Boyack KW, Klavans R, Börner K (2005) Mapping the backbone of science. *Scientometrics* 64(3): 351-374.

Boyack KW, Newman D, Duhon RJ, Klavans R, Patek M, Biberstine JR, Schijvenaars B, Skupin A, Ma N, Börner K. Clustering more than two million biomedical publications: comparing the accuracies of nine text-based similarity approaches. *PLoS One*. 2011 Mar 17;6(3):e18029. doi: 10.1371/journal.pone.0018029.

Burgun A, Bodenreider O. Methods for exploring the semantics of the relationships between co-occurring UMLS concepts. *Stud Health Technol Inform.*2001;84(Pt 1):171-5.

Cohen, T, Whitfield, GK, Schvaneveldt, RW, Mukund, K, Rindflesch, T. (2010). EpiphaNet: An Interactive Tool to Support Biomedical Discoveries. *Journal of Biomedical Discovery and Collaboration*, 5, 21–49.

Chen, C. *The Fitness of Information: Quantitative Assessments of Critical Evidence*. John Wiley & Sons, NY, 2014.

D'Souza JL, Smalheiser NR. Three journal similarity metrics and their application to biomedical journals. *PLoS One*. 2014 Dec 23;9(12):e115681. doi: 10.1371/journal.pone.0115681.

Kastin A, Rindflesch TC, Hristovski D. Large-scale structure of a network of co-occurring MeSH terms: statistical analysis of macroscopic properties. *PLoS ONE* 9 (7): e102188.

Lee M, Wang W, Yu H. Exploring supervised and unsupervised methods to detect topics in biomedical text. *BMC Bioinformatics*. 2006 Mar 16;7:140.

Leydesdorff L, Goldstone RL (2014) Interdisciplinarity at the journal and specialty level: The changing knowledge bases of the journal *Cognitive Science*. *J Assoc Inf Sci Technol* 65(1): 164-177.

Mishra S, Torvik VI. (2014) Measures Of Novelty And Growth For Bibliometrics. <https://www.ideals.illinois.edu/handle/2142/49962>

Pedersen T, Pakhomov SVS, Patwardhan S, Chute CG. Measures of semantic similarity and relatedness in the biomedical domain. *Journal of Biomedical Informatics*. 2007;40(3):288-299.

Srinivasan P, Hristovski D. Distilling conceptual connections from MeSH co-occurrences. *Stud Health Technol Inform*. 2004;107(Pt 2):808-12.

Theodosiou T, Vizirianakis IS, Angelis L, Tsiftaris A, Darzentas N. MeSHy: Mining unanticipated PubMed information using frequencies of occurrences and concurrences of MeSH terms. *J Biomed Inform*. 2011 Dec;44(6):919-26.

Torvik VI, Smalheiser NR (2009) Author name disambiguation in MEDLINE. *ACM Transactions on Knowledge Discovery from Data (TKDD)* 3(3): 11.

Torvik VI, Weeber M, Swanson DR, Smalheiser NR (2005) A probabilistic similarity metric for MEDLINE records: a model for author name disambiguation. *J Assoc Inf Sci Technol* 56(2): 140-158.

Workman TE, Fiszman M, Cairelli MJ, Nahl D, Rindfleisch TC. Spark, an application based on Serendipitous Knowledge Discovery. *J Biomed Inform*. 2015 Dec 28. pii: S1532-0464(15)00294-4. doi: 10.1016/j.jbi.2015.12.014.

Workman TE, Roseblat G, Fiszman M, Rindfleisch TC. A literature-based assessment of concept pairs as a measure of semantic relatedness. *AMIA Annu Symp Proc*. 2013 Nov 16;2013:1512-21.

Uzzi B, Mukherjee S, Stringer M, Jones B. Atypical combinations and scientific impact. *Science* 2013 Oct 25; 468-72.

Zhou J, Shui Y, Peng S, Li X, Mamitsuka H, Zhu S. MeSHSim: An R/Bioconductor package for measuring semantic similarity over MeSH headings and MEDLINE documents. *J Bioinform Comput Biol*. 2015 Dec;13(6):1542002. doi: 10.1142/S0219720015420020.

Table 1. Comparison of article-based and author-based MeSH term pair odds ratios against human raters' judgments of semantic relatedness.

<u>Physicians</u>	<u>Coders</u>	<u>CUI1</u>	<u>CUI2</u>	<u>mesh1</u>	<u>mesh2</u>	<u>article odds ratio</u>	<u>author odds ratio</u>
4	4	C0035078	C0035078	Renal Insufficiency	Renal Insufficiency		
3	3.3	C0156543	C0000786	no exact match	Abortion, Spontaneous		
3.3	3	C0018787	C0027061	Heart	Myocardium	27.5	4.2246
3	2.8	C0038454	C0021308	Stroke	Infarction	1.16	1.7916
3	2.2	C0011253	C0036341	Delusions	Schizophrenia	35	5.2131
2.7	2	C0175895	C0009814	Vascular Calcification	Constriction, Pathologic	2.84	0
2.7	1.8	C0027627	C0001418	Neoplasm Metastasis	Adenocarcinoma	11.6	4.5045
3	1.4	C0018802	C0034063	Heart Failure	Pulmonary Edema	12.6	3.8184
1.7	1.4	C0034069	C0242379	Pulmonary Fibrosis	Lung Neoplasms	8.86	3.327
2.3	1.3	C0011991	C0344375	Diarrhea	no exact match		
2.3	1.3	C0026269	C0004238	Mitral Valve Stenosis	Atrial Fibrillation	28.5	7.9298
2	1.3	C0006118	C0151699	Intracranial Hemorrhages	Hemorrhages	5.74	3.8165
1.7	1.2	C0003232	C0020517	Brain Neoplasms Anti-Bacterial Agents	Hypersensitivity Myocardial	0.68	0.9068
1.7	1.2	C0034065	C0027051	Pulmonary Embolism	Infarction	4.75	2.3836
2	1.1	C0007286	C0029408	Carpal Tunnel Syndrome	Osteoarthritis	4.65	4.2669
2	1.1	C0003873	C0409974	Arthritis, Rheumatoid	Lupus Erythematosus, Systemic	18.8	3.6328
2	1	C0702166	C0039142	Acne Vulgaris	Syringes	0	0.7958
2	1	C0011849	C0020538	Diabetes Mellitus	Hypertension	5.97	1.8314
1.7	1	C0010137	C0086511	Cortisone	Arthroplasty, Replacement, Knee	0.05	0.2433
1.3	1	C0206698	C0009378	Cholangiocarcinoma	Colonoscopy	0.44	8.0176
1.3	1	C0333997	C0007107	Giant Lymph Node Hyperplasia	Laryngeal Neoplasms	0	3.9563
1	1	C0003615	C0029456	Appendicitis	Osteoporosis	0	0.7581
1	1	C0011581	C0007642	Depressive Disorder	Cellulitis	0	0.2698
1	1	C0020473	C0027627	Neoplasm Hyperlipidemias	Metastasis	0.08	1.142
1	1	C0026769	C0033975	Multiple Sclerosis	Psychotic Disorders	0.73	1.0165
1	1	C0030920	C0027092	Peptic Ulcer	Myopia	0.09	0.163
1	1	C0034887	C0003483	Colonic Polyps	Aorta	0	1.2344
1	1	C0042345	C0224701	Varicose Veins	Medial Collateral	0	0.4605

1	1	C0043352	C0023891	Xerostomia	Ligament, Knee Liver Cirrhosis, Alcoholic	0	2.0231
---	---	----------	----------	------------	---	---	--------

Pedersen et al (2007) compiled a list of 29 UMLS concept (CUI) pairs annotated by physicians or medical coders, on a 1 to 4 scale of semantic similarity. We mapped these to the corresponding MeSH term pairs as far as possible and displayed their article-based and author-based odds ratios.

Table 2. The top 20 MeSH terms most similar to “Clergy” [MeSH], ranked by article-based odds ratio.

<i>Rank</i>	<i>MeSH Term</i>	<i>Article Count</i>	<i>Article Co-Occurrence</i>	<i>Article Odds Ratio</i>	<i>Author Co-Occurrence</i>	<i>Author Odds Ratio</i>
1	Catholicism	7383	403	73.9721	489	11.9222
2	Pastoral Care	3105	426	73.0453	541	28.6092
3	Chaplaincy Service, Hospital	924	180	68.9655	230	30.6748
4	Religion and Medicine	9795	216	43.1310	419	9.7469
5	Spirituality	4549	121	31.1054	188	9.2904
6	Religion and Psychology	5439	122	27.4899	288	10.0404
7	Christianity	6287	170	26.7632	353	10.4056
8	Religion	11340	156	25.9395	376	6.2134
9	Protestantism	693	50	20.6782	79	14.5488
10	Professional Role	7739	90	20.3712	150	4.2735
11	Theology	1141	49	19.0661	78	10.2821
12	Child Abuse, Sexual	8060	76	17.5115	112	3.0210
13	Judaism	2255	45	17.4014	133	9.7722
14	Counseling	26810	108	16.6821	308	3.6347
15	Value of Life	5338	56	15.6863	137	5.3399
16	Anecdotes as Topic	4470	46	14.4201	67	3.4561
17	Terminally Ill	5155	44	12.7536	136	5.3030
18	Attitude	37996	112	12.2351	397	2.7826

19	Euthanasia, Passive	5808	57	12.1899	142	4.7142
20	Ethics	9353	51	12.0796	161	3.9591

Shown are the top 20 MeSH terms that co-occurred with “Clergy” [MeSH], article count = 1641, ranked by article odds ratio. Author-based co-occurrences and odds ratios are also shown.

Table 3. Top 20 MeSH terms most related to “Tennis” [MeSH] by article and by author odds ratios.

<i>Rank</i>	<i>MeSH Term 2</i>	<i>Article Odds Ratio</i>	<i>Rank</i>	<i>MeSH Term 2</i>	<i>Author Odds Ratio</i>
1	Athletic Injuries	34.1242	1	Golf	29.4118
2	Sports	31.6081	2	Racquet Sports	26.8496
3	Tennis Elbow	20.4491	3	Baseball	25.2618
4	Biomechanical Phenomena	20.1603	4	Gymnastics	21.0877
5	Shoulder Joint	19.4338	5	Basketball	20.7207
6	Athletic Performance	18.4908	6	Tennis Elbow	20.6044
7	Motor Skills	15.8241	7	Sports Equipment	19.3218
8	Baseball	15.4834	8	Weight Lifting	17.0541
9	Competitive Behavior	14.7309	9	Track and Field	16.4690
10	Elbow Joint	14.3665	10	Hockey	16.4425
11	Golf	12.6683	11	Fractures, Stress	15.1724
12	Task Performance and Analysis	12.6183	12	Shoulder Impingement Syndrome	14.6710
13	Elbow	12.0805	13	Football	14.5765
14	Range of Motion, Articular	11.5265	14	Athletic Injuries	14.5104
15	Tendinopathy	11.2257	15	Ergometry	14.4312
16	Running	11.0825	16	Cumulative Trauma Disorders	14.1844
17	Cumulative Trauma Disorders	10.6751	17	Soccer	13.9590
18	Tendon Injuries	10.6642	18	Tendinopathy	13.7232
19	Soccer	10.4322	19	Acromion	12.7377
20	Physical Education and Training	9.8885	20	Jogging	12.6743

Columns at left show the top 20 MeSH terms that co-occurred with “Tennis” ranked by article odds ratio. At right, top 20 ranked by author odds ratio.

Table 4. Top 20 MeSH terms most related to “Abbreviations as Topic” [MeSH] by article and by author odds ratios.

Rank	MeSH Term 2	Article Odds Ratio	Rank	MeSH Term 2	Author Odds Ratio
1	Terminology as Topic	28.2360	1	Dictionaries as Topic	8.8602
2	Medication Errors	14.5379	2	MEDLINE	8.3068
3	Nursing Assessment	11.9482	3	Subject Headings	7.5937
4	Periodicals as Topic	10.9649	4	Weights and Measures	7.4758
5	Drug Prescriptions	10.2136	5	Medical Subject Headings	7.1207
6	Weights and Measures	9.1896	6	Natural Language Processing	6.8958
7	Writing	9.1089	7	Metric System	6.7002
8	Medical Records	7.7367	8	Abstracting and Indexing as Topic	6.6596
9	MEDLINE	7.4530	9	Unified Medical Language System	6.1576
10	Language	7.2093	10	Databases, Bibliographic	6.0713
11	Communication	6.9483	11	International System of Units	5.9347
12	Joint Commission on Accreditation of Healthcare Organizations	6.8768	12	Dictionaries, Medical	5.8901
13	Natural Language Processing	6.3139	13	Names	5.2069
14	Abstracting and Indexing as Topic	6.1107	14	Wit and Humor as Topic	4.5648
15	Nursing Records	5.9940	15	Vocabulary, Controlled	4.2828
16	Safety Management	5.8633	16	Hypermedia	4.2093
17	Publishing	5.6080	17	Reminder Systems	4.0059
18	Names	5.2402	18	Patient Identification Systems	3.8644
19	Information Storage and Retrieval	5.1120	19	Peer Review, Research	3.8040
20	Handwriting	5.0601	20	National Library of Medicine (U.S.)	3.6824

Columns at left show the top 20 MeSH terms that co-occurred with “Abbreviations as Topic” ranked by article odds ratio. At right, top 20 ranked by author odds ratio.