Genomics of primary chemoresistance and remission induction failure in pediatric and adult acute myeloid leukemia

Fiona C. Brown¹, Paolo Cifani¹, Esther Drill², Jie He³, Eric Still¹, Shan Zhong³, Sohail Balasubramanian³, Dean Pavlick³, Bahar Yilmazel³, Kristina M. Knapp⁴, Todd A. Alonzo^{5,6}, Soheil Meshinchi^{5,7}, Richard M. Stone⁸, Steven M. Kornblau⁹, Guido Marcucci¹⁰, Alan S. Gamis^{5,11}, John C. Byrd¹², Mithat Gonen², Ross L. Levine¹³ and Alex Kentsis^{1,14}

¹ Molecular Pharmacology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, United States

² Department of Biostatistics and Epidemiology, Memorial Sloan Kettering Cancer Center, New York, NY, United States

³ Foundation Medicine, Cambridge, MA, United States

⁴ Leukemia Center, Memorial Sloan Kettering Cancer Center, New York, NY, United States

⁵ Children's Oncology Group, Monrovia, CA, United States

⁶ Department of Biostatistics, University of Southern California, Los Angeles, CA, United States

⁷ Fred Hutchinson Cancer Research Center and the University of Washington School of Medicine, Seattle, WA, United States

⁸ Division of Hematologic Malignancies, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA, USA

⁹ Department of Leukemia, The University of Texas and MD Anderson Cancer Center, Houston, TX, United States

¹⁰ Gehr Family Leukemia Center, City of Hope Comprehensive Cancer Center, Duarte, CA, United States

¹¹ Division of Hematology-Oncology/, Children's Mercy Hospitals and Clinics, Kansas City, MO, United States

¹² Division of Hematology, Department of Internal Medicine, The Ohio State University, Columbus, OH, United States

¹³ Human Oncology and Pathogenesis Program, Memorial Sloan Kettering Cancer Center and Weill Medical College of Cornell University, New York, NY, United States

¹⁴ Department of Pediatrics, Memorial Sloan Kettering Cancer Center and Weill Medical College of Cornell University, New York, NY, United States

Correspondence: Alex Kentsis, MD, PhD, Memorial Sloan Kettering Cancer Center, 1275 York Ave, New York, NY, USA 10065, Phone number: 1-646-888-2593, Email: kentsisresearchgroup@gmail.com

Key Points

- Targeted gene sequencing of 670 genes in adult and pediatric AML
- Profiling of 107 primary AML samples identifies new genomic alterations for primary chemoresistance

Abstract

Despite intense efforts, the cure rates of children and adults with AML remain unsatisfactory in large part due to resistance to chemotherapy. Whilst cytogenetic risk stratification proved valuable in identifying causes of therapy failure and disease relapse, cytogenetically normal AML remains the most prevalent disease type, with significant heterogeneity of clinical outcomes, including primary chemoresistance. Using targeted sequencing of 670 genes recurrently mutated in hematologic malignancies, we investigated the genetic basis of primary chemotherapy resistance and remission induction failure of 107 primary cases obtained at diagnosis from children and adults with cytogenetically normal AML. Comparative analysis revealed mutations of *SETBP1*, *ASXL1* and *RELN* to be significantly enriched at diagnosis in primary induction failure as compared to remission cases. In addition, this analysis revealed novel genomic alterations not previously described in AML, as well as distinct genes that are significantly overexpressed in therapy resistant AML. However, identified gene mutations were sufficient to explain only a minority of cases of primary induction failure. Thus, additional genetic or molecular mechanisms must cause primary chemoresistance in pediatric and adult acute myeloid leukemias.

Introduction

For patients with acute myeloid leukemia (AML), failure to achieve complete remission after induction therapy or relapse after complete remission represents the major barrier to cure for both children and adults. Despite recent efforts to improve risk stratification and to incorporate targeted therapies into AML care, relapse rates remain approximately 30% and 50% for children and adults with AML, respectively¹⁻³. Whilst leukemia cytogenetics at the time of diagnosis has proven to be the most useful prognostic biomarker of clinical outcomes and therapy stratification, patients with cytogenetically normal (CN) AML encompass the largest group for whom prognostication and therapy selection are hindered by the lack of prognostic markers and effective therapeutic targets.

Genomic profiling has been used to stratify patients with CN AML, identifying recurrent somatic gene mutations, such as those in *NPM1*, *TET2*, *ASXL1*, and *DNMT3A*, that are significantly associated with clinical outcomes^{4, 5}. Recent studies using whole-genome sequencing have assessed the mutational landscape of relapsed AML in adults identifying two major patterns of clonal evolution in response to chemotherapy: 1) persistence of the dominant pre-leukemic clone with mutations acquired at relapse, and 2) expansion of a diagnostic sub-clone, emphasizing the importance of therapeutically targeting specific molecular mechanisms that contribute to chemotherapy resistance⁶⁻⁸. Similar observations were recently made using genomic studies of pediatric AML, with specific variant allele frequencies representing potential prognostic indicators⁹. Clonal persistence after induction chemotherapy appears to be associated with the increased risk of relapse¹⁰, but whether these mutant clones directly cause chemoresistance or reflect age-related clonal hematopoiesis remains to be determined¹¹. Importantly, the molecular basis of primary chemotherapy resistance, and thus fundamental causes of chemoresistance in general, remain poorly understood. Here, we used targeted gene sequencing of the majority of genes known to be recurrently mutated in hematologic

malignancies to identify diagnostic gene mutations that are associated with primary chemoresistance in adult and pediatric CN AML.

Study design

Patient Selection

Specimens were collected from patients treated at the Memorial Sloan Kettering Cancer Center, MD Anderson Cancer Center, Dana-Farber Cancer Institute, Ohio State University Comprehensive Cancer Center, and Children's Oncology Group (COG) participating institutions. All patients provided informed consent, were enrolled on respective institutional research protocols and treated using current institutional remission induction regimens (Table 1). Primary chemoresistance was defined based on the presence of at least 5% AML blasts by morphologic assessment of bone marrow aspirates obtained after two cycles of induction chemotherapy, as assessed by the respective institutional or central pathologic review.

Targeted Gene Sequencing

Mononuclear cells isolated from bone marrow aspirates obtained at diagnosis were purified by Ficoll gradient centrifugation, and enriched for leukemia cells by negative immunomagnetic selection against CD3, CD14, CD19 and CD235a given the lack of expression of these markers by the majority of AML cells¹². Targeted DNA and RNA capture and sequencing of 670 genes (405 genes in DNA and 265 genes in RNA) known to be recurrently mutated in hematologic malignancies were performed using the Foundation One Heme sequencing platform. Extraction of DNA and RNA, library preparation and Illumina sequencing were carried out as previously described¹³. Sequencing data were analyzed to identify genomic alterations including base substitutions, insertions, deletions, and gene fusions¹³. Differential gene expression analysis was performed based on per million processing regions (PPM) gene expression measurements, as median normalized to all other AML specimens analyzed to

date¹³. All statistical comparisons were corrected for multiple hypothesis testing using the Bonferroni correction, and statistical significance was calculated as indicated.

Results and Discussion

We were able to obtain genomic profiles of 107 CN AML specimens in total, collected from 55 adults and 52 children, balanced for outcome groups, gender and age (Table 1). No significant differences were observed in the total number of coding mutations between the primary failure of induction chemotherapy and complete remission outcome groups for both pediatric and adult cohorts (Supplemental Table 1). Consistent with previous studies, the most frequently mutated genes among adults were *DNMT3A* (49%), *FLT3* (42%) and *NPM1* (42%), and for pediatric patients *FLT3* (54%), *NRAS* (27%), and *WT1* (25%), which were not significantly associated with either outcome group (Figure 1). Mutations of *DNMT3A* and *NPM1* were relatively rare in pediatric (2%) as compared to adult cases ($p < 0.001$, permutation test). In contrast, *CEBPA* mutations were more frequently observed in pediatric as compared to adult CN AML (21% versus 2%, respectively, $p < 0.001$, permutation test).

Genomic profiling also identified 15 gene rearrangements among the 21 specimens for which diagnostic material of adequate quality was available for RNA capture (Supplemental Table 1). Cryptic rearrangements of known genes in AML were detected, including *KMT2A* (*MLL*) and *NUP98-NSD1*. The latter was observed in only pediatric patients from this study, consistent with previous reports^{14, 15}. We identified several novel cryptic gene rearrangements including intergenic deletions involving *NOTCH1/EDF1* and *GATA2/DNAJB8*, as well as translocations leading to the rearrangements of *SQSTM1-NUP214* and *IGH-BCL2*. Mutations of these genes have been reported in hematologic malignancies, for example the *SQSTM1-NUP214* gene fusion in cases of T-cell acute lymphoblastic leukemias (ALL), but not in AML¹⁶. In addition, our study identified potential novel coding mutations in *MAP3K1* (*MEKK*), *MAP3K6*

(*MEKK6*), *EP300* (p300), and *WDR90* (Supplemental Table 1). These findings emphasize the value of genomic profiling to identify potential pathogenic lesions that may elude conventional cytogenetic analyses and diagnosis based on morphologic or immunophenotypic studies.

We assessed whether any gene mutations were significantly associated with primary chemoresistance as compared to complete remission among the adult and pediatric patients in the studied cohorts. This identified mutations of *SETBP1* (8 out of 50 versus 1 out of 57, $p = 0.01$) and *ASXL1* (7 out of 50 versus 1 out of 57, $p = 0.02$), as well as *RELN* (8 out of 50 versus 2 out of 57, $p = 0.04$, permutation test). Mutations of *SETBP1* and *ASXL1* have been reported to be associated with inferior outcomes and increased risk of relapse in patients with AML^{17, 18}. Relapse-associated *SETBP1* mutations in AML commonly cause missense mutations in the SKI domain (p.858-871). Our analysis identified the missense p.D868N mutation in one CN AML specimen, as well as additional missense mutations (p.E183K, p.A374T, p.A408T, p.M502I, p.D1161H, p.T1547N and p.S1590C) of *SETBP1*, suggesting that alternative mechanisms of *SETBP1* dysregulation may exist in AML. Mutations of *RELN* have been reported in early T-cell precursor (ETP) ALL, a stem cell-like leukemia with high rates of intrinsic chemotherapy resistance and genetic features that are shared with subtypes of AML¹⁹. Our findings of *RELN* mutations associated with primary chemoresistance in CN AML raise the possibility that these leukemias may comprise a distinct biologic entity.

In addition, using available RNA sequencing and gene expression data, we assessed whether genes included in our analysis were significantly altered in expression in the induction failure as compared to complete remission cohorts (Supplemental Table 2). This analysis revealed significant overexpression of genes encoding the DNA damage repair and response regulators *ATM*, *ATR*, *PSIP1* (*LEDGF*), and *EMSY* (*C11ORF30*), endoplasmic reticulum stress response regulator *HERPUD1*, transcription factors *CBFB* and *MEF2C*, kinases *AKT3* and *MAP3K7* (*TAK1*), metabolic enzymes *LDHA* and *ACSL6*, clathrin *CLTC*, and immune regulator

CTLA4 ($p < 0.05$, t-test). Additional profiling and functional studies will be necessary to establish their prevalence and pathogenicity in AML. Indeed, recent studies have implicated overexpression of *MEF2C* with inferior outcomes in pediatric AML²⁰, and overexpression of *MAP3K7* has been observed to enhance AML cell survival²¹. These results suggest that other mechanisms such as those involving epigenetic or functional molecular alterations of cell signaling may also mediate chemotherapy response and failure.

In summary, we have assessed the mutational landscape of cytogenetically normal AML at diagnosis in adults and children to determine the genetic basis of induction failure and primary chemoresistance. In spite of careful selection of cytogenetically normal specimens, our study identified numerous cryptic gene rearrangements, including those that have potential therapeutic implications. Likewise, we found novel gene alterations including lesions that have been observed in non-myeloid hematologic malignancies, some of which may be pathogenic. In addition, our analysis revealed significant overexpression of genes that are associated with primary chemoresistance. Though mutations of *SETBP1*, *ASXL1* and *RELN* appear to be associated with primary chemoresistance and induction failure in our study, their prevalence was relatively low. We did not identify single gene mutations that are significantly associated with primary chemoresistance and induction failure in the majority of the patients studied. It is possible that additional genetic lesions such as those not included in the current target gene panel¹³, or alternatively, combinations of genetic alterations that are below the statistical power of our study may cause primary chemoresistance and failure in AML, which may be elucidated by larger and more comprehensive genomic profiling studies. Finally, it remains to be determined whether functional molecular profiling such as that achieved using gene expression, epigenetic and proteomic analyses may reveal more universal biomarkers and therapeutic targets to prospectively identify and therapeutically block chemotherapy resistance in AML²²⁻²⁵.

Acknowledgements

We thank Alejandro Gutierrez for critical discussions. This work was supported by the NIH R21 CA188881, K08 CA160660, P30 CA008748, U10 CA180899, Burroughs Wellcome Fund, Alex's Lemonade Stand Foundation, Gabrielle's Angel Foundation, and the Josie Robertson Investigator Program.

Authorship

FB and AK wrote the manuscript and performed analysis. FB, PC, ES performed laboratory experiments. JH, SZ, SB, DP and BY performed analysis. KK, TA, SM, RS, SK, GM, JB compiled patient samples. ED, MG and JH performed statistical analysis. AK and RL designed the study. RL contributed clinical and genomic expertise.

Conflict-of-interest disclosure: JH, SZ, SB, DP and BY are employees and equity holders of Foundation Medicine Inc. RL is a consultant for Foundation Medicine Inc.

Correspondence: Alex Kentsis, MD, PhD, Memorial Sloan Kettering Cancer Center, 1275 York Ave, New York, NY, USA 10065, Phone number: 1-646-888-2593, Email: kentsisresearchgroup@gmail.com

References

1. de Rooij, J.D., C.M. Zwaan, and M. van den Heuvel-Eibrink, Pediatric AML: From Biology to Clinical Management. *J Clin Med*, 2015. **4**(1): p. 127-49.
2. Breems, D.A., et al., Prognostic index for adult patients with acute myeloid leukemia in first relapse. *J Clin Oncol*, 2005. **23**(9): p. 1969-78.
3. Mangan, J.K. and S.M. Luger, Salvage therapy for relapsed or refractory acute myeloid leukemia. *Ther Adv Hematol*, 2011. **2**(2): p. 73-82.
4. Kihara, R., et al., Comprehensive analysis of genetic alterations and their prognostic impacts in adult acute myeloid leukemia patients. *Leukemia*, 2014. **28**(8): p. 1586-95.
5. Patel, J.P., et al., Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. *N Engl J Med*, 2012. **366**(12): p. 1079-89.
6. Ding, L., et al., Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. *Nature*, 2012. **481**(7382): p. 506-10.
7. Sood, R., et al., Somatic mutational landscape of AML with inv(16) or t(8;21) identifies patterns of clonal evolution in relapse leukemia. *Leukemia*, 2016. **30**(2): p. 501-4.
8. Corces-Zimmerman, M.R., et al., Preleukemic mutations in human acute myeloid leukemia affect epigenetic regulators and persist in remission. *Proc Natl Acad Sci U S A*, 2014. **111**(7): p. 2548-53.
9. Farrar, J.E., et al., Genomic profiling of pediatric acute myeloid leukemia reveals a changing mutational landscape from disease diagnosis to relapse. *Cancer Res*, 2016.
10. Klcio, J.M., et al., Association Between Mutation Clearance After Induction Therapy and Outcomes in Acute Myeloid Leukemia. *JAMA*, 2015. **314**(8): p. 811-22.
11. Wong, T.N., et al., Rapid expansion of preexisting nonleukemic hematopoietic clones frequently follows induction therapy for de novo AML. *Blood*, 2016. **127**(7): p. 893-7.

12. Creutzig, U., et al., Clinical significance of surface antigen expression in children with acute myeloid leukemia: results of study AML-BFM-87. *Blood*, 1995. **86**(8): p. 3097-108.
13. He, J., et al., Integrated genomic DNA/RNA profiling of hematologic malignancies in the clinical setting. *Blood*, 2016.
14. Shiba, N., et al., NUP98-NSD1 gene fusion and its related gene expression signature are strongly associated with a poor prognosis in pediatric acute myeloid leukemia. *Genes Chromosomes Cancer*, 2013. **52**(7): p. 683-93.
15. Hollink, I.H., et al., NUP98/NSD1 characterizes a novel poor prognostic group in acute myeloid leukemia with a distinct HOX gene expression pattern. *Blood*, 2011. **118**(13): p. 3645-56.
16. Gorello, P., et al., SQSTM1-NUP214: a new gene fusion in adult T-cell acute lymphoblastic leukemia. *Haematologica*, 2010. **95**(12): p. 2161-3.
17. Thol, F., et al., SETBP1 mutation analysis in 944 patients with MDS and AML. *Leukemia*, 2013. **27**(10): p. 2072-5.
18. Meggendorfer, M., et al., SETBP1 mutations occur in 9% of MDS/MPN and in 4% of MPN cases and are strongly associated with atypical CML, monosomy 7, isochromosome i(17)(q10), ASXL1 and CBL mutations. *Leukemia*, 2013. **27**(9): p. 1852-60.
19. Zhang, J., et al., The genetic basis of early T-cell precursor acute lymphoblastic leukaemia. *Nature*, 2012. **481**(7380): p. 157-63.
20. Laszlo, G.S., et al., High expression of myocyte enhancer factor 2C (MEF2C) is associated with adverse-risk features and poor outcome in pediatric acute myeloid leukemia: a report from the Children's Oncology Group. *J Hematol Oncol*, 2015. **8**(1): p. 115.
21. Bosman, M.C., et al., The TAK1-NF-kappaB axis as therapeutic target for AML. *Blood*, 2014. **124**(20): p. 3130-40.

22. Rijal, S., et al., Inositol polyphosphate 4-phosphatase II (INPP4B) is associated with chemoresistance and poor outcome in AML. *Blood*, 2015. **125**(18): p. 2815-24.
23. Alizadeh, A.A., et al., Prediction of survival in diffuse large B-cell lymphoma based on the expression of 2 genes reflecting tumor and microenvironment. *Blood*, 2011. **118**(5): p. 1350-8.
24. Chan, S.M., et al., Isocitrate dehydrogenase 1 and 2 mutations induce BCL-2 dependence in acute myeloid leukemia. *Nat Med*, 2015. **21**(2): p. 178-84.
25. Chen, Y., et al., CXCR4 downregulation of let-7a drives chemoresistance in acute myeloid leukemia. *J Clin Invest*, 2013. **123**(6): p. 2395-407.

Table 1: Demographic features of induction failure and induction remission cohorts

Characteristics	Pediatric		Adult	
	Remission	Failure	Remission	Failure
Number, N (%)	25 (48)	27 (52)	32 (58)	23 (42)
Gender, % female	52	44	47	39
Age (years), median (range)	11.5 (2.8-16.8)	10.4 (0.6-16.8)	48.4 (19-84)	54.7 (16-73)
Therapy, N (%)				
ADE	25 (48)	27 (52)	0	0
IDA + HDAC	0	0	10 (31.2)	4 (17.4)
7+3 DNR and AraC	0	0	11 (34.4)	14 (60.9)
FAI	0	0	4 (12.5)	1 (4.3)
BID - FA	0	0	2 (6.25)	2 (8.7)
FLAG - IDA	0	0	4 (12.5)	0 (0)
IDA + AraC	0	0	1 (3.1)	1 (4.3)
Clofa + IDA + AraC	0	0	0	1 (4.3)

Abbreviations: ADE, cytarabine, daunomycin and etoposide-based regimen; IDA, idarubicin; HDAC, high-dose cytarabine; DNR, daunorubicin; BID-FA, twice daily fludarabine + cytarabine; FAI, fludarabine, cytarabine, idarubicin; FLAG-IDA, fludarabine, cytarabine, idarubicin and granulocyte colony stimulating factor; Clofa, clofarabine.

Figure 1: Mutational spectrum of pediatric and adult AML with primary chemotherapy resistance and induction failure. Tile plot of the genomic alterations identified in DNA/RNA targeted sequencing. Data are shown for 107 CN AML specimens, collected from 52 and 57 adult and pediatric specimens respectively, grouped by clinical outcome.

Supplemental Table 1: Genomic alterations.

Supplemental Table 2: Gene expression analysis.

Pediatric

Adult

Figure 1. Mutational spectrum of pediatric and adult AML with primary chemotherapy resistance and induction failure