

1 P2Y2 purinergic receptor is induced following human cytomegalovirus infection
2 and its activity is required for efficient viral replication

3
4

5

6 Saisai Chen, Thomas Shenk, Maciej T. Nogalski[#]

7

8

9

10 Department of Molecular Biology

11 Princeton University

12 Princeton, NJ 08544-1014 USA

13

14

15

16

17 **Running Title:** Role of purinergic receptors in HCMV infection

18

19 [#]Corresponding author: nogalski@princeton.edu

20

21

22

23

24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Abstract

Human cytomegalovirus (HCMV) manipulates many aspects of host cell biology to create an intracellular milieu optimally supportive of its replication and spread. The current study reveals a role for purinergic signaling in HCMV infection. The levels of several components of the purinergic signaling system, including the P2Y2 receptor, were altered in HCMV-infected fibroblasts. P2Y2 receptor RNA and protein are strongly induced following infection. Pharmacological inhibition of receptor activity or knockdown of receptor expression markedly reduced the production of infectious HCMV progeny. When P2Y2 activity was inhibited, the accumulation of most viral RNAs tested and viral DNA was reduced. In addition, the level of cytosolic calcium within infected cells was reduced when P2Y2 signaling was blocked. The HCMV-coded UL37x1 protein was previously shown to induce calcium flux from the smooth endoplasmic reticulum to the cytosol, and the present study demonstrates that P2Y2 function is required for this mobilization. We conclude that P2Y2 supports the production of HCMV progeny, possibly at multiple points within the viral replication cycle that interface with signaling pathways induced by the purinergic receptor.

47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69

Importance

HCMV infection is ubiquitous and can cause life-threatening disease in immunocompromised patients, debilitating birth defects in newborns, and has been increasingly associated with a wide range of chronic conditions. Such broad clinical implications result from the modulation of multiple host cell processes. This study documents that cellular purinergic signaling is usurped in HCMV-infected cells and that the function of this signaling axis is critical for efficient HCMV infection. Therefore, we speculate that blocking P2Y2 receptor activity has the potential to become an attractive novel treatment option for HCMV infection.

70

Introduction

71 Human cytomegalovirus (HCMV) is a beta herpesvirus that infects a large percentage of
72 the adult population worldwide. Infection in immunocompetent people is typically
73 asymptomatic. In contrast, HCMV is a leading opportunistic pathogen in immunosuppressed
74 individuals [1-4] and is a major infectious cause of birth defects [5, 6].

75 HCMV causes broad cellular effects that likely contribute to the diverse pathologies
76 associated with infection. One mechanism utilized by the virus to change the biology of infected
77 cells is via the regulation of expression levels and activities of cell surface proteins [7, 8]. Here,
78 we describe the role of purinergic receptors during HCMV infection. Purinergic receptors are
79 ubiquitous cell surface receptors that are activated by extracellular adenosine di- (ADP) and tri-
80 (ATP) phosphates (P2 receptors) or adenosine (P1 receptors). P2 purinergic receptors are further
81 divided into ionotropic P2X and metabotropic P2Y families. P2X receptors are ATP-gated ion
82 channels and P2Y receptors are G protein-coupled receptors that are activated by adenine and
83 uridine nucleotides or nucleotide sugars. Seven subtypes of P2X receptors (P2X1-7) and eight of
84 P2Y receptors (P2Y1, P2Y2, P2Y4, P2Y6, P2Y11, P2Y12, P2Y13, and P2Y14) have been
85 identified [9, 10].

86 Signaling via the P2Y receptors triggers the activation of a heterotrimeric G protein,
87 which leads to the activation of effector protein phospholipase C (PLC), and generation of
88 diacylglycerol (DAG) and inositol 1,4,5-triphosphate (IP₃)[11]. DAG stimulates protein
89 kinase C (PKC)[12]. The activation of PKC has been shown to increase the expression of early
90 growth response protein 1 (Egr-1) [13], a transcription factor critical for DNA synthesis,
91 proliferation, and migration of fibroblasts and other cells [14]. IP₃ mobilizes cytosolic Ca²⁺ from
92 the smooth endoplasmic reticulum (SER). It has been observed that in lung fibroblasts, P2Y2 is

93 the only purinergic receptor subtype that, when activated, causes the mobilization of intracellular
94 Ca^{2+} [15]. P2Y2-mediated intracellular Ca^{2+} increases have been implicated in promoting the
95 proliferation and migration of hepatocellular carcinoma cells in mice [16]. Moreover, P2Y2
96 signaling was found to stimulate HIV-1 viral fusion through the activation of proline-rich
97 tyrosine kinase 2 (Pyk2) [17].

98 P2X receptors differ notably from their P2Y counterparts in ligand selectivity. While
99 P2Y receptors recognize a wide range of agonists, P2X receptors are activated only by ATP.
100 Moreover, P2X receptors are assembled as trimeric proteins. Specifically, P2X5 mainly
101 functions as heterotrimers involving P2X1, P2X2, or P2X4 [18]. The expression of P2X5
102 receptors is normally restricted to the trigeminal mesencephalic nucleus of the brainstem,
103 sensory neurons, cervical spinal cord, and some blood vessels [19]. P2X5 receptor expression in
104 atypical locations has been linked to cancer [20-22]. Activation of the P2X5 receptor causes an
105 influx of cations (Na^+ , K^+ , and Ca^{2+}) across the plasma membrane [23].

106 Knowing that purinergic receptors regulate cellular calcium levels and that an increase in
107 calcium levels has been observed following HCMV infection [24], it is intriguing to speculate
108 that members of the family of purinergic receptors could be important factors during HCMV
109 infection. It has been reported previously that a few members of the family of purinergic
110 receptors exhibited higher expression in HCMV-infected cells, however their molecular role
111 during infection has not been found [25, 26]. The regulation of Ca^{2+} release is critical for viral
112 DNA synthesis and the production of infectious progeny [27]. It has been found that the HCMV
113 immediate early protein pUL37x1 induces the mobilization of Ca^{2+} from SER to the cytosol [28].
114 Altering the release of Ca^{2+} from SER affects the activity of Ca^{2+} -dependent ER chaperones,
115 resulting in the accumulation of unfolded proteins and contributing to the unfolded protein

116 response [29]. In addition, the Ca^{2+} -dependent protein kinase, $\text{PKC}\alpha$, is activated following
117 infection, leading to the production of large (1-5 μm diameter) cytoplasmic vesicles at late times
118 during infection [30]. The presence of these vesicles correlates with the efficient accumulation of
119 enveloped virions. The release of Ca^{2+} from SER during infection is also expected to influence
120 Ca^{2+} -dependent processes that occur in the mitochondria. For instance, Ca^{2+} uptake can stimulate
121 aerobic metabolism and enhance ATP production [31, 32]. Ca^{2+} can also induce the activity of
122 Ca^{2+} /calmodulin-dependent protein kinase kinase (CaMKK), which activates 5' AMP-activated
123 protein kinase (AMPK). AMPK activity has been shown to support HCMV-induced changes to
124 the infected cell metabolome, as well as to be necessary for HCMV DNA synthesis and the
125 expression of viral late genes [33, 34]. Interestingly, blocking viral DNA synthesis by inhibiting
126 viral DNA polymerase, pUL54, can serve as an effective therapeutic measure for HCMV
127 infection [35]. Additionally, cellular phosphoinositide 3-kinase (PI3-K) and p38 kinase activities
128 were found to be required for viral DNA replication and the production of infectious progeny
129 [36, 37]. As both PI3-K and p38 are downstream factors of the purinergic receptor signaling axis
130 [38, 39], it is conceivable that purinergic receptors can have an effect on viral DNA synthesis.
131 Finally, since Ca^{2+} plays a central role in mediating apoptosis, regulating the concentration of
132 intracellular Ca^{2+} may affect the infected cell's sensitivity to apoptotic stimuli [40, 41].

133 Although the viral pUL37x1 protein has already been implicated in mediating Ca^{2+}
134 release during HCMV infection, there may also be other pathways that affect intracellular Ca^{2+}
135 levels. Therefore, we speculated that the purinergic receptor-mediated signaling may work
136 alongside pUL37x1 in regulating Ca^{2+} release into the cytosol, influencing the efficiency of
137 HCMV infection. In fact, one study using mouse fibroblasts showed that extracellular ATP and
138 UTP could stimulate the mobilization of Ca^{2+} from intracellular stores by an IP_3 -mediated

139 pathway downstream of P2Y2 signaling that is independent of PKC activation [42]. To test this
140 notion, we used pharmacological agents and siRNA technology to study the consequences of
141 inhibiting purinergic receptors during HCMV infection. We provide evidence that P2Y2
142 inhibition interferes with the expression of viral genes and the release of infectious progeny,
143 whereas P2X5 inhibition enhances HCMV yield. P2Y2-mediated signaling affects HCMV
144 infection at the stage of viral DNA synthesis and significantly contributes to the regulation of
145 intracellular Ca²⁺ homeostasis.

146

147 **Methods and Materials**

148 **Cells, viruses and drugs.** Human foreskin fibroblasts (HFF) and human lung fibroblasts
149 (MRC5) were cultured in Dulbecco's Modified Eagle Medium (DMEM; Sigma-Aldrich)
150 supplemented with 10% fetal bovine serum (FBS). Penicillin and streptomycin were added to the
151 media for all experiments except those involving siRNA transfection.

152 HFF cells stably expressing the viral protein, pUL37x1, were made using the pLVX-
153 EF1 α promotor lentivirus packaging plasmid (Clontech). To insert AD169 UL37x1 between
154 EcoRI-BamHI sites of the pLVX-EF1 plasmid, the sequence was amplified by PCR from
155 BADwt, a BAC containing the AD169 strain [43]. Lentivirus particles were generated using
156 sequence-confirmed plasmids containing UL37x1 or GFP as a control. Following lentiviral
157 treatment, cells expressing pUL37x1 or GFP were selected using 2 μ g/ml puromycin, and protein
158 expression was evaluated by Western blot using mouse monoclonal antibody 4B6-B [28].

159 Previously described HCMV strains TB40/E, TB40-GFP, ADwt and ADsubUL37x1
160 HCMV strains were used in these studies [43, 44]. For some experiments, TB40/E virus was
161 inactivated by irradiation (50 J/m²) in a Stratalinker (Stratagene) [45]. A pUL37x1-deficient

162 derivative of AD_{wt}, AD_{subUL37x1} [43], lacks the AD169 genomic sequence from 169,144 to
163 169,631 and includes kanamycin resistance and *LacZ* markers [46]. All viral stocks were purified
164 by centrifugation through a 20% D-sorbitol cushion containing 50mM Tris·HCl, 1mM MgCl₂,
165 pH 7.2, and resuspended in DMEM. Infections were performed by treating cells with viral
166 inoculum for 2 h, followed by removal of inoculum and washing with phosphate-buffered saline
167 (PBS) before applying with fresh media.

168 Virus titers were determined based on viral IE1 expression on a reporter plate containing
169 fibroblasts [34, 47]. At 24 hpi, cells were assayed by indirect immunofluorescence using a
170 primary antibody against the viral IE1 protein [48] and secondary goat anti-mouse antibody
171 conjugated to Alexa Fluor 488 (Molecular Probes). Nuclei were stained using Hoescht dye
172 (Thermo Fisher Scientific). IE1-positive nuclei were quantified using the Operetta High-Content
173 Imaging System (PerkinElmer). In some cases, titers were determined by TCID₅₀ analysis, and
174 calculations were performed according to the Reed and Muench formula [49]. All virus stocks
175 were titered by the TCID₅₀ method.

176 Kaempferol (Sigma-Aldrich) was dissolved in DMSO and stored at 4°C until use.
177 PPADS tetrasodium salt (Torcis) was dissolved in dimethyl water and stored at -20°C until use.
178 Cytotoxicity assays for drugs were performed at 96 h post treatment using the CellTiter 96[®]
179 AQueous One Solution Cell Proliferation Assay (Promega, Madison, WI), according to the
180 manufacturer's protocol. Cell viability was measured based on absorbance at 490 nm using a
181 SpectraMax Plus spectrometer (Molecular Devices).

182
183 **siRNA knockdown experiments.** siRNAs targeting P2Y2 (siP2Y2), P2X5 (siP2X5), ENPP4
184 (siENPP4) or ENTPD2 (siENTPD2) were designed and purchased from Life Technologies.
185 HFFs were grown to ~80% confluence in DMEM supplemented with 10% FBS and then

186 transfected with siRNA using Lipofectamine[®] RNAiMAX Reagent (Life Technologies). Cells
187 transfected with non-specific, scrambled siRNA (siSc) (Life Technologies) served as a negative
188 control. Following a 24 h-incubation, cells were washed with PBS and either infected with
189 TB40/E-GFP virus or mock infected.

190

191 **Assay for viral entry.** For drug assays, kaempferol, PPADS, or solvent was applied to 95-100%
192 confluent HFFs for 1 h, and for siRNA analyses, cells were transfected with siP2Y2, siP2X5 or
193 siSc for 24 h. Cells were then washed with PBS and infected at a multiplicity of 1 TCID₅₀/cell or
194 mock infected for 1 h at 4°C. Virus that had not penetrated cells was removed with a low-pH
195 citrate buffer (40 mM citric acid, 10 mM KCl, 135 mM NaCl, pH 3.0) [50]. Cells were then
196 incubated at 37°C in DMEM containing 10% FBS. Cells were either fixed in cold MeOH and
197 immunostained for viral IE1 protein at 24 hpi, as described above, or viral DNA was isolated 1 h
198 after the reaction temperature was raised to 37°C, using the QIAamp DNA Mini kit (Qiagen) and
199 quantified by quantitative polymerase chain reaction (qPCR) performed in duplicate using SYBR
200 Green master mix (Applied Biosystems) using the QuantStudio™ 6 Flex Real-Time PCR System
201 (Applied Biosystems). Primers for the viral UL123 locus were used (Table 1). Fold change was
202 calculated using the $\Delta\Delta C_t$ method and GAPDH was used as an internal control [51].

203

204 **Nucleic acid and protein analyses.** For RNA-Seq analysis of protein-coding RNAs, MRC-5
205 human lung fibroblasts were infected with the AD169 strain of HCMV. RNA from mock- and
206 HCMV-infected cells was collected at 48 hpi and isolated using the miRNeasy kit (Qiagen.).
207 RNA quality was analyzed using the Bioanalyzer 2100 (Agilent Technologies). TruSeq Stranded
208 Total RNA Library Prep Kit with Ribo-Zero (Illumina) was used to generate cDNA libraries as
209 per the manufacturer's instructions.. Briefly, cytosolic ribosomal RNA (rRNA) was depleted

210 from total RNA using biotinylated probes that selectively bind rRNA species. The resulting
211 rRNA-depleted RNA underwent fragmentation, reverse transcription, end repair, 3'-end
212 adenylation, adaptor ligation and subsequent PCR amplification and SPRI bead purification
213 (Beckman Coulter). The unique barcode sequences were incorporated in the adaptors for
214 multiplexed high-throughput sequencing. The final product was assessed for its size distribution
215 and concentration using BioAnalyzer High Sensitivity DNA Kit (Agilent) and Kapa Library
216 Quantification Kit (Kapa Biosystems). The libraries were pooled, denatured and loaded onto a
217 HiSeq Rapid Paired-Read flow cell (Illumina), which was subjected to 2X100 cycles of
218 sequencing by an Illumina HiSeq 2500 (Illumina). Illumina CASAVA pipeline Version 1.8 was
219 used to extract de-multiplexed sequencing reads. Sequencing reads were normalized using the
220 trimmed mean of M-values method (TMM), mapped to the human reference genome and
221 transcripts annotated to which reads have been mapped. Mapped reads were counted and the
222 differential gene expression between triplicate samples from mock- and HCMV-infected cells
223 was computed based on counts per million (CPM). Ingenuity Pathway Analysis cloud software
224 (Qiagen) was used to group the genes identified as significantly up- or down-regulated into gene
225 ontologies and to determine which cellular networks were the most significantly regulated in
226 HCMV-infected cells.

227 For quantitative reverse transcription PCR (qRT-PCR) analysis of RNA, RNA was
228 extracted from samples collected in QIAzol lysis reagent using the miRNeasy kit (Qiagen).
229 Complimentary DNA (cDNA) was made from 1 µg of total RNA with oligo dT and MultiScribe
230 reverse transcriptase (Applied Biosystems), according to the manufacturer's protocol. Reverse
231 transcription reactions were run at 25°C for 10 min, 48°C for 30 min, and 90°C for 5 min.
232 Primers used in the study are listed in Table 1. qPCR was performed in duplicates on equal

233 volumes of cDNA using SYBR Green master mix (Applied Biosystems) on the QuantStudio™ 6
234 Flex Real-Time PCR System (Applied Biosystems). Transcript levels were analyzed using the
235 $\Delta\Delta C_t$ method and GAPDH was used as an internal control [51]. Error ranges are reported as
236 standard error of the mean (SEM).

237 For quantification of viral DNA, total DNA was isolated from HCMV-infected cells or
238 media at 96 hpi, and viral DNA was quantified by qPCR using primers specific for UL123 gene
239 and a standard curve created by serially diluting *BAD_{wt}* DNA. To calculate the particle-to-
240 infectious unit ratio, media collected at 96 hpi was titered and viral DNA was isolated from
241 DNase I-treated (10U/mL; 30 min. at 37C) virions to calculate viral DNA copy number/mL as a
242 measure of virus particles.

243 For protein analysis, fibroblasts were harvested using lysis buffer (50 mM Tris-HCl, pH
244 7.5, 5 mM EDTA, 100 mM NaCl, 1% Triton X-100, 0.1% SDS, and 10% glycerol). Samples
245 were mixed with 6x SDS sample buffer (325 mM Tris pH 6.8, 6% SDS, 48% glycerol, 0.03%
246 bromophenol blue containing 9% 2-mercaptoethanol). Equal protein amounts of the different
247 samples were separated by electrophoresis (SDS-PAGE) and transferred to ImmunoBlot
248 polyvinylidene difluoride (PVDF) membranes (BioRad Laboratories). Western blot analyses
249 were performed using primary antibodies recognizing P2Y2 (H-70; Santa Cruz Biotechnology),
250 P2X5 (Sigma-Aldrich), β -actin-HRP (Abcam), or pUL37x1-specific mouse monoclonal antibody
251 4B6-B [28]. Goat anti-mouse and donkey anti-rabbit (GE Healthcare Biosciences) conjugated
252 with horseradish peroxidase (HRP) were used as secondary antibodies. Western blots were
253 developed using WesternSure ECL Detection Reagents (Licor).

254
255 **Intracellular calcium assay.** HFFs were cultured to 95% confluency, drug-treated and infected
256 at a multiplicity of ~ 1 with *AD_{wt}*, *AD_{sub}UL37x1* viruses or mock infected. At 20 hpi, cells were

257 washed and incubated in media containing 2 μ M Fluo-4 AM dye (Life Technologies) for 30 min.
258 at 37°C. After washing with PBS, images were captured using the Operetta High-Content
259 Imaging System (PerkinElmer) and the fluorescent signal was measured based on 10 cells per
260 experimental arm using ImageJ software [52]. Background fluorescence was subtracted and fold
261 change was determined relative to control samples.

262

263 **Results**

264 **HCMV infection elevates the steady-state levels of several purinergic receptors**

265 To investigate the differential expression of purinergic receptors during infection, we
266 performed whole RNA sequencing of MRC-5 fibroblasts infected with the AD169 strain of
267 HCMV. RNA from mock and infected cells was analyzed in three independent experiments
268 where samples were collected at 48 hpi, allowing us to analyze RNAs during the late phase of
269 the viral replication cycle. Several components involved in purinergic receptor signaling are
270 expressed at significantly higher levels in the infected cells, including P2Y2, P2X5, ENPP4 and
271 ENTPD2, and several others were reduced (Table 2).

272 To confirm the results obtained from RNA-Seq and extend the findings to a different
273 fibroblast cell population and virus strain, we used qRT-PCR to investigate the expression levels
274 of the RNAs identified in our RNA-seq analysis. HFFs were infected with the TB40/E-GFP
275 strain of HCMV at a multiplicity of 3 TCID₅₀/cell. At 48 hpi, HCMV-infected cells expressed
276 76.41 ± 20.10 times more P2Y2, 2.74 ± 0.54 times more P2X5, 9.98 ± 3.22 times more ENPP4
277 and 45.55 ± 16.54 times more ENTPD2 RNA than mock-infected controls (Fig. 1).

278 We also examined the kinetics of P2Y2 and P2X5 receptor expression throughout the
279 viral replication cycle. HFFs were infected at a multiplicity of 3 TCID₅₀/cell, and samples were

280 collected after various time intervals. P2Y2 and P2X5 RNAs were assayed by qRT-PCR and
281 receptor proteins were assayed by Western blot. P2Y2 RNA was elevated at 24 hpi and increased
282 as the infection progressed (Fig. 2A). P2Y2 protein became highly abundant at 48 hpi and its
283 levels stayed elevated through the last time point assayed at 120 hpi (Fig. 2C). These results also
284 qualitatively match the mass spectroscopy-based analysis of cell surface proteins, which
285 determined that P2Y2 protein is 3-, 4- and 5-fold more abundant at 24, 48, and 72 hpi,
286 respectively, compared to uninfected cells [8]. In the case of P2X5, its transcript levels increased
287 up to 48 hpi and then decreased later in infection (Fig. 2B). This corresponded with increasing
288 levels of P2X5 protein between 24 and 72 hpi, which then markedly decreased later in infection
289 (Fig. 2C).

290 Taken together, data from both RNA-Seq and qRT-PCR analyses confirm that HCMV
291 infection in fibroblasts causes an increase in the expression of the cellular purinergic receptors
292 P2Y2 and P2X5, as well as the cell surface ectonucleotidases ENPP4 and ENTPD2.

293

294 **Viral gene expression is necessary for the normal modulation of P2Y2 and P2X5 expression**

295 Since the overexpression of these receptors was seen during the early and late phases of
296 infection, we wished to determine whether the binding of the virion to the fibroblast cell surface
297 was sufficient to induce their upregulation. To do so, we used UV-irradiated TB40/E virus that
298 can bind to and enter cells but cannot express its genes. HFFs were infected with the untreated
299 virus or UV-irradiated virus at a multiplicity of 3 TCID₅₀/cell, samples were collected at 24 and
300 48 hpi and receptor transcript levels were quantified by qRT-PCR. Compared to cells infected
301 with untreated TB40/E virus, those infected with the UV-irradiated virus exhibited 90% lower
302 expression of P2Y2 RNA (Fig. 3A) but 40% higher expression of P2X5 (Fig. 3B). This finding

303 suggests that viral binding and entry is not sufficient to produce the increased levels of P2Y2
304 transcript normally seen during infection. However, viral gene expression appears not to be
305 required for the upregulation of P2X5 expression.

306

307 **P2Y2 and P2X5 receptors have opposite effects on HCMV yield**

308 To determine whether the upregulated expression of these purinergic receptors and
309 ectonucleotidases is related to HCMV pathogenesis, we assessed the effects of inhibiting their
310 expression or activity on the production of infectious progeny.

311 First, we employed siRNAs to knock down their expression. Specific siRNAs designed to
312 target the P2Y2 (siP2Y2), P2X5 (siP2X5), ENPP4 (siENPP4) and ENTPD2 (siENTPD2)
313 transcripts were used to inhibit the expression of these four genes. We verified that the siRNAs
314 reduced expression of their targeted transcripts by using qRT-PCR (Fig. 4A). HFFs were
315 transfected with the selected siRNAs for 24 h before HCMV infection at a multiplicity of 3
316 TCID₅₀/ml. At 120 hpi, the media was collected and viral titer assayed. Knockdown of the P2Y2
317 (~15% of its normal level) or P2X5 receptor (~10% of its normal level) reduced the viral yield
318 by ~95% or increased the yield by almost 400%, respectively (Fig. 4B), compared to viral titer of
319 the media collected from scrambled siRNA (siSc)-treated cells. In contrast, siRNA-mediated
320 knockdown of ENPP4 or ENTPD2 had minimal effects on the amount of infectious progeny
321 released by the infected cells (Fig. 4B).

322 Seeing the significant effect of P2Y2 and P2X5 knockdown on HCMV yield, we focused
323 further study on the roles of these two receptors. To investigate the kinetics of viral release
324 during infection, HFFs were transfected with selected siRNAs for 24 h before HCMV infection
325 at a multiplicity of 3 TCID₅₀/cell. Differences in the amount of virus released from cells

326 transfected with siP2Y2 (reduced virus yield) or siP2X5 (increased virus yield) relative to
327 control cells transfected with siSc were first observed at 72 hpi and gradually increased up to 120
328 hpi (Fig. 4C).

329 We next employed pharmacological perturbations to confirm the roles of the receptors
330 during HCMV infection. Kaempferol is a selective P2Y2 receptor antagonist [17, 53]. PPADS
331 has ~10-fold higher affinity for blocking P2X5 than other P2X family members [23, 54]. We
332 used kaempferol and PPADS at 50 μ M in our studies, because the drugs are commonly used to
333 treat fibroblasts at concentrations of 20-100 μ M [55-58] and we detected little toxicity when
334 uninfected fibroblasts were treated at doses ranging from 0-400 μ M (Fig. 5A, B). To test the
335 effects of inhibiting P2Y2 and P2X5 receptor activity on the release of viral progeny, HFFs were
336 pretreated with drug for 1 h prior to HCMV infection at a multiplicity of 3 TCID₅₀/cell. After
337 allowing 2 h for viral entry into cells, cells were washed and supplemented with media
338 containing either kaempferol or PPADS. Media with drug was replaced every 24 h until media
339 samples were collected at 96 hpi and assayed for infectious virus. Kaempferol decreased the viral
340 yield by more than 99%, whereas PPADS nearly tripled the yield (Fig. 5C). Therefore, we
341 conclude that elevated P2Y2 receptor activity normally facilitates infection whereas the P2X5
342 receptor plays an antiviral role.

343
344 **Inhibiting P2Y2 and P2X5 purinergic receptors does not affect the efficiency of viral entry**
345 **into fibroblasts.**

346 We next aimed to elucidate the point during infection at which the two purinergic
347 receptors play a role. A possible explanation for the observed changes in viral yield upon
348 inhibiting the activity or expression of P2Y2 and P2X5 is that the cell surface purinergic

349 receptors may function in the initial entry of the virus into fibroblasts. Although this notion did
350 not fit well with the kinetics of receptor expression following HCMV infection (Fig. 2),
351 purinergic signaling was shown to be involved in the entry step for HIV-1 infection [17, 59].
352 Therefore, we tested this possibility using both pharmacological antagonists and siRNA-
353 mediated expression knockdown. First, we assessed the efficiency of viral entry into cells that
354 have been pre-treated with kaempferol or PPADS compared to solvent controls. Cells were
355 treated with drugs and after 1 h, cells were washed and HCMV-infected at a multiplicity of 1
356 TCID₅₀/cell for 1 h at 4°C to allow virus attachment to the cell surface. Then, the viral particles
357 that did not attach were removed and cells were incubated at 37°C, allowing the attached virus to
358 penetrate the cells. After 24 h, the cells were fixed and immunostained for IE1 protein. Pre-
359 treatment with either drug did not significantly alter the percentage of cells expressing IE1
360 protein compared to solvent controls (Fig. 6A). To confirm these results, we assessed the
361 efficiency of viral entry in cells transfected with siP2Y2, siP2X5 or siSc as a control. Cells were
362 transfected with the appropriate siRNA for 24 h before HCMV infection at a multiplicity of 1
363 TCID₅₀/cell. After 1 h at 4°C, unattached virions were removed and cultures were shifted to
364 37°C. One hour later, samples were collected and viral DNA copy number was quantified by
365 qPCR. Again, results showed no significant differences in fold change of viral DNA copies
366 between the siP2Y2 or siP2X5 transfected cells and the siSc controls (Fig. 6B). We conclude that
367 inhibiting P2Y2 and P2X5 activity or expression does not affect viral entry into HFFs.

368

369 **Inhibiting P2Y2 reduces the accumulation of viral transcripts**

370 To probe when and how the cellular P2Y2 receptor might affect viral gene expression
371 subsequent to virus entry, we analyzed viral gene expression in siSc- and siP2Y2-treated cells at

372 various times during the viral replication cycle. qRT-PCR was used to quantify transcript levels
373 for representatives of the three main classes of viral genes. The data show that infection in cells
374 transfected with siP2Y2 resulted in a reduction of immediate early (UL123, UL122), early
375 (UL26, UL54), and late (UL69, UL82, UL99) viral transcripts compared to siSc controls (Fig.
376 7A-B, D-H). Viral RNAs were most dependent on P2Y2 during the late phase of infection,
377 consistent with the timing of P2Y2 RNA accumulation (Fig. 2). Intriguingly, one viral transcript,
378 immediate early UL37x1, was expressed at higher levels in cells transfected with siP2Y2 than in
379 control cells (Fig. 7C). UL37x1 has been implicated in the mobilization of intracellular Ca^{2+}
380 during infection [28, 30].

381

382 **P2Y2 receptor affects the efficiency of viral DNA synthesis**

383 One of the strongest inhibitory effects of P2Y2 knockdown was seen on the accumulation
384 of RNA coding for the viral DNA polymerase, UL54 (Fig. 7E). Given its direct role in viral
385 DNA replication [60], we tested whether P2Y2 could have an impact on viral DNA (vDNA)
386 accumulation. Total DNA was isolated from HCMV-infected control and P2Y2-deficient cells
387 and vDNA copy numbers were measured by qPCR using primers specific to UL123. There were
388 about 5-fold fewer copies of vDNA in P2Y2-deficient cells than in control cells (Fig. 8A). This
389 supports the hypothesis that P2Y2 inhibition prevents viral DNA accumulation by blocking viral
390 DNA synthesis. To confirm this result, we also compared the vDNA copy numbers in released
391 viral particles between siP2Y2-treated cells and siSc-treated controls. Similarly, when viral
392 genomic DNA was isolated from viral particles and quantified, there were also nearly 5 times
393 fewer copies of vDNA released from P2Y2-deficient cells compared to control cells (Fig. 8B).
394 To confirm that lower viral DNA synthesis is the predominant explanation for the decrease in

395 infectious particles found in the media collected from P2Y2-deficient cells (Fig. 4B, C and 5C),
396 we decided to test the alternative hypothesis that progeny released from P2Y2 knockdown cells
397 are less infectious by investigating if there are any effects of P2Y2 inhibition on the infectivity of
398 virions. Accordingly, the ratios of vDNA copy number to infectious unit were calculated, and
399 there was no significant difference between the siP2Y2-treated cells and siSc controls (Fig. 8C).
400 Therefore, the reduction in HCMV yield observed in the P2Y2 knockdown condition is caused
401 by reduced viral DNA synthesis and release of viral particles, and not by reduced infectivity of
402 the virions released from P2Y2-deficient cells. Taken together, it appears that P2Y2 is critical for
403 viral DNA synthesis. Inhibiting the P2Y2 receptor reduces viral DNA synthesis and causes a
404 drastic reduction in viral titer.

405

406 **P2Y2 and UL37x1 cooperate to regulate intracellular Ca²⁺ during infection**

407 One of the most striking findings in our analysis of the role of purinergic receptors on
408 viral gene expression during HCMV infection is the fact that of all the viral genes studied, only
409 one was upregulated in the siP2Y2-treated cells compared to siSc-treated control cells (Fig. 7C).
410 This was the transcript of the viral immediate early UL37x1 gene. Previously, UL37x1 has been
411 shown to be involved in the release of Ca²⁺ from SER stores to increase cytosolic Ca²⁺
412 concentrations during infection [28, 30]. Interestingly, P2Y2 is a G-protein coupled receptor that,
413 when activated, leads to the downstream hydrolysis of PIP₂ into IP₃, which is also known to
414 cause the release of Ca²⁺ ions from SER stores [11, 61]. Because of the potential involvement of
415 P2Y2 in Ca²⁺ homeostasis following infection, we investigated the effects of inhibiting P2Y2
416 expression on intracellular Ca²⁺ levels. To do so, fibroblasts were infected with either wild-type
417 AD_{wt} virus or AD_{subUL37x1} virus lacking the UL37x1 gene at a multiplicity of 1 TCID₅₀/cell

418 or mock infected, and cultures were treated 2 h later with kaempferol or solvent control. Using a
419 multiplicity of 1 TCID₅₀/cell allowed us also to measure Ca²⁺ levels in infected and uninfected
420 cells in the same experimental well. At 20 hpi, we used the fluorescence-based fluo-4 AM
421 calcium assay to compare the changes in intracellular Ca²⁺ concentrations. We determined that
422 wild type virus caused ~2.5-fold increase in Ca²⁺ levels compared to uninfected cells (Fig. 9A).
423 We postulated that if P2Y2 contributes to elevated levels of Ca²⁺ in infected cells, then we should
424 see a partial decrease of Ca²⁺ levels in cells either infected with ADwt and treated with
425 kaempferol or infected with AD_{sub}UL37x1. Instead, we saw that in both instances, Ca²⁺ was
426 reduced almost to the level found in uninfected cells (Fig. 9A). Interestingly, Ca²⁺ levels did not
427 decrease further in cells infected with AD_{sub}UL37x1 and treated with kaempferol. Therefore,
428 these data imply that P2Y2 and pUL37x1 work together to elevate Ca²⁺ levels during HCMV
429 infection and that the lack of either component prevents cells from increasing intracellular
430 calcium levels during infection.

431 To further test this hypothesis, we investigated the effect of UL37x1 expression on Ca²⁺
432 homeostasis in HFF cells without the background of infection. For this purpose, a UL37x1-
433 expressing lentiviral pLVX-EF1 plasmid was created. HFF cells were transduced with the vector,
434 and subjected to puromycin selection. Two cellular clones were obtained [pLVX-UL37x1(1) and
435 pLVX-UL37x1(2)] that exhibited stable expression of pUL37x1. The cells were treated with
436 kaempferol or solvent control for 1 h. Then, Ca²⁺ levels were measured using the fluo-4 AM
437 assay and compared to levels in wild type HFF cells. No significant changes in Ca²⁺ levels were
438 observed between samples (Fig. 9B), which supports our conclusion that increased expression of
439 pUL37x1 in the absence of P2Y2 overexpression is not sufficient to stimulate the robust increase
440 in calcium seen in HCMV-infected cells.

441 As a control, cells transduced with pLVX-UL37x1 or pLVX-GFP plasmids were
442 analyzed by Western blot assay for the level of pUL37x1 expression. The assay confirmed the
443 pUL37x1 overexpression only in pLVX-UL37x1 plasmid-transduced HFF cells (Fig. 9C).
444 Because inhibiting the P2Y2 receptor activity decreases the extent to which intracellular Ca^{2+} is
445 mobilized during infection, we propose that P2Y2 signaling plays a critical role in controlling
446 intracellular Ca^{2+} during HCMV infection.

447 Since previous studies as well as our current study have shown that the viral pUL37x1
448 protein is involved in the mobilization of intracellular Ca^{2+} from calcium stores in the SER and
449 knowing that P2Y2 expression was upregulated early in infection, we aimed to further elucidate
450 the relationship between the viral UL37x1 and cellular P2Y2. Specifically, we assessed whether
451 the upregulation of P2Y2 expression during viral infection may be a downstream effect of the
452 immediate early kinetics of UL37x1 expression [62]. Since the upregulation of P2Y2 transcript is
453 most evident around 48hpi, it is conceivable that the viral pUL37x1 protein increases the
454 expression of P2Y2 and that increased P2Y2 activity further increases intracellular Ca^{2+} levels
455 for the remainder of the infection. In order to test this idea, HFFs were infected with either the
456 *ADsubUL37x1* virus or the wild type *ADwt* virus. Samples were collected at 24 hpi. P2Y2,
457 UL54, UL123 and UL37x1 transcript levels were quantified by qRT-PCR. If the hypothesis that
458 P2Y2 upregulation is dependent on pUL37x1 expression were true, then P2Y2 transcripts would
459 be significantly lower in cells infected with *ADsubUL37x1* than *ADwt* virus. However, although
460 P2Y2 expression was appeared somewhat lower in cells infected with *ADsubUL37x1* compared
461 to those infected with *ADwt*, the difference was not significant (Fig. 10A). Interestingly, it was
462 also noted that unlike siP2Y2-treated cells, *ADsubUL37x1*-infected cells did not exhibit lower
463 UL54 RNA levels compared to *ADwt*-infected cells (Fig. 10B). The same expression levels of

464 UL123 in cells infected with either *ADwt* or *ADsubUL37x1* confirm that the cells received an
465 equal amount of virus (Fig. 10C). As expected, UL37x1 expression was only evident in *ADwt*-
466 and not in *ADsubUL37x1*-infected cells (Fig. 10D). Both clonal populations of cells expressing
467 pUL37x1 were characterized by a robust expression of pUL37x1, but there was only a minimally
468 higher level of P2Y2 protein in these cells compared to controls expressing GFP (Fig. 9C).
469 Therefore, these data suggest that viral pUL37x1 is likely not a significant upstream activator of
470 P2Y2 expression.

471 Taken together, these findings suggest that both cellular P2Y2 and viral pUL37x1
472 cooperate to modulate intracellular Ca^{2+} within HCMV-infected cells.

473

474

Discussion

475 The purpose of this study was to investigate the role of purinergic signaling during
476 HCMV infection in human fibroblasts. RNA-Seq analysis identified several members of the
477 purinergic receptor network whose expression was modulated upon infection (Table 2). qRT-
478 PCR analysis confirmed that infection increased the cellular P2Y2, P2X5, ENPP4 and ENTPD2
479 transcripts. Elevated expression of purinergic receptors during HCMV infection has been
480 reported previously. In myeloid cells, HCMV infection increases the expression of the P2Y5
481 receptor [63]. In endothelial cells, HCMV infection has been associated with higher levels of the
482 P2Y2, P2Y1, and P2X7 receptors, as well as ecto-5' nucleotidase CD73 [25, 26]. Yet, the impact
483 of these altered expression patterns on the efficiency of HCMV infection has not been
484 investigated. We now report that siRNA targeting ENPP4 and ENTPD2 transcripts did not affect
485 viral yield (Fig. 4B). In contrast, inhibiting the activity or expression of P2Y2 and P2X5
486 receptors had a robust effect on the amount of infectious progeny released from infected cells

487 (Fig. 4B, C). These findings suggest that, at least in fibroblasts, purinergic signaling through the
488 P2Y2 and P2X5 receptors is important in HCMV infection independent from the activity of
489 ectonucleotidases.

490 Inhibiting P2Y2 and P2X5 by using pharmacological compounds or siRNA technology
491 drastically altered HCMV yield. It is noteworthy that, even though both receptors are activated
492 by ATP, they had completely opposite effects on the production of HCMV progeny. Compared
493 to control treated cells, infection in P2Y2-deficient cells led to a drastic reduction (~95% lower)
494 in the number of infectious virions released (Fig. 4B, C), which demonstrated that P2Y2 supports
495 HCMV replication. On the other hand, infection in P2X5-deficient cells resulted in enhanced (4x
496 higher) viral release compared to control treated cells (Fig. 4B, C), suggesting that P2X5
497 antagonizes HCMV replication.

498 Our analysis of P2Y2 and P2X5 expression kinetics supported the view that these
499 receptors are independently regulated and have different functions during HCMV infection.
500 While active viral gene expression is necessary for the upregulation of P2Y2 expression (Fig.
501 3A), it is not for the increased expression of P2X5 (Fig. 3B). Moreover, P2Y2 gradually
502 increased in expression from 2 to 96 hpi, as seen at both RNA and protein levels (Fig. 2A, C),
503 P2X5 expression increased up to 48 hpi (RNA) or 72 hpi (protein) and then decreased
504 immediately afterwards (Fig. 2B, C). The gradual increase in P2Y2 expression is consistent with
505 a supportive role for the receptor throughout the viral replication cycle. Conversely, the initial
506 spike in the levels of the inhibitory P2X5 receptor might be antagonized by a virus-mediated
507 mechanism that suppresses its expression later after infection.

508 How do the P2Y2 and P2X5 receptors modulate HCMV replication? The P2X5
509 purinergic receptor is a non-selective cation channel that is gated by extracellular ATP. It is

510 typically found on cells of the skin, gut, bladder, thymus, skeletal muscle, and spinal cord and is
511 thought to play important roles in cell differentiation [64]. For instance, it has been shown to be
512 involved in the differentiation of skin keratinocytes and mucosal epithelial cells during turnover
513 in the gut and bladder [19, 65]. P2X5 receptors have also been observed to play various roles in
514 regulating osteoblastic differentiation and proliferation, triggering differentiation of skeletal
515 muscle satellite cells, and the differentiation of human fetal epidermis [66, 67]. Although the
516 expression of P2X5 receptors have been associated with differentiating and proliferating cell
517 types, the exact mechanism by which its signaling affects these cellular processes is still unclear.
518 Thus it is difficult to speculate on how a block to its activity might influence HCMV replication.

519 P2Y2 is a G-protein coupled receptor that can be activated by extracellular ATP and
520 UTP. HCMV infection of P2Y2 knockdown cells exhibited reduced expression of most viral
521 genes tested, including UL54, the viral DNA polymerase catalytic subunit (Fig. 7E). Reduced
522 pUL54 levels would be expected to reduce key downstream events in the viral replication cycle.
523 Indeed, P2Y2-deficient cells exhibited reduced accumulation of viral DNA (Fig. 8A) and late
524 viral RNAs (Fig. 7F-H). Although it is likely that the receptor has additional effects on HCMV
525 replication, its effect on viral DNA accumulation, followed by inhibition of late RNAs that
526 depend on active DNA replication for their synthesis, can explain the low viral yield from P2Y2-
527 deficient cells.

528 There are several plausible mechanisms by which purinergic signaling via the P2Y2
529 receptor can affect viral DNA replication. It could affect DNA replication indirectly, through
530 inhibition of an upstream viral function that we have not yet measured, or it could act directly at
531 DNA replication. One potential mechanism is through p38 mitogen-activated protein kinases
532 (MAPKs). Extracellular ATP and UTP can activate the MAPK kinase (MKK) 3/6-p38-MAPK

533 cascade via the P2Y2 receptor in glomerular mesangial cells [68]. p38 activation results from
534 increased activity of MKK3/6, which are downstream effectors of P2Y2 signaling [69]. It has
535 been reported that active p38 plays a critical role in HCMV viral DNA replication in infected
536 human fibroblasts [69]. Thus, inhibiting P2Y2 could decrease MKK3/6 activity, which might
537 then reduce p38 activation and viral DNA replication. This role for P2Y2 action within infected
538 cells remains speculative, because the exact mechanism by which p38 regulates viral DNA
539 replication via the UL54 polymerase has not yet been characterized.

540 In addition to its impact on p38 MAPK, the P2Y2 receptor has a well-established role in
541 regulating cellular calcium levels [15, 16, 23]. Extracellular nucleotides act via the P2Y2
542 receptor to induce intracellular Ca^{2+} mobilization from SER in a variety of cell types [70-72].
543 The increase in intracellular Ca^{2+} can be blocked by PLC inhibitors and by low molecular weight
544 heparin, indicating the involvement of IP_3 -sensitive intracellular Ca^{2+} stores, which is known to
545 be downstream of P2Y2-mediated signaling [73]. As calcium homeostasis was found to be
546 critical for an efficient HCMV infection [27, 28], it was intriguing to speculate that P2Y2
547 inhibition may have important consequences for HCMV replication due to disruption of
548 intracellular calcium regulation. Treating infected cells with drugs that disrupt SER Ca^{2+}
549 homeostasis inhibits the production of infectious progeny due to retarded accumulation of late
550 gene products [27], patterns that were observed during HCMV infection in P2Y2 knockdown
551 cells.

552 The first indication that P2Y2-dependent Ca^{2+} regulation may be important during
553 HCMV infection came from analyzing differential expressions of viral genes in wild type cells
554 and cells treated with siP2Y2. Among all tested viral transcripts only UL37x1 RNA levels were
555 upregulated in P2Y2-deficient cells (Fig. 7C). It is known that the HCMV immediate early

556 protein pUL37x1 induces the mobilization of Ca^{2+} from the SER to the cytosol [28]. In addition,
557 the Ca^{2+} -dependent protein kinase, PKC α , is activated following infection, leading to the
558 production of large (1-5 μm diameter) cytoplasmic vesicles late after infection [30]. The
559 presence of these vesicles correlated with the efficient accumulation of enveloped virions.
560 Regulating intracellular Ca^{2+} levels may also influence viral DNA replication. Ca^{2+} can induce
561 the activity of Ca^{2+} /calmodulin-dependent protein kinase kinase (CaMKK), which activates
562 AMPK. AMPK activity has been shown to be necessary for HCMV DNA synthesis and the
563 expression of viral late genes [33, 34]. In addition, it has been found that extracellular
564 nucleotides stimulate the PI3-K/Akt pathway through P2Y2-mediated signaling involving Ca^{2+}
565 influx, CaM, and Src [74]. It was previously reported that treating infected fibroblasts with an
566 inhibitor of PI3-K activity caused inhibition of viral DNA replication and a 4-log decrease in
567 viral titers [36]. Although intriguing, it is more likely that P2Y2-mediated inhibition of viral
568 UL54 is independent from its effects on intracellular Ca^{2+} . Cells infected with AD_{sub}UL37x1,
569 which lacks the UL37x1 gene, show reduced intracellular Ca^{2+} levels (Fig. 9A), but,
570 nevertheless, do not show reduced UL54 RNA levels (Fig. 10B). Hence the effect P2Y2 on viral
571 DNA accumulation may be distinct from its effects on intracellular Ca^{2+} concentrations.

572 To broaden our understanding on a potential role of P2Y2 in Ca^{2+} levels and the
573 relationship between P2Y2 and UL37x1 during infection, we tested the effect of kaempferol, a
574 P2Y2 antagonist, on intracellular Ca^{2+} in cells infected with wild-type HCMV (AD_{wt}) versus the
575 pUL37x1-deficient mutant virus (AD_{sub}UL37x1). As shown previously [28], AD_{wt}-infected
576 cells exhibited increased free Ca^{2+} compared to uninfected cells, whereas AD_{sub}UL37x1-
577 infected cells had Ca^{2+} levels similar to those in uninfected cells (Fig. 9A). Interestingly,
578 kaempferol treatment also lowered Ca^{2+} levels in AD_{wt}-infected cells to its level in uninfected

579 cells and similar Ca^{2+} levels were measured in kaempferol-treated, *ADsubUL37x1*-infected cells
580 (Fig. 9A). Additionally, when we tested Ca^{2+} levels in uninfected cells that overexpress UL37x1,
581 no elevated levels of Ca^{2+} were observed (Fig. 9B). While P2Y2-deficient cells were
582 characterized by increased expression of UL37x1 (Fig. 7C), neither *ADsubUL37x1*-infected
583 cells nor cells overexpressing UL37x1 affected P2Y2 RNA or protein levels (Fig.10A, 9C).
584 Taken together, these results strongly suggest that P2Y2 has a critical role in regulating Ca^{2+}
585 levels following HCMV infection. Both P2Y2 and pUL37x1 are indispensable for maintaining
586 favorable intracellular Ca^{2+} levels during infection. It is not yet clear how the viral and cellular
587 gene products cooperate in this process.

588 Aside from disrupting viral DNA replication and Ca^{2+} homeostasis, a third mechanism by
589 which P2Y2 activation may play a role in HCMV infection is through affecting the nuclear
590 egress of the viral capsid. It has been reported that PKC, a downstream effector of P2Y2
591 signaling, plays an important role in destabilizing the nuclear lamina by phosphorylating several
592 types of nuclear lamin [75]. The phosphorylation and reorganization of the nuclear lamins is an
593 essential part of viral nuclear egress [75]. It is conceivable, then, that a block to nuclear egress of
594 HCMV capsids could contribute to the reduction in infectious virus produced following
595 inhibition of P2Y2 (Fig. 4, 5).

596 It has been reported that during hypoxic conditions, fibroblasts release ATP, which
597 activates P2Y2 to regulate cellular DNA synthesis via extracellular signal-regulated kinases 1
598 and 2 (ERK_{1/2}) induced Egr-1 activation [14]. Additionally, it has been found that in human
599 cardiac fibroblasts, ATP-mediated activation of the P2Y2 receptor leads to cell cycle progression
600 and cell proliferation [76]. Moreover, P2Y2 activation results in the activation of PKC, which
601 acts in concert with ERK and PI3-K/PKC pathways to induce c-Fos protein expression and HeLa

602 cell proliferation [77]. Since P2Y2 is dramatically overexpressed in HCMV-infected cells, it
603 would be interesting to understand how the expected upregulation of cellular DNA synthesis and
604 cell cycle progression is avoided in HCMV-infected cells in order to maintain the cell cycle
605 arrest characteristic of infection [78, 79].

606 Signaling via the P2Y2 receptor also may influence cell migration. P2Y2 activation
607 increases MCF-7 breast cancer cell migration via the MEK-ERK_{1/2} signaling pathway [80].
608 Specifically, extracellular ATP can activate MAPKs through the P2Y2/PLC/PKC/ERK signaling
609 pathway to induce the translocation of ERK_{1/2} into the nucleus [13]. Also, fibroblasts appear to
610 require PKC activation, which is downstream of P2Y2 signaling, in order to respond to
611 hyaluronan stimulation with increased locomotion [81]. Taken together, it is conceivable that the
612 overexpression of P2Y2 receptors during HCMV infection may lead to increased migration of
613 the host cell. This not only has potential to facilitate dissemination of the virus within its infected
614 host, it may also contribute to the likely oncomodulatory activity of HCMV [82].

615 In summary, our results show that purinergic signaling through the P2Y2 and P2X5
616 receptors plays a critical role within infected cells and set the stage for additional investigations
617 on the impact of this receptor family in HCMV biology. Further, our observation that a
618 pharmacological block to P2Y2 dramatically reduces viral yield raises the possibility that P2Y2
619 antagonists, if well tolerated, could prove to be attractive candidates for new HCMV therapies.

620

621 **Acknowledgments:**

622 This work was supported by a grant from the National Institutes of Health (AI112951).
623 M.N. was supported by a fellowships from the American Cancer Society (PF-14-116-01-MPC).

624

625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644

References

1. Sia, I.G., et al., *Cytomegalovirus (CMV) DNA load predicts relapsing CMV infection after solid organ transplantation*. J Infect Dis, 2000. **181**(2): p. 717-20.
2. George, M.J., et al., *The independent role of cytomegalovirus as a risk factor for invasive fungal disease in orthotopic liver transplant recipients*. Boston Center for Liver Transplantation CMVIG-Study Group. Cytogam, MedImmune, Inc. Gaithersburg, Maryland. Am J Med, 1997. **103**(2): p. 106-13.
3. Kanj, S.S., et al., *Cytomegalovirus infection following liver transplantation: review of the literature*. Clin Infect Dis, 1996. **22**(3): p. 537-49.
4. Gerna, G., et al., *Human cytomegalovirus end-organ disease is associated with high or low systemic viral load in preemptively treated solid-organ transplant recipients*. New Microbiol, 2012. **35**(3): p. 279-87.
5. Tabata, T., et al., *Human cytomegalovirus infection interferes with the maintenance and differentiation of trophoblast progenitor cells of the human placenta*. J Virol, 2015. **89**(9): p. 5134-47.
6. Lanzieri, T.M., et al., *Cytomegalovirus infection among infants in California neonatal intensive care units, 2005-2010*. J Perinat Med, 2014. **42**(3): p. 393-9.
7. Gudleski-O'Regan, N., et al., *Increased expression of LDL receptor-related protein 1 during human cytomegalovirus infection reduces virion cholesterol and infectivity*. Cell Host Microbe, 2012. **12**(1): p. 86-96.

- 645 8. Weekes, M.P., et al., *Quantitative temporal viromics: an approach to investigate host-*
646 *pathogen interaction*. Cell, 2014. **157**(6): p. 1460-72.
- 647 9. North, R.A., *Molecular physiology of P2X receptors*. Physiol Rev, 2002. **82**(4): p. 1013-
648 67.
- 649 10. Fischer, W., et al., *Conserved lysin and arginin residues in the extracellular loop of*
650 *P2X(3) receptors are involved in agonist binding*. Eur J Pharmacol, 2007. **576**(1-3): p. 7-
651 17.
- 652 11. Ralevic, V. and G. Burnstock, *Receptors for purines and pyrimidines*. Pharmacol Rev,
653 1998. **50**(3): p. 413-92.
- 654 12. Nishizuka, Y., *Intracellular signaling by hydrolysis of phospholipids and activation of*
655 *protein kinase C*. Science, 1992. **258**(5082): p. 607-14.
- 656 13. Chang, S.J., et al., *Extracellular ATP activates the PLC/PKC/ERK signaling pathway*
657 *through the P2Y2 purinergic receptor leading to the induction of early growth response 1*
658 *expression and the inhibition of viability in human endometrial stromal cells*. Cell Signal,
659 2008. **20**(7): p. 1248-55.
- 660 14. Gerasimovskaya, E.V., et al., *Extracellular ATP is an autocrine/paracrine regulator of*
661 *hypoxia-induced adventitial fibroblast growth. Signaling through extracellular signal-*
662 *regulated kinase-1/2 and the Egr-1 transcription factor*. J Biol Chem, 2002. **277**(47): p.
663 44638-50.

- 664 15. Homolya, L., et al., *Nucleotide-regulated calcium signaling in lung fibroblasts and*
665 *epithelial cells from normal and P2Y₂ receptor (-/-) mice*. J Biol Chem, 1999. **274**(37):
666 p. 26454-60.
- 667 16. Xie, R., et al., *The P2Y₂ nucleotide receptor mediates the proliferation and migration of*
668 *human hepatocellular carcinoma cells induced by ATP*. J Biol Chem, 2014. **289**(27): p.
669 19137-49.
- 670 17. Seror, C., et al., *Extracellular ATP acts on P2Y₂ purinergic receptors to facilitate HIV-1*
671 *infection*. J Exp Med, 2011. **208**(9): p. 1823-34.
- 672 18. Compan, V., et al., *P2X₂ and P2X₅ subunits define a new heteromeric receptor with*
673 *P2X₇-like properties*. J Neurosci, 2012. **32**(12): p. 4284-96.
- 674 19. Groschel-Stewart, U., et al., *Localisation of P2X₅ and P2X₇ receptors by*
675 *immunohistochemistry in rat stratified squamous epithelia*. Cell Tissue Res, 1999.
676 **296**(3): p. 599-605.
- 677 20. Greig, A.V., et al., *Expression of purinergic receptors in non-melanoma skin cancers and*
678 *their functional roles in A431 cells*. J Invest Dermatol, 2003. **121**(2): p. 315-27.
- 679 21. Calvert, R.C., et al., *Immunocytochemical and pharmacological characterisation of P2-*
680 *purinoceptor-mediated cell growth and death in PC-3 hormone refractory prostate*
681 *cancer cells*. Anticancer Res, 2004. **24**(5A): p. 2853-9.
- 682 22. Overes, I.M., et al., *Expression of P2X₅ in lymphoid malignancies results in LRH-1-*
683 *specific cytotoxic T-cell-mediated lysis*. Br J Haematol, 2008. **141**(6): p. 799-807.

- 684 23. Bo, X., et al., *Pharmacological and biophysical properties of the human P2X5 receptor*.
685 *Mol Pharmacol*, 2003. **63**(6): p. 1407-16.
- 686 24. Fortunato, E.A., et al., *Exploitation of cellular signaling and regulatory pathways by*
687 *human cytomegalovirus*. *Trends Microbiol*, 2000. **8**(3): p. 111-9.
- 688 25. Zandberg, M., et al., *Infection of human endothelium in vitro by cytomegalovirus causes*
689 *enhanced expression of purinergic receptors: a potential virus escape mechanism?*
690 *Transplantation*, 2007. **84**(10): p. 1343-7.
- 691 26. Kas-Deelen, A.M., et al., *Cytomegalovirus infection increases the expression and activity*
692 *of ecto-ATPase (CD39) and ecto-5'nucleotidase (CD73) on endothelial cells*. *FEBS Lett*,
693 2001. **491**(1-2): p. 21-5.
- 694 27. Isler, J.A., T.G. Maguire, and J.C. Alwine, *Production of infectious human*
695 *cytomegalovirus virions is inhibited by drugs that disrupt calcium homeostasis in the*
696 *endoplasmic reticulum*. *J Virol*, 2005. **79**(24): p. 15388-97.
- 697 28. Sharon-Friling, R., et al., *Human cytomegalovirus pUL37x1 induces the release of*
698 *endoplasmic reticulum calcium stores*. *Proc Natl Acad Sci U S A*, 2006. **103**(50): p.
699 19117-22.
- 700 29. Zhao, L. and S.L. Ackerman, *Endoplasmic reticulum stress in health and disease*. *Curr*
701 *Opin Cell Biol*, 2006. **18**(4): p. 444-52.

- 702 30. Sharon-Friling, R. and T. Shenk, *Human cytomegalovirus pUL37x1-induced calcium flux*
703 *activates PKC α , inducing altered cell shape and accumulation of cytoplasmic*
704 *vesicles*. Proc Natl Acad Sci U S A, 2014. **111**(12): p. E1140-8.
- 705 31. Hajnoczky, G., et al., *Decoding of cytosolic calcium oscillations in the mitochondria*.
706 Cell, 1995. **82**(3): p. 415-24.
- 707 32. Jouaville, L.S., et al., *Regulation of mitochondrial ATP synthesis by calcium: evidence*
708 *for a long-term metabolic priming*. Proc Natl Acad Sci U S A, 1999. **96**(24): p. 13807-12.
- 709 33. McArdle, J., N.J. Moorman, and J. Munger, *HCMV targets the metabolic stress response*
710 *through activation of AMPK whose activity is important for viral replication*. PLoS
711 Pathog, 2012. **8**(1): p. e1002502.
- 712 34. Terry, L.J., et al., *Human kinome profiling identifies a requirement for AMP-activated*
713 *protein kinase during human cytomegalovirus infection*. Proc Natl Acad Sci U S A, 2012.
714 **109**(8): p. 3071-6.
- 715 35. Wiebusch, L., M. Truss, and C. Hagemeyer, *Inhibition of human cytomegalovirus*
716 *replication by small interfering RNAs*. J Gen Virol, 2004. **85**(Pt 1): p. 179-84.
- 717 36. Johnson, R.A., et al., *Human cytomegalovirus up-regulates the phosphatidylinositol 3-*
718 *kinase (PI3-K) pathway: inhibition of PI3-K activity inhibits viral replication and virus-*
719 *induced signaling*. J Virol, 2001. **75**(13): p. 6022-32.

- 720 37. Johnson, R.A., et al., *Domain mapping of the human cytomegalovirus IE1-72 and*
721 *cellular p107 protein-protein interaction and the possible functional consequences.* J Gen
722 Virol, 1999. **80** (Pt 5): p. 1293-303.
- 723 38. Buzzi, N., et al., *Extracellular ATP activates MAP kinase cascades through a P2Y*
724 *purinergic receptor in the human intestinal Caco-2 cell line.* Biochim Biophys Acta,
725 2009. **1790**(12): p. 1651-9.
- 726 39. Bian, S., et al., *P2X7 integrates PI3K/AKT and AMPK-PRAS40-mTOR signaling*
727 *pathways to mediate tumor cell death.* PLoS One, 2013. **8**(4): p. e60184.
- 728 40. Nakamura, T., et al., *Benidipine, a long-acting calcium-channel blocker, prevents the*
729 *progression to end-stage renal failure in a rat mesangioproliferative glomerulonephritis.*
730 Nephron, 2000. **86**(3): p. 315-26.
- 731 41. Pinton, P., et al., *The Ca²⁺ concentration of the endoplasmic reticulum is a key*
732 *determinant of ceramide-induced apoptosis: significance for the molecular mechanism of*
733 *Bcl-2 action.* EMBO J, 2001. **20**(11): p. 2690-701.
- 734 42. Gonzalez, G.A. and M.R. Montminy, *Cyclic AMP stimulates somatostatin gene*
735 *transcription by phosphorylation of CREB at serine 133.* Cell, 1989. **59**(4): p. 675-80.
- 736 43. Yu, D., et al., *Construction of a self-excisable bacterial artificial chromosome containing*
737 *the human cytomegalovirus genome and mutagenesis of the diploid TRL/IRL13 gene.* J
738 Virol, 2002. **76**(5): p. 2316-28.

- 739 44. Sinzger, C., et al., *Cloning and sequencing of a highly productive, endotheliotropic virus*
740 *strain derived from human cytomegalovirus TB40/E*. J Gen Virol, 2008. **89**(Pt 2): p. 359-
741 68.
- 742 45. Gaspar, M. and T. Shenk, *Human cytomegalovirus inhibits a DNA damage response by*
743 *mislocalizing checkpoint proteins*. Proc Natl Acad Sci U S A, 2006. **103**(8): p. 2821-6.
- 744 46. Murphy, E., et al., *Coding potential of laboratory and clinical strains of human*
745 *cytomegalovirus*. Proc Natl Acad Sci U S A, 2003. **100**(25): p. 14976-81.
- 746 47. Koyuncu, E., et al., *Saturated very long chain Fatty acids are required for the production*
747 *of infectious human cytomegalovirus progeny*. PLoS Pathog, 2013. **9**(5): p. e1003333.
- 748 48. Zhu, H., Y. Shen, and T. Shenk, *Human cytomegalovirus IE1 and IE2 proteins block*
749 *apoptosis*. J Virol, 1995. **69**(12): p. 7960-70.
- 750 49. Reed, L.J. and H. Muench, *A simple method of estimating fifty percent endpoints*. The
751 American Journal of Hygiene, 1938. **27**: p. 493-497.
- 752 50. Highlander, S.L., et al., *Neutralizing monoclonal antibodies specific for herpes simplex*
753 *virus glycoprotein D inhibit virus penetration*. J Virol, 1987. **61**(11): p. 3356-64.
- 754 51. Livak, K.J. and T.D. Schmittgen, *Analysis of relative gene expression data using real-*
755 *time quantitative PCR and the 2^{(-Delta Delta C(T))} Method*. Methods, 2001. **25**(4): p.
756 402-8.
- 757 52. Rasband, W.S., *ImageJ*. 1997-2010, U.S. National Institutes of Health: Bethesda,
758 Maryland, USA.

- 759 53. Kaulich, M., et al., *Flavonoids — novel lead compounds for the development of P2Y2*
760 *receptor antagonists*. Drug Development Research, 2003. **59**(1): p. 72-81.
- 761 54. North, R.A. and A. Surprenant, *Pharmacology of cloned P2X receptors*. Annu Rev
762 Pharmacol Toxicol, 2000. **40**: p. 563-80.
- 763 55. Yoon, H.Y., et al., *Kaempferol inhibits IL-1beta-induced proliferation of rheumatoid*
764 *arthritis synovial fibroblasts and the production of COX-2, PGE2 and MMPs*. Int J Mol
765 Med, 2013. **32**(4): p. 971-7.
- 766 56. Granatiero, V., et al., *Reduced mitochondrial Ca(2+) transients stimulate autophagy in*
767 *human fibroblasts carrying the 13514A>G mutation of the ND5 subunit of NADH*
768 *dehydrogenase*. Cell Death Differ, 2016. **23**(2): p. 231-41.
- 769 57. Janssen, L.J., et al., *ATP stimulates Ca(2+)-waves and gene expression in cultured*
770 *human pulmonary fibroblasts*. Int J Biochem Cell Biol, 2009. **41**(12): p. 2477-84.
- 771 58. Mezzaroma, E., et al., *The inflammasome promotes adverse cardiac remodeling*
772 *following acute myocardial infarction in the mouse*. Proc Natl Acad Sci U S A, 2011.
773 **108**(49): p. 19725-30.
- 774 59. Hazleton, J.E., J.W. Berman, and E.A. Eugenin, *Purinergic receptors are required for*
775 *HIV-1 infection of primary human macrophages*. J Immunol, 2012. **188**(9): p. 4488-95.
- 776 60. Kerry, J.A., et al., *Multiple regulatory events influence human cytomegalovirus DNA*
777 *polymerase (UL54) expression during viral infection*. J Virol, 1996. **70**(1): p. 373-82.

- 778 61. Klein, I., et al., [*Molecular genetic study of medullary thyroid cancer*]. Orv Hetil, 1999.
779 **140**(49): p. 2739-46.
- 780 62. Adair, R., et al., *Human cytomegalovirus temporally regulated gene expression in*
781 *differentiated, immortalized retinal pigment epithelial cells*. J Clin Virol, 2006. **35**(4): p.
782 478-84.
- 783 63. Slobedman, B., et al., *Impact of human cytomegalovirus latent infection on myeloid*
784 *progenitor cell gene expression*. J Virol, 2004. **78**(8): p. 4054-62.
- 785 64. Ruan, H.Z. and G. Burnstock, *The distribution of P2X5 purinergic receptors in the*
786 *enteric nervous system of mouse*. Cell Tissue Res, 2005. **319**(2): p. 191-200.
- 787 65. Lee, H.Y., M. Bardini, and G. Burnstock, *Distribution of P2X receptors in the urinary*
788 *bladder and the ureter of the rat*. J Urol, 2000. **163**(6): p. 2002-7.
- 789 66. Hoebertz, A., et al., *Expression of P2 receptors in bone and cultured bone cells*. Bone,
790 2000. **27**(4): p. 503-10.
- 791 67. Greig, A.V., et al., *Purinergic receptors are part of a functional signaling system for*
792 *proliferation and differentiation of human epidermal keratinocytes*. J Invest Dermatol,
793 2003. **120**(6): p. 1007-15.
- 794 68. Huwiler, A., et al., *Extracellular nucleotides activate the p38-stress-activated protein*
795 *kinase cascade in glomerular mesangial cells*. Br J Pharmacol, 2000. **129**(3): p. 612-8.

- 796 69. Johnson, R.A., S.M. Huong, and E.S. Huang, *Activation of the mitogen-activated protein*
797 *kinase p38 by human cytomegalovirus infection through two distinct pathways: a novel*
798 *mechanism for activation of p38*. J Virol, 2000. **74**(3): p. 1158-67.
- 799 70. Berra-Romani, R., et al., *The mechanism of injury-induced intracellular calcium*
800 *concentration oscillations in the endothelium of excised rat aorta*. J Vasc Res, 2012.
801 **49**(1): p. 65-76.
- 802 71. Sugamoto, Y., K. Hirai, and T. Tokoro, *P2Y2 receptor elevates intracellular calcium*
803 *concentration in rabbit eye suprachoroid*. J Med Dent Sci, 1999. **46**(2): p. 83-92.
- 804 72. Shahidullah, M. and W.S. Wilson, *Mobilisation of intracellular calcium by P2Y2*
805 *receptors in cultured, non-transformed bovine ciliary epithelial cells*. Curr Eye Res,
806 1997. **16**(10): p. 1006-16.
- 807 73. Viana, F., et al., *Calcium signalling through nucleotide receptor P2Y2 in cultured human*
808 *vascular endothelium*. Cell Calcium, 1998. **24**(2): p. 117-27.
- 809 74. Katz, S., et al., *Activation of the PI3K/Akt signaling pathway through P2Y(2) receptors*
810 *by extracellular ATP is involved in osteoblastic cell proliferation*. Arch Biochem
811 Biophys, 2011. **513**(2): p. 144-52.
- 812 75. Milbradt, J., et al., *Novel mode of phosphorylation-triggered reorganization of the*
813 *nuclear lamina during nuclear egress of human cytomegalovirus*. J Biol Chem, 2010.
814 **285**(18): p. 13979-89.

- 815 76. Chen, J.B., et al., *Adenosine-5'-triphosphate up-regulates proliferation of human cardiac*
816 *fibroblasts*. Br J Pharmacol, 2012. **166**(3): p. 1140-50.
- 817 77. Muscella, A., et al., *Activation of P2Y2 receptor induces c-FOS protein through a*
818 *pathway involving mitogen-activated protein kinases and phosphoinositide 3-kinases in*
819 *HeLa cells*. J Cell Physiol, 2003. **195**(2): p. 234-40.
- 820 78. Bresnahan, W.A., et al., *Human cytomegalovirus inhibits cellular DNA synthesis and*
821 *arrests productively infected cells in late G1*. Virology, 1996. **224**(1): p. 150-60.
- 822 79. Lu, M. and T. Shenk, *Human cytomegalovirus infection inhibits cell cycle progression at*
823 *multiple points, including the transition from G1 to S*. J Virol, 1996. **70**(12): p. 8850-7.
- 824 80. Chadet, S., et al., *The activation of P2Y2 receptors increases MCF-7 breast cancer cells*
825 *migration through the MEK-ERK1/2 signalling pathway*. Carcinogenesis, 2014. **35**(6): p.
826 1238-47.
- 827 81. Hall, C.L., et al., *Fibroblasts require protein kinase C activation to respond to*
828 *hyaluronan with increased locomotion*. Matrix Biol, 2001. **20**(3): p. 183-92.
- 829 82. Dziurzynski, K., et al., *Consensus on the role of human cytomegalovirus in glioblastoma*.
830 Neuro Oncol, 2012. **14**(3): p. 246-55.
831

Gene	Forward Primer (5'→3')	Reverse Primer (5'→3')
UL123	TGC TGT GCT GCT ATG TCT TAG AGG	TTG GTT ATC AGA GGC CGC TTGG
UL122	TGA CCG AGG ATT GCA ACGA	CGG CAT GAT TGA CAG CCTG
UL37x1	TCC CGC CTT GGT TAA GA	CGA GTT CTC ACC GTC AAT TA
UL26	CCA GCA GCT TCC AGT ATT C	ACC TGG ATC TGC CCT ATC
UL54	TGC TTT CGT CGG TGC TCT CTA AG	TGT GCG GCA GGT TAG ATT GACG
UL69	ACG AGT GTC AGA ACG AGA TGT GC	TGA AAC GAT AGG GTG CCA ACG C
UL82	AGA CGT CGA AGC GGT AAC AAC G	AGT CGT CAA GGC TCG CAA AGA C
UL99	ACG ACA ACA TCC CTC CGA CTTC	TCT GTT GCC GCT CCT CGT TATC
P2Y2	CCA CCT GCC TTC TCA CTA GC	TGG GAA ATC TCA AGG ACT GG
P2X5	CCT GCG GAG AGA GAA CTT GG	AGT TGA ATT TGG GGA AAC GGA TG
ENPP4	TTT GGT TGC CGA TGA AGG CT	CGT GGG CAG CTA GAA ATG GA
ENPTD2	TCA ATC CAG CTC CTT GAA CC	TCC CCA GTA CAG ACC CAG AC
GAPDH	CAA GAG CAC AAG AGG AAG AGA G	CTA CAT GGC AAC TGT GAG GAG

TABLE 1. Primers used in qPCR assays.

Gene	Description	Log ₂ (FC)
P2Y2	Purinergic receptor, P2Y subtype 2	7.49
ENPP4	Ectonucleotide pyrophosphatase/phosphodiesterase 4	4.20
P2X5	Purinergic receptor, P2X subtype 5	3.21
ENTPD2	Ectonucleoside triphosphate diphosphohydrolase 2	3.18
P2X1	Purinergic receptor, P2X subtype 1	2.78
ENTPD8	Ectonucleoside triphosphate diphosphohydrolase 8	2.65
PANX2	Pannexin channel 2	2.28
ENPP5	Ectonucleotide pyrophosphatase/phosphodiesterase 5	1.41
ENPP6	Ectonucleotide pyrophosphatase/phosphodiesterase 6	-3.66
ENPP7P10	Ectonucleotide pyrophosphatase/phosphodiesterase 7 pseudogene 10	-3.58
P2X6	Purinergic receptor, P2X subtype 6	-2.76
ENPP7P4	Ectonucleotide pyrophosphatase/phosphodiesterase 7 pseudogene 4	-2.17
P2X7	Purinergic receptor, P2X subtype 7	-1.97
NT5E	Ecto-5' nucleotidase (CD73)	-1.68
ENPP7P12	Ectonucleotide pyrophosphatase/phosphodiesterase 7 pseudogene 12	-1.55

TABLE 2. Differential expression of members of the purinergic receptor network in HCMV-infected MRC-5 fibroblasts at 48 hpi compared to mock-infected controls. Log₂ fold change (FC) was calculated from sequence reads determined from triplicate biological samples.




FIG 1. HCMV infection increases the transcript levels of P2Y2, P2X5, ENPP4, and ENTPD2 at 48 hpi. HFFs were infected with TB40/E-GFP virus (MOI=3) or mock infected for 2 h. Samples were collected at 48 hpi, transcripts were quantified by qRT-PCR and GAPDH was used as an internal control. Data are represented as fold change compared to mock-infected cells. Results were averaged between two biological replicates and error bars represent standard error of the mean (SEM).




FIG 2. Kinetics of P2Y2 and P2X5 expression in HCMV-infected HFFs. (A, B) P2Y2 and P2X5 RNA levels are modulated by infection. HFFs were infected with TB40/E-GFP (MOI=3) or mock infected. RNA and protein samples were collected at various times after infection, transcripts were quantified by qRT-PCR and GAPDH was used as an internal control. Results are shown as fold change compared to mock-infected cells. Data were averaged for two biological replicates and error bars represent SEM. **(C) P2Y2 and P2X5 proteins are modulated by infection.** Proteins were subjected to Western blot analyses using antibodies specific for P2Y2 and P2X5. Actin was monitored as a loading control.




FIG 3. UV-irradiated HCMV failed to increase P2Y2 but did increase P2X5 RNA levels. Untreated or UV-irradiated TB40/E virus was applied to HFFs (MOI=3). Samples were collected at 48 hpi (P2Y2) or 24 hpi (P2X5), and transcripts were quantified by qRT-PCR. GAPDH was assayed as an internal control. Results are shown as fold change compared to untreated control virus condition. Data were averaged between two biological replicates and error bars represent SEM.






FIG 4. siRNA-mediated knockdown of P2Y2 or P2X5 affects HCMV yield. HFFs were transfected with siP2Y2, siP2X5, siENPP4, siENTPD2, or scrambled siRNA (siSc) as a control. After 24 h, cells were infected with TB40/E-GFP virus (MOI=3). **(A) siRNA validation.** Samples were collected at 48 hpi and qRT-PCR was used to measure levels of P2Y2, P2X5, ENPP4 and ENTPD2 transcripts. GAPDH was used as an internal control. The data are represented as fold change compared to siSc control. The mean fold change for three biological replicates are shown and error bars represent SEM. **(B) Effect of siRNA-mediated knockdowns on virus yield at 120 hpi.** TCID₅₀/ml values were determined and fold change was calculated relative to siSc control. Results show the average fold change from two biological replicates and error bars represent SEM. **(C) Virus growth kinetics following siRNA-mediated knockdown.** Samples were collected after various time intervals. Viral titers were determined by applying the infectious media to a reporter plate of HFFs and immunostaining for IE1 protein 24 h later. The data are presented as percent of IE1-positive cells averaged from two biological replicates and error bars represent SEM.




FIG 5. Kaempferol and PPADS have opposite effects on HCMV yield. (A, B) Drug toxicity assays. HFFs were treated with various concentrations of (A) kaempferol and (B) PPADS. The media with drugs was replenished every 24 h. At 96 h, cell viability was measured using the CellTiter 96[®] AQ_{ueous} Assay. Results are shown as average absorbance from two biological replicates and error bars represent SEM. **(C) Drugs modulate the yield of HCMV.** HFFs were infected with TB40/E-GFP virus (MOI=3) or mock infected for 2 h and treated with either kaempferol (50 μM) or PPADS (50 μM). The drugs were replaced every 24 h until media samples were collected at 96 hpi. Viral yield was determined by monitoring expression of IE1 protein at 24 h after infection. Results were averaged between two biological replicates and error bars represent SEM.




FIG 6. P2Y2 and P2X5 receptors do not affect HCMV entry into HFFs. HFFs were either **(A)** pre-treated with either kaempferol (50 μ M) or PPADS (50 μ M), or solvent control for 1 h, or **(B)** transfected with siP2Y2, siP2X5, or siSc as a control for 24 h. Then, they were infected with TB40/E-GFP virus (MOI=1) or mock-infected for 1 h at 4°C. Cells were washed with citrate buffer and incubated at 37°C. Viral entry was assayed by either **(A)** immunostaining for IE1 protein at 24 hpi or **(B)** qPCR quantification of intracellular viral DNA at 1 hpi. Results are reported as fold change (FC) compared to the solvent control or siSc control conditions. Data for two technical replicates were averaged and error bars represent SEM.




FIG 7. P2Y2 receptor regulates levels of viral gene expression. HFFs were transfected with siP2Y2 or siSc as a control. After 24 h, cells were infected with TB40/E-GFP virus (MOI=3) or mock infected, and RNA samples were collected after various time intervals. Viral transcript levels were determined by qRT-PCR with GAPDH serving as an internal control. The data are shown as fold change compared to transcript levels at 2 hpi, calculated from two technical replicates. Error bars represent SEM.




FIG 8. P2Y2 receptor affects viral DNA accumulation. HFFs were transfected with siP2Y2 or siSc as a control. After 24 h, cells were infected with TB40/E-GFP virus (MOI=3) or mock infected. **(A) P2Y2 KD reduces intracellular vDNA.** At 96 hpi, total DNA was isolated from siSc- or siP2Y2-treated, HCMV-infected cells and the level of intracellular viral DNA was measured by qPCR. **(B) P2Y2 KD reduces extracellular vDNA.** At 96 hpi, total DNA was isolated from media of siSc- or siP2Y2-treated, HCMV-infected cells and the level of extracellular viral DNA was measured by qPCR. **(C) P2Y2 KD does not change the particle-to-infectious unit ratio.** The infectivity of virus in media collected at 96 hpi was titered and viral DNA was isolated from virions present in the media and quantified by qPCR to calculate a viral DNA copy number-to-infectious unit ratio. The results are depicted as fold change of this ratio. Data were averaged between two biological replicates and error bars represent SEM.




FIG 9. P2Y2 and UL37x1 work cooperatively to regulate Ca^{2+} in HCMV-infected cells. (A) Effects of kaempferol on Ca^{2+} levels in ADwt vs. ADsubUL37x1 infected cells. HFFs were infected with either ADwt or ADsubUL37x1 (MOI=1) or mock infected for 2 h and treated with kaempferol (50 μ M) or solvent control. Fluo-4 AM dye was applied at 20 hpi. (B) Effects of kaempferol on Ca^{2+} levels in wild type vs. UL37x1 overexpressing cells. Wild type HFF (WT HFF) cells or HFF cells overexpressing UL37x1 [pLVX-UL37x1(1) and pLVX-UL37x1(2)] were treated with kaempferol (50 μ M) or solvent control. Fluo-4 AM dye was applied 1 h later. Pictures of cells were taken using the Operetta High-Content Imaging System and the fluorescent signal was measured using ImageJ software based on 10 cells per experimental arm. Results are presented as the intensity of fluorescent signal in arbitrary units [AU]. (C) P2Y2 and UL37x1 protein in UL37x1 overexpressing cells. Protein samples were collected from two clones of MRC5 fibroblasts overexpressing UL37x1 [pLVX-UL37x1 (1) and pLVX-UL37x1 (2)] and control cells expressing GFP. Proteins were separated on SDS-PAGE and Western blot analysis was performed using antibodies recognizing P2Y2, UL37x1 and actin as a loading control.




FIG 10. P2Y2 overexpression in HCMV-infected HFFs is not regulated by viral UL37x1. HFFs were infected with either *ADwt* or *ADsubUL37x1* virus (MOI=3) or mock-infected for 2 h. Samples were collected at 24 hpi. (A) P2Y2, (B) UL54, (C) UL123 and (D) UL37x1 transcript levels were determined by qRT-PCR with GAPDH serving as an internal control. Results are shown as fold change compared to mock-infected or *Adwt*-infected controls. Data are averaged across three biological replicates and error bars represent SEM.