Corticosteroid and progesterone transactivation of mineralocorticoid receptors from Amur sturgeon and tropical gar

Akira Sugimoto¹, Kaori Oka¹, Rui Sato¹, Shinji Adachi²,
Michael E. Baker^{3,*}, Yoshinao Katsu^{1,4,*}

¹Graduate School of Life Science, Hokkaido University, Sapporo, Japan; ²Faculty of Fisheries Sciences, Hokkaido University, Hakodate, Japan; ³Department of Medicine, University of California, San Diego, CA, USA; ⁴Department of Biological Sciences, Hokkaido University, Sapporo, Japan

*Corresponding authors:

Michael E. Baker: mbaker@ucsd.edu

Yoshinao Katsu; ykatsu@sci.hokudai.ac.jp

Abstract

The response to a panel of steroids by the mineralocorticoid receptor (MR) from Amur sturgeon and tropical gar, two basal ray-finned fish, expressed in HEK293 cells was investigated. Half-maximal responses (EC50s) for transcriptional activation of sturgeon MR by 11-deoxycorticosterone, corticosterone, 11-deoxycortisol, cortisol and aldosterone, and progesterone were between 13 pM and 150 pM. For gar MR, EC50s were between 8 pM and 55 pM. Such low EC50s support physiological regulation by these steroids of the MR in sturgeon and gar. Companion studies with human MR and zebrafish MR found higher EC50s compared to EC50s for sturgeon and gar MR, with EC50s for zebrafish MR closer to gar and sturgeon MR than was human MR. For zebrafish MR, EC50s were between 75 pM and 740 pM; for human MR, EC50s were between 65 pM and 2 nM. In addition to progesterone, spironolactone and 19nor-progesterone were agonists for all three fish MRs, in contrast to their antagonist activity for human MR, which is hypothesized to involve serine-810 in human MR because all three steroids are agonists for a mutant human Ser810Leu-MR. Paradoxically, sturgeon, gar and zebrafish MRs contain a serine corresponding to serine-810 in human MR. Our data suggests alternative mechanism(s) for progesterone, spironolactone and 19nor-progesterone as MR agonists in these three ray-finned fishes and the need for caution in applying data for progesterone signaling in zebrafish to human physiology.

1 INTRODUCTION

2 The mineralocorticoid receptor (MR) is a transcription factor that belongs to the
3 nuclear receptor family, a diverse group of transcription factors that also includes
4 receptors for androgens (AR), estrogens (ER), glucocorticoids (GR) and progestins (PR),
5 and other small lipophilic ligands, such as thyroid hormone and retinoids [1-5]. The
6 MR and GR are descended from a common corticosteroid receptor (CR), which are
7 present in lampreys and hagfish [5-7]. Several corticosteroids (Figure 1), including
8 aldosterone (Aldo), cortisol (F), 11-deoxycortisol (S), corticosterone (B) and
9 11-deoxycorticosterone (DOC), as well as progesterone (Prog), are transcriptional
10 activators of Atlantic sea lamprey CR and hagfish CR [6]. Among these steroids, Aldo,
11 the main physiological activator of the MR in human and other terrestrial vertebrates
12 [8-11], had the lowest half-maximal response (EC50) for transcriptional activation of
13 the CR. This strong response to Aldo is surprising because Aldo is not found in either
14 lamprey or hagfish serum [6]. S, which along with DOC is present in Atlantic sea
15 lamprey serum, has been found to have mineralocorticoid activity in lamprey [12, 13].

16 Distinct MR and GR genes first appear in cartilaginous fishes (Chondrichthyes),
17 such as sharks, rays and skates [6, 14]. Carroll et al. [14] determined EC50s of several
18 corticosteroids for skate MR; EC50s were 70 pM for Aldo, 30 pM for DOC, 90 pM for
19 B, 1 nM for F and 22 nM for S. In teleosts, which comprise about 95% of known
20 ray-fish species (*Actinopterygii*), corticosteroid activation of the MR has been
21 investigated for cichlid [15], trout [16], carp [17], midshipman fish [18] and zebrafish
22 [19], with Aldo, F and DOC being the principal steroids that were studied. Although
23 Aldo has not been found in teleost fish [20], Aldo has a low EC50 for teleost MRs,
24 similar to that found for Aldo activation of lamprey CR and skate MR. DOC also has
25 a low EC50 for teleost MRs, and DOC has been proposed as mineralocorticoid in fish
26 [16, 21-25]. F also has been proposed to be ligand for teleost fish MR [22, 24, 25].
27 The response of the teleost MRs to B and S, which are found in fish [25, 26], has been
28 studied only in trout, in which the EC50s are 10 nM for B and 3.7 nM for S [16].
29 Interestingly, spironolactone (spiron) and Prog are agonists for trout MR [16], in
30 contrast to their antagonist activity for human MR [16, 27]. Spiron also is an agonist
31 for zebrafish MR [19]. Together, these studies indicate that several corticosteroid(s)
32 are potential transcriptional activators of teleost MRs [22, 25, 28, 29].

33 An important gap in our understanding of the evolution of selectivity of
34 ray-finned fish MRs for steroids is the absence of data on the MR in Chondrostei
35 (sturgeons, paddlefishes, reedfishes, bichirs) and Holostei (bowfins, gars), which
36 evolved before a fish-specific genome duplication occurred after the split of the

- 1 Acipenseriformes (sturgeons) and the Semionotiformes (gars) from the lineage leading
- 2 to teleost fish, but before the divergence of Osteoglossomorpha (Figure 2) [30-32].
- 3

- 4
- 5
- 6
- 7
- 8
- 9
- 10

Figure 1. Structures of potential steroid regulators of fish MR.

Aldo, the physiological ligand for terrestrial vertebrate MRs, is not found in fish [20]. F and DOC have been proposed to be mineralocorticoids in teleosts [22, 25]. S is a ligand for corticosteroid receptor in lamprey [12]. Progesterone is an antagonist for human MR [27].

Figure 2. Phylogenetic relationship of sturgeon and gar MRs to other vertebrates.

To investigate the relationship of sturgeon and gar to other fish, we constructed a phylogenetic tree of the steroid-binding domains on MRs in sturgeon, gar, selected teleosts, elasmobranchs and tetrapods. The phylogenetic tree was constructed using the maximum likelihood with JTT+G model with 1000 bootstrap replications, which are shown as percentages at the nodes of the tree.

Our interest in the evolution of steroid hormone action [4, 5, 33, 34] prompted us to investigate transcriptional activation of the MR from Amur sturgeon, *Acipenser schrenckii*, and tropical gar, *Atractosteus tropicus* by a broad panel of corticosteroids (Aldo, F, B, DOC, S), Prog, spiron and 19nor-progesterone (19norP), a steroid that has not previously been studied for activation of fish MR. To gain further insight into the evolution of steroid specificity, we compared our results with companion studies of zebrafish and human MRs. In agreement with studies of teleost MRs, we find that Aldo and DOC have the lowest EC50 (highest activity) for sturgeon and gar MRs. However, we also find that S, B, F, and Prog have low EC50s, consistent with these steroids also having a physiological role as ligands for these MRs. Spiron and 19norP also activated sturgeon and gar MRs. In comparison, zebrafish MR has a strong response to Aldo and DOC and a good response to B, F, S Prog, spiron and 19norP, while human MR has strong response to Aldo, DOC and B and a good response to F and S, and a weak response to Prog and 19norP and no response to spiron. The weak response to Prog and 19norP and absence of a response to spiron by human MR is in agreement with other studies [27, 35, 36]. The strong response to 19norP, Prog and spiron of sturgeon, gar and zebrafish MR is perplexing because the basis for the low response to to these steroids by human MR is thought to be due to the presence of Ser-810 on α -helix 5 [27, 37, 38]. 19norP, Prog and spiron are agonists for

human MR with Ser810Leu mutation [27, 37, 38]. Sturgeon, gar and zebrafish MRs contain a serine corresponding to serine-810 in human MR, suggesting the presence of an alternative mechanism for these steroids acting as MR agonists in these three ray-finned fishes, as well as the need to apply caution in interpreting data on Prog activity in zebrafish to human physiology.

MATERIALS AND METHODS

Animals and chemical reagents

Amur sturgeon and tropical gar were obtained as described previously [32]. All experimental procedures involving live fish followed the policies and guidelines of the Hokkaido University Animal Care and Use Committee. Aldosterone (Aldo) (CAS 52-39-1), corticosterone (B) (CAS 50-22-6), cortisol (F) (CAS 50-23-7), 11-deoxycortisol (S) (CAS 152-58-9), 11-deoxycorticosterone (DOC) (CAS 64-85-7), progesterone (Prog) (CAS 57-83-0), 19nor-progesterone (19norP) (CAS 472-54-8), spironolactone (Spiron) (CAS 52-01-7), 5 α -dihydrotestosterone (DHT) (CAS 521-18-6), and 17 β -estradiol (E2) (CAS 50-28-2) were purchased from Sigma-Aldrich. For the reporter gene assays, all hormones were dissolved in dimethyl-sulfoxide (DMSO) and the final concentration of DMSO in the culture medium did not exceed 0.1%.

Molecular cloning of mineralocorticoid receptors

Two conserved amino acid regions, GCHYGV and LYFAPD of vertebrate MRs were selected and degenerate oligonucleotides were used as primers for PCR. First-strand cDNA was synthesized from 2 μ g of total RNA isolated from the liver after amplification, and an additional primer set (CKVFFK and LYFAPD) was used for the second PCR. The amplified DNA fragments were subcloned with TA-cloning plasmid pGEM-T Easy vector, sequenced using a BigDye terminator Cycle Sequencing-kit with T7 and SP6 primers, and analyzed on the 3130 Genetic Analyzer (Applied Biosystems). The 5'- and 3'-ends of the mineralocorticoid receptor cDNAs were amplified by rapid amplification of the cDNA end (RACE) using a SMART RACE cDNA Amplification kit.

Database and sequence analysis

MRs for phylogenetic analysis were collected with Blast searches of GenBank. A phylogenetic tree for MRs was constructed by the Neighbor-Joining Method [39] after sequences were aligned by MUSCLE [40] using several fish, frog, alligator, chicken, rat, mouse, human MRs. Maximum likelihood (ML) analysis was conducted using the JTT+G model. Statistical confidence for each branch in the tree was evaluated by the bootstrap method [41] with 1000 replications. We used the MEGA5 program [42] for these analyses.

1

2 **Construction of plasmid vectors**

3 Full-coding regions of mineralocorticoid receptors were amplified by PCR with KOD
4 DNA polymerase. PCR products were gel-purified and ligated into pcDNA3.1 vector. Mouse
5 mammary tumor virus-long terminal repeat (MMTV-LTR) was amplified from pMSG vector by
6 PCR, and inserted into pGL3-basic vector containing the *Photinus pyralis* luciferase gene. All
7 constructs were verified by sequencing.

8

9 **Transactivation Assay**

10 Human embryonic kidney 293 (HEK293) cells were used in the reporter gene assay.
11 Transfection and reporter assays were carried out as described previously [33], except that we
12 used PEI-max as transfection reagent [43]. All transfections were performed at least three times,
13 employing triplicate sample points in each experiment. The values shown are mean $\pm$ SEM
14 from three separate experiments, and dose-response data and EC50 were analyzed using
15 GraphPad Prism.

16

17 **Statistical methods**

18 Results are presented as mean $\pm$ SE (SEM) from three separate experiments. All
19 multi-group comparisons were performed using one-way ANOVA followed by Bonferroni test.
20 Dose-response data and EC50 were analyzed using GraphPad Prism. $P < 0.05$ was considered
21 statistically significant.

22

23 **RESULTS**

24 **Isolation of mineralocorticoid receptors from sturgeon and gar**

25 We cloned sturgeon MR cDNA containing an open reading frame encoding 953 amino
26 acids (GenBank accession LC149818)], and gar MR cDNA containing an open reading frame
27 encoding 987 amino acids (GenBank accession LC149819). Sturgeon and gar MR sequences
28 can be divided into four domains (Figure 3). The overall amino acid identity between these two
29 MRs was 72%, with particularly high sequence identities for the DBD (100%) and LBD (89%)
30 (Figure 3). Comparison of sturgeon MR with five other species (human, chicken, alligator,
31 *Xenopus*, and zebrafish) revealed that sturgeon MR had identities of 44-36% in A/B
32 domains, 100-95% in DBDs, 67-47% in D domains, and 90-74% in LBDs (Figure 3).

33

34 **Phylogenetic analysis of ancient fish corticoid receptors**

35 To investigate the evolutionary position of gar and sturgeon MR in relationship to other
36 fish MRs and tetrapods, we collected MR sequences from several teleosts, skates and elephant

shark and selected terrestrial vertebrates. Consistent with the evolution of Acipenseriformes and Holostei, phylogenetic analysis places sturgeon and gar MRs close to the base of ray-finned fish (Figure 2).

Figure 3. Comparisons of functional domains in sturgeon, gar, zebrafish, *X. laevis*, alligator, chicken and human MRs.

Comparison of the domains in sturgeon MR, gar MR, zebrafish MR, *X. laevis* MR, alligator MR, and human MR. The functional A/B domain, C domain, D domain and E domain are schematically represented with the numbers of amino acid residues at each domain boundary indicated. The percentage of amino acid identity between domains is depicted. GenBank accession numbers are: LC149818 for sturgeon MR; LC149819 for gar MR; NM_001100403 for zebrafish MR; NM_001090605 for *Xenopus* MR; AB701406 for alligator MR; and NM_000901 for human MR.

Strong response to 3-keto-steroids by sturgeon and gar mineralocorticoid receptors

We examined steroid-inducible transcriptional activation of gar and sturgeon MRs using the MMTV-driven reporter construct [33, 44]. For comparison, we also examined transcriptional activation of human MR and zebrafish MR. At 1 nM, Aldo, B, S, DOC, F and Prog were strong inducers of luciferase activation by gar MR and sturgeon MR and by zebrafish

MR, with the exception of Prog which had a lower signal. These MRs show little stimulation by 1 nM DHT and E2 (Figure 4). At 1 nM, Aldo, B, DOC were strong transcriptional activators of human MR, which was activated to a lesser extent by S, and weakly activated by Prog (Figure 4).

Steroid Concentration = 1 nM

Figure 4. Ligand-specificities of fish and human MRs.

Full-length sturgeon MR (A), gar MR (B), zebrafish (C), and human MR (D) were expressed in HEK293 cells with an MMTV-luciferase reporter. Cells were treated with 10^{-8} M Aldo, F, B, S, DOC, Prog, 5 α -dihydrotestosterone (DHT), 17 β -estradiol (E2) or vehicle alone (DMSO). Results are expressed as means $\pm$ SEM, n=3. Y-axis indicates fold-activation compared to the activity of control vector with vehicle (DMSO) alone as 1.

The agonist activity of Prog for sturgeon, gar and zebrafish MRs and for trout MR [16] and the evidence that spiron is an agonist for trout [16] and zebrafish MRs [19] stimulated us to examine transcriptional activation of sturgeon and gar MRs by spiron, an agonist for human MR. We also investigated transcriptional of sturgeon, gar and zebrafish MRs by 19norP, which, like Prog and spiron, is an agonist for human Ser810Leu-MR [27]. As shown in Figure 5, spiron

1 and 19norP are agonists for sturgeon, gar and zebrafish MRs.

2

3

4

5 **Figure 5. Transcriptional activation by 19nor-progesterone and spironolactone of sturgeon,**
6 **gar, zebrafish and human MRs.**

7 Full-length sturgeon MR (A), gar MR (B), zebrafish MR (C), and human MR (D) were
8 expressed in HEK293 cells with an MMTV-luciferase reporter. Cells were treated with 1 nM or
9 10 nM Aldo, 19norP or spiron or vehicle alone (DMSO). Results are expressed as means $\pm$
10 SEM, N=3 and represent fold-activation compared to the control vector with vehicle.

11

12 We examined concentration-dependent activation of gar, sturgeon, zebrafish, and
13 human MRs by Aldo, F, B, DOC, S and Prog (Figure 6, Table 1). Both gar and sturgeon MRs
14 had similar low EC50s, which varied from 7.7 pM to 150 pM for these steroids. For each
15 steroid, the EC50s for gar MR were a little lower than for sturgeon MR.

1
2 **Figure 6. Concentration-dependent transcriptional activities of fish and human MRs.**
3 Concentration-response profiles of full-length sturgeon MR (A and B), gar MR (C and D),
4 zebrafish MR (E and F), and human MR (G and H) for various steroids. HEK293 cells were
5 transiently transfected with the MMTV-containing vector together with an MR expression vector.
6 Cells were incubated with increasing concentrations of Aldo, B, and S (A, C, E, and G) or Prog,

F, and DOC (B, D, F, and H) (10^{-13} to 10^{-6} M). Data are expressed as a ratio of steroid to vehicle (DMSO). Each column represents the mean of triplicate determinations, and vertical bars represent the mean $\pm$ SEM.

Table 1. EC50 activities for 3-keto-steroid transcriptional activation of sturgeon, gar, zebrafish and human MRs

	Aldo	DOC	B	S	F	Prog
Sturgeon MR	2.7×10^{-11}	1.3×10^{-11}	4.8×10^{-11}	8.2×10^{-11}	1.5×10^{-10}	7.0×10^{-11}
Gar MR	1.7×10^{-11}	7.7×10^{-12}	3.1×10^{-11}	2.6×10^{-11}	5.3×10^{-11}	1.8×10^{-11}
Zebrafish MR	8.8×10^{-11}	7.4×10^{-11}	3.3×10^{-10}	5.0×10^{-10}	5.9×10^{-10}	7.4×10^{-10}
Human MR	6.5×10^{-11}	1.7×10^{-10}	2.2×10^{-10}	2.0×10^{-9}	6.5×10^{-10}	-

In comparison, EC50s of Aldo, B and F were similar for zebrafish and human MR and a little higher than their EC50s for sturgeon and gar MR. EC50s of DOC, S and Prog for zebrafish MR were higher than their EC50s for sturgeon and gar MR, but lower than the EC50s for human MR. Prog had a lower, but still significant, maximal activation for zebrafish MR while 100 nM Prog had little activation of human MR. Overall all corticosteroids and Prog had EC50s that would be consistent with a physiological role in transcription of the MR in sturgeon, gar and zebrafish (Table 1, Figure 6).

In human MR, Ser-810 and Ala-773 are important in the low transcriptional activity of Prog. Prog, spiron and 19norP can activate human MR with selective mutations at either Ser-810 or Ala-773 [27, 37, 38]. For example, at 1 nM, prog, spiron and 19norP are agonist for a Ser810Leu mutant MR [27, 37, 38]. We extracted the sequence of helices 3-5, which contain Ser-810 and Ala-773, from sturgeon, gar and zebrafish MR (Figure 7) and other teleosts. All of these ray-finned fish contain a serine and alanine that aligns with Ser-810 and Ala-773 in human MR.

Figure 7. Alignment of vertebrate MRs to Serine-810 and Alanine-773 in helices 3-5 in

1 **human MR.**

2 Human Ser-810 and Ala-773 are conserved in ray-finned fish MRs. Skate MR, elephant shark
3 MR, lamprey CR and hagfish CR contain a methionine corresponding to human Ser-810.
4 Lamprey CR and hagfish CR contain a cysteine corresponding to Ala-773 in human MR.
5 Amino acids that are identical to amino acids in human MR are denoted by (-).
6

7 **DISCUSSION**

8 Several corticosteroids are physiological activators of the MR in cartilaginous fishes,
9 ray-finned fishes and tetrapods. Interestingly, Aldo, the mineralocorticoid for terrestrial
10 vertebrates, first appears in lungfish [45]. Nevertheless, Aldo is a potent activator of the
11 lamprey CR [6], which is ancestral to the MR [5-7, 11]. Interestingly, F, DOC, B and S and
12 Prog also are transcriptional activators of the CR in lamprey and hagfish [6], with only S, thus far,
13 found to have mineralocorticoid activity in lamprey [12, 13]. In skate, which has separate MR
14 and GR genes, Aldo, F, DOC and B are strong transcriptional activators of the MR [14]. F,
15 DOC, B, S and Prog are found in teleosts [26] and F and DOC have been proposed to be
16 transcriptional activators of teleost MRs [4, 15-19, 21, 22, 24, 25].

17 Absent, until now, was information about the response to corticosteroids of MRs in
18 sturgeon and gar, two basal fish that fill in the gap between elasmobranchs and teleosts (Figure 2).
19 Here we report that sturgeon MR and gar MR have EC50s below 1 nM for Aldo, F, DOC, B, S
20 and Prog. Interestingly, we find that zebrafish MR also has a similar strong response to these
21 corticosteroids and Prog. Moreover, spiron and 19norP are agonists for sturgeon, gar and
22 zebrafish MRs. This low selectivity for 3-keto-steroids (Figure 1) that can activate these fish
23 MRs resembles the response to these steroids by lamprey and hagfish CR [6] and skate MR [14].
24 Thus, this strong response of the MR to a broad panel of 3-keto-steroids was conserved after the
25 third whole-genome duplication at the base of the teleosts [30-32, 46].

26 In contrast, human MR is more selective for 3-keto-steroids with higher EC50s for S and
27 Prog. Our data showing weak activation by Prog and 19norP of human MR is in agreement
28 with other studies [35-37]. The strong response to Prog, spiron and 19norP of ray-finned fish
29 MR is interesting in the light of the report by Geller et al. [27] that human MR with a Ser810Leu
30 mutation was activated by 1 nM Prog, 19norP and spiron. Mutagenesis studies and structural
31 analyses of the MR-Leu810 mutant led to the hypothesis that Leu-810 on α -helix 5 has
32 stabilizing van der Waals contacts with Ala-773 on α -helix 3 [27, 37, 38] to explain the strong
33 transcriptional activation by Prog, 19norP and spiron. This serine and alanine are conserved in
34 and sturgeon and gar MRs, as well as in zebrafish MR (Figure 7) [4] and other teleost MRs
35 indicating that other mechanism(s) can lead to a strong response of sturgeon, gar and zebrafish
36 MR to Prog, 19norP and spiron. Activation by Prog of zebrafish MR is of concern because

zebrafish is an established model system for studying gene regulation in teleosts, as well as providing insights into human physiology [47]. Prog activation of zebrafish MR may confound data that focuses on activation of the PR. Prog may also be an agonist for the MR in medaka and other teleosts that have a serine and alanine that correspond to Ser-810 and Ala-773 in human MR.

Mechanisms for regulation of steroid activation of ray-finned fish MR

The strong response of zebrafish MR, as well as sturgeon and gar MRs, to five corticosteroids, Prog, 19norP and spiron requires one or more mechanisms to provide steroid-specific regulation of transcriptional activation of these ray-finned fish MRs. At this time, such mechanisms in gar, sturgeon and zebrafish MRs or other ray-finned fish MRs are poorly understood. Clues for possible mechanisms may be found from insights into regulation of mammalian MRs [4, 11, 48-53]. One possibility is an important mechanism in epithelial cells for regulating access of F and B to mammalian MR by tissue specific expression of 11 β -HSD2, which selectively converts F and B, respectively, to cortisone (E) and 11-dehydrocortisone (A), two inactive steroids. Aldo is inert to 11 β -HSD2, allowing Aldo to occupy the MR in epithelial cells in which 11 β -HSD2 inactivates F and B [11, 51, 52]. 11 β -HSD2 is found in ray-finned fish [54, 55], including sturgeon (Accession: ALE30175) and gar (Accession: XP_006641583). Expression of 11 β -HSD2 in MR-containing tissues provides a mechanism to exclude F and B from the MR. DOC, S and Prog, which have low EC₅₀s in gar, sturgeon and zebrafish, lack an 11 β -hydroxyl group and are inert to 11 β -HSD2.

Other regulatory mechanisms of the response of the MR to 3-keto-steroids include tissue-selective synthesis of 3-keto-steroids [5, 56-58], selective sequestration of 3-keto-steroids to plasma proteins [48, 53, 59], steroid-specific conformational changes that regulate MR binding of co-activators [60-64], effects of inter-domain interactions between the NTD and the LBD [19, 50, 64, 65] and post-translational modifications, such as phosphorylation, and SUMOylation [49, 66, 67].

Authors Contributions

A.S., K.O., R.S., and S.A. carried out the research. M.E.B. and Y.K. conceived and designed the experiments and wrote the paper. All authors gave final approval for publication. We have no competing interests.

Acknowledgments

We thank colleagues in our laboratories. We also thank Drs. Arlette Hernandez-Franyutti and Mari Carmen for providing gar tissues.

1

2 **Funding**

3 K.O. was supported by the Japan Society for the Promotion of Science (JSPS) Research
4 Fellowships for Young Scientists. This work was supported in part by Grants-in-Aid for
5 Scientific Research 23570067 and 26440159 (YK) from the Ministry of Education, Culture,
6 Sports, Science and Technology of Japan.

7

8 **References**

- 9 1 Lu, N. Z., Wardell, S. E., Burnstein, K. L., Defranco, D., Fuller, P. J., Giguere,
10 V., Hochberg, R. B., McKay, L., Renoir, J. M., Weigel, N. L., Wilson, E. M., McDonnell,
11 D. P. and Cidlowski, J. A. (2006) International Union of Pharmacology. LXV. The
12 pharmacology and classification of the nuclear receptor superfamily: glucocorticoid,
13 mineralocorticoid, progesterone, and androgen receptors. *Pharmacological reviews*. **58**,
14 782-797
- 15 2 Markov, G. V., Tavares, R., Dauphin-Villemant, C., Demeneix, B. A., Baker,
16 M. E. and Laudet, V. (2009) Independent elaboration of steroid hormone signaling
17 pathways in metazoans. *Proc Natl Acad Sci U S A*. **106**, 11913-11918
- 18 3 Bridgham, J. T., Eick, G. N., Larroux, C., Deshpande, K., Harms, M. J.,
19 Gauthier, M. E., Ortlund, E. A., Degnan, B. M. and Thornton, J. W. (2010) Protein
20 evolution by molecular tinkering: diversification of the nuclear receptor superfamily
21 from a ligand-dependent ancestor. *PLoS biology*. **8**
- 22 4 Baker, M. E., Funder, J. W. and Kattoula, S. R. (2013) Evolution of hormone
23 selectivity in glucocorticoid and mineralocorticoid receptors. *J Steroid Biochem Mol*
24 *Biol*. **137**, 57-70
- 25 5 Baker, M. E., Nelson, D. R. and Studer, R. A. (2015) Origin of the response to
26 adrenal and sex steroids: Roles of promiscuity and co-evolution of enzymes and steroid
27 receptors. *J Steroid Biochem Mol Biol*. **151**, 12-24
- 28 6 Bridgham, J. T., Carroll, S. M. and Thornton, J. W. (2006) Evolution of
29 hormone-receptor complexity by molecular exploitation. *Science*. **312**, 97-101
- 30 7 Thornton, J. W. (2001) Evolution of vertebrate steroid receptors from an
31 ancestral estrogen receptor by ligand exploitation and serial genome expansions. *Proc*
32 *Natl Acad Sci U S A*. **98**, 5671-5676
- 33 8 Funder, J. W. (2012) Aldosterone and mineralocorticoid receptors: a personal
34 reflection. *Molecular and cellular endocrinology*. **350**, 146-150
- 35 9 Hawkins, U. A., Gomez-Sanchez, E. P., Gomez-Sanchez, C. M. and
36 Gomez-Sanchez, C. E. (2012) The ubiquitous mineralocorticoid receptor: clinical

1 implications. *Curr Hypertens Rep.* **14**, 573-580

2 10 Martinerie, L., Munier, M., Le Menuet, D., Meduri, G., Viengchareun, S. and

3 Lombes, M. (2013) The mineralocorticoid signaling pathway throughout development:

4 expression, regulation and pathophysiological implications. *Biochimie.* **95**, 148-157

5 11 Rossier, B. C., Baker, M. E. and Studer, R. A. (2015) Epithelial sodium

6 transport and its control by aldosterone: the story of our internal environment revisited.

7 *Physiological reviews.* **95**, 297-340

8 12 Close, D. A., Yun, S. S., McCormick, S. D., Wildbill, A. J. and Li, W. (2010)

9 11-deoxycortisol is a corticosteroid hormone in the lamprey. *Proc Natl Acad Sci U S A.*

10 **107**, 13942-13947

11 13 Roberts, B. W., Didier, W., Rai, S., Johnson, N. S., Libants, S., Yun, S. S. and

12 Close, D. A. (2014) Regulation of a putative corticosteroid,

13 17,21-dihydroxypregn-4-ene,3,20-one, in sea lamprey, *Petromyzon marinus*. *Gen Comp*

14 *Endocrinol.* **196**, 17-25

15 14 Carroll, S. M., Bridgham, J. T. and Thornton, J. W. (2008) Evolution of

16 hormone signaling in elasmobranchs by exploitation of promiscuous receptors.

17 *Molecular biology and evolution.* **25**, 2643-2652

18 15 Greenwood, A. K., Butler, P. C., White, R. B., DeMarco, U., Pearce, D. and

19 Fernald, R. D. (2003) Multiple corticosteroid receptors in a teleost fish: distinct

20 sequences, expression patterns, and transcriptional activities. *Endocrinology.* **144**,

21 4226-4236

22 16 Sturm, A., Bury, N., Dengreville, L., Fagart, J., Flouriot, G., Rafestin-Oblin, M.

23 E. and Prunet, P. (2005) 11-deoxycorticosterone is a potent agonist of the rainbow trout

24 (*Oncorhynchus mykiss*) mineralocorticoid receptor. *Endocrinology.* **146**, 47-55

25 17 Stolte, E. H., de Mazon, A. F., Leon-Koosterziel, K. M., Jesiak, M., Bury, N.

26 R., Sturm, A., Savelkoul, H. F., van Kemenade, B. M. and Flik, G. (2008) Corticosteroid

27 receptors involved in stress regulation in common carp, *Cyprinus carpio*. *J Endocrinol.*

28 **198**, 403-417

29 18 Arterbery, A. S., Fergus, D. J., Fogarty, E. A., Mayberry, J., Deitcher, D. L.,

30 Lee Kraus, W. and Bass, A. H. (2011) Evolution of ligand specificity in vertebrate

31 corticosteroid receptors. *BMC Evol Biol.* **11**, 14

32 19 Pippal, J. B., Cheung, C. M., Yao, Y. Z., Brennan, F. E. and Fuller, P. J. (2011)

33 Characterization of the zebrafish (*Danio rerio*) mineralocorticoid receptor. *Molecular*

34 *and cellular endocrinology.* **332**, 58-66

35 20 Jiang, J. Q., Young, G., Kobayashi, T. and Nagahama, Y. (1998) Eel (*Anguilla*

36 *japonica*) testis 11 β -hydroxylase gene is expressed in interrenal tissue and its product

1 lacks aldosterone synthesizing activity. *Molecular and cellular endocrinology*. **146**,
2 207-211

3 21 Baker, M. E. (2003) Evolution of glucocorticoid and mineralocorticoid
4 responses: go fish. *Endocrinology*. **144**, 4223-4225

5 22 Bury, N. R. and Sturm, A. (2007) Evolution of the corticosteroid receptor
6 signalling pathway in fish. *Gen Comp Endocrinol*. **153**, 47-56

7 23 Baker, M. E., Chandsawangbhuwana, C. and Ollikainen, N. (2007) Structural
8 analysis of the evolution of steroid specificity in the mineralocorticoid and
9 glucocorticoid receptors. *BMC Evol Biol*. **7**, 24

10 24 Sakamoto, T., Mori, C., Minami, S., Takahashi, H., Abe, T., Ojima, D., Ogoshi,
11 M. and Sakamoto, H. (2011) Corticosteroids stimulate the amphibious behavior in
12 mudskipper: potential role of mineralocorticoid receptors in teleost fish. *Physiol Behav*.
13 **104**, 923-928

14 25 Takahashi, H. and Sakamoto, T. (2013) The role of 'mineralocorticoids' in
15 teleost fish: relative importance of glucocorticoid signaling in the osmoregulation and
16 'central' actions of mineralocorticoid receptor. *Gen Comp Endocrinol*. **181**, 223-228

17 26 Milla, S., Wang, N., Mandiki, S. N. and Kestemont, P. (2009) Corticosteroids:
18 Friends or foes of teleost fish reproduction? *Comparative biochemistry and physiology*.
19 Part A, *Molecular & integrative physiology*. **153**, 242-251

20 27 Geller, D. S., Farhi, A., Pinkerton, N., Fradley, M., Moritz, M., Spitzer, A.,
21 Meinke, G., Tsai, F. T., Sigler, P. B. and Lifton, R. P. (2000) Activating
22 mineralocorticoid receptor mutation in hypertension exacerbated by pregnancy. *Science*.
23 **289**, 119-123

24 28 Gilmour, K. M. (2005) Mineralocorticoid receptors and hormones: fishing for
25 answers. *Endocrinology*. **146**, 44-46

26 29 Prunet, P., Sturm, A. and Milla, S. (2006) Multiple corticosteroid receptors in
27 fish: from old ideas to new concepts. *Gen Comp Endocrinol*. **147**, 17-23

28 30 Jaillon, O., Aury, J. M., Brunet, F., Petit, J. L., Stange-Thomann, N., Mauceli,
29 E., Bouneau, L., Fischer, C., Ozouf-Costaz, C., Bernot, A., Nicaud, S., Jaffe, D., Fisher,
30 S., Lutfalla, G., Dossat, C., Segurens, B., Dasilva, C., Salanoubat, M., Levy, M., Boudet,
31 N., Castellano, S., Anthouard, V., Jubin, C., Castelli, V., Katinka, M., Vacherie, B.,
32 Biemont, C., Skalli, Z., Cattolico, L., Poulain, J., De Berardinis, V., Cruaud, C., Duprat,
33 S., Brottier, P., Coutanceau, J. P., Gouzy, J., Parra, G., Lardier, G., Chapple, C.,
34 McKernan, K. J., McEwan, P., Bosak, S., Kellis, M., Volff, J. N., Guigo, R., Zody, M. C.,
35 Mesirov, J., Lindblad-Toh, K., Birren, B., Nusbaum, C., Kahn, D., Robinson-Rechavi,
36 M., Laudet, V., Schachter, V., Quetier, F., Saurin, W., Scarpelli, C., Wincker, P., Lander,

- 1 E. S., Weissenbach, J. and Roest Crollius, H. (2004) Genome duplication in the teleost
2 fish *Tetraodon nigroviridis* reveals the early vertebrate proto-karyotype. *Nature*. **431**,
3 946-957
- 4 31 Kasahara, M., Naruse, K., Sasaki, S., Nakatani, Y., Qu, W., Ahsan, B., Yamada,
5 T., Nagayasu, Y., Doi, K., Kasai, Y., Jindo, T., Kobayashi, D., Shimada, A., Toyoda, A.,
6 Kuroki, Y., Fujiyama, A., Sasaki, T., Shimizu, A., Asakawa, S., Shimizu, N., Hashimoto,
7 S., Yang, J., Lee, Y., Matsushima, K., Sugano, S., Sakaizumi, M., Narita, T., Ohishi, K.,
8 Haga, S., Ohta, F., Nomoto, H., Nogata, K., Morishita, T., Endo, T., Shin, I. T., Takeda,
9 H., Morishita, S. and Kohara, Y. (2007) The medaka draft genome and insights into
10 vertebrate genome evolution. *Nature*. **447**, 714-719
- 11 32 Katsu, Y., Kohno, S., Hyodo, S., Ijiri, S., Adachi, S., Hara, A., Guillelte, L. J.,
12 Jr. and Iguchi, T. (2008) Molecular cloning, characterization, and evolutionary analysis
13 of estrogen receptors from phylogenetically ancient fish. *Endocrinology*. **149**,
14 6300-6310
- 15 33 Oka, K., Hoang, A., Okada, D., Iguchi, T., Baker, M. E. and Katsu, Y. (2015)
16 Allosteric role of the amino-terminal A/B domain on corticosteroid transactivation of
17 gar and human glucocorticoid receptors. *J Steroid Biochem Mol Biol*. **154**, 112-119
- 18 34 Katsu, Y., Kohno, S., Oka, K. and Baker, M. E. (2016) Evolution of
19 corticosteroid specificity for human, chicken, alligator and frog glucocorticoid receptors.
20 *Steroids*. **113**, 38-45
- 21 35 Rupprecht, R., Reul, J. M., van Steensel, B., Spengler, D., Soder, M., Berning,
22 B., Holsboer, F. and Damm, K. (1993) Pharmacological and functional characterization
23 of human mineralocorticoid and glucocorticoid receptor ligands. *Eur J Pharmacol*. **247**,
24 145-154
- 25 36 Fagart, J., Wurtz, J. M., Souque, A., Hellal-Levy, C., Moras, D. and
26 Rafestin-Oblin, M. E. (1998) Antagonism in the human mineralocorticoid receptor. *The*
27 *EMBO journal*. **17**, 3317-3325
- 28 37 Bledsoe, R. K., Madauss, K. P., Holt, J. A., Apolito, C. J., Lambert, M. H.,
29 Pearce, K. H., Stanley, T. B., Stewart, E. L., Trump, R. P., Willson, T. M. and Williams,
30 S. P. (2005) A ligand-mediated hydrogen bond network required for the activation of the
31 mineralocorticoid receptor. *The Journal of biological chemistry*. **280**, 31283-31293
- 32 38 Fagart, J., Huyet, J., Pinon, G. M., Rochel, M., Mayer, C. and Rafestin-Oblin,
33 M. E. (2005) Crystal structure of a mutant mineralocorticoid receptor responsible for
34 hypertension. *Nature structural & molecular biology*. **12**, 554-555
- 35 39 Saitou, N. and Nei, M. (1987) The neighbor-joining method: a new method
36 for reconstructing phylogenetic trees. *Molecular biology and evolution*. **4**, 406-425

1 40 Edgar, R. C. (2004) MUSCLE: multiple sequence alignment with high
2 accuracy and high throughput. *Nucleic acids research*. **32**, 1792-1797

3 41 Felsenstein, J. (1985) Confidence-Limits on Phylogenies - an Approach Using
4 the Bootstrap. *Evolution*. **39**, 783-791

5 42 Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S.
6 (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood,
7 evolutionary distance, and maximum parsimony methods. *Molecular biology and*
8 *evolution*. **28**, 2731-2739

9 43 Kadlecova, Z., Nallet, S., Hacker, D. L., Baldi, L., Klok, H. A. and Wurm, F.
10 M. (2012) Poly(ethyleneimine)-mediated large-scale transient gene expression:
11 influence of molecular weight, polydispersity and N-propionyl groups. *Macromol*
12 *Biosci*. **12**, 628-636

13 44 Oka, K., Kohno, S., Urushitani, H., Guillelte, L. J., Jr., Ohta, Y., Iguchi, T. and
14 Katsu, Y. (2013) Molecular cloning and characterization of the corticoid receptors from
15 the American alligator. *Molecular and cellular endocrinology*. **365**, 153-161

16 45 Joss, J. M. P., Arnoldreed, D. E. and Balment, R. J. (1994) The Steroidogenic
17 Response to Angiotensin-II in the Australian Lungfish, *Neoceratodus-Forsteri*. *J Comp*
18 *Physiol B*. **164**, 378-382

19 46 Inoue, J., Sato, Y., Sinclair, R., Tsukamoto, K. and Nishida, M. (2015) Rapid
20 genome reshaping by multiple-gene loss after whole-genome duplication in teleost fish
21 suggested by mathematical modeling. *Proc Natl Acad Sci U S A*. **112**, 14918-14923

22 47 Tokarz, J., Moller, G., de Angelis, M. H. and Adamski, J. (2013) Zebrafish and
23 steroids: what do we know and what do we need to know? *J Steroid Biochem Mol Biol*.
24 **137**, 165-173

25 48 Funder, J. and Myles, K. (1996) Exclusion of corticosterone from epithelial
26 mineralocorticoid receptors is insufficient for selectivity of aldosterone action: in vivo
27 binding studies. *Endocrinology*. **137**, 5264-5268

28 49 Pascual-Le Tallec, L. and Lombes, M. (2005) The mineralocorticoid receptor:
29 a journey exploring its diversity and specificity of action. *Mol Endocrinol*. **19**,
30 2211-2221

31 50 Fuller, P. J., Yao, Y., Yang, J. and Young, M. J. (2012) Mechanisms of ligand
32 specificity of the mineralocorticoid receptor. *J Endocrinol*. **213**, 15-24

33 51 Odermatt, A. and Kratschmar, D. V. (2012) Tissue-specific modulation of
34 mineralocorticoid receptor function by 11beta-hydroxysteroid dehydrogenases: an
35 overview. *Molecular and cellular endocrinology*. **350**, 168-186

36 52 Chapman, K., Holmes, M. and Seckl, J. (2013) 11beta-hydroxysteroid

1 dehydrogenases: intracellular gate-keepers of tissue glucocorticoid action. *Physiological*
2 *reviews.* **93**, 1139-1206

3 53 Sivukhina, E. V. and Jirikowski, G. F. (2014) Adrenal steroids in the brain:
4 role of the intrinsic expression of corticosteroid-binding globulin (CBG) in the stress
5 response. *Steroids.* **81**, 70-73

6 54 Kusakabe, M., Nakamura, I. and Young, G. (2003) 11beta-hydroxysteroid
7 dehydrogenase complementary deoxyribonucleic acid in rainbow trout: cloning, sites of
8 expression, and seasonal changes in gonads. *Endocrinology.* **144**, 2534-2545

9 55 Baker, M. E. (2010) 11Beta-hydroxysteroid dehydrogenase-type 2 evolved
10 from an ancestral 17beta-hydroxysteroid dehydrogenase-type 2. *Biochemical and*
11 *biophysical research communications.* **399**, 215-220

12 56 Nebert, D. W. and Russell, D. W. (2002) Clinical importance of the
13 cytochromes P450. *Lancet.* **360**, 1155-1162

14 57 Payne, A. H. and Hales, D. B. (2004) Overview of steroidogenic enzymes in
15 the pathway from cholesterol to active steroid hormones. *Endocr Rev.* **25**, 947-970

16 58 Taves, M. D., Plumb, A. W., Sandkam, B. A., Ma, C., Van Der Gugten, J. G.,
17 Holmes, D. T., Close, D. A., Abraham, N. and Soma, K. K. (2015) Steroid profiling
18 reveals widespread local regulation of glucocorticoid levels during mouse development.
19 *Endocrinology.* **156**, 511-522

20 59 Breuner, C. W. and Orchinik, M. (2002) Plasma binding proteins as mediators
21 of corticosteroid action in vertebrates. *J Endocrinol.* **175**, 99-112

22 60 Bledsoe, R. K., Montana, V. G., Stanley, T. B., Delves, C. J., Apolito, C. J.,
23 McKee, D. D., Consler, T. G., Parks, D. J., Stewart, E. L., Willson, T. M., Lambert, M.
24 H., Moore, J. T., Pearce, K. H. and Xu, H. E. (2002) Crystal structure of the
25 glucocorticoid receptor ligand binding domain reveals a novel mode of receptor
26 dimerization and coactivator recognition. *Cell.* **110**, 93-105

27 61 Li, Y., Suino, K., Daugherty, J. and Xu, H. E. (2005) Structural and
28 biochemical mechanisms for the specificity of hormone binding and coactivator
29 assembly by mineralocorticoid receptor. *Mol Cell.* **19**, 367-380

30 62 Hultman, M. L., Krasnoperova, N. V., Li, S., Du, S., Xia, C., Dietz, J. D., Lala,
31 D. S., Welsch, D. J. and Hu, X. (2005) The ligand-dependent interaction of
32 mineralocorticoid receptor with coactivator and corepressor peptides suggests multiple
33 activation mechanisms. *Mol Endocrinol.* **19**, 1460-1473

34 63 Khan, S. H., Awasthi, S., Guo, C., Goswami, D., Ling, J., Griffin, P. R.,
35 Simons, S. S., Jr. and Kumar, R. (2012) Binding of the N-terminal region of coactivator
36 TIF2 to the intrinsically disordered AF1 domain of the glucocorticoid receptor is

1 accompanied by conformational reorganizations. The Journal of biological chemistry.
2 **287**, 44546-44560
3 64 Fuller, P. J. (2015) Novel interactions of the mineralocorticoid receptor.
4 Molecular and cellular endocrinology. **408**, 33-37
5 65 Pippal, J. B., Yao, Y., Rogerson, F. M. and Fuller, P. J. (2009) Structural and
6 functional characterization of the interdomain interaction in the mineralocorticoid
7 receptor. Mol Endocrinol. **23**, 1360-1370
8 66 Bhargava, A. and Pearce, D. (2004) Mechanisms of mineralocorticoid action:
9 determinants of receptor specificity and actions of regulated gene products. Trends
10 Endocrinol Metab. **15**, 147-153
11 67 Faresse, N. (2014) Post-translational modifications of the mineralocorticoid
12 receptor: How to dress the receptor according to the circumstances? J Steroid Biochem
13 Mol Biol. **143**, 334-342
14