Better together: a transboundary approach to brown bear monitoring in the Pyrenees

Blaise Piédallu (1) – Corresponding author

Pierre-Yves Quenette (2)

Ivan Afonso Jordana (3)

Nicolas Bombillon (2)

Adrienne Gastineau (2)

Ramón Jato (4)

Christian Miquel (5)

Pablo Muñoz (6)

Santiago Palazón (7)

Jordi Solà de la Torre (8)

Olivier Gimenez (1)

(1) CEFÉ, CNRSUMR 5175, Université de Montpellier, Université Paul-Valéry Montpellier, EPHE, 1919 Route de Mende, 34293 Montpellier Cedex 5, France

(2) Office National de la Chasse et de la Faune Sauvage, CNERA PAD-Equipe Ours, Impasse de la Chapelle, 31800 Villeneuve-de-Rivière, France

(3) Medi Ambient, Conselh Generau d'Aran, Plaça dera Libertat, 16, 25530 Vielha, Spain

(4) Departamento de Producción y Asistencias Técnicas, Sociedad Aragonesa de Gestión Agroambiental, S. L. U., Ctra HUV-Fornillos, 22195 Fornillos de Apiés, Huesca, Spain

(5) Laboratoire d'Ecologie Alpine, CNRS UMR 5553, Université Joseph Fourier, BP 53, F-38041 Grenoble Cedex 9, France

- 23 (6) Jefe de la Sección de Hábitats, Gobierno de Navarra
- 24 (7) Fauna and Flora Service, Department of Territory and Sustainability. Dr. Roux, 80. 08017
- 25 Barcelona, Spain
- 26 (8) Unitat de Fauna, Departament de Medi Ambient, C/ Prat de la Creu 62-64, AD500 Andorra
- 27 la Vella, Principality of Andorra
- 28 Corresponding author personal information :
- 29 Centre d'Ecologie Fonctionnelle et Evolutive
- 30 1919 Route de Mende
- 31 34293 Montpellier Cedex 5
- 32 blaise.piedallu@cefe.cnrs.fr
- 33 +334 67 61 33 14

34 Abstract

35 Human administrative borders have no effect on wild animals, and the vast home ranges of large carnivores
 36 often cause them to live simultaneously on the territory of two or more countries or jurisdictions with
 37 different management policies. Here, we investigate the importance of transboundary population monitoring
 38 using as a case study the Pyrenean brown bear population (*Ursus arctos*) that lives in France, Andorra and
 39 Spain. Using capture-recapture models and the Pollock's robust design, we estimated abundance and
 40 demographic parameters using data collected separately in France and Spain and a dataset gathered from
 41 joint monitoring on both sides of the border. As expected, the abundance estimates from French (from 11
 42 bears in 2008 to 13 in 2014) or Spanish (from 4 bears in 2008 to 9 in 2014) data only were lower than
 43 abundance obtained from both sides of the border (from 11 in 2008 to 18 in 2014). The joint monitoring
 44 dataset also highlighted the importance of individual detection heterogeneity that, if ignored, would lead to
 45 underestimation. Our results reinforce the importance of transboundary cooperation when dealing with
 46 animal populations with territory spanning two or more administrative jurisdictions for collecting reliable
 47 scientific data and providing relevant abundance estimation to take sound management decisions.

48 *Keywords: capture-recapture, elusive species, robust design, transboundary monitoring, Ursus arctos*

49 Introduction

50 Large carnivore populations in Europe are recovering from years of decrease after having been highly
 51 threatened or locally extinct, leading to larger population sizes and wider distributions (Chapron et al. 2014).
 52 These populations have large home ranges despite interspecific (Gittleman and Harvey 1982) or intraspecific
 53 (Mattisson et al. 2013; Nilsen et al. 2005) variations, and their recovery increases the likelihood of crossing
 54 the administrative border between two countries, or between two distinct jurisdictions within the same
 55 country. Therefore, the same large carnivore population may overlap several territories where the inhabitants
 56 have different or conflicting opinions on their monitoring or management (Bischof and Swenson 2012; Bull
 57 et al. 2009; Piédallu et al. 2016).

58 The conservation status of a population is commonly estimated based on its abundance and distribution,
 59 which in turn may help to assess its viability and to predict its range in the future. However, while
 60 transboundary large carnivore populations are extremely common, transboundary conservation policies are

still rarely implemented (Gervasi et al. 2015; Linnell and Boitani 2012; Trouwborst 2010). Moreover, population monitoring is usually performed independently between the countries on one side and the other of the boundary. These countries in turn perform abundance estimation based solely on the data gathered on their own territory, leading to a discrepancy between management policy and ecological reality (Linnell and Boitani 2012; Trouwborst 2010). This approach might cause significant errors when trying to assess the abundance of a species, due to the possibility for a single animal to be seen across different territories (Bischof et al. 2016).

Here, we aimed at illustrating the value of collaborative monitoring at a transboundary scale, using brown bears (*Ursus arctos*) in the Pyrénées as a case study. Brown bears are widely distributed in the Northern hemisphere, being spread over North America (Waits et al. 1998), Europe (Swenson et al. 2011), and Asia (Hirata et al. 2013). The species as a whole is not endangered. However, some populations can be both small and isolated from others, leading to an important risk of local extinction (Linnell et al. 2008; Swenson et al. 2011). Among those endangered populations is the Pyrenean brown bear population, which resides in the mountains between Southwestern France and Northeastern Spain. Despite being reinforced by the reintroduction of individuals from Slovenia in 1996-97 (3 bears) and 2006 (5 bears) with another one scheduled for 2016 (1 bear), its viability is still considered to be low (Chapron et al. 2010) due to its small size, its fragmentation in different cores, and heavy inbreeding that might negatively impact its growth in a medium- or long-term future. Moreover, there are conflicting attitudes among local inhabitants over bear presence (Piédallu et al. 2016).

Combining data collected through camera trapping, systematic search-effort protocols and opportunistic observations in Spain and France, we estimated abundance and demographic parameters for the Pyrenean brown bear population using Pollock's robust design capture-recapture models (Kendall et al. 1997). We reproduced the same analyses mimicking a lack of collaboration between the two countries and compared the results. Overall, we demonstrate the benefits of cooperation across the border in the monitoring and management of a large carnivore population.

Material & Methods

1. Study area and bear population

This study was performed in the Pyrenees Mountains, on both sides of the border, which included Southwestern France, the North of the Spanish Autonomous communities of Catalonia, Aragon and Navarre, and the principality of Andorra (Figure 1). Bears that live in the Pyrenees descend from individuals that were translocated here from Slovenia in 1996-97 (2 females and 1 male) and 2006 (4 females and 1 male), although a hybrid between a Slovenian bear and a now-deceased Pyrenean female is still alive. The relocations were performed to save the population that was in critical danger of extinction in 1995, with only 5 individuals remaining, including a single adult female. The current population is split in two unconnected population cores (Camarra et al. 2015): one in the Western Pyrenees, over France, Aragon and Navarre, and one in Central-Eastern Pyrenees, over France, Catalonia, Aragon and Andorra.

2. Data collection and monitoring

Systematic monitoring was performed in France, Catalonia and Andorra from 2008 to 2014, consisting in fixed itineraries along which agents of governmental wildlife looked for bear tracks with help on the French side of the Pyrenees from volunteers of the Brown Bear Network, whose members include 135 professional members (from various agencies) and 228 amateur members (who are not affiliated to any agency). In Catalonia, two teams of 10 professional members handle systematic monitoring with help from Forestry Agents. Visits of these itineraries were repeated at regular time intervals between April and November, with at least a monthly visit (Figure 1). Due to the very low number of bears on their territories, there was no systematic monitoring in Aragon and Navarre. In all areas, data was also obtained through opportunistic means, which included all sightings or tracks found outside of systematic monitoring and validated by the local agencies, or damages caused on livestock (sheep in particular) or beehives. Since 2008, both countries have worked to standardize their monitoring protocols in the framework decided by the Transborder Monitoring Group for Bears in Pyrenees (GSTOP).

The most common bear tracks included hair samples (with 4-5 hair traps being scattered along the itineraries to improve the chances of getting a sample), scats, footprints, claw marks, photographs, films, attacks on livestock, sightings and opened ant-hills. Hair traps were added on trees along the itineraries to increase the chances to obtain hair samples. We discarded the tracks that did not allow the identification of an individual, mostly due to a lack of a good enough genetic sample. We also paid a particular attention to estimate the

interval in which the bear might have left the track (see Supplementary materials). Genetic samples (hairs and faeces) were analyzed using a Polymerase Chain Reaction (PCR) multitube approach with four repeats for each sample to avoid the risk of misidentification due to the low DNA quantity of some hair samples (Miquel et al. 2006; Taberlet et al. 1997). 13 microsatellites markers and 1 marker for sex were targeted by the PCR in order to identify the bear individuals. The results were then aggregated between all sources.

We assigned to each individual its age class at first capture (Cub: up to 1 year old, Juvenile: up to 3 years old, Adult: 4 years old and more) and its gender. The population was assumed geographically closed, i.e. no emigration or immigration between this population and another one outside the Pyrenees.

3. Abundance estimation

We used the Pollock's robust design (Kendall et al. 1997) to estimate abundance while accounting for imperfect detection of individuals and temporary emigration. The robust design approach uses repeated captures in a short timeframe, called secondary occasions, within a single time step (a so-called primary occasion). The population is considered to be closed (without demographic processes such as births, deaths, immigrations and emigrations) between two consecutive secondary occasions, hence allowing the robust estimation of abundance corrected for imperfect detection, and open (with births, deaths, emigrations and immigrations) between two consecutive primary occasions. In this study, we used years as primary occasions of capture (7 in total, from 2008 to 2014) and the months from May to September as secondary occasions (5 in total). We chose these secondary occasions because no births occur in this time interval, and we did not include the first months of the year because very young cubs have a high risk of dying.

We built three capture-recapture datasets: one for French data (including Andorra), one for Spanish data (Catalonia, Aragon, Navarre), and one for the Combined (France and Spain) data. We tested three possible types of temporary emigration: no emigration, a random emigration (where the probability of being a temporary emigrant only depends on the status of the individual at a given capture occasion) and a Markovian emigration (where the probability of being a temporary emigrant depends on the status of the individual at a given time step and at the previous time step). We tested 4 possible effects on survival: no effect, a sex effect, an age effect, and an additive effect of sex and age. Finally, 6 possible effects on detection were considered: no effect, a sex effect, a time effect (where the probability of detection changed

between primary occasions – years, in our study), a mixture effect (with two distinct classes or individuals to account for detection heterogeneity, e.g. Cubaynes et al. (2010)), and the additive effects of sex and time and sex and mixture. In total, we tested 72 different models (Supplementary materials).

We used the Akaike Information Criterion corrected for small sample size (AICc) to perform model selection (Burnham and Anderson 2002). For each dataset (France, Spain, Combined), we kept the models with a AICc weight > 0.01 . We also model-averaged parameter estimates to account for uncertainty in selecting one single ‘best’ model. To obtain 95% confidence intervals on abundance, we used a non-parametric bootstrap in which individual capture histories from each dataset were resampled with replacement a hundred times (Buckland et al. 1997).

We used the program MARK (White and Burnham 1999) called from software R (RCoreTeam 2013) with the ‘RMark’ package (Laake 2013).

Results

1. Descriptive statistics

7517 bear tracks were gathered in the Pyrenees between 2008 and 2014. The number of tracks that were usable in a robust design framework for the French, Spanish and Combined dataset by both being accurate enough and collected between May and September (see Supplementary data) is shown in Table 1, and amounted for 15 to 17% of the total tracks.

2. Model selection with the French dataset

The models that were selected for the French dataset mostly included a mixture effect on detection (Table 2A), whether alone or with an additive effect of sex. Survival was linked either to no effect, or to an effect of sex. However, no specific type of emigration (none, random or Markovian) was preferentially selected. The three best models (AICc weight > 0.1) included a sex effect along with individual heterogeneity on detection, and no effect on survival. No time effect on detection was found in the best models, and only the 12th best had an age effect on survival.

3. Model selection with the Spanish dataset

All models selected for the Spanish dataset included emigration, whether it was random or Markovian (Table 2B), with the four best-ranked models having a random emigration. Models without emigration had a significantly lower weight. In contrast with the French dataset, the heterogeneity of detection (mixture effect) was seldom kept in the best models. Instead, most of the best models either had no effect or a sex effect for both survival and detection. The four best models (AICc weight >0.08) had a random emigration, and either no effect of a sex effect on both survival and detection.

4. Model selection with the Combined dataset

Like in the French dataset, all the best models for the Combined dataset had a mixture effect on detection, whether it was alone or with an additive effect of sex (Table 2C). No clear pattern emerged for emigration, although the models with Markovian emigration tended to have a lower weight. The results were similarly contrasted for the effects on survival, with sex being kept in 8 of the 15 models and age in 5 of 15.

5. Abundance estimates

In Spain, population increased to stabilize around 10 individuals, but showed a steep fall in 2013 before recovering in 2014 (Figure 2), which may indicate individuals that were temporarily unavailable. The French dataset pictured a very small increase between 2008 and 2014. The Combined dataset showed a trend similar to the Spanish dataset, with an increase between 2008 and 2014 and a temporary fall in 2013 mostly due to the lack of individuals detected on the Spanish side of the Pyrenees.

6. Demographic parameter estimates

The survival estimates were higher in France than in Spain (Table 3). There was a notable difference between male and female survival in the Combined dataset ($F = 0.81 \pm 0.051$, $M = 0.933 \pm 0.041$). In the France and Combined datasets, two classes of individuals were identified detection-wise, with one class being easily detected (~0.7-0.8) and one being harder to detect (~0.3-0.4). In both cases, the detection probability of females ended up being slightly inferior to the detection probability of males.

In the Spain dataset, emigration was much higher than in the other datasets. This might be explained by the very few individuals found in 2013 that were probably treated by the models as temporary emigrating. Moreover, all models selected for the Spain dataset included a random or Markovian emigration, while the

193 France and Combined datasets also selected models with no emigration, which might have lowered the
194 estimated values when averaged.

195 **Discussion**

196 1. Abundance and heterogeneity in the detection process

197 The results for the French and Combined datasets highlighted two classes of individuals with significantly
198 different detection. A previous study on wolves highlighted the importance of individual detection
199 heterogeneity (IDH) when estimating abundance of large carnivore populations (Cubaynes et al. 2010). IDH
200 in the Pyrenean brown bears might stem from intraspecific home range disparities (McLoughlin et al. 2000)
201 making it more likely to find tracks of individuals who move a lot. The individual that was detected the most
202 often, Pyros, is also the male with the highest reproductive success, and the three most commonly seen males
203 (Pyros, Balou and Nere) were the three oldest living males in 2014. Another factor that might cause IDH is
204 the efficiency of human agents when looking for bear tracks. Some Pyrenean bears display a stable spatial
205 behavior over the years (Camarra et al. 2015), making their movements predictable in time. The personality
206 differences displayed by bears (Fagen and Fagen 1996) might allow the agents to become better at finding
207 tracks.

208 2. Sex and age effect on survival

209 A result that was consistent in all datasets was the lack of an age effect on survival. This is unexpected since
210 young bears generally suffer from higher mortality rates (Bunnell and Tait 1985). However, because our
211 analysis only included data ranging from May to September, we excluded the first four or five months of life
212 for the bear cubs, during which their survival rate is at its lowest, meaning that bear cub deaths might have
213 simply gone unnoticed due to the absence of the cubs from the dataset.

214 The presence of an effect of sex on survival, with higher mortality rates for females, seemed to contradict a
215 previous study performed on grizzly bears where mortality was higher among males (McLellan et al. 1999).
216 However, in this study, hunting explained the bias towards male mortality, since death from natural causes
217 was more common among females. Because bear hunting is banned in the Pyrenees, the results between the
218 two studies are actually consistent.

3. Estimates of the demographic parameters

Female bears survival in the Combined dataset was consistent with previous studies performed in other populations (De Barba et al. 2010; Garshelis et al. 2005; Wielgus and Bunnell 1994). Male bears survival in the same dataset was consistent with a previous study on a population without hunting pressure (De Barba et al. 2010). The emigration rate was significantly higher when using only the Spanish dataset compared to the French and Combined datasets, which could indicate that bears travel more from the Spanish to the French side of the Pyrenees. This important emigration rate on the Spanish side might also result from demographic stochasticity, stemming from the small size of the population and the very low number of individuals identified in Spain in 2013.

4. Robust design as a tool to estimate population size for cryptic species

Rare populations can either be defined by the low number of individuals, or by the fact that the population belongs to elusive, hard-to-observe species with large home ranges and low density (McDonald 2004). The Pyrenean brown bear population falls under both cases, and as such estimating its demographic parameters is made even more difficult. We confirmed however that robust design capture-recapture models allowed estimating several population parameters (Smith et al. 2013). In our case, the robust design study was combined with the use of multiple data sources (Boulanger et al. 2008; Gervasi et al. 2012) to obtain enough captures so we could build capture-recapture datasets with a sufficient number of observations to be exploited.

Abundance estimates were different from those calculated yearly by the GSTOP, which found a significant increase in the minimal population size between 2008 and 2014, going from 17 to 31 individuals (Camarra et al. 2015). This discrepancy can be explained by the method used by the agencies, which use both a Minimal Kept Population Size (EMR) and a Minimal Detected Population Size (EMD). Every year, the EMD can be used to correct the previous year's EMR if some bears were not detected, due to the fact that the population is assumed geographically closed. In comparison, our robust design framework includes temporary emigration, which means that a bear that is not found during an entire year will not be included in the total population size. Moreover, to use the robust design framework, we excluded tracks that were difficult to date, and those that fell outside of the secondary occasions (May to September), which left some individuals

identified by the GSTOP (Camarra et al. 2015) out of our database (see Supplementary materials). Agency estimates performed so far always included the individuals that were found dead in their yearly counts (Camarra et al. 2015), while a robust design model would only include them if the death occurred after the end of the primary occasion. Finally, the gender of a single individual born in 2012 was not determined, and as such that bear was excluded from our analyses. These factors put together account for the differences with the counts performed by the GSTOP.

5. Implications for bear abundance estimates

Even though different methods are used to estimate brown bear population size such as distance sampling and capture-recapture methods (Solberg et al. 2006), once the analyses are performed, most brown bear populations over the world are described by their total number of individuals, whether it is in Italy (Gervasi et al. 2008), in Sweden (Kindberg et al. 2011) or in the USA (Boulanger et al. 2008). Some other carnivore populations, however, are described with different indicators, such as the number of potential family groups (Andren et al. 2002) or the number of females with cubs of the year (Palomero et al. 2007). Considering the importance of the minimum population size for its viability (Shaffer 1981), we suggest that in complement to the minimal population size used by the GSTOP to describe the Pyrenean brown bears, it is important to develop an alternate counting method, especially since the heavy inbreeding (Camarra et al. 2015; Chapron et al. 2009) means that the genetic effective size (Palstra and Ruzzante 2008) of the Pyrenean brown bear population is lower than the “real” population size would let us to believe. Brown bear females in Europe (Swenson et al. 2007) usually start reproducing at the age of 4 or 5 with an interbirth interval of 2 years at least (Schwartz et al. 2003). Therefore, we suggest also describing the population by using the number of females with cubs of the year (Palomero et al. 2007) or the total number of 4+ years old females in the population, instead of limiting ourselves to the total number of bears. This would give a much more accurate insight of the viability of the population, which is an important tool to use when dealing with the conservation of endangered populations (Beissinger and Westphal 1998).

6. Importance of transboundary management

A transboundary management strategy helps to avoid significant errors in the estimates, most importantly the overestimation that occurs when a single individual is accounted for in one or more political jurisdiction due

to its mobility(Bischof et al. 2016). The sum of yearly abundance in France and Spain was always larger than abundance from the Combined dataset, hence confirming this assumption. Even though administrative borders might coincide with a difference in local priorities (Moilanen and Arponen 2011), such as the Scandinavian lynx (*Lynx lynx*) population which is much more heavily hunted in Norway than it is in Sweden (Swenson and Andrén 2005),a transboundary approach allows for more biologically logical management (Linnell and Boitani 2012). Differing management policies between two neighboring jurisdictions can have consequences on a large carnivore population, e.g. by creating source/sink dynamics (Swenson and Andrén 2005)that will in turn affect demography (Robinson et al. 2008). Here, we showed that it was possible to build a transborder model despite differences in monitoring methods between the different jurisdictions.

Conclusion

Perceived errors in abundance estimates might lead to skepticism towards the results provided by scientists, which can be followed by a negative evolution of a conservation conflict (Redpath et al. 2013). While accurate scientific data is not sufficient to solve the contention surrounding some large carnivore populations, it is a necessary step to advance towards mitigating conflicts by allowing the identification of human and ecological impacts (Redpath et al. 2013). It is especially important in the case of the Pyrenean brown bear, where the recent relocation of a male Slovenian bear in Catalonia could lead to an increase of the conflict, playing on the attitude differences between Pyrenean residents (Piédallu et al. 2016). In this context, reliable scientific data will be needed to avoid widening the gap between supporters and opponents of bear presence in the French-Spanish mountains. As large carnivore populations recover in Europe (Chapron et al. 2014), transboundary monitoring, however complex it might be (Kark et al. 2015),could become an essential first step towards harmonious and efficient transboundary management(Bischof et al. 2016; Kark et al. 2015), as it will be easier to implement than transboundary policies(Kark et al. 2015), and could facilitate the development of efficient solutions to ingrained conservation conflicts (Redpath et al. 2013).

298 Tables & Figures

299 **Table 1: Number of bear tracks found yearly**

		2008	2009	2010	2011	2012	2013	2014	Total	%Total Tracks	%ID'd Tracks
France	All year	99	58	159	189	228	203	162	1098	22.80%	74.34%
	May-Sep	80	52	123	130	157	161	121	824	17.11%	55.79%
	Apr-Nov	94	56	154	178	208	199	147	1036	21.52%	70.14%
Spain	All year	37	54	90	145	101	48	93	568	21.02%	63.18%
	May-Sep	24	41	65	124	66	29	77	426	15.77%	47.39%
	Apr-Nov	36	51	80	142	101	48	91	549	20.32%	61.07%
Total	All year	136	112	249	334	329	251	255	1666	22.16%	70.12%
	May-Sep	104	93	188	254	223	190	198	1250	16.63%	52.61%
	Apr-Nov	130	107	234	320	309	247	238	1585	21.09%	66.71%

300

301 Number of bear tracks per primary occasion of capture/Year. Two sets of secondary occasions are depicted:
 302 May-Sep (from May to September) and Apr-Nov (from April to November). The Total and Identified
 303 number of tracks in each dataset are shown in Table 1 of the Appendix section.

304

305 **Table 2: Model selection for each of the three datasets**

(a) France							
#	Survival	Detection	Emigration	npar	AICc	Weight	Deviance
6	0	Sex + Mix	None	12	649.3	0.27	636.1
30	0	Sex + Mix	Markovian	14	650.2	0.17	632.4
54	0	Sex + Mix	Random	13	651.1	0.11	635.6
12	Sex	Sex + Mix	None	13	651.2	0.10	635.7
28	0	Mix	Markovian	13	651.6	0.08	636.1
4	0	Mix	None	11	652.7	0.05	641.7
60	Sex	Sex + Mix	Random	14	653.1	0.04	635.3
34	Sex	Mix	Markovien	14	653.5	0.03	635.7
52	0	Mix	Random	12	653.8	0.03	640.6
10	Sex	Mix	None	12	654.5	0.02	641.2
26	0	Sex	Markovian	12	655.4	0.01	642.2
18	Age	Sex + Mix	None	19	655.5	0.01	625.9
58	Sex	Mix	Random	13	655.6	0.01	640.2
36	Sex	Sex + Mix	Markovian	15	655.8	0.01	635.7
(b) Spain							
#	Survival	Detection	Emigration	npar	AICc	Weight	Deviance
49	0	0	Random	10	447.8	0.25	420.9
50	0	Sex	Random	11	448.1	0.21	418.9
55	Sex	0	Random	11	449.6	0.10	420.3
56	Sex	Sex	Random	12	450.0	0.08	418.4
25	0	0	Markovian	11	450.1	0.08	420.8
26	0	Sex	Markovian	12	450.5	0.07	418.8
52	0	Mix	Random	12	451.4	0.04	419.7
31	Sex	0	Markovian	12	451.9	0.03	420.3
54	0	Sex + Mix	Random	13	452.3	0.03	418.2
32	Sex	Sex	Markovian	13	452.4	0.03	418.3
58	Sex	Mix	Random	13	453.3	0.02	419.2
28	0	Mix	Markovian	13	453.7	0.01	419.7
(c) Combined							
#	Survival	Detection	Emigration	npar	AICc	Weight	Deviance
12	Sex	Sex + Mix	None	13	725.0	0.21	724.5
6	0	Sex + Mix	None	12	725.2	0.19	726.9
58	Sex	Mix	Random	13	726.2	0.11	725.8
52	0	Mix	Random	12	726.4	0.10	728.1
18	Age	Sex + Mix	None	19	727.6	0.06	713.6
72	Sex + Age	Sex + Mix	Random	21	727.8	0.05	709.1
34	Sex	Mix	Markovian	14	728.4	0.04	725.8
28	0	Mix	Markovian	13	728.5	0.04	728.1
24	Sex + Age	Sex + Mix	None	20	728.6	0.03	712.3
10	Sex	Mix	None	12	728.9	0.03	730.7
64	Age	Mix	Random	19	729.2	0.02	715.3
4	0	Mix	None	11	729.4	0.02	733.3
36	Sex	Sex + Mix	Markovian	15	729.4	0.02	724.5
30	0	Sex + Mix	Markovian	14	729.6	0.02	726.9
70	Sex + Age	Mix	Random	20	730.2	0.02	713.9

306

307 List of the best models (Weight > 0.01) selected for each one of the three datasets ((a) France (b) Spain (c)

308 Total). #: Number of the model. Survival, Detection, Emigration: effects tested on each parameter in the

309 model. 0: No effect on the parameter. npar: number of parameters. Note: whenever model averaging was

310 impossible due to the different nature of the models (with or without a mixture effect), we only selected the
311 category with the highest combined weight.

Table 3: Model-averaged parameter estimates

Dataset	Bear Class	Survival		Detection		Emigration		Immigration	
France	Female class 1	0.90	± 0.07	0.84	± 0.13	0.08	± 0.08	0.50	± 0.46
	Female class 2			0.36	± 0.05				
	Male class 1	0.92	± 0.06	0.88	± 0.10				
	Male class 2			0.45	± 0.06				
Spain	Female	0.83	± 0.06	0.48	± 0.04	0.27	± 0.09	0.25	± 0.14
	Male	0.85	± 0.07	0.52	± 0.04				
Total	Female class 1	0.88	± 0.05	0.73	± 0.13	0.07	± 0.05	0.10	± 0.14
	Female class 2			0.32	± 0.05				
	Male class 1	0.93	± 0.04	0.81	± 0.05				
	Male class 2			0.41	± 0.06				

Model-averaged parameter estimates for each of the datasets. Bear Class represents the different individual classes identified by the mixture effect.

317 **Fig. 1: Map of bear monitoring in the French and Spanish Pyrenees**

318
319 Bear monitoring in the French and Spanish Pyrenees. Thin lines: systematic itineraries. Dots: camera traps.
320 The top-right mini-map indicates French counties and Spanish province – only the westernmost French
321 county is part of the Aquitaine-Poitou-Charentes region, all the others are part of the Languedoc-Roussillon-
322 Midi-Pyrénées region. In France, data monitoring is handled by the Game and Wildlife Agency. In Spain,
323 monitoring is handled separately between the three Autonomous communities: in Catalonia, the Fauna and
324 Flora Service of the Department of Territory and Sustainability is in charge with a local delegation to the
325 General Council in the Val d’Aran; in Aragon, it is the Environmental Services Department that oversees
326 bear management, and in Navarre, it is the Public Society for Environmental Management of Navarre. In
327 Andorra, monitoring is handled by the Natural Heritage Department.

328 **Figure 2: Population size estimation for each dataset**

329

330 Population size for the Pyrenean brown bears between 2008 and 2014 estimated using the three different
 331 datasets (France only, Spain only, and Combined).

332 **References**

- 333 Andren H et al. (2002) Estimating total lynx Lynx lynx population size from censuses of family groups
334 Wildlife Biology 8:299-306
- 335 Beissinger SR, Westphal I (1998) On the use of demographic models of population viability in endangered
336 species management Journal of Wildlife Management 62:821-841
- 337 Bischof R, Brøseth H, Gimenez O (2016) Wildlife in a Politically Divided World: Insularism Inflates
338 Estimates of Brown Bear Abundance Conservation Letters 9:122-130 doi:10.1111/conl.12183
- 339 Bischof R, Swenson JE (2012) Linking noninvasive genetic sampling and traditional monitoring to aid
340 management of a trans-border carnivore population Ecological Applications 22:361-373
- 341 Boulanger J, Kendall KC, Stetz JB, Roon DA, Waits LP, Paetkau D (2008) Multiple Data Sources Improve
342 DNA-Based Mark-Recapture Population Estimates Of Grizzly Bears Ecological Applications
343 18:577-589
- 344 Buckland ST, Burnham KP, Augustin NH (1997) Model Selection: An Integral Part of Inference Biometrics
345 53:603-618
- 346 Bull J, Nilsen EB, Mysterud A, Milner-Gulland EJ (2009) Survival on the Border: A Population Model to
347 Evaluate Management Options for Norway's Wolves Canis lupus Wildlife Biology 15:412-424
348 doi:10.2981/08-010
- 349 Bunnell FL, Tait DEN (1985) Mortality Rates of North American Bears Arctic 38:316-323
- 350 Burnham KP, Anderson DR (2002) Model Selection and Multimodel Inference: A Practical Information-
351 Theoretic Approach, 2nd Edition. Springer-Verlag, New York
- 352 Camarra JJ, Sentilles J, Bombillon N, Gastineau A, Quenette PY (2015) Suivi de l'ours brun dans les
353 Pyrénées françaises (Sous-populations occidentale et centrale) - Rapport annuel, Année 2015. Office
354 National de la Chasse et de la Faune Sauvage,
- 355 Chapron G, Gimenez O, Quenette P-Y (2010) Demographic parameters and viability of the Pyrenean brown
356 bear population.
- 357 Chapron G et al. (2014) Recovery of large carnivores in Europe's modern human-dominated landscapes
358 Science 346:1517-1519 doi:10.1126/science.1257553

- 359 Chapron G, Wielgus R, Quenette PY, Camarra JJ (2009) Diagnosing mechanisms of decline and planning
360 for recovery of an endangered brown bear (*Ursus arctos*) population PloS one 4:e7568
361 doi:10.1371/journal.pone.0007568
- 362 Cubaynes S et al. (2010) Importance of accounting for detection heterogeneity when estimating abundance:
363 the case of French wolves Conservation biology : the journal of the Society for Conservation
364 Biology 24:621-626 doi:10.1111/j.1523-1739.2009.01431.x
- 365 De Barba M, Waits LP, Garton EO, Genovesi P, Randi E, Mustoni A, Groff C (2010) The power of genetic
366 monitoring for studying demography, ecology and genetics of a reintroduced brown bear population
367 Mol Ecol 19:3938-3951 doi:10.1111/j.1365-294X.2010.04791.x
- 368 Fagen R, Fagen JM (1996) Individual Distinctiveness in Brown Bears, *Ursus arctos* L Ethology 102:212-226
369 doi:10.1111/j.1439-0310.1996.tb01119.x
- 370 Garshelis DL, Gibeau M, Herrero S (2005) Grizzly bear demographics in and around Banff National Park
371 and Kananaskis Country, Alberta Journal of Wildlife Management 69:277-297
- 372 Gervasi V, Brøseth H, Nilsen EB, Ellegren H, Flagstad Ø, Linnell JDC (2015) Compensatory immigration
373 counteracts contrasting conservation strategies of wolverines (*Gulo gulo*) within Scandinavia
374 Biological Conservation 191:632-639 doi:10.1016/j.biocon.2015.07.024
- 375 Gervasi V et al. (2008) A preliminary estimate of the Apennine brown bear population size based on hair-
376 snag sampling and multiple data source mark–recapture Huggins models *Ursus* 19:105-121
- 377 Gervasi V, Ciucci P, Boulanger J, Randi E, Boitani L (2012) A multiple data source approach to improve
378 abundance estimates of small populations: The brown bear in the Apennines, Italy Biological
379 Conservation 152:10-20 doi:10.1016/j.biocon.2012.04.005
- 380 Gittleman JL, Harvey PH (1982) Carnivore Home-Range Size, Metabolic Needs and Ecology Behavioral
381 Ecology and Sociobiology 10:57-63
- 382 Hirata D et al. (2013) Molecular phylogeography of the brown bear (*Ursus arctos*) in Northeastern Asia
383 based on analyses of complete mitochondrial DNA sequences Molecular biology and evolution
384 30:1644-1652 doi:10.1093/molbev/mst077
- 385 Kark S, Tulloch A, Gordon A, Mazar T, Bunnefeld N, Levin N (2015) Cross-boundary collaboration: key to
386 the conservation puzzle Current Opinion in Environmental Sustainability 12:12-24
387 doi:10.1016/j.cosust.2014.08.005

388 Kendall WL, Nichols JD, Hines JE (1997) Estimating Temporary Emigration Using Capture-Recapture Data
389 With Pollock's Robust Design Ecology 78:563-578

390 Kindberg J, Swenson JE, Ericsson G, Bellemain E, Miquel C, Taberlet P (2011) Estimating population size
391 and trends of the Swedish brown bear *Ursus arctos* population Wildlife Biology 17:114-123
392 doi:10.2981/10-100

393 Laake JL (2013) RMark: An R Interface For Analysis Of Capture-Recapture Data With MARK. Alaska
394 Fisheries Science Center,

395 Linnell JD, Salvatori V, Boitani L (2008) Guidelines for Population Level Management Plans for Large
396 Carnivores. European Commission,

397 Linnell JDC, Boitani L (2012) Building biological realism into wolf management policy: the development of
398 the population approach in Europe *Hystrix*, the Italian Journal of Mammalogy 23:80-91
399 doi:10.4404/hystrix-23.1-4676

400 Mattisson J et al. (2013) Home range size variation in a recovering wolf population: evaluating the effect of
401 environmental, demographic, and social factors *Oecologia* 173:813-825 doi:10.1007/s00442-013-
402 2668-x

403 McDonald LL (2004) Sampling rare populations. In: Thompson WL (ed) Sampling Rare and Elusive
404 Species. Island Press, Washington, DC, pp 11-42

405 McLellan B et al. (1999) Rates and Causes of Grizzly Bear Mortality in the Interior Mountains of British
406 Columbia, Alberta, Montana, Washington and Idaho *Journal of Wildlife Management* 63:911-920
407 doi:10.2307/3802805

408 McLoughlin PD, Ferguson SH, Messier F (2000) Intraspecific Variation in Home Range Overlap with
409 Habitat Quality: A Comparison among Brown Bear Populations *Evolutionary Ecology* 14:39-60
410 doi:10.1023/a:1011019031766

411 Miquel C, Bellemain E, Poillot C, BessiÈRe J, Durand A, Taberlet P (2006) Quality indexes to assess the
412 reliability of genotypes in studies using noninvasive sampling and multiple-tube approach *Molecular*
413 *Ecology Notes* 6:985-988 doi:10.1111/j.1471-8286.2006.01413.x

414 Moilanen A, Arponen A (2011) Administrative regions in conservation: Balancing local priorities with
415 regional to global preferences in spatial planning *Biological Conservation* 144:1719-1725
416 doi:10.1016/j.biocon.2011.03.007

417 Nilsen EB, Herfindal I, Linnell JDC (2005) Can intra-specific variation in carnivore home-range size be
418 explained using remote-sensing estimates of environmental productivity? *Ecoscience* 12:68-75
419 doi:10.2980/i1195-6860-12-1-68.1

420 Palomero G, Ballesteros F, Nores C, Blanco JC, Herrero J, Garcia-Serrano A (2007) Trends in number and
421 distribution of brown bear females with cubs-of-the-year in the Cantabrian Mountains, Spain *Ursus*
422 18:145-157

423 Palstra FP, Ruzzante DE (2008) Genetic estimates of contemporary effective population size: what can they
424 tell us about the importance of genetic stochasticity for wild population persistence? *Molecular*
425 *Ecology* 17:3428-3447 doi:10.1111/j.1365-294X.2008.03842.x

426 Piédallu B et al. (2016) Spatial variation in public attitudes towards brown bears in the French Pyrenees
427 *Biological Conservation* 197:90-97 doi:10.1016/j.biocon.2016.02.027

428 RCoreTeam (2013) R: A Language and Environment for Statistical Computing. R Foundation for Statistical
429 Computing,

430 Redpath SM et al. (2013) Understanding and managing conservation conflicts *Trends in ecology & evolution*
431 28:100-109 doi:10.1016/j.tree.2012.08.021

432 Robinson HS, Wielgus R, Cooley HS, Cooley SW (2008) Sink Populations in Carnivore Management:
433 Cougar Demography and Immigration in a Hunted Population *Ecological Applications* 18:1028-
434 1037

435 Schwartz CC et al. (2003) Reproductive maturation and senescence in the female brown bear *Ursus* 14:109-
436 119

437 Shaffer ML (1981) Minimum Population Sizes for Species Conservation *BioScience* 31:131-134

438 Smith HC, Pollock K, Waples K, Bradley S, Bejder L (2013) Use of the robust design to estimate seasonal
439 abundance and demographic parameters of a coastal bottlenose dolphin (*Tursiops aduncus*)
440 population *PloS one* 8:e76574 doi:10.1371/journal.pone.0076574

441 Solberg KH, Bellemain E, Drageset O-M, Taberlet P, Swenson JE (2006) An evaluation of field and non-
442 invasive genetic methods to estimate brown bear (*Ursus arctos*) population size *Biological*
443 *Conservation* 128:158-168 doi:10.1016/j.biocon.2005.09.025

444 Swenson JE, Adamic M, Huber D, Stokke S (2007) Brown bear body mass and growth in northern and
445 southern Europe *Oecologia* 153:37-47 doi:10.1007/s00442-007-0715-1

446 Swenson JE, Andrén H (2005) A tale of two countries: large carnivore depredation and compensation
447 schemes in Sweden and Norway. In: Woodroffe R, Thirgood S, Rabinowitz A (eds) People and
448 Wildlife, Conflict or Co-existence? Cambridge University Press, pp 323-339

449 Swenson JE, Taberlet P, Bellemain E (2011) Genetics and conservation of European brown bears *Ursus*
450 *arctos* Mammal Review 41:87-98 doi:10.1111/j.1365-2907.2010.00179.x

451 Taberlet P et al. (1997) Noninvasive genetic tracking of the endangered Pyrenean brown bear population
452 Molecular Ecology 6:869-876

453 Trouwborst A (2010) Managing the Carnivore Comeback: International and EU Species Protection Law and
454 the Return of Lynx, Wolf and Bear to Western Europe Journal of Environmental Law 22:347-372
455 doi:10.1093/jel/eqq013

456 Waits LP, Talbot SL, Ward RH, Shields GF (1998) Mitochondrial DNA Phylogeography of the North
457 American Brown Bear and Implications for Conservation Conservation Biology 12:408-417
458 doi:10.1111/j.1523-1739.1998.96351.x

459 White GC, Burnham KP (1999) Program MARK: survival estimation from populations of marked animals
460 Bird Study 46:S120-S139 doi:10.1080/00063659909477239

461 Wielgus R, Bunnell FL (1994) Dynamics of a small, hunted Brown Bear *Ursus Arctos* population in
462 Southwestern Alberta, Canada Biological Conservation 67:161-166

463

464