1 Title

2 “Same Difference”: Comprehensive evaluation of three DNA methylation measurement
3 platforms.

4 Authors

5 Thadeous J Kacmarczyk¹, Mame P. Fall¹, Xihui Zhang¹, Yuan Xin¹, Yushan Li¹, Alicia Alonso¹,
6 Doron Betel^{1,2}

7 1. Department of Medicine, Division of Hematology/Oncology, Epigenomics Core Facility, Weill
8 Cornell Medicine, New York, New York

9 2. Institute for Computational Biomedicine, Weill Cornell Medicine, New York, New York

10

11 Corresponding author: Thadeous J Kacmarczyk, thk2008@med.cornell.edu

12 Abstract

13 **Background:** DNA methylation in CpG context is fundamental to the epigenetic regulation of
14 gene expression in high eukaryotes. Disorganization of methylation status is implicated in many
15 diseases, cellular differentiation, imprinting, and other biological processes. Techniques that enrich
16 for biologically relevant genomic regions with high CpG content are desired, since, depending on
17 the size of an organism's methylome, the depth of sequencing required to cover all CpGs can be
18 prohibitively expensive. Currently, restriction enzyme based reduced representation bisulfite
19 sequencing and its modified protocols are widely used to study methylation differences. Recently,
20 Agilent Technologies and Roche NimbleGen have ventured to both reduce sequencing costs and
21 capture CpGs of known biological relevance by marketing in-solution custom-capture hybridization
22 platforms. We aimed to evaluate the similarities and differences of these three methods

23 considering each targets approximately 10-13% of the human methylome.

24 **Results:** Overall, the regions covered per platform were as expected: targeted capture based
 25 methods covered >95% of their designed regions whereas the restriction enzyme-based method
 26 covered >70% of the expected fragments. While the total number of CpG loci shared by all
 27 methods was low, ~30% of any platform, the methylation levels of CpGs common across platforms
 28 were concordant. Annotation of CpG loci with genomic features revealed roughly the same
 29 proportions of feature annotations across the three platforms. Targeted capture methods
 30 encompass similar amounts of annotations with the restriction enzyme based method covering
 31 fewer promoters (~9%) and shores (~8%) and more unannotated loci (7-14%).

32 **Conclusions:** Although all methods are largely consistent in terms of covered CpG loci and cover
 33 similar proportions of annotated CpG loci, the restriction based enrichment results in more
 34 unannotated regions and the commercially available capture methods result in less off-target
 35 regions. Quality of DNA is very important for restriction based enrichment and starting material
 36 can be low. Conversely, quality of the starting material is less important for capture methods, and
 37 at least twice the amount of starting material is required. Pricing is marginally less for restriction
 38 based enrichment, and number of samples to be prepared is not restricted to the number of
 39 samples a kit supports. The one advantage of capture libraries is the ability to custom design
 40 areas of interest. The choice of the technique should be decided by the number of samples, the
 41 quality and quantity of DNA available and the biological areas of interest since comparable data
 42 are obtained from all platforms.

43

44 **Keywords:** Bisulfite sequencing, DNA methylation, Methylome capture, RRBS, 5mC, CpG

45 Background

46 DNA cytosine methylation in the form of 5-methylcytosine (5mC) in CpG context is an epigenetic
 47 marker that is important for regulation of gene expression. Changes in CpG methylation are
 48 implicated in many diseases, and proper methylation patterns are required for normal
 49 development [1]. Large scale studies such as ENCODE [2] and the Human Epigenomics
 50 Roadmap [3] have performed extensive profiling of 5mC in various cell lines and tissues revealing
 51 a rich and dynamic landscape of 5mC patterns in the human genome. Given the importance of
 52 these markers to cellular development and contribution to disease, a number of approaches have
 53 been developed for detecting the methylation status of cytosines [4], with bisulfite sequencing
 54 (BS-seq) being widely used to provide single-base quantitative measurement of 5mC (and 5-
 55 hydroxymethylcytosine, 5hmC). Bisulfite sequencing refers to massively parallel sequencing after
 56 chemical deamination of cytosines (C) to uracils (U), followed by polymerase chain reaction
 57 (PCR). The deamination of cytosines is accomplished by the use of sodium bisulfite, and this pre-
 58 treatment preserves both the methyl modification in 5mC and the 5-hydroxymethyl modification in
 59 5hmC [5]. The benchmark in methylome coverage is whole genome bisulfite sequencing (WGBS),
 60 which at 30X sequencing coverage, accounts for ~94% of all cytosines in the genome with 99.8%
 61 of them being CpG loci [6]. However, different WGBS library preparation protocols can bias region
 62 coverage. Since no method completely covers the methylome, and biologically relevant CpGs
 63 have been identified in known genomic features [1,7], developing focused assays considering
 64 these features is in demand with the caveat that these approaches will leave gaps in the
 65 methylome potentially excluding important CpGs.

66
 67 There are several methods for acquiring DNA methylation levels and we investigated the
 68 characteristics of three currently widely used platforms: i) enrichment by enzymatic digestion

(MspI) enhanced reduced representation bisulfite sequencing (ERRBS)[8], ii) capture based Agilent SureSelect Methyl-Seq (SSMethylSeq), and iii) capture based Roche NimbleGen SeqCap Epi CpGiant (CpGiant).

In this paper we present an analysis of the methylation pattern of the human lung fibroblast IMR-90 cell line profiled by each of the platform protocols, using two technical replicate libraries for ERRBS, and two libraries each for SSMethylSeq and CpGiant, one at the manufacturer's suggested concentration and one at a reduced concentration (Table 1 and Additional file 1: Table S1). The capture libraries differ only in concentration of input material and are treated effectively as technical replicates. All libraries were sequenced to equivalent depth and compared to a library made using WGBS.

Materials and Methods

Cell growth and DNA preparation

IMR90 cells (American Type Culture Collection, Manassas, VA cat # CCL-186) were provided by Dan Hasson (Mount Sinai School of Medicine, New York, NY). DNA from 5×10^7 cells was purified using the Gentra Puregene DNA kit according to manufacturer protocol (cat # 158389, Qiagen Valencia, CA). DNA was resuspended in TE, quantified using fluorometric quantification (Qubit 2.0 ThermoFisher Scientific Waltham, MA) and quality was assessed by running on a 1% agarose gel.

ERRBS (digestion-based enhanced reduced representation bisulfite sequencing)

Two ERRBS libraries were prepared as described in Garrett-Bakelman, et al [8]. Briefly, 75 ng of DNA was digested with the methylation insensitive MspI enzyme ($C^{\wedge}CGG$). After end-repair, A-

tailing, and adapter ligation with Illumina TruSeq adapters, the region corresponding to 84-334bp was size-selected as two fractions. Each fraction was subjected to overnight bisulfite conversion (55 cycles of 95°C for 30 sec, 50°C for 15 min) using EZ DNA methylation kit (Cat # D5002, Zymo Research, Irvine CA). Purified bisulfite converted DNA was PCR amplified using TruSeq primers (Illumina Inc. San Diego, CA) for 18 cycles of denaturing, annealing and extension/elongation steps using Roche FastStart (cat # 03 553 361 001) – 94°C for 20 secs, 65°C for 30 secs, 72°C for 1 min, followed by 72°C for 3 min. The resulting libraries were normalized to 2nM and pooled at the same molar ratio. Samples were clustered at 6.5pM on a V3 paired-end read flow cell and sequenced for 100 cycles on an Illumina HiSeq 2500.

Agilent SureSelect Methyl-Seq (SSMethylSeq)

Two libraries were made according to the company's specifications using 3ug and 1ug of DNA. DNA was sonicated using a Covaris S220 sonicator (Covaris, Woburn, MA) to obtain products of 150-200bp. DNA was then end-repaired, A-tailed and ligated with methylated adapters to create pre-capture DNA libraries. DNA Libraries were then hybridized to the RNA SureSelect Human methyl-seq capture library at 65°C for 16 hrs. Hybridized products were purified by capture with Streptavidin beads and then subjected to bisulfite conversion (64°C for 2.5hr) using the Zymo EZ DNA Gold kit (Cat # D5005, Zymo Research, Irvine CA). The bisulfite treated libraries were PCR amplified for 8 cycles with Agilent Taq to get the required amount of DNA library and then indexed by another 6 cycles of PCR amplification to create the final libraries. Libraries were clustered at 11pM on a V3 paired-end read flow cell and sequenced for 100 cycles on an Illumina HiSeq 2500.

Roche NimbleGen SeqCap Epi CpGiant (CpGiant)

Two libraries were made using 1ug and 0.25ug of starting material, according to the company's specifications. DNA was sonicated using a Covaris S220 sonicator (Covaris, Woburn, MA) to

114 obtain products of 180-220bp. DNA was then end-repaired, A-tailed and ligated with methylated
 115 indexed-adapters to create pre-capture DNA libraries. The pre-capture libraries were bisulfite
 116 converted at 54°C for 1 hour using the Zymo EZ DNA Lightning kit (Cat # D5030, Zymo Research,
 117 Irvine CA). The bisulfite treated pre-capture libraries were PCR amplified with HiFi HotSart Uracil+
 118 polymerase (Cat# KK2802, Kapa Biosystems, Wilmington, MA). The amplified, bisulfite converted
 119 sample libraries were then hybridized to the probe pool of fully-, partially- and un-methylated
 120 cytosines from both strands of DNA oligos at 42°C for 72 hrs. Hybridized products were purified by
 121 capture with Capture Beads and PCR amplified for 15 cycles to create the final libraries. Libraries
 122 were clustered at 12pM on a V3 paired-end read flow cell and sequenced for 100 cycles on an
 123 Illumina HiSeq 2500.

124 **Whole Genome Bisulfite Sequencing (WGBS)**

125 Briefly, 100 ng of DNA were bisulfite converted using EZ DNA Methylation-Gold Kit (cat # D5005,
 126 Zymo Research Corporation, Irvine, CA) and the single stranded DNA obtained processed for
 127 library construction using the EpiGnome Methyl-Seq kit (Cat. # EGMK81324) as per manufacturer
 128 protocol (Illumina Madison, Madison, WI). The DNA was made double stranded by the use of
 129 5'tagged random hexamers and subsequently 3' tagged with terminal tagging oligo. The di-tagged
 130 DNA was enriched using 10 cycles of PCR, with PCR indexed-primers compatible with Illumina
 131 sequencing. The library was clustered at 7 pM on a V3 paired-end read flow cell and sequenced
 132 for 100 cycles on an Illumina HiSeq 2500.

133 **Computational analysis**

134 Illumina's CASAVA 1.8.2 was used to generate fastq files from basecalls. Raw fastq reads were
 135 processed by a custom pipeline that consists of: 1) filtering raw fastq reads for pass filter reads,
 136 2) trimming adapter sequence by FLEXBAR [9], 3) genomic alignments performed using Bismark

[10] and Bowtie [11] to reference human genome hg19, and 4) methylation calling by a custom PERL script [8]. Custom analysis scripts were written in R (version 3.3.0 [12]), including packages: Bioconductor - Biobase version 2.32.0 [13], GenomicRanges package version 1.24.0 [14], Beeswarm package version 0.2.3 [15], Affy package version 1.50.0 [16], BSgenome.Hsapiens.UCSC.hg19 package version 1.4.0 [17], PreprocessCore package version 1.34.0 [18], UpSetR package version 1.2.0 [19], and VennDiagram package version 1.6.17 [20], were used to perform the analysis and are available at GitHub (<https://github.com/thk2008/methylseqplatformcomparison>). Sequencing data and methylation results are available to download from GEO (GSE83595)

Results

Region coverage

Targeted capture techniques (SSMethylSeq and CpGiant) have a designed set of genomic regions and therefore, a predicted set of CpGs covered. SSMethylSeq is specifically designed to capture CpGs from a single DNA strand, where the other platform capture CpGs from both DNA strands. The ERRBS protocol is considered targeted to the extent that the restriction digest produces consistent genomic fragments, sizes from 84-334bp are isolated during the library preparation gel extraction step. Note that since the DNA for WGBS is randomly sheared and coverage depends upon sequencing depth, there are no predicted regions per se and consequently WGBS is excluded from the region coverage analysis. Regions are considered overlapping if any two regions overlap by at least 10 bases. Overall, SSMethylSeq and MspI 84-334b have similar region length proportions, while CpGiant has fewer regions and more variable lengths (Figure 1A and Additional file 1: Table S2). Additionally, the two capture platforms cover similar regions, and similar amount of different regions from MspI 84-334b (Figure 1B).

160

161 **Description of analysis**

162 Even though the platforms are different, the data are all bisulfite sequencing data, thus the
 163 procedure for processing the data is the same. First we evaluated sequencing and alignment
 164 consistency. Next we compared strand symmetry of methylation values, since the SSMethylSeq
 165 platform is designed to cover predominantly one strand. Then we evaluated several measures to
 166 quantify properties of each platform relating to the target regions that are covered, methylation
 167 levels of cytosines in CpG context and coverage of genomic regions. Target region analysis
 168 includes the number of CpGs covered, the fraction of target regions covered, the coverage of
 169 target region CpGs, the overlap of CpGs across all platforms, and the concordance of methylation
 170 levels of overlapping CpGs. Similarly, genomic region evaluation is comprised of the number and
 171 fraction of annotated CpGs by genomic feature (i.e. CpG islands or shores – 2kb flanking the
 172 islands) and by gene part (i.e. promoter, exon, intron), and the overlap, coverage, and
 173 concordance of genomic annotations. We used the median absolute deviation (MAD), a robust
 174 measure of variability insensitive to outliers, to estimate the statistical dispersion in methylation
 175 level comparisons.

176

177 **Sample preparation and sequencing**

178 The general library preparation protocol for ERRBS is to digest input DNA with the methylation
 179 insensitive MspI enzyme followed by addition of adapters and barcodes, enrichment by gel size
 180 selection for fragments 84-334bp, bisulfite conversion, and amplification by PCR. The capture
 181 methods of SSMethylSeq and CpGiant are similar in that targeted capture of genomic regions is
 182 performed by hybridization to probes thereby providing a direct measure of CpG methylation at
 183 predefined regions. However, the methods differ in two important aspects. First, the SSMethylSeq

platform captures and measures methylation from only one DNA strand. Second, in CpGiant's methodology bisulfite conversion is performed before hybridization to oligonucleotides whereas in SSMethylSeq bisulfite conversion is performed prior to genomic capture. Hence, CpGiant's probe design must capture the various combinations of DNA sequences that can result from bisulfite conversion. In contrast to the targeted approaches WGBS is unbiased and relies on bisulfite breakage of genomic DNA. All libraries were made from human lung fibroblast IMR90 cell line DNA and prepared as described in the materials and methods with prominent differences outlined in Table 1.

Table 1: Protocols comparison

	ERRBS	SS-MethylSeq	CpGiant	WGBS
DNA requirement	75ng >40kb	1, 3ug	0.25,1ug	100 ng >40kb
DNA processing	MspI digestion to completion followed by fractionation of 84-334bp	Sonication to 150-200bp	Sonication to 180-220bp	Single stranded and fragmented during bisulfate conversion
Enrichment method	Size fractionation of 84-334bp sizes	Hybridization to RNA capture library stranded design	Hybridization to oligo probes containing fully-, partially- and un-methylated cytosines from both strands	none
Bisulfite conversion step	Post-adaptor ligation	Post-hybridization capture	Pre-hybridization capture	Pre-adaptor ligation
Zymo Research Bisulfite Conversion kit	EZ-DNA Methylation (50°C, 55cycles)	EZ-DNA Methylation Gold (64°C, 2.5hr)	EZ-Methylation Lightning (54°C, 1hr)	EZ DNA Methylation-Gold
Total PCR amplification cycles	18 Post enrichment and bisulfite conversion	16 Post enrichment and bisulfite conversion (8 for amplification and 6 for Indexing)	28 12 post bisulfite conversion & 16 post enrichment	10 Post bisulfite conversion
DNA Polymerase (uracil tolerant)	FastStart Taq (Roche)	Taq2000 (Agilent Technologies)	HiFi HotSart Uracil+ (Kapa Biosystems)	FailSafe Enzyme (Epicentre)
Predicted number of targeted CpG	6.6M	3.7M	5.6M	56M

sites				
Relative price (library preparation and fixed coverage)	13.5 %	15.5%	15.5%	100%

Sequencing and alignment characteristics

Generally, all platforms produced similar sequencing results with no noticeable bias or reduced quality scores. The number of clusters and number of pass filter reads produced a typically consistent number of usable reads for all samples (254M +/- 37M), see Additional file 1: Table S3 for more details.

Consistent with other bisulfite converted samples the number of uniquely mapped reads was ~ 70.8% sd=5.2% (Figure 2A). Across all platforms, a mean of 24.2%, s= 2.2% of reads were not aligned and ambiguously mapped reads showed a larger proportion for ERRBS (~12.1%), than WGBS (4.6%), SSMethylSeq (1.4%), or CpGiant (1.4%) (Figure 2A). These observations indicate the propensity of designed platforms to avoid repetitive regions or paralogous regions in contrast to ERRBS where no such selection is possible.

Strand methylation symmetry

Maintenance of symmetric methylation patterns across complementary CpG sites is required in order to preserve methylation patterns across cellular divisions. In principle, measuring methylation levels on one strand is sufficient to infer the cellular methylation state. Indeed, the SSMethylSeq platform is designed to capture only one strand in contrast to all other platforms that capture from both strands (Figure 2B). We note that this is different from measuring hemi-methylation states, which refers to different methylation between two parental alleles. This

requires genomic phasing information and is not considered in this analysis. To validate that methylation is indeed strand symmetric we compared the methylation values between complementary CpG sites (Figure 3). For ERRBS and CpGiant about 40% of the CpGs were complementary and analysis of SSMethylSeq was done with the ~2% of the probes that were complementary. We found strong agreement in methylation values in all samples and all platforms supporting the notion of symmetric methylation (Figure 3, mean MAD=0.28, sd=0.06). ERRBS contains a small set of discordant CpG sites (Figure 3A, B), where methylation values are inconsistent among complementary sites ($\Delta > 99$, 3.8% of sites ERRBS_A, 1.0% in ERRBS_B at 10X coverage). This discordance is a consequence of the ERRBS library preparation where the MspI staggered cut sites are *de novo (in vitro)* filled with C' and G' to generate blunt ends. As a result, methylated CpGs at the restriction sites (i.e. at the ends of sequences fragments) are discordant. To correct this artificial discordance, it's common for single end sequencing to discard CpG values at MspI cut sites. We observed additional disparity occurring in paired-end sequencing where the sequenced fragments are shorter than the overall sequencing length where sequencing reads from opposite strands are overlapping. Since the fraction of discordant sites is small, we have kept them in the analysis. However, In order to maintain equal evaluation of CpG capture and methylation levels among all platforms, all subsequent analyses are based on CpG-units, which are defined as CpG sites with ≥ 10 spanning reads regardless of which strand the reads are mapped (i.e. if two CpG sites are complementary they are combined into one unit and then filtered for ≥ 10 x coverage).

233

234 **CpG-unit coverage and target region coverage.**

235 We evaluated the number of CpG-units covered and the mean coverage of all CpG-units that
236 were sequenced at 10X depth or more. ERRBS, SSMethylSeq and CpGiant platforms cover on

average 3.8M CpG-units, sd=0.2M at mean coverage of 114.27, sd=23.6 (Figure 4A,B and Additional file 1: Figures S1 and S2). In contrast, WGBS sequenced at ~400M paired-end reads covered 14.4M CpG-units (~50% of the 28M total CpG-units in the genome) at a mean coverage of 19x, demonstrating that achieving comparable coverage by WGBS is significantly more costly than targeted platforms. Therefore, these results confirm that while WGBS profiles a large portion of CpG-units, the targeted platforms provide a cost effective way to interrogate a limited, yet potentially most informative, set of CpG-units at considerably reduced cost.

Next, we evaluated the extent of coverage in the targeted regions and the fraction of their CpGs identified. Here, a region is considered as having coverage if at least one CpG-unit is sufficiently covered within the region. We observed that ERRBS covers on average 79.8% of its predicted regions 84-334bp, whereas CpGiant and SSMethylSeq cover, on average, 98.7% and 95.3% of their expected capture regions, respectively (Figure 4A). While region coverage and the coverage of CpG-units within the targeted regions are high, we observed that roughly 20%-40% of the CpG-units covered are outside the targeted regions (Figure 4A). In the case of ERRBS this is expected since restriction digestion and gel isolation can be variable. In the case of capture methods, this may indicate either cross-hybridization of the probes to other genomic locations or hybridization of longer fragments. Since there are typically several CpG-units within a region, we looked at the distribution of the fraction of CpG-units covered per region and found that nearly all CpG-units in targeted regions are covered although for both SSMethylSeq and ERRBS a number of CpGs in targeted regions are not represented presumably due to coverage bias (Figure 4C).

CpG-unit overlap and methylation levels concordance

Overall, the average number of common CpG-units covered by all three platforms is ~30% +/- 1% (Figure 4D). When comparing the pairwise overlap of shared CpG-units across all samples, we

observe high overlap between intra-platform replicates with median number of shared CpG-units ~3.58M, median overlap 92.1% and high methylation level concordance with mean MAD = 0.23 and sd = 0.05 (Additional file 1: Figure S3 and Table S4). Among inter-platform samples we observed less overlap; median number of shared CpG-units ~1.5M, median overlap 39.1% (Figure 5 upper triangle, Additional file 1: Figure S3 and Table S4). Methylation levels of common CpG-unit's across all platforms are highly concordant indicating that methylation level measurements are consistent and reproducible with average MAD = 0.41 and sd = 0.25 (Figure 5 lower triangle, Additional file 1: Figure S3 and Table S4). Inter-platform concordance was slightly lower than intra-platform concordance with mean MAD = 0.46 and sd = 0.27 (Figure 5 lower triangle, Additional file 1: Figure S3 and Table S4). These results demonstrate that while the platforms differ in their capture approaches and bisulfite conversion kits (Table 1) these differences are not biasing methylation measurements. The differences among the platforms, therefore, are largely restricted to variations in targeted regions and not in methylation measurements.

275

276 **Coverage of genomic feature regions by CpG-units**

Next we were interested in what genomic features are covered by each platform and the degree of coverage. Analogous to the previous analysis, here we define a region as being one of exon, intron, promoter, CpGi, or shore, and coverage is a genomic region that contains at least one detected CpG-unit. It should be noted that the genomic feature annotations are not mutually exclusive and that some CpG-units are annotated by more than one category. Naturally, the targeted platforms focus on genomic regions known to play important roles in epigenetic regulation such as promoter regions, CpG islands, and CpG shores and ERRBS covers the same regions, but to a lesser degree (Figure 6A). Moreover, each platform covers similar proportions of

285 CpG-units in particular genomic region (Figure 6B). Conversely, we looked at the number of CpG-
 286 units that have an annotation and observed similar trends for the three platforms with ERRBS
 287 having a larger proportion of unannotated CpG-units (~27%, Figure 6C). Overall, no platform is
 288 significantly enriched for particular genomic region although ERRBS is slightly less represented in
 289 promoters, CpG islands and CpG shores, while having more representation of unannotated CpG-
 290 units.

291

292 **Overlap of CpG-units annotated with a genomic feature**

293 We evaluated the overlap of annotated CpG-units to determine whether any platform is enriched
 294 for a particular genomic feature (e.g. exons, introns, promoters, CpG islands and shores). Again, it
 295 should be mentioned that the genomic feature annotations are not mutually exclusive and that
 296 some CpG-units are annotated by more than one category. However, a CpG-unit is counted as
 297 annotated if it has one or more annotations. Comparing the overlap in annotations across
 298 platforms, we see a similar grouping as above with the methylation levels, intra-platform overlap is
 299 high (mean overlap 93.7% sd=2.9%), and inter-platform overlap is lower (mean 55.1% sd=15.5%)
 300 (Figure 7 and Additional file 1: Figures S4-S9). Thus, we observed similar proportions of genomic
 301 region coverage across all platforms, but lower overlap of annotated CpG-units suggesting
 302 coverage of different loci within those regions.

303

304 **Discussion**

305 We performed a systematic analysis of the characteristics of three prominent platforms for
 306 measuring DNA methylation levels: ERRBS, Agilent SureSelect Methyl-Seq (SSMethylSeq),
 307 Roche NimbleGen SeqCap Epi CpGiant (CpGiant), and WGBS with the goal of identifying

whether one method outperforms the others for any particular property.

We assessed the expected region coverage for SSMethylSeq and CpGiant and MspI (for ERRBS) and observed that, overall, the CpG sites covered by each platform are largely distinct from each other. SSMethylSeq and CpGiant cover roughly 70% of the same CpGs, while MspI has only ~30% overlap with either capture. However, each platform covers a large fraction of its targeted regions. SSMethylSeq and CpGiant cover greater than 95% of their designed capture regions and ERRBS covers at least 77% of its expected fragments. Furthermore, methylation levels of overlapping CpG-units are highly concordant.

Finally, each platform covered roughly the same proportions of genomic features (CpG islands, shores, promoters, exons, and introns), but profiled different CpG sites within those regions. In addition to the differences in the targeted CpG positions, there are also differences in the library preparation protocols. ERRBS can be performed using small amounts of starting material (as little as 75ngs), whereas SSMethylSeq requires 1ug and CpGiant 0.25ug of starting DNA (although those are expected to reduce with further optimization by the vendors). ERRBS, or digestion-based methods, can cover certain genomic regions that are not amenable to capture. ERRBS provides comparable data to the other platforms, although there is more variability among the profiled CpGs. Capture platforms are more precise and can be customized for profiling specific genomic regions of interest. In addition, capture platforms are the only methods available for methylation profiling of low quality, degraded DNA although library preparation is more labor intensive and efficiency of capture depends on shearing.

Conclusions

Epigenetic state is a fundamental element of cellular development and regulation. Therefore, accurate, reproducible and cost effective approaches for profiling DNA methylation are important for advancements in biomedical research. While the cost of sequencing continues to decrease,

332 reaching sufficient coverage for reliable measurement of CpG methylation by WGBS is still
 333 prohibitive. We conclude from our comparative study that capture-based approaches provide
 334 comparable results and cover more precisely their intended targets than ERRBS, which is a
 335 restriction enzyme based approach. They also provide the added flexibility of designing custom
 336 capture for surveying regions of interest. On the other hand, digestion based protocols are
 337 currently more cost effective and may be the only option for clinical samples where the amount of
 338 input DNA is limited. In the absence of any prior knowledge about the genomic regions of interest
 339 for a particular study, the choice of methylation profiling platform should be guided by cost and
 340 amount and quality of input DNA.

341

342 **List of abbreviations**

343 5mC: 5-methylcytosine, BS-seq: bisulfite sequencing, CpGiant: Roche NimbleGen SeqCap Epi
 344 CpGiant, CpGi: CpG island, ERRBS: enhanced reduced representation bisulfite sequencing,
 345 GEO: gene expression omnibus, MAD: median absolute deviation, PCR: polymerase chain
 346 reaction, SSMethylSeq: Agilent SureSelect Methyl-Seq, WGBS: whole-genome bisulfite
 347 sequencing

348

349 **Declarations**

350 *Ethics approval and consent to participate*

351 Not applicable

352 *Consent for publication*

353 Not applicable

354 *Availability of data and material*

355 The data that support the findings of this study are available in NCBI GEO
 356 (<http://www.ncbi.nlm.nih.gov/geo/>) under Accession Number GSE83595. The code for the analysis
 357 is available from the code hosting platform GitHub
 358 (<https://github.com/thk2008/methylseqplatformcomparison>)

359 *Competing interests*

360 The authors declare that they have no competing interests.

361 *Funding*

362 DB is funded by grants from the Starr Consortium (I8-A8-132) and Tri-SCI (2013-036).

363 *Authors' contributions*

364 DB, AA Conceived and designed the experiments. MPF, XZ, YL performed the experiments. TJK
 365 performed the bioinformatic analysis and prepared the figures. TJK, DB, AA, analyzed the data.
 366 TJK, DB drafted the manuscript. TJK, DB, AA, MPF reviewed the manuscript. All authors read and
 367 approved the final manuscript.

368 *Acknowledgements*

369

370

371 *Additional file*

372 Additional file1. **Table S1.** Library input details. **Table S2.** Target region properties and CpGs
 373 covered. **Table S3.** Sequencing details. **Figure S1.** Number of CpG-units covered. **Figure S2.**

374 Mean and median coverage per CpG-unit. **Figure S3.** Intra- and Inter- platform CpG-unit overlap
 375 and methylation levels concordance. **Table S4.** Intra- and Inter- platform details. **Figure S4.**
 376 Overlap of exon annotation of CpG-units as UpSet plot. **Figure S5.** Overlap of intron annotation of
 377 CpG-units as UpSet plot. **Figure S6.** Overlap of promoters annotation of CpG-units as UpSet plot.
 378 **Figure S7.** Overlap of CpG island annotation of CpG-units as UpSet plot. **Figure S8.** Overlap
 379 CpG shores annotation of CpG-units as UpSet plot. **Figure S9.** Overlap of unannotated CpG-units
 380 as UpSet plot.

381

382 References

- 383 1. Jones PA. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat*
 384 *Rev Genet.* 2012;13(7):484–92.
- 385 2. The ENCODE Project Consortium. The ENCODE (ENCyclopedia Of DNA Elements)
 386 Project. *Science.* 2004;306(5696):636–40.
- 387 3. Consortium RE, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, et al. Integrative
 388 analysis of 111 reference human epigenomes. *Nature.* 2015;518(7539):317–30.
- 389 4. Harris RA, Wang T, Coarfa C, Nagarajan RP, Hong C, Downey SL, et al. Comparison of
 390 sequencing-based methods to profile DNA methylation and identification of monoallelic
 391 epigenetic modifications. *Nat Biotechnol.* 2010;28(10):1097–105.
- 392 5. Frommer M, McDonald LE, Millar DS, Collis CM, Watt F, Grigg GW, et al. A genomic
 393 sequencing protocol that yields a positive display of 5-methylcytosine residues in individual
 394 DNA strands. *Proc Natl Acad Sci USA.* 1992;89(5):1827–31.
- 395 6. Lister R, Pelizzola M, Dowen RH, Hawkins RD, Hon G, Tonti-Filippini J, et al. Human DNA
 396 methylomes at base resolution show widespread epigenomic differences. *Nature.*
 397 2009;462(7271):315–22.
- 398 7. The BLUEPRINT consortium. Quantitative comparison of DNA methylation assays for
 399 biomarker development and clinical applications. *Nat Biotech.* 2016;34:726–737
- 400 8. Garrett-Bakelman FE, Sheridan CK, Kacmarczyk TJ, Ishii J, Betel D, Alonso A, et al.
 401 Enhanced reduced representation bisulfite sequencing for assessment of DNA methylation
 402 at base pair resolution. *J Vis Exp.* 2015;(96):e52246.
- 403 9. Dodt M, Roehr JT, Ahmed R, Dieterich C. FLEXBAR-Flexible Barcode and Adapter

- Processing for Next-Generation Sequencing Platforms. Biology (Basel). 2012;1(3):895–905.
10. Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. Bioinformatics . 2011;27(11):1571–2.
11. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. Genome Biol . 2009;10(3):R25.
12. R Core Team. R: A Language and Environment for Statistical Computing . Vienna, Austria; 2016. Available from: <https://www.r-project.org/>
13. Huber, W., Carey, J. V, Gentleman, R., et al. Orchestrating high-throughput genomic analysis with Bioconductor. Nat Methods . 2015;12(2):115–21.
14. Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, et al. Software for Computing and Annotating Genomic Ranges. {PLoS} Comput Biol . 2013;9(8).
15. Eklund A. beeswarm: The Bee Swarm Plot, an Alternative to Stripchart . 2016. Available from: <https://cran.r-project.org/package=beeswarm>
16. Gautier L, Cope L, Bolstad BM, Irizarry RA. affy--analysis of Affymetrix GeneChip data at the probe level. Bioinformatics. 2004;20(3):307–15.
17. Team TBD. BSgenome.Hsapiens.UCSC.hg19: Full genome sequences for Homo sapiens (UCSC version hg19). 2014.
18. Bolstad BM. preprocessCore: A collection of pre-processing functions. 2016. Available from: <https://github.com/bmbolstad/preprocessCore>
19. Gehlenborg N. UpSetR: A More Scalable Alternative to Venn and Euler Diagrams for Visualizing Intersecting Sets. 2016. Available from: <https://cran.r-project.org/package=UpSetR>
20. Chen H. VennDiagram: Generate High-Resolution Venn and Euler Plots. 2016. Available from: <https://cran.r-project.org/package=VennDiagram>

Figure legends

Figure 1. Length and overlap of design regions and MspI predicted regions.

A) Boxplot showing the distributions of targeted region's lengths. SSMethylSeq and MspI 84-334b (ERRBS) show similar region proportions, while CpGiant regions are generally longer and are more variable in length. **B)** Venn diagrams showing the pairwise overlap comparison of each of the platform's regions and the total number of regions. Circle size is proportional to the size of the

435 set. The two capture platforms have a higher degree of common regions with each other than
436 either one with MspI 84-344b.

437

438 **Figure 2. Read alignment and strand parity.**

439 **A)** Percent alignment of uniquely aligned reads (green), ambiguously mapped reads (grays),
440 reads with no alignment (pink) and rejected reads (blue). **B)** The fraction of CpG's covered at
441 $\geq 10\times$ coverage grouped by strand, plus strand (blue) and the minus strand (pink). The strand
442 specific protocol of SSMethylSeq platform is evident by the high proportion of reads mapped to
443 the negative strand.

444

445 **Figure 3: Strand symmetry of methylation values MA-plots.**

446 MA-plots of the log average of the methylation levels (A) on the x-axis and log ratio of the
447 methylation levels (M) on the y-axis, between complementary CpG positions. Median absolute
448 deviation (MAD) values are used to evaluate the agreement in methylation levels. The bimodal
449 nature of methylation patterns (mostly unmethylated or methylated) is reflected in the high density
450 at both ends of the x-axis. The artificially discordant sites introduced during ERRBS library
451 preparation are identified as increased density off the center line at the low methylation values (at
452 $A < 0$ range) in panels A, B. Panels are: **A)** ERRBS_A, **B)** ERRBS_B, **C)** SSMethylSeq_A, **D)**
453 SSMethylSeq_B, **E)** CpGiant_A, **F)** CpGiant_B, and **G)** WGBS.

454

455 **Figure 4. Platform CpG-unit region coverage and CpG-unit overlap.** **A)** Number of CpG-units
456 identified in targeted and off target regions by each platform. WGBS (orange) covers $\sim 14.4\text{M}$
457 CpG-units however there is no notion of targeted regions in this platform. The targeted platforms
458 predicted total CpG-units are depicted as gray bar and coverage of the CpG-units in the predicted
459 regions (a.k.a “on-target”) are shown in blue. CpG-units outside the predicted set (off-target) are

shown in red bars. The red square points and right scale represent the percent recovery of targeted CpG-units. **B)** Distribution of coverage per CpG-unit. **C).** Density plots of the percent of the dataset's regions (color scale) with the fraction of CpG-units covered per region. These plots demonstrate that the reduced number of recovered CpG-units in ERRBS relative to the other platforms (red squares in panel A) are attributed to increased number of missed CpG-units shown as increased density at 0 region. **D)** Triple Venn diagram showing the proportion of overlap of CpG-units for the three platforms; size is not proportional to dataset.

Figure 5. Inter-platform CpG-unit overlap and methylation levels concordance. The upper triangle shows the total number of CpG-units (blue) and number of overlapping CpG-units between any two samples (red) with the percent overlap indicated. The lower triangle shows MA-plots of common CpG-units between any two samples, where M is the log ratio of the methylation levels and A is the average log methylation level from the two platforms. There are blue clouds that at log scale show the variance in methylation at low levels. See Additional file 1: Figure S3 for all pairwise comparisons.

Figure 6: Summary of coverage and representation of annotated genomic regions by each platform. **A)** The fraction of genomic feature covered by sample CpG-units where at least one CpG-unit is covered in a region. **B)** The fraction of a region's CpG-units covered, and **C)** The fraction of CpG-units annotated with a genomic feature. CpGi (CpG island).

Figure 7: CpG-unit annotations overlap. Overlaps of all CpG-unit annotations across all platforms. The UpSet visualization technique [19] for set intersections is displayed as matrix layout. Horizontal bars on the lower left indicate the total number of annotated CpG-units in the set. Dark circles in the matrix (lower right) indicate sets that are part of the intersection. Bars in the

485 main plot area (upper right) indicate the number of intersecting CpG-units for the sets represented
486 by the dark circles. Roughly 33% of the annotations are common to all three suggesting that while
487 there may be similar proportions of CpG-unit annotations (i.e. they may be covering similar
488 regions), they are covering different loci within those regions.

Figure 1

A

B

Region overlap

Figure 2

Figure 3

Figure 4

A

B

C

Percent of target regions with percent CpG-unit coverage

D

Figure 5

ERRBS_A

ERRBS_A::SSMethylSeq_A_opt

SSMethylSeq_A

ERRBS_A::CpGiant_A_opt

SSMethylSeq_A_opt::CpGiant_A_opt

CpGiant_A

Figure 6

A

Region coverage

B

Region CpG-unit coverage

C

Sample CpG-units annotations

Figure 7

Annotation overlap