

1 **MtrA is an essential regulator that coordinates antibiotic production**
2 **and development in *Streptomyces* species**

3

4 Nicolle F. Som^{1#}, Daniel Heine^{2#}, John T. Munnoch¹, Neil A. Holmes¹, Felicity Knowles¹,
5 Govind Chandra², Ryan F. Seipke^{1,3}, Paul A. Hoskisson⁴, Barrie Wilkinson^{2*} and Matthew I
6 Hutchings^{1*}

7

8 # Authors contributed equally; * Correspondence

9

10 ¹School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich,
11 United Kingdom. NR4 7TJ.

12

13 ²Department of Molecular Microbiology, John Innes Centre, Norwich Research Park, Norwich,
14 United Kingdom. NR4 7TJ.

15

16 ³ School of Molecular & Cellular Biology, Astbury Centre for Structural Molecular Biology,
17 University of Leeds, Leeds, LS2 9JT, UK

18

19 ⁴Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, 161,
20 Cathedral Street, Glasgow, G4 0RE, UK

21

22

23

24

25 **Abstract.** *Streptomyces* bacteria make numerous secondary metabolites, including half of all
26 known antibiotics. Production is coordinated with their complex life cycles but the regulators
27 that coordinate development with antibiotic biosynthesis are largely unknown. This is
28 important to understand because most *Streptomyces* secondary metabolites are not produced
29 under laboratory conditions and unlocking the 'cryptic' biosynthetic gene clusters (BGCs) is a
30 major focus for antibiotic discovery. Here we characterise the highly conserved actinobacterial
31 response regulator MtrA in *Streptomyces* species. MtrA is essential and regulates cell cycle
32 progression in *Mycobacterium tuberculosis*. We show that MtrA is also essential in
33 *Streptomyces venezuelae* where it controls genes required for DNA replication and cell
34 division. MtrA also directly regulates the expression of genes in >70% of its BGCs and
35 artificially activating MtrA switches on the production of antibiotics in *S. coelicolor* and *S.*
36 *venezuelae*. We propose that MtrA coordinates antibiotic production and development in
37 *Streptomyces* species.

38

39 **Introduction.**

40 *Streptomyces* secondary metabolites account for two thirds of all known antibiotics and
41 numerous other compounds that are used in human medicine as anticancer, anti-parasitic,
42 antiviral and immunosuppressant drugs. Discovery of these natural products peaked in the
43 1950s but there has been a resurgence of interest in the 21st century, driven by genome
44 sequencing and the increasing threat of drug resistant infections (Katz & Baltz 2016). Despite
45 their importance however, we still have a poor understanding of how *Streptomyces* bacteria
46 control the production of their secondary metabolites. This is important because $\geq 75\%$ of their
47 secondary metabolite biosynthetic gene clusters (BGCs) are not expressed in laboratory
48 culture and activating cryptic BGCs could facilitate the discovery of new antibiotics and other
49 useful compounds (Hosaka et al. 2009; Rutledge & Challis 2015).

50 The major way in which bacteria sense and respond to their environment is through
51 two-component systems and several have been implicated in the regulation of antibiotic
52 production in *Streptomyces* species (Hsiao & Kirby 2009). Two component systems typically

53 consist of a bifunctional sensor kinase and a cognate response regulator (Salazar & Laub
54 2015). The sensor kinase perceives an extracellular signal and activates its cognate response
55 regulator through a two-step phosphorylation. The phosphorylated regulator (RR~P) brings
56 about a response to the original signal, usually by modulating target gene expression. In the
57 absence of signal, the bifunctional sensor kinase dephosphorylates its response regulator to
58 keep the response switched off (Capra & Laub 2012). The delicate balance of kinase and
59 phosphatase activities is crucial in modulating the activity of the response regulator and its
60 target genes during the bacterial cell cycle (Salazar & Laub 2015). Cross-talk between two
61 component systems is rare in wild-type cells but removal of a sensor kinase can result in
62 constitutive activation of its response regulator by a non-cognate sensor kinase or by the
63 cellular pool of acetyl phosphate (Hutchings, Hong & Buttner 2006). Similarly, altering a sensor
64 kinase to block its phosphatase activity can result in a response regulator that cannot be
65 dephosphorylated and is rendered constitutively active (Salazar & Laub 2015).

66 Here we report characterisation of the highly conserved actinobacterial response
67 regulator MtrA in the model organism *Streptomyces venezuelae* (Hoskisson & Hutchings
68 2006). MtrA was first identified as an essential regulator in *Mycobacterium tuberculosis*
69 (Mycobacterium tuberculosis regulator A) (Zahrt & Deretic 2000). *M. tuberculosis* MtrA (TB-
70 MtrA) regulates the expression of *dnaA* and *dnaN*, which are essential for DNA replication,
71 and sequesters the origin of DNA replication, *oriC*, in dividing cells (Purushotham et al. 2015).
72 TB-MtrA also regulates the expression of cell division genes and interacts directly with the
73 DnaA protein, which presumably prevents the initiation of DNA replication (Purushotham et al.
74 2015). The MtrB sensor kinase is activated following localisation to the site of cell division
75 through interaction with FtsI and DivIVA (Wag31) and these data have led to a model in which
76 oscillations in TB-MtrA~P levels play a key role in cell cycle progression by repressing DNA
77 replication and activating cell division (Purushotham et al. 2015; Plocinska et al. 2012;
78 Plocinska et al. 2014). An accessory lipoprotein called LpqB modulates the activity of MtrB in
79 mycobacteria and an *M. smegmatis* Δ *lpqB* mutant grows as *Streptomyces*-like filaments
80 (Nguyen et al. 2010). In the closely related *Corynebacterium glutamicum*, Cg-MtrA controls

81 genes involved in cell wall remodelling and the osmotic stress response (Brocker, Mack & Bott
82 2011a; Krämer 2009) but deletion of the *mtrAB* genes gives rise to elongated cells which are
83 indicative of a defect in cell division (Möker et al. 2004).

84 *Streptomyces* bacteria are filamentous saprophytes which grow through the soil as
85 branching substrate mycelium that extends at the hyphal tips. Nutrient starvation triggers the
86 production of reproductive aerial hyphae that accelerate DNA replication, generating up to 200
87 copies of the chromosome in each aerial hypha, before undergoing cell division to form chains
88 of unigenomic spores. Aerial hyphae production and sporulation are coordinated with the
89 production of bioactive secondary metabolites, including antibiotics. *S. coelicolor* has
90 traditionally been used to study development and antibiotic production because it makes
91 pigmented antibiotics and spores but it only differentiates into aerial hyphae and spores when
92 grown on solid agar. *S. venezuelae* has recently emerged as an alternative model for studying
93 development because it completes a full developmental life cycle in liquid growth medium in
94 ~20 hours (Bush et al. 2015). Here we present evidence that MtrA is essential for the growth
95 of *S. venezuelae* and that it directly regulates the expression of genes involved in DNA
96 replication, cell division and secondary metabolism, several of which (*dnaA*, *dnaN*, *oriC* and
97 *wblE/whiB1*) are also regulated by TB-MtrA. We performed ChIP-seq throughout the *S.*
98 *venezuelae* life-cycle and found that the DNA binding activity of Sv-MtrA displays a biphasic
99 plasticity such that it is active during vegetative growth and sporulation, but inactive during
100 formation of aerial hyphae. Sv-MtrA also binds to sites spanning ~85% of the BGCs in *S.*
101 *venezuelae* and directly modulates expression of target genes in at least 75% of these BGCs.
102 ChIP-seq in *S. coelicolor* revealed that Sc-MtrA binds to many of the same cell division targets
103 and to genes spanning ~70% of its BGCs. Key developmental genes bound by MtrA in both
104 streptomycetes include *bldM*, *ftsZ*, *ssgA*, *smc*, *smeA*, *whiI*, *whiB*, *whiD* and *wblE*. Deletion of
105 *mtrB* constitutively activates MtrA and its target genes and leads to constitutive production of
106 antibiotics in *S. coelicolor* and *S. venezuelae*. Finally, we show that introducing TB-MtrA and
107 gain of function TB-MtrA and Sv-MtrA proteins into *S. venezuelae* also activates antibiotic

108 production. We propose that manipulating the activity of MtrA could be a route to the activation
109 of cryptic BGCs in all *Streptomyces* species.

110

111 **Results**

112 **MtrA is essential in *Streptomyces venezuelae*.**

113 To investigate the function of the MtrAB two component system in *S. venezuelae* we deleted
114 either *mtrA* or *mtrB* (Gust et al. 2003). *mtrB* was deleted easily but all attempts to delete *mtrA*
115 were unsuccessful until we introduced a second copy of *mtrA* *in trans*. The same result was
116 reported for TB-MtrA and the authors concluded that it is essential in *M. tuberculosis* (Zahrt &
117 Deretic 2000). It follows that if MtrA is essential in these bacteria then it must be active in the
118 absence of MtrB, otherwise deleting *mtrB* would also be lethal. We propose that deletion of
119 *mtrB* leads to constitutive phosphorylation of MtrA, by the cellular pool of acetyl phosphate or
120 another sensor kinase. MtrA~P has been shown to be the active form for DNA binding in both
121 *C. glutamicum* and *M. tuberculosis* (Fol et al. 2006; Brocker, Mack & Bott 2011b). We
122 observed a similar result with *Streptomyces coelicolor* VanRS, where VanR and its target
123 genes are constitutively active in a $\Delta vanS$ mutant (Hutchings, Hong & Buttner 2006).
124 Surprisingly, deletion of *mtrB* has no effect on growth rate or development in liquid cultures
125 but in agar grown cultures, deletion of $\Delta mtrB$ results in a sporulation defect in the centre of the
126 colonies (Figures S1 and S2).


127

128 **MtrA activity changes during the life cycle of *S. venezuelae*.**

129 Published microarray data for *S. venezuelae* NRRL B-65442 (Bibb et al. 2012) (ArrayExpress
130 accession E-MEXP-3612), shows that *mtrA* expression levels are highest during vegetative
131 growth at 10 and 12 hours and drop approximately 2-fold at 14-20 hours growth (Figure S3).
132 RNA sequencing at 12, 16 and 20 hours (accession number = GSE81104), shows a dip in
133 transcript levels at 16 hours (aerial growth) and an increase following sporulation, at 20 hours
134 (Figure S4). Thus, the *mtrA* gene appears to be most highly transcribed during active growth
135 and following sporulation. We did not detect any significant change in MtrA protein levels

136 during the life cycle using immunoblotting (Figure S3) and attempts to compare levels of
137 phosphorylated and non-phosphorylated MtrA using Phostag were unsuccessful (not shown).
138 To examine MtrA activity during the *S. venezuelae* life cycle, we performed ChIP-seq
139 throughout development. We introduced an expression construct for MtrA-3xFlag *in trans*
140 under the control of its own promoter and then deleted the native *mtrA* gene. Survival of this
141 strain shows that MtrA-3xFlag is functional and there are no significant difference in growth
142 rates between this strain and the wild-type (Figures S5 and S6). We performed ChIP-seq on
143 the wild-type (control) and the MtrA-3xFlag strain at 8, 10, 12, 14, 16, 18 and 20 hours to
144 examine the DNA binding activity of MtrA at key stages of the developmental life cycle
145 (accession number GSE84311, Table S1). The 8-12h time points correspond to the active
146 growth of substrate hyphae, between 14-18 hours they produce aerial hyphae and undergo
147 DNA replication and cell division to form spores and at 20 hours the culture is made up of
148 immature and mature spore chains. We mapped the reads to the *S. venezuelae* NRRL B-
149 65442 genome sequence (Genbank accession CP018074). Analysis of the ChIP-seq data
150 using a P value > 0.05 revealed that only one target is enriched at all time points, the promoter
151 region of the *ectABCD* operon, which is the BGC for the secondary metabolites ectoine and
152 5' hydroxyectoine (5HE). We could not detect ectoine or 5HE in the wild-type or $\Delta mtrB$ mutant
153 using LCMS with authentic standards so we assume that MtrA~P represses *ectABCD*
154 expression throughout the life cycle. Analysis of all the ChIP-seq datasets using $P > 0.05$
155 shows that the pattern of binding of MtrA is dynamic throughout the lifecycle with peak binding
156 activity seen at 10 and 20 hours, coinciding with mid-exponential growth and within spores
157 (Figure S7). This is more consistent with *mtrA* expression levels observed in the dRNA-seq
158 experiment than in the microarray data (Figures S3-4). This may be because the microarray
159 experiments were performed in a different laboratory so growth rates may vary whereas we
160 performed the RNA-seq experiments under identical conditions. However, these data show
161 that the DNA binding activity of MtrA changes during the life cycle and mapping ChIP-seq
162 reads at targets such as the *mtrA* promoter provides a good illustration of this biphasic activity
163 (Figure 1). Differential RNA sequencing also shows that the *mtrAB-lpqB* genes are transcribed

164 as a single leaderless transcript and that MtrA binds immediately upstream of the -35 region,
165 suggesting positive autoregulation. There is a second putative transcript start site at -79bp,
166 which is upstream of the MtrA binding site, suggesting there may be two promoters driving
167 expression of *mtrAB-lpqB* (Figure 1).
168



169

Figure 1. MtrA autoregulates its own expression. ChIP-seq during the *Streptomyces venezuelae* life cycle shows that MtrA has highest activity in actively growing mycelium (10 and 12 hours) and in spores (20 hours). The peaks shown are at the *mtrAB-lpqB* operon promoter region which has two transcript start sites at +1 (P1) and -79 (P2). The position of the MtrA ChIP peak suggests it activates P1 and blocks P2.

170

171 **Identifying the MtrA binding site.** To identify an MtrA consensus binding sequence we
172 performed MEME analysis (Bailey et al. 2015) on three MtrA target promoters identified by
173 ChIP-seq and bound by purified MtrA using *in vitro* electrophoretic mobility shift assays
174 (EMSAs). These analyses suggested that MtrA recognises and binds to an AT-rich 7 bp motif.
175 We then took 50 bp of sequence from beneath each peak at the 14h and the 16/18h time
176 points and ran them through MEME to identify conserved motifs. This analysis suggests that

177 MtrA binds a direct repeat of the AT rich motif separated by a 5bp spacer (Figure 2). However,
178 many of the sequences enriched in the MtrA ChIP-seq dataset do not contain this motif and
179 we hypothesise that MtrA binds target sites directly and also by interacting with other
180 transcription factors. Immunoprecipitation experiments using anti-Flag beads pulled down four
181 different regulators in the MtrA-3xFlag cultures that are not immunoprecipitated in the wild-
182 type control. The genes encoding these regulators (*sven15_0243*, *sven15_2691*,
183 *sven15_3571* and *sven15_4644*) are also targets for MtrA (Table S1). *Sven15_3571* is DnaA,
184 which initiates DNA replication and acts as a transcription factor in bacteria. TB-MtrA also
185 interacts with DnaA although the effects of this interaction are not known (Purushotham et al.
186 2015). Thus, our data suggests (but does not prove) that MtrA forms complexes with other
187 DNA binding proteins and this may explain why there are so many enrichment peaks in the
188 20-hour ChIP-seq dataset. Interaction between regulatory proteins is not unprecedented in *S.*
189 *venezuelae*. The response regulator BldM modulates one set of target genes as a homodimer
190 and another set by forming heterodimers with the WhiI response regulator (Bassam et al.
191 2014). The developmental regulators WhiA and WhiB also proposed to interact and have
192 identical ChIP-seq regulons (Bush et al. 2016).



193

194 **MtrA regulates the expression of DNA replication and cell division genes.**

195 ChIP-seq shows that MtrA activity changes during the *S. venezuelae* life cycle (Figure 1). RNA
196 sequencing of wild-type and $\Delta mtrB$ strains shows that expression of *mtrA* is 3-fold upregulated
197 in the $\Delta mtrB$ mutant (Table S2) which is consistent with activation by MtrA~P. Given the
198 essentiality of MtrA and its altered activities at different stages of the life cycle we predicted
199 that it could play a key role in regulating cell cycle progression in *S. venezuelae*, as in *M.*
200 *tuberculosis* (Purushotham et al. 2015; Plocinska et al. 2014). Indeed, ChIP-seq shows that
201 several key developmental genes are bound by MtrA in *S. venezuelae* and *S. coelicolor* (see
202 Tables S1 and S2 for complete lists).

203


Target	P value	Binding site
Sven15_0205 (<i>ectA</i>)	4.23e-5	<u>GCGTTACCCAT</u> CCGTTA TAAACGGTCAT
Sven15_0880 (<i>cmlF</i>)	4.23e-5	ATT CGCCATCTACGGTTC CGGGCTGGCC
Sven15_3571 (<i>dnaA</i>)	1.51e-5	CACAATCTTT CCGTTCT GTCCTTACCT

204

Figure 2. MtrA binds to 6-5-6bp direct repeat motif. The table shows three MtrA ChIP targets that are bound *in vitro* by purified MtrA protein. They are the promoter regions of *ectABCD* operon (ectoine BGC), *cmlF* (chloramphenicol transporter) and *dnaA* (transcription factor and initiator of DNA replication). MEME analysis identified a conserved 7bp motif (coloured) that is present as a 6-5-6bp repeat motif at the *ectA* promoter (second half underlined). MEME analysis of different subsets of ChIP targets identified the same direct repeat motif (shown for the targets identified at 14 hours and 16 and 18 hours).

205 Targets common to both TB- and Sv-MtrA are the promoter regions of *wbIE*, *dnaA* and *dnaN*,
 206 and the *oriC* region between *dnaA* and *dnaN* (Figure 3A). RNA-seq shows that deletion of
 207 *mtrB* does not affect the expression levels of *dnaA* or *dnaN* under the conditions used but
 208 deletion of *mtrB* causes a two-fold increase in *wbIE* transcript levels, suggesting MtrA-
 209 dependent activation. MtrA also directly represses *adpA* expression, which is down-regulated
 210 nearly 4-fold in the $\Delta mtrB$ mutant (Table S2). AdpA controls key developmental genes and
 211 represses DNA replication in *Streptomyces* species (Wolański et al. 2012; Wolański et al.
 212 2011). *wbIE* encodes a WhiB-like transcription factor of unknown function that is homologous
 213 to WhiB1 in *M. tuberculosis*. Both *wbIE* and *whiB1* are essential (Stapleton et al. 2012). Sv-
 214 MtrA also binds upstream of the cell division genes *ssgA* and *ssgB*, which are upregulated in
 215 the $\Delta mtrB$ mutant, and *ftsZ* which is unaffected by loss of MtrB, probably because this gene
 216 is subject to complex regulation (Figure 3B and Table S2).



217

Figure 3. MtrA binds to targets required for DNA replication and cell division in *S. venezuelae*. Top. MtrA ChIP peaks upstream of *dnaA*, which encodes the initiator of DNA replication, and *dnaN* which encodes the DNA polymerase beta clamp subunit and at the origin of DNA replication, *oriC*. Bottom. MtrA ChIP peaks upstream of *ftsZ*, which encodes the Z ring forming FtsZ protein and *ssgA*, whose product helps localise FtsZ to the sites of cell division during sporulation.

218

219 SsgAB localise the divisome marker protein FtsZ to the correct positions in aerial hyphae to
220 mark the sites of cell division, prior to sporulation (Schwedock et al. 1997; Willemse et al.
221 2011). Sv-MtrA also regulates expression of the *smeA-ssfA* operon, which is elevated
222 between 2- and 4-fold in the $\Delta mtrB$ mutant. SmeA targets the DNA pump SffA to the cell
223 division septa (Bush et al. 2015). MtrA also binds upstream of (and activates some) key
224 developmental regulators, including *whiB*, *whiD*, *whiI* and *bldM* (Table S2) (Bassam et al.
225 2014; Molle et al. 2000; Bush et al. 2016). Many of these Sv-MtrA target genes are also
226 repressed by BldD (in complex with c-di-GMP), a master regulator of *Streptomyces*
227 development. Sv-MtrA does not regulate *bldD* expression but it does regulate the expression
228 of genes encoding proteins that metabolise cyclic di-GMP, which is required for BldD-
229 mediated repression (Tschowri et al. 2014) (Table S2).

230

231 **MtrA regulates global BGC expression in *Streptomyces venezuelae*.**


232 ChIP-seq shows that MtrA binds to sites spanning 28 of the 31 BGCs predicted by antiSMASH
233 3.0 in the *S. venezuelae* genome (Weber et al. 2015) (Table S3). The only BGCs with no MtrA
234 enrichment are those encoding biosynthesis of the desferrioxamine siderophores, the WhiE
235 polyketide spore pigment and a putative insecticidal complex. Of these three clusters, the
236 WhiE BGC is upregulated in the $\Delta mtrB$ mutant suggesting indirect regulation by MtrA, probably
237 via BldM (Bassam et al. 2014). The other two BGCs are unaffected by loss of MtrB (Table
238 S3). Of the 28 BGCs that are bound by MtrA, nine have genes that are positively regulated by
239 MtrA, 10 have genes that are negatively regulated by MtrA and three are subject to both
240 positive and negative regulation by MtrA at individual genes within the gene cluster (Table
241 S3). The other six BGCs have sites that are enriched in MtrA ChIP-seq but expression of the
242 genes nearest the ChIP-seq peaks are not affected by deletion of *mtrB* under the conditions
243 we used (Table S3). We predict the *ectABCD* operon is repressed by MtrA because the *ectA*
244 promoter is bound by MtrA at all time points. Consistent with this, we cannot detect ectoine or
245 5HE in the wild-type or $\Delta mtrB$ strains. Since most of the *S. venezuelae* BGCs are

246 uncharacterised we know little about their cluster specific regulation, or their natural products.
247 It should also be noted that the antiSMASH predictions we have used here likely include
248 additional flanking genes that may not be part of the BGC so the actual number of BGCs
249 bound by MtrA may be slightly lower.

250

251 **MtrA activates chloramphenicol production in *S. venezuelae*.**

252 Phenotypic screening of the $\Delta mtrB$ mutant revealed that a cryptic antibacterial is activated by
253 removing MtrB whereas over-expressing *mtrA* in wild-type *S. venezuelae* has no effect (Figure
254 S8). This is probably because MtrA~P levels only increase in the absence of MtrB. *S.*
255 *venezuelae* encodes the biosynthetic pathway for the antibiotic chloramphenicol, which
256 according to previous studies is off in the wild-type strain (Fernández-Martínez et al. 2014).
257 CHIP-seq shows that Sv-MtrA binds in the intergenic region between the divergent *cmIF*
258 (*sven15_0879*) and *cmIN* (*sven15_0880*) genes (Figure 4) and expression of *cmIN* is 6-fold
259 upregulated in the $\Delta mtrB$ mutant, consistent with direct activation by MtrA. CmlN is an efflux
260 permease, predicted to export chloramphenicol (Fernández-Martínez et al. 2014). HPLC
261 confirmed that chloramphenicol is produced in the $\Delta mtrB$ mutant but we also detected very
262 low levels in the wild-type strain suggesting the cluster is not completely silent (Figure 4).
263 Cultivation for 24 hours in biological and technical triplicates confirmed an increased
264 production of chloramphenicol in the $\Delta mtrB$ mutant with a mean concentration of 0.41 mg/L
265 which is >30 times higher than the wild-type strain (0.013 mg/L) or wild-type over-expressing
266 *mtrA* (0.010 mg/L).




267

Figure 4. MtrA controls chloramphenicol production. Top. MtrA ChIP peaks at the chloramphenicol (Cm) BGC show two significant peaks between *cmIN* and *cmIF* and a smaller peak upstream of the regulator *cmIR* which is $P < 0.05$. Middle. Expression data for wild-type and $\Delta mtrB$ strains at 14 and 20 hours shows *cmIF* is 6-fold upregulated in $\Delta mtrB$ suggesting it is activated by MtrA. All the other Cm genes are downregulated in the mutant which may be due to negative feedback. Bottom left. HPLC shows the Cm standard and extracts of wild-type, $\Delta mtrB$, wild-type plus empty pMS82 vector, and wild-type plus pMS82 expressing Sv-Y99C MtrA, TB-MtrA and TB-Y102C MtrA. Right. Zooming in shows that Sv-Y99C MtrA induces production of low levels of Cm production.

268 **Deleting *mtrB* has a global effect on the *S. venezuelae* metabolome.**

269 While loss of MtrB clearly leads to increased production of chloramphenicol, we were
270 interested to know if there are any other effects on the metabolome. We cultivated the wild
271 type strain and three independently isolated $\Delta mtrB$ mutants in biological and technical
272 triplicates and analysed the extracts by UPLC/HRMS using an untargeted metabolomics
273 approach. Runs were aligned to compensate for between-run variation and a peak-picking
274 algorithm was applied to allow for the immaculate matching of each feature (a discrete *m/z*
275 value and its retention time) among all runs. Following normalisation, features could be
276 compared quantitatively and their putative identity proposed based on their high-resolution
277 MS-signal. Comparing the level of metabolite signals, it appeared that all $\Delta mtrB$ mutants
278 showed increased production of a considerable portion of their putative secondary
279 metabolites. To display multidimensional data, we used Principle Component Analysis (PCA)
280 (Figure 5). Each sphere in the 3D Plot represents one dataset obtained from a particular
281 UPLC-HRMS run. Data from the $\Delta mtrB$ mutant strains clearly group together, and are distinct
282 from data obtained from the wild type, while variations within each group are comparably
283 small. The 3D Plot therefore shows consistent and global changes in the metabolome upon
284 loss of MtrB (Figure 5).

285



286

Figure 5. Loss of MtrB results in a global shift in the *S. venezuelae* metabolome.

Principle Component Analysis on the wild-type (blue dots) and triplicate samples of the $\Delta mtrB$ strain (red, green and orange dots). Data from $\Delta mtrB$ mutant strains clearly group together, and are distinct from data obtained from the wildtype while variations within each group are comparably small.

287

288 MtrA directly activates antibiotic production in *Streptomyces coelicolor*.

289 MtrA is conserved in all sequenced *Streptomyces* genomes so we reasoned that it might
290 activate antibiotic production in other streptomycetes. To test this, we deleted *mtrB* in the
291 model organism *Streptomyces coelicolor* which is probably the best characterised species in
292 terms of its secondary metabolite BGCs (Challis 2013). The 16S rDNA phylogenetic tree of
293 the family *Streptomycetaceae* shows that *S. venezuelae* (clade 40) is highly divergent from *S.*
294 *coelicolor* (clade 112) so we reasoned that if deleting *mtrB* switches on antibiotic production
295 in these distantly related species it may be universal to all streptomycetes (Labeda et al. 2011).
296 The *S. coelicolor* $\Delta mtrB$ mutant forms small colonies that are delayed in sporulation but normal
297 growth and differentiation are restored by reintroducing *mtrB* *in trans* (Figures S9-10). The
298 $\Delta mtrB$ mutant over-produces the red antibiotic undecylprodigiosin and the blue antibiotic


299 actinorhodin and we also detected significant amounts of metacycloprodigiosin, a potent
300 anticancer compound encoded by the undecylprodigiosin BGC but not previously reported
301 from *S. coelicolor* (Figure 6). Other secondary metabolites are clearly down-regulated in the
302 $\Delta mtrB$ mutant, including the desferrioxamine B and E siderophores and germicidin A. We
303 performed ChIP-seq in *S. coelicolor* because its BGCs and their natural products are the most
304 well defined of any *Streptomyces* species (Bentley et al. 2002; Challis 2013; van Keulen &
305 Dyson 2014). The results show that MtrA binds to sites spanning 21 of its 29 BGCs (Tables
306 S4 and S5). MtrA binds upstream of the genes encoding the actinorhodin activator ActII-4 and
307 the undecylprodigiosin activator RedZ (Figure S11). MtrA does not bind to the desferrioxamine
308 BGC (*desABCD*) in *S. coelicolor* (Table S4) but does bind upstream of *sco4394*, which
309 encodes DesR, an iron dependent repressor of desferrioxamine biosynthesis (Figure S11). A
310 single type III polyketide gene, *SCO7221*, is responsible for germicidin biosynthesis (Song et
311 al. 2006) and it is not regulated by MtrA, so the effect on germicidin biosynthesis must be
312 indirect.

313

314 ***M. tuberculosis* MtrA activates chloramphenicol production in *S. venezuelae*.**

315 Modulating MtrA activity by deleting *mtrB* to switch on antibiotic production is time consuming
316 and impractical, particularly when applied to multiple *Streptomyces* strains. To test whether
317 gain-of-function MtrA proteins might be used instead we constructed *Streptomyces* expression
318 vectors for wild-type TB-MtrA, gain-of-function Y102C TB-MtrA (Satsangi et al. 2013) and a
319 Y99C Sv-MtrA variant which has the equivalent change to Y102C TB-MtrA (Figure S12). We
320 were unable to delete the native *mtrA* gene in strains carrying these constructs but all three
321 induced chloramphenicol production in the wild-type strain (Figure 4). Y99C Sv-MtrA switches
322 on the production of relatively small amounts of chloramphenicol in wild-type *S. venezuelae*
323 but the TB-MtrA proteins switch on much higher levels of production, with the strain expressing
324 Y102C-TB-MtrA making more chloramphenicol than the $\Delta mtrB$ mutant (Figure 4). These
325 results suggest that expression of different MtrA proteins is a useful avenue to explore in terms
326 of activating BGCs in *Streptomyces* species and could be useful tools for the community. The

327 MtrA expression constructs described in this work are available from AddGene (IDs 85988-
328 94).
329



330

Figure 6. Deleting *mtrB* switches on antibiotic production in *S. coelicolor*. (a) HPLC on extracts of wild-type *S. coelicolor* M145 cultures shows production of desferrioxamines and germicidin. (b) HPLC on extracts of an *S. coelicolor* $\Delta mtrB$ mutant shows production of actinorhodin and undecylprodigiosin is switched on while the production of the desferrioxamine siderophores and germicidin are switched off.

331

332 Discussion.

333 This work describes the first detailed analysis of the highly conserved actinobacterial response
334 regulator MtrA in *Streptomyces* species. As with most response regulators, MtrA activity is
335 controlled through phosphorylation and dephosphorylation by its cognate sensor kinase, MtrB.
336 Our data, and previously published work on MtrA in mycobacteria and corynebacteria, support
337 a model in which removing MtrB results in constitutive phosphorylation and activation of MtrA
338 which then modulates the expression of its target genes in the cell. In *S. coelicolor* and *S.*
339 *venezuelae* these include key developmental genes and there is a significant overlap with the
340 regulon controlled by the master repressor of development, BldD, which is activated for DNA
341 binding by the secondary messenger cyclic-di-GMP (Tschowri et al. 2014). Exactly where

342 MtrA fits in to the hierarchy of developmental regulation is not yet clear and there is much work
343 to be done to determine the role of the MtrAB two-component system in the cell biology of
344 *Streptomyces* species. However, our data show that MtrA has an essential role in *S.*
345 *venezuelae* and shares some functions with *M. tuberculosis* MtrA. It will be interesting to
346 compare the complete overlap when the TB-MtrA regulon is published (Purushotham et al.
347 2015). Loss of MtrB has minor effects on growth and sporulation in *S. venezuelae* and it will
348 be interesting to deplete MtrA activity in future studies to examine the effects on growth,
349 development and secondary metabolite production. Here we have focused on the role of MtrA
350 in regulating secondary metabolism and how it might be exploited to activate cryptic BGCs.
351 The effects on antibiotic production appear to be direct in both *S. coelicolor* and *S. venezuelae*
352 because MtrA activates expression of the chloramphenicol transporter CmlN and activation of
353 MtrA switches on chloramphenicol production. It also binds to the promoters of the cluster
354 specific activators ActII-4 and RedZ in *S. coelicolor* and activates the products under their
355 control, actinorhodin and undecylprodigiosin, respectively. In a developmental time course it
356 is remarkable that MtrA binds to sites spanning 27 of the 31 BGCs in *S. venezuelae* and
357 directly affects the expression of target genes in at least 22 of these BGCs. In vegetatively
358 growing *S. coelicolor* MtrA binds to sites spanning 21 out of 29 BGCs. We propose that MtrA
359 is a master regulator of secondary metabolism in *Streptomyces* species. Consistent with this,
360 unbiased metabolomics analysis of *S. venezuelae* shows that deleting *mtrB* results in a global
361 change in the metabolome. Given that secondary metabolite BGCs are often subject to
362 complex and multilevel regulation, and given the energy costs associated with making these
363 natural products, it is perhaps not surprising that we do not see obvious over-production of
364 other compounds in the $\Delta mtrB$ strains. We predict that deletion of known BGCs in the $\Delta mtrB$
365 mutants and / or depletion of *mtrA* transcripts will have positive effects on the production of
366 additional secondary metabolites.

367 In summary, our study has revealed an important role for MtrA, a previously
368 uncharacterised but essential response regulator, in the life cycles of streptomycetes.
369 Significant work remains to be done to elucidate its role in *Streptomyces* species but our data

370 suggest it might play a similar role to CtrA in *Caulobacter crescentus*, acting as a master
371 regulator of the cell cycle alongside other master regulators such as BldD. Perhaps even more
372 important is the fact that it coordinates production of antibiotics and other natural products with
373 the developmental programme. Rapid progress has been made recently in elucidating the
374 roles of such developmental regulators, using next generation sequencing techniques and *S.*
375 *venezuelae* as a model system (Bush et al. 2015). We have shown that *S. venezuelae* can
376 also be used as a model to study global effects on secondary metabolism and to elucidate the
377 regulatory cascades that link secondary metabolism and differentiation. Understanding these
378 genetic circuits is essential if we are to unlock the full potential of these bacteria and their
379 secondary metabolites.

380

381 **Materials and Methods**

382 **Strains, plasmids and primers.** The bacterial strains, plasmids and cosmids and primers
383 used in this study are listed in Tables S6-8. The Sv- and TB-MtrA expression vectors have
384 been deposited with AddGene (ID 85988-94). *S. venezuelae* NRRL B-65442 is deposited in
385 the USDA Agricultural Research Services (ARS) Culture Collection
386 (<http://nrri.ncaur.usda.gov/cgi-bin/usda/prokaryote/report.html?nrriCodes=B%2d65442>).

387 Plasmid stocks were prepared using Qiagen miniprep kits and cosmids were prepared as
388 described previously (Kieser et al. 2000). Genes were deleted using the ReDirect PCR
389 targeting method (Gust et al. 2003) and an *S. venezuelae* NRRL B-65442 cosmid library
390 provided by Professor Mark Buttner at the John Innes Centre, Norwich. All of the expression
391 constructs used in this work were made by Genscript by synthesising the *mtrA* alleles with or
392 without 3' tags and then cloning into the required vectors. Liquid cultures of *E. coli* were
393 routinely grown shaking at 220 rpm in Lennox Broth at 37°C. Liquid cultures of *S. coelicolor*
394 or *S. venezuelae* were grown in Mannitol Yeast Extract Malt Extract (MYM) at 30°C, shaking
395 at 220 rpm. Cultures grown on solid MYM agar were grown at 30°C. Spore stocks of *S.*
396 *coelicolor* were prepared from cultures grown on Soya Flour + Mannitol (SFM) agar. All media
397 recipes have been published previously and *Streptomyces* spores were prepared as

398 described (Kieser et al. 2000). To determine the developmental growth in liquid culture *S.*
399 *venezuelae* and mutant strains were grown, shaking in 35ml MYM in 250ml conical flasks
400 containing springs at 30°C at 220rpm. A spore inoculum sufficient to reach an OD₆₀₀ of 0.35
401 after 8 hours of growth was added to 35ml of MYM media in 250 ml flasks containing springs.
402 The culture densities were measured at OD₆₀₀. Development in liquid cultures was monitored
403 using an GXML3000B microscope from GX optical. Pictures of agar plate grown colonies were
404 taken with a Zeiss SVII stereo microscope. SEM images were taken at the bioimaging facility
405 at the John Innes Centre.

406

407 **Purification of MtrA-His.** The *mtrA* gene was cloned into pETDuet1 (Novagene) to express
408 the protein with a C-terminal hexa-His tag and purified using a batch method. Cell pellets were
409 resuspended in 25ml lysis buffer (75mM Tris-HCl pH8, 20mM NaCl, 0.1% Triton X100, 50 µl
410 10mg/ml lysozyme, 3 x Pierce EDTA-free Protease Inhibitor Mini Tablets (Thermo Scientific)
411 in 1L) and incubated for 30 minutes at room temperature. The cell lysate was sonicated 2 x
412 40 seconds at 50Hz with 1 minute in between sonication steps. Cell debris was removed by
413 centrifugation at 18,000rpm for 20 minutes at 4°C in Beckman Coulter Avanti[®] J-20 high
414 performance centrifuge using a JLA-25-50 rotor (Beckman Coulter). The supernatant was
415 transferred in a fresh 50ml Falcon tube and 350µl of Ni-NTA agarose beads (Qiagen) were
416 added and incubated under gentle agitation for 1 hour at 4°C. The Ni-NTA agarose beads
417 were spun down gently (maximum of 1200rpm, 4°C) and the supernatant was discarded. The
418 Ni-NTA beads were resuspended in 2ml wash buffer (75mM Tris-HCl pH8, 200mM NaCl, 10%
419 glycerol, 10mM MgCl₂, 0.1mM DTT, 20mM 2-mercaptoethanol in 1L) and transferred in
420 Polypropylene columns (1ml, Qiagen). The beads in the column were washed with 20ml wash
421 buffer. The protein was eluted from the beads with 2.5ml elution buffer (wash buffer plus
422 350mL imidazole in 1L). Polyclonal antiserum was raised by Cambridge Research
423 Biochemicals. Immunoblotting was performed as described previously (Hutchings, Hong,
424 Leibovitz, et al. 2006).

425

426 **DNA binding studies.** Chromatin Immunoprecipitation followed by sequencing (ChIP-seq)
427 was performed on *S. coelicolor* M145 $\Delta mtrA$ + MtrA-3xFlag and *S. venezuelae* NRRL B-65442
428 $\Delta mtrA$ + MtrA-3xFlag grown in liquid MYM medium (Kieser et al. 2000). Note that *S. coelicolor*
429 M145 grows vegetatively and does not differentiate under these growth conditions whereas
430 *S. venezuelae* NRRL B-65442 undergoes a full life cycle in 20 hours. Samples were taken at
431 16 and 20 hours for *S. coelicolor* and at 8, 10, 12, 14, 16, 18 and 20 hours for *S. venezuelae*.
432 ChIP-seq and electrophoretic mobility shift assays (EMSAs) were performed as described
433 previously (Crack et al. 2015) using probes generated by PCR with 6-Fam labelled primers
434 from Integrated DNA Technologies (Table S8). ChIP-seq Gels were visualised using a
435 Typhoon FLA 9500 laser scanner (GE Healthcare) with LBP/BPB1 emission filter, Exmax
436 495nm Exmax 576nm, at 50 μ M resolution. ChIP-seq data was analysed as described
437 previously (Bush et al. 2013) and peaks were visually inspected using integrated genome
438 browser (Nicol et al. 2009). Binding sites were identified using MEME (Bailey et al. 2015).

439

440 **RNA-sequencing.** Duplicate wild-type and $\Delta mtrB$ cultures were grown for 14 and 20 hours in
441 MYM medium and RNA was prepared as described previously (Munnoch et al. 2016).
442 Libraries were prepared and sequenced by Vertis Biotechnologie and analysed using CLC
443 Genomics Workbench.

444

445 **LCMS and PCA.**

446 Analytical HPLC was performed on an HPLC 1100 system (Agilent Technologies) connected
447 to a Gemini® 3 μ m NX-C18 110 Å, 150×4.6 mm column (Phenomenex) and on a Synergi™
448 4 μ m Fusion-RP 80 Å LC column 250×10 mm. UPLC-MS for metabolic profiling was performed
449 on a Nexera X2 liquid chromatograph (LC-30AD) LCMS system (Shimadzu) connected to an
450 autosampler (SIL-30AC), a Prominence column oven (CTO-20AC) and a Prominence photo
451 diode array detector (SPD-M20A). A Kinetex® 2.6 μ m C18 100 Å, 100×2.1 mm column
452 (Phenomenex) was used. The UPLC-System was connected with a LCMS-IT-TOF Liquid
453 Chromatograph mass spectrometer (Shimadzu). UPLC-HRMS Data was acquired on an

454 Acquity UPLC system (Waters Corporation) equipped with an ACQUITY UPLC[®] BEH 1.7 μ m
455 C18, 1.0 \times 100 mm column (Waters Corporation) and connected to a Synapt G2-Si high
456 resolution mass spectrometer (Waters Corporation). A gradient between mobile phase A (H₂O
457 with 0.1 % formic acid) and mobile phase B (acetonitrile with 0.1 % formic acid) at a flow rate
458 of 80 μ L/min was used. Initial conditions were 1 % B for 1 min, ramped to 90 % B within
459 6 minutes, ramped to 100 % B within 0.5 min, held for 0.5 min, returned to 1 % B within 0.1 min
460 and held for 1.9 min. MS spectra were acquired with a scan time of one second in the range
461 of m/z = 150 - 1200 in positive MSe-Resolution mode. The following instrument parameters
462 were used: capillary voltage of 3 kV, sampling Cone 40, source offset: 80, source temperature
463 of 120 °C, desolvation temperature 350 °C, desolvation gas flow 800 L/h. A solution of sodium
464 formate was used for calibration. A solution of leucine encephalin (H₂O/MeOH/formic acid:
465 49.95/49.95/0.1) was used as lock mass and was injected every 15 sec. The lock mass has
466 been acquired with a scan time of 0.3 sec and 3 scans were averaged each time. The lock
467 mass (m/z = 556.2766) has been applied during data acquisition. For processing
468 metabolomics data we used the Software Progenesis QI (Waters). All solvents for analytical
469 HPLC and UPLC-MS were obtained commercially at least in HPLC grade from Fisher
470 Scientific and were filtered prior to use. Formic acid (EZInfo 3.0 (MKS Umetrics AB) was used
471 for plotting PCA data.

472

473 **References**

- 474 Bailey, T.L. et al., 2015. The MEME Suite. *Nucleic Acids Research*, 43(W1), pp.W39–49.
- 475 Bassam, Al, M.M. et al., 2014. Response regulator heterodimer formation controls a key
476 stage in *Streptomyces* development. *PLoS genetics*, 10(8), p.e1004554.
- 477 Bentley, S.D. et al., 2002. Complete genome sequence of the model actinomycete
478 *Streptomyces coelicolor* A3(2). *Nature*, 417(6885), pp.141–147.
- 479 Bibb, M.J. et al., 2012. Expression of the chaplin and rodlin hydrophobic sheath proteins in
480 *Streptomyces venezuelae* is controlled by σ (BldN) and a cognate anti-sigma factor, RsbN.

- 481 *Molecular Microbiology*, 84(6), pp.1033–1049.
- 482 Brocker, M., Mack, C. & Bott, M., 2011a. Target genes, consensus binding site, and role of
483 phosphorylation for the response regulator MtrA of *Corynebacterium glutamicum*. *Journal of*
484 *Bacteriology*, 193(5), pp.1237–1249.
- 485 Brocker, M., Mack, C. & Bott, M., 2011b. Target Genes, Consensus Binding Site, and Role
486 of Phosphorylation for the Response Regulator MtrA of *Corynebacterium glutamicum*.
487 *Journal of Bacteriology*, 193(5), pp.1237–1249.
- 488 Bush, M.J. et al., 2015. c-di-GMP signalling and the regulation of developmental transitions
489 in streptomycetes. *Nature Reviews Microbiology*, 13(12), pp.1–12.
- 490 Bush, M.J. et al., 2013. Genes Required for Aerial Growth, Cell Division, and Chromosome
491 Segregation Are Targets of WhiA before Sporulation in *Streptomyces venezuelae*. *mBio*,
492 4(5), pp.e00684–13.
- 493 Bush, M.J. et al., 2016. Genome-Wide Chromatin Immunoprecipitation Sequencing Analysis
494 Shows that WhiB Is a Transcription Factor That Cocontrols Its Regulon with WhiA To Initiate
495 Developmental Cell Division in *Streptomyces*. *mBio*, 7(2), pp.e00523–16.
- 496 Capra, E.J. & Laub, M.T., 2012. Evolution of Two-Component Signal Transduction Systems.
497 *Annual Review Microbiology*, 66(1), pp.325–347.
- 498 Challis, G.L., 2013. Exploitation of the *Streptomyces coelicolor* A3(2) genome sequence for
499 discovery of new natural products and biosynthetic pathways. *Journal of Industrial*
500 *Microbiology & Biotechnology*. 41(2), pp219-32.
- 501 Crack, J.C. et al., 2015. NsrR from *Streptomyces coelicoloris* a Nitric Oxide-Sensing [4Fe-
502 4S] Cluster Protein with a Specialized Regulatory Function. *Journal of Biological Chemistry*,
503 15;290(20), pp12689-704.
- 504 Fernández-Martínez, L.T. et al., 2014. New Insights into Chloramphenicol Biosynthesis in
505 *Streptomyces venezuelae* ATCC 10712. *Antimicrobial Agents and Chemotherapy*, 58(12),

506 pp.7441–7450.

507 Fol, M. et al., 2006. Modulation of *Mycobacterium tuberculosis* proliferation by MtrA, an
508 essential two-component response regulator. *Molecular Microbiology*, 60(3), pp.643–657.

509 Gust, B. et al., 2003. PCR-targeted *Streptomyces* gene replacement identifies a protein
510 domain needed for biosynthesis of the sesquiterpene soil odor geosmin. *Proceedings of the*
511 *National Academy of Sciences USA*, 100(4), pp.1541–1546.

512 Hosaka, T. et al., 2009. Antibacterial discovery in actinomycetes strains with mutations in
513 RNA polymerase or ribosomal protein S12. *Nature Biotechnology*, 27(5), pp.462–464.

514 Hoskisson, P.A. & Hutchings, M.I., 2006. MtrAB-LpqB: a conserved three-component
515 system in actinobacteria? *Trends in Microbiology*, 14(10), pp.444–449.

516 Hsiao, N.-H. & Kirby, R., 2009. Two-component signal transduction systems in
517 *Streptomyces* and related organisms studied using DNA comparative microarray analysis.
518 *Antonie van Leeuwenhoek*, 95(3), pp.189–206.

519 Hutchings, M.I., Hong, H.-J. & Buttner, M.J., 2006. The vancomycin resistance VanRS two-
520 component signal transduction system of *Streptomyces coelicolor*. *Molecular Microbiology*,
521 59(3), pp.923–935.

522 Hutchings, M.I., Hong, H.-J., Leibovitz, E., et al., 2006. The sigma(E) cell envelope stress
523 response of *Streptomyces coelicolor* is influenced by a novel lipoprotein, CseA. *Journal of*
524 *Bacteriology*, 188(20), pp.7222–7229.

525 Katz, L. & Baltz, R.H., 2016. Natural product discovery: past, present, and future. *Journal of*
526 *Industrial Microbiology & Biotechnology*, 43(2-3), pp.1–22.

527 Kieser, T. et al., 2000. *Practical Streptomyces Genetics*. Published by the John Innes
528 Foundation. ISBN 0-7084-0623-8

529 Krämer, R., 2009. Osmosensing and osmosignaling in *Corynebacterium glutamicum*. *Amino*
530 *Acids*, 37(3), pp.487–497.

- 531 Labeda, D.P. et al., 2011. Phylogenetic study of the species within the family
532 Streptomycetaceae. *Antonie van Leeuwenhoek*. 101(1), pp73-104
- 533 Molle, V. et al., 2000. WhiD and WhiB, Homologous Proteins Required for Different Stages
534 of Sporulation in *Streptomyces coelicolor* A3(2). *Journal of Bacteriology*, 182(5), pp.1286–
535 1295.
- 536 Möker, N. et al., 2004. Deletion of the genes encoding the MtrA-MtrB two-component system
537 of *Corynebacterium glutamicum* has a strong influence on cell morphology, antibiotics
538 susceptibility and expression of genes involved in osmoprotection. *Molecular Microbiology*,
539 54(2), pp.420–438.
- 540 Munnoch, J.T. et al., 2016. Characterization of a putative NsrR homologue in *Streptomyces*
541 *venezuelae* reveals a new member of the Rrf2 superfamily. *Scientific Reports*, 6, 31597.
- 542 Nguyen, H.T. et al., 2010. A lipoprotein modulates activity of the MtrAB two-component
543 system to provide intrinsic multidrug resistance, cytokinetic control and cell wall homeostasis
544 in *Mycobacterium*. *Molecular Microbiology*, 76(2), pp.348–364.
- 545 Nicol, J.W. et al., 2009. The Integrated Genome Browser: free software for distribution and
546 exploration of genome-scale datasets. *Bioinformatics*, 25(20), pp.2730–2731.
- 547 Plocinska, R. et al., 2014. *Mycobacterium tuberculosis* MtrB Sensor Kinase Interactions with
548 FtsI and Wag31 Proteins Reveal a Role for MtrB Distinct from That Regulating MtrA
549 Activities. *Journal of Bacteriology*, 196(23), pp.4120–4129.
- 550 Plocinska, R. et al., 2012. Septal Localization of the *Mycobacterium tuberculosis* MtrB
551 Sensor Kinase Promotes MtrA Regulon Expression. *Journal of Biological Chemistry*,
552 287(28), pp.23887–23899.
- 553 Purushotham, G. et al., 2015. *Mycobacterium tuberculosis* oriC sequestration by MtrA
554 response regulator. *Molecular Microbiology*, 98(3), pp.586–604.
- 555 Rutledge, P.J. & Challis, G.L., 2015. Discovery of microbial natural products by activation of

- 556 silent biosynthetic gene clusters. *Nature Reviews Microbiology*, 13(8), pp.509–523.
- 557 Salazar, M.E. & Laub, M.T., 2015. Temporal and evolutionary dynamics of two-component
558 signaling pathways. *Current Opinion in Microbiology*, 24, pp.7–14.
- 559 Satsangi, A.T. et al., 2013. Mycobacterium tuberculosis MtrAY102C is a gain-of-function
560 mutant that potentially acts as a constitutively active protein. *Tuberculosis (Edinb)*, 93,
561 pp.S28–S32.
- 562 Schwedock, J. et al., 1997. Assembly of the cell division protein FtsZ into ladder-like
563 structures in the aerial hyphae of *Streptomyces coelicolor*. *Molecular Microbiology*, 25(5),
564 pp.847–858.
- 565 Song, L. et al., 2006. Type III polyketide synthase beta-ketoacyl-ACP starter unit and
566 ethylmalonyl-CoA extender unit selectivity discovered by *Streptomyces coelicolor* genome
567 mining. *Journal of the American Chemical Society*, 128(46), pp.14754–14755.
- 568 Stapleton, M.R. et al., 2012. Mycobacterium tuberculosis WhiB1 represses transcription of
569 the essential chaperonin GroEL2. *Tuberculosis (Edinb)*, 92(4), pp.328–332.
- 570 Tschowri, N. et al., 2014. Tetrameric c-di-GMP Mediates Effective Transcription Factor
571 Dimerization to Control *Streptomyces* Development. *Cell*, 158(5), pp.1136–1147.
- 572 van Keulen, G. & Dyson, P.J., 2014. Production of Specialized Metabolites by *Streptomyces*
573 *coelicolor* A3(2). In *Advances in Applied Microbiology*. Elsevier, pp. 217–266.
- 574 Weber, T. et al., 2015. antiSMASH 3.0—a comprehensive resource for the genome mining of
575 biosynthetic gene clusters. *Nucleic Acids Research*, 43(W1), pp.W237–43.
- 576 Willemse, J., Mommaas, A.M. & Wezel, G.P., 2011. Constitutive expression of *ftsZ* overrides
577 the *whi* developmental genes to initiate sporulation of *Streptomyces coelicolor*. *Antonie van*
578 *Leeuwenhoek*, 101(3), pp.619–632.
- 579 Wolański, M. et al., 2011. The level of AdpA directly affects expression of developmental
580 genes in *Streptomyces coelicolor*. *Journal of Bacteriology*, 193(22), pp.6358–6365.

581 Wolański, M., Jakimowicz, D. & Zakrzewska-Czerwińska, J., 2012. AdpA, key regulator for
582 morphological differentiation regulates bacterial chromosome replication. *Open Biology*,
583 2(7), p.120097.

584 Zahrt, T.C. & Deretic, V., 2000. An essential two-component signal transduction system in
585 *Mycobacterium tuberculosis*. *Journal of Bacteriology*, 182(13), pp.3832–3838.

586

587 **Acknowledgements**

588 We thank Kim Findlay for performing the Scanning Electron Microscopy at the John Innes
589 Centre Bioimaging Facility, Mark Buttner for the gift of *S. venezuelae* NRRL B-65442, cosmid
590 Sv-6-A04, access to microarray data prior to publication and helpful discussions, Matt Bush
591 for helpful discussions and critical reading of the manuscript and Elaine Patrick for excellent
592 technical support. This research was supported by a BBSRC PhD studentship to NS, a UEA-
593 funded PhD studentship to FK, MRC Milstein award G0801721 to MIH, Norwich Research
594 Park translational funding, NPRONET Proof of Concept funding and Natural Environment
595 Research Council responsive mode grants NE/M015033/1 and NE/M014657/1 to MIH and
596 BW and NE/M001415/1 to PAH.

597

598 **Author contributions**

599 FK and RFS constructed the *S. coelicolor* mutants and FK undertook phenotype analysis, NS
600 constructed all the *S. venezuelae* strains and constructs, analysed phenotypes and performed
601 ChIP-seq experiments, JTM performed RNA-seq and dRNA-seq experiments, JTM and GC
602 analysed the ChIP- and RNA-seq datasets, NH made constructs and prepared strain extracts
603 for LCMS, DH purified and analysed all natural products including LCMS and PCA analyses,
604 PAH, MIH and BW conceived the study and all the authors analysed data and wrote the
605 manuscript.

606