Running title: CRITICAL TEMPERATURE LIMITS XYLOGENESIS

Critical minimum temperature limits xylogenesis and maintains treelines on the Tibetan Plateau

Xiaoxia Li^{1,2}, Eryuan Liang^{1,2*}, Jozica Grigar³, Sergio Rossi⁴, Katarina Cufar⁵ & Aaron M. Ellison⁶

¹*Key Laboratory of Alpine Ecology and Biodiversity, Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100085, China;* ²*CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing 100101, China;* ³*Slovenian Forestry Institute, Department of Yield and Silviculture, Vecna pot 2, SI-1000 Ljubljana, Slovenia;* ⁴*University of Quebec in Chicoutimi, Département des Sciences Fondamentales, 555, Boulevard de l'Université, Chicoutimi (QC) G7H2B1, Canada ;* ⁵*University of Ljubljana, Biotechnical Faculty, Department of Wood Science and Technology, Jamnikarjeva 101, SI-1000 Ljubljana, Slovenia* ⁶*Harvard Forest, Harvard University, 324 North Main St., Petersham, MA 01366, USA;*

* Correspondence:

Dr. Eryuan Liang
Institute of Tibetan Plateau Research, Chinese Academy of Sciences
Building 3, Courtyard 16, Lin Cui Road, Chaoyang District
Beijing 100101, China
Phone: 0086 10-84097069
Fax: 0086 10-84097079
Email: liangey@itpcas.ac.cn

ABSTRACT

Physiological and ecological mechanisms that define treelines are still debated. It is suggested that the absence of trees above the treeline is caused by the low temperature that limits growth. Thus, we raise the hypothesis that there is a critical minimum temperature (CT_{min}) preventing xylogenesis at treeline. We tested this hypothesis by examining weekly xylogenesis across three and four growing seasons in two natural Smith fir (*Abies georgei* var. *smithii*) treeline sites on the south-eastern Tibetan Plateau. Despite differences in the timing of cell differentiation among years, minimum air temperature was the dominant climatic variable associated with xylem growth; the critical minimum temperature (CT_{min}) for the onset and end of xylogenesis occurred at 0.7±0.4 °C. A process-based-modeled chronology of tree-ring formation using this CT_{min} was consistent with actual tree-ring data. This extremely low CT_{min} permits Smith fir growing at treeline to complete annual xylem production and maturation and provides both support and a mechanism for treeline formation.

Key- words: cambial activity; climate; growing season; growth limitation hypothesis; threshold temperatures; xylem; treeline formation

INTRODUCTION

The explanations for treeline formation focus on limitations of available resources (Stevens & Fox 1991; Susiluoto *et al.* 2010), establishment sites (e.g., Smith *et al.* 2003), or time available for growth (e.g., Körner 1998), although this ecophysiological causes still remain debated (Tranquillini 1979; Holtmeier 2003; Körner 2003; Cui *et al.* 2005). Based on notable similarities in temperatures at treelines (Körner & Paulsen 2004), Körner (1998) proposed that low temperatures limit the time available for meristematic growth and cell division (Körner 1998). This hypothesis has been supported by phenomenological data. For example, treeline trees tend to have higher amounts of non-structural carbohydrates than trees growing at lower elevation, suggesting that treeline are limited more by growth processes than by photosynthesis and carbon assimilation (e.g. Hoch & Körner 2003; Shi *et al.* 2008; Fajardo *et al.* 2012). In parallel, dendroclimatic studies have identified a signal of reduced growth during periods with low temperatures at treelines in cold and humid areas (e.g., Mäkinen *et al.* 2000; Oberhuber 2004; Frank & Esper 2005; Ettinger *et al.* 2011).

Physiological manifestations of the growth limitation hypothesis include a constraint on the production of new cells by meristems below a critical minimum temperature (CT_{min}) (Körner 1998) and a trade-off between taking maximal advantage of the length of the growing season while avoiding cellular damage due to early (fall, winter) or late (winter, spring) freezing events (Chuine & Beaubien 2001; Kollas *et al.* 2014). Such a trade-off would suggest a narrow thermal window for the onset and cessation of xylem formation at treeline. Indeed, recent studies focused on temporal dynamics of xylem formation in various tree species at treeline (Rossi *et al.* 2007; Seo *et al.* 2008; Gruber *et al.* 2009; Moser *et al.* 2010;

Lenz *et al.* 2012). One study reported that a gradual increase in temperature (heat sum) was associated with the onset of cambial activity (Seo *et al.* 2008), whereas another estimated a CT_{min} of 5.6 –8.5 °C for xylogenesis at the altitudinal treeline in the Eastern Alps (Rossi *et al.* 2007). Separating gradual (heat-sum) and threshold (CT_{min}) effects on xylogenesis at treeline has not yet been accomplished.

A mechanistic model can provide a deeper understanding on a climatic control on tree growth dynamics. The process-based forward model, such as, Vaganov-Shashkin (VS) model, was used to simulate the climatic control on conifer tree-ring growth (Vaganov *et al.* 1999; Anchukaitis *et al.* 2006; Evans *et al.* 2006). The critical temperature for cambial activity is a key parameter to model tree growth despite that its observed data are less available as a model input.

Our observations at the upper Smith fir (*Abies georgei* var. *smithii*) treeline of the south-eastern Tibetan Plateau, including a decade of uninterrupted *in situ* micrometeorological measurements and weekly collection of microcores containing cambium and forming wood during three consecutive growing seasons provides an opportunity to examine both gradual and threshold effects of temperature on xylogenesis at a natural alpine treeline. Specifically, we tested the potential for thermal control of xylogenesis to be a mechanism underlying the growth limitation hypothesis by (1) identifying the timing and dynamics of xylem formation in Smith fir growing at treeline as a function of climatic factors; and (2) detecting a plausible CT_{min} for xylogenesis. Previous studies have found that its growth near treeline is constrained by the minimum temperature in summer (Bräuning & Mantwill 2004; Liang *et al.* 2009). The onset of bud swelling and needle unfolding in Smith

fir is delayed by 3.5 days per 100 m increase in elevation (Wang *et al.* 2013), indicating a thermal limitation of tree phenology. Therefore, we hypothesized that minimum temperature limits xylem formation and that a threshold minimum temperature controls the timing of the onset and ending of xylem formation. If this potential CT_{min} is reasonable, as a primary temperature parameter, it will be expected to be used in Vaganov-Shashkin (VS) model of tree-ring formation to simulate regional tree growth at Smith fir treeline on the south-eastern Tibetan Plateau.

MATERIALS AND METHODS

Study sites and tree selection

The study focused on the natural alpine treeline of Smith fir (*Abies georgei* var. *smithii*) growing on the eastern side of the Sygera Mountains (29° 10' – 30° 15' N, 93° 12' – 95° 35' E) on the south-eastern Tibetan Plateau (Liang *et al.* 2011). The south-eastern Tibetan Plateau is characterized by a cold and humid climate, and has the highest natural treeline (up to 4900 m a.s.l.) in the Northern Hemisphere (Miehe *et al.* 2007). Smith fir (*Abies georgei* var. *smithii*) is one of the dominant treeline species in this region. This tree is a shade-requiring conifer and the upper treeline depends on topographic aspect and ranges from 4,250 to 4,400 m a.s.l. We studied two sites at open-canopy treelines: Site 1, was at 4360 m a.s.l. on an east-facing slope, and Site 2, was at 4250 m a.s.l. on a south-east-facing slope. The sites were 200 m apart, on slopes < 15° *Rhododendron aganniphum* var. *schizopeplum* dominated the understory. The coverage of Smith fir was < 20% and the podzolic soils had an average pH value of 4.5.

At each site, five dominant trees were selected in April 2007. These trees had a mean age of 201 ± 24 and 117 ± 14 years, and mean diameters at 1.3-m above ground of 34 ± 4 and 44 ± 7 cm in Sites 1 and 2, respectively. Because repeated sampling could cause severe wounding that could modify xylem formation, another five trees per site with homogeneous diameters were chosen for the samplings in 2009 and 2010. Trees with polycormic stems, partially dead crowns, reaction wood, or other evident damage were avoided.

Meteorological data

An automatic weather station (Campbell Scientific, CR1000) was installed in November 2006 in an open area above the treeline ($29^{\circ}39' \text{ N}$, $94^{\circ}42' \text{ E}$, 4390 m a.s.l.) at a linear distance of ≈ 150 m and 200 m from Sites 1 and 2, respectively. Measurements of air (3-m above ground) and soil temperature (at 10-, 20- and 40-cm depths), precipitation, snow fall, and soil water content (at 10-, 20-, and 40-cm depths) were collected at 30-minute intervals. These data were used to compute daily averages, minima, and maxima of each variable.

Microcoring and histological analyses

Xylem formation was studied from 2007 until 2010 at Site 1 and from 2007 to 2009 at Site 2. One microcore (15-mm long, 2-mm diameter) was collected from each tree weekly from May until October around the stem at breast height using a Trephor tool. Immediately after removal from the trees, the microcores were fixed in a formalin-ethanol-acetic acid (FAA) solution. The microcores contained innermost phloem, cambium, developing xylem, and at least three previous xylem growth rings. In the laboratory, the microcores were dehydrated

with successive immersions in a graded series of ethanol and *d*-limonene, then embedded in paraffin. Transverse sections (9-12 µm in thickness) were cut from the samples with a Leica RM 2245 rotary microtome using Feather N35H knives (Osaka, Japan). Sections were stained with a mixture of safranin (0.5 % in 95 % ethanol) and astra blue (0.5 % in 95% ethanol) and observed with a Nikon Eclipse 800 light microscope under bright field and polarized light to identify the phases of differentiation of the developing xylem cells. In cross-section, cambial cells were characterized by thin cell walls and small radial diameters (Deslauriers *et al.* 2003). Newly-formed xylem cells in the phase of cell enlargement contained protoplasts, had thin primary walls, and a radial diameter at least twice the size of the cambial cells. The onset of cell-wall thickening was determined by birefringence in the cell walls under polarized light. Mature cells had completely red-stained walls and empty lumen. For each sample, the total current xylem cell number was determined by counting the number of cells undergoing enlargement, cell-wall thickening, and the number of mature cells along three radial files (Deslauriers *et al.* 2003).

Data standardization and estimating the rate of xylem formation

The data were standardized to compensate the variation in the number of xylem cells along the tree circumference. The total cell number of the previous years was counted on three radial files per sample and used for standardization. The standardized number of cells nc_{ij} number in the i^{th} phase of the j^{th} sample was calculated as:

$$nc_{ij} = n_{ij} (a_m / a_j) \quad (1)$$

where n_{ij} is the number of cells in the current year, a_m is the mean number of cells of the

previous ring of all j -samples and a_j is the mean number of cells of the previous ring in each j -sample .

We modelled the dynamics of xylem formation by fitting a Gompertz function to the number of xylem cells that were produced through time:

$$y = A \exp[-e^{(\beta - \kappa t)}] \quad (2)$$

where y is the weekly cumulative sum of tracheids, t is the time of the year computed as day of the year, A is an asymptote (constant) and β and κ are constants reflecting the x -intercept placement and rate of change, respectively (Deslauriers *et al.* 2003). Model parameters were estimated using the Origin software package (Version 8.5, OriginLab Corporation, Northampton, MA, USA).

Estimation of the onset and ending of xylem formation

We used directly observations of cell differentiation to identify the onset, ending, and duration of xylem formation from counts of the number of cells in three radial files per tree. In spring, xylem formation was considered to have started when at least one tangential row of cells was observed in the enlarging phase. Because of the weekly resolution of the monitoring, we used the occurrence of 1-2 enlarging xylem cells along any of the checked three radial files as an indicator the xylem formation had begun (Li *et al.* 2013). In late summer, when cells were no longer observed in the wall thickening and lignification phase, xylem formation was considered to have ended. The duration of xylogenesis was estimated as the number of days between the dates of onset and ending of xylem formation; we report the average among trees for each studied site and year.

Comparisons between sites in onset, duration, and ending of differentiation in the developing xylem ring, were done with generalized linear models (GLM). Homoscedasticity was checked using Shapiro-Wilk and Levene tests.

Identifying CT_{min}

Logistic regression (LOGISTIC procedure in SPSS 16.0) was used to model the probability of xylogenesis as a function of air temperature. Xylem cell production was coded zero (not occurring) or one (occurring). CT_{min} was estimated as that temperature when the probability of ongoing xylem growth equalled 0.5 (Rossi et al. 2008). CT_{min} s also were calculated for 5% and 95% of the maximum number of xylem cells. For each tree and year, the model was fitted with three respective daily temperature series (mean, absolute minimum, and absolute maximum). Model verification included the likelihood-ratio χ^2 , Wald's χ^2 for regression parameter and goodness of fit, and Hosmer-Lemeshow $\hat{C}$ for possible lack of fit. None of the models were excluded because of a lack of fit. CT_{min} s were compared between sites and years using analysis of variance (ANOVA) models. Model validation was performed by comparing the observations with the predicted values when using estimated CT_{min} s.

Climate- growth relationships

We used two approaches to identify relationships between intra-annual xylem growth and climatic variables during four growing seasons. One approach consisted of computation of Pearson's correlation coefficients between xylem cell production and weather data for weekly intervals. Weather data here include daily mean, daily absolute minimum, daily absolute

maximum temperatures, growing degree-days (GDD) > 5°C, and sums of precipitation.

Intra-annual xylem growth may be controlled both by endogenous (e.g. hormonal regulation) and exogenous factors (e.g., climate). To analyse the climatic effect, a common approach was used to remove the endogenous growth trend by fitting a growth curve, and to estimate the growth departure, calculated as the dimensionless ratio between observed and expected growth (Fritts 1976). This ratio (hereafter called ‘growth index’) was calculated as the number of tracheids produced during the week divided by the expected values estimated using the Gompertz function (Zhai *et al.* 2012). To account for possible effects of time-lags, daily weather data were averaged (temperature) or summed (precipitation) weekly from 1 to 10 d prior to each sampling date (referred to as P1 to P10). To minimize the effects of temporal autocorrelation, correlation coefficients were calculated on first-order differences for both datasets.

Tree-ring modeling

We used the Vaganov-Shashkin (VS) model to simulate tree-ring growth at the Smith fir treelines in the Sygera Mountains. The VS model estimates xylem formation and its internal characteristics based on equations relating daily temperature, precipitation, and sunlight to the kinetics of xylem development (Vaganov *et al.* 2006). It assumes that climatic influences are directly but nonlinearly related to tree-ring characteristics through controls on the rates of cambial activity processes. To date, it has been successfully used to simulate and evaluate the relationships between climate and tree-ring formation under a variety of environmental conditions in many different regions (Anchukaitis *et al.* 2006; Evans *et al.* 2006; George *et al.*

2008; Shi *et al.* 2008; Zhang *et al.* 2011; Gou *et al.* 2013). Variables used as input for the VS model included soil moisture, depth of root system, temperature sum for initiation of growth, soil water drainage rate, and maximum daily precipitation falling into soil were taken from field observations. For the parameter of minimum temperature, we used the abovementioned CT_{min}. Model fit was evaluated against an actual high-quality tree-ring width chronology from Smith fir treeline in the Sygera Mts., which was developed and used for paleoclimatic reconstructions in this region (Liang *et al.* 2009). The best estimate of physiological CT_{min} was found by iteration and comparison between simulated and observed chronologies (1960 – 2006).

Finally, a single simulated tree-ring width chronology was created for the Smith fir treeline in the Sygera Mts. based on daily climate data from Nyingchi meteorological stations (3,000 m a.s.l.). To account for the altitude differences between Nyingchi and the study sites, we extended the time series of daily temperatures at the treeline back to 1960 based on a linear regression of the Nyingchi data and our own micrometeorological data ($r^2 \geq 0.89$, 2007-2010).

RESULTS

Micrometeorological conditions at the upper treeline

The sampling sites at the upper treeline were cold and humid. In spite of a difference of 110 m in elevation and different topographical aspects of the two treeline sites, they had fairly similar temperatures (Supporting Information Fig. S1). Annual temperatures (2007 – 2010) ranged from 0.1 to 0.9 °C, while growing-season (June-September) temperatures ranged from

6.4 to 7.1 °C (Fig. 1). On average, the annual precipitation was 951 mm, of which 62 % fell during the monsoon season (June to September). Snowfall mainly occurred from November to May. Due to snowmelt and increased precipitation, soil moisture content increased rapidly from the beginning of April and remained above 30 % from early May until November, and finally decreased to near 0 in late November and early December. The year 2008 was characterized by heavy spring snowfall and had the latest snowmelt and soil thawing during the four studied years (Fig. 1).

Xylem formation

The onset of xylem formation occurred from late May to early June and differed significantly among years ($F = 15.73$, $P < 0.001$). The onset of xylem formation was observed 4 - 9 days later in 2008 than in the other years, at both sites (Fig. 2 a). No difference was found in onset of xylogenesis between sites ($F = 2.31$, $P > 0.05$). Xylem formation ended between the beginning and the end of September and differed significantly among years ($F = 10.42$, $P < 0.005$), and 1-2 weeks later in 2010 at site 1 (Fig. 2 b).

Overall, the duration of xylem formation lasted from 109 to 125 days (Fig. 2 c), with no significant differences detected between sites ($F = 3.80$, $P > 0.05$). Conversely, there were significant variations among years ($F = 4.71$, $P < 0.05$). From 2007 to 2009, the average period between the onset and ending of xylem formation was 107 days, whereas an average of 118 days was required to complete xylem formation in 2010.

Relationship between climate and xylem formation

Weekly cumulative xylem production was fit well by the Gompertz function ($0.96 \leq r^2 \leq 0.98$; Supporting Information Table S1 and Fig. S2). Intra-annual xylem cell production was significantly and positively correlated with daily minimum and mean air temperatures and $GDD > 5^\circ\text{C}$ at both sites (Fig. 3 a, b). However, only minimum temperature was significantly correlated with growth indices after removing the growth trends (Fig. 3 c, d). At site 1, positive correlations between growth indices and minimum temperature were found for time lags of 0 – 3 days ($r = 0.34$, $P < 0.05$), whereas the corresponding time lags were 7 – 10 days at site 2 ($r = 0.42$, $P < 0.05$). No significant correlations were found between xylem cell production or growth index and precipitation from P0 to P10.

CT_{min}

The critical minimum air temperature (CT_{min}) at which there was a 0.5 probability that xylem formation was ongoing is shown in Fig. 4 and Table 1 for Site 1 (2007 – 2010) and Site 2 (2007 – 2009). Based on direct observations of xylogenesis, the values respectively for minimum, mean, and maximum temperatures of 0.6, 4.0, and 9.3°C were estimated for the onset of xylogenesis, while the corresponding values for the ending of xylem differentiation were 0.7, 3.9, and 9.0 °C. There were no differences among critical temperatures for the onset and ending of xylogenesis (ANOVA, $P > 0.05$), with values of 0.7 ± 0.4 , 3.9 ± 0.5 , and $9.1 \pm 0.6^\circ\text{C}$ for the minimum, mean and maximum temperatures, respectively. No significant differences were found between the two sites in terms of the estimated air temperature thresholds for the onset and ending of xylogenesis (ANOVA, $P > 0.05$). The mean air temperature during the period of xylem formation based on weekly observations at both sites

was $6.8 \pm 0.4^{\circ}\text{C}$.

Tree-ring modeling

Applying the Vaganov-Shashkin (VS) model starting with an estimated $\text{CT}_{\min} = 0.7^{\circ}\text{C}$, the best estimate of physiological $\text{CT}_{\min}$ was 0.9°C (Table 2, Fig. 5). The correlation between observations and predictions varied slightly at $\text{CT}_{\min}$ of $0.3\text{--}0.9^{\circ}\text{C}$, while it decreased rapidly for values of $\text{CT}_{\min} > 1^{\circ}\text{C}$ (Fig. 6). Overall, significant, positive correlations were found between the modeled and real chronologies when $\text{CT}_{\min}$ varied within the range of $0.7 \pm 0.4^{\circ}\text{C}$ ($r = 0.62$, $P < 0.01$).

DISCUSSION

The importance of temperature for xylem formation during and after its onset has been demonstrated repeatedly (e.g., Oribe *et al.* 2001; Gričar *et al.* 2006; Rossi *et al.* 2008). These and other data suggest that air temperature, not soil temperature, directly limits xylem formation at high latitudes and altitudes (Rossi *et al.* 2007; Lenz *et al.* 2012; Lupi *et al.* 2012). Minimum temperature is assumed to be an important driver of tree species range limits (Körner 2003; Kollas *et al.* 2014), and so a critical minimum temperature ($\text{CT}_{\min}$) with fairly narrow bounds should exist for the onset and ending of xylogenesis. However, long-term monitoring of xylem formation at natural treelines is limited, which has precluded assessment of $\text{CT}_{\min}$ for xylem formation by direct observations.

Effects of climate on xylem formation

As predicted, minimum air temperature strongly limited xylem formation of Smith fir at the upper treeline on the south-eastern Tibetan Plateau. This finding accords with those from dendroclimatological analysis in the same study area (Liang *et al.* 2009) and xylem formation studies at high latitudes and altitudes (e.g., Rossi *et al.* 2008; Gruber *et al.* 2009). The importance of minimum air temperature may be related to the timing of cell differentiation, which perhaps occurs mainly during the night, when the temperature is lower (Hosoo *et al.* 2002, Steppe *et al.* 2015). Controlled experiments in Hamburg, Germany also showed that night temperatures could directly influence xylem cell expansion of *Podocarpus latifolius* (Dünisch 2010). According to Körner (2003), cell doubling time, which is highest and fairly constant at temperatures of 10-25°C, approaches infinity at 1-2°C, suggesting a minimum temperature limit on cell division. The simulated ring-width chronologies produced by VS model of tree-ring formation also exhibit similar positive correlations with the minimum temperature during summer (Supporting Information Fig. S3, $P < 0.01$). CT_{min} is thus expected to limit xylogenesis of Smith fir at the treeline.

Critical temperatures for xylem formation

Despite the variance in timing and duration of xylem formation during our four years of observations, minimum, average, and maximum temperatures for the onset and ending of xylogenesis were narrowly bounded with average values of 0.7, 3.9, and 9.1 °C, respectively. Most studies to date have indicated that xylogenesis in conifers growing in cold climates can take place when the daily minimum temperatures $\geq 4 - 5$ °C (e.g. Rossi *et al.* 2008; Boulouf Lugo *et al.* 2012). However, based on the presented 4-year observations of xylogenesis and

uninterrupted *in situ* micrometeorological measurements directly at the treeline, we found that the CT_{min} for xylogenesis in Smith fir is as low as 0.7 °C. In particular, based on this CT_{min} , the modeled chronology produced by VS model is consistent with actual tree-ring data, suggesting that minimum temperature could be considered as a significant driver of xylem growth. Such a low CT_{min} may have evolved to provide sufficient time to complete xylogenesis at alpine treelines. The length of the growing season for stem growth diminishes with altitude and reaches a minimum at the alpine treeline. According to some authors (Rossi *et al.* 2008; Körner 2012), a tree can only survive when the length of growing seasons is at least 3 months and the mean growing season air temperature is 6.4.°C, since each of them critically constrains the growth and development of trees. At Smith fir treelines in south-eastern Tibet, the duration of xylem growth of 115 days provided by a $CT_{min} < 1$ °C, together with a mean air temperature of 6.8 °C during the growing season extended by this low CT_{min} meets these prerequisites for tree growth and development.

The dates of snow melting and soil thawing also are thought to be critical for the onset of xylogenesis and could therefore determine the annual xylem production (Vaganov *et al.* 1999). At our treeline sites, the onset of xylem growth occurred 7 - 30 days after soil thawing in spring, which coincided with the surpassing of CT_{min} . This temporal lag might be related to a strategy of avoiding the risk of freezing damage during early cell development.

The growth limitation hypothesis predicts that the absence of trees above the treeline is attributable to critical minimum temperature for growth (Körner 1998). Treeline trees often have slower growth rates and higher non-structural carbohydrate levels than trees at lower altitudes (e.g. Hoch & Körner 2003; Shi *et al.* 2008; Fajardo *et al.* 2012), suggesting a carbon

sink rather than carbon limitation. However, some authors have argued that tree populations with the highest non-structural carbohydrate concentrations may be the most carbon limited in terms of growth (Li *et al.* 2008; Wiley & Helleker 2012). Although our observations of xylogenesis cannot differentiate between carbon limitation and a carbon sink in Smith fir, the significant effect of a narrowly bounded CT_{min} on xylem cell production provides a physiological, rather than an ecological, mechanism for the growth limitation hypothesis.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (41525001, 41130529, 41601204). International cooperation was supported by the bilateral project between China and Slovenia (BI-CN/09-11-012) and COST Action (FP1106, STReESS). AME's participation in the project was supported by the Chinese Academy of Sciences President International Fellowship Initiative for Visiting Scientists, Grant no. 2016VBA074. We thank the Southeast Tibet Station for Alpine Environment Observation and Research, Chinese Academy of Sciences for the fieldwork and monitoring; Yongxiang Zhang from National Climate Centre, Beijing, China for her support on the VS model and Martin Cregeen for additional language editing.

REFERENCES

- Anchukaitis K. J., Evans M. N., Kaplan A. , Vaganov E. A., Hughes M. K., Grissino-Mayer H. D. & Cane M. A. (2006) Forward modeling of regional scale tree-ring patterns in the south-eastern United States and the recent influence of summer drought. *Geophysical Research Letters* **33**, L04705.
- Boulouf Lugo J., Deslauriers A. & Rossi S. (2012) Duration of xylogenesis in black spruce

lengthened between 1950 and 2010. *Annals of Botany* **110**, 1099-1108.

Bräuning, A., & Mantwill B. (2004) Summer temperature and summer monsoon history on the Tibetan plateau during the last 400 years recorded by tree rings. *Geophysical Research Letters* **31**, L24205.

Camarero, J. J., Olano J. M. & Parras A. (2010) Plastic bimodal xylogenesis in conifers from continental Mediterranean climates. *New Phytologist* **185**, 471-480.

Chuine, I. & Beaubien E. G. (2001) Phenology is a major determinant of tree species range. *Ecology Letters* **4**, 500-510.

Cui, H., Liu H. & Dai J. (2005) Study on mountain ecology and alpine timberline. Beijing, Science Press.

Dünisch, O. (2010) Low night temperatures cause reduced tracheid expansion in *Podocarpus latifolius*. *IAWA Journal* **31**, 245-255.

Deslauriers, A. & Morin H. (2005) Intra-annual tracheid production in balsam fir stems and the effect of meteorological variables. *Trees - Structure and Function* **19**, 402-408.

Deslauriers, A., Morin, H. & Begin, Y. (2003) Cellular phenology of annual ring formation of *Abies balsamea* in the Quebec boreal forest (Canada). *Canadian Journal of Forest Research* **33**, 190-200.

Ettinger, A. K., Ford K. R. & HilleRisLambers J. (2011) Climate determines upper, but not lower, altitudinal range limits of Pacific Northwest conifers. *Ecology* **92**, 1323-1331.

Evans, M. N., Reichert B. K., Kaplan A., Anchukaitis K. J., Vaganov E. A., Hughes M. K. & Cane M. A. (2006) A forward modeling approach to paleoclimatic interpretation of tree-ring data. *Journal of Geophysical Research-Biogeosciences* **111**, G03008

395 Fajardo, A., Piper F. I. & Hoch G. (2013) Similar variation in carbon storage between
396 deciduous and evergreen treeline species across elevational gradients. *Annals of Botany*
397 **112**, 623-631.

398 Frank, D. & Esper J. (2005) Characterization and climate response patterns of a
399 high-elevation, multi-species tree-ring network in the European Alps.
400 *Dendrochronologia* **22**, 107-121.

401 Fritts, H. (1976) Tree rings and climate. Academic Press, New York.

402 George, S. S., Meko D. M. & Evans M. N. (2008) Regional tree growth and inferred summer
403 climate in the Winnipeg River basin, Canada, since AD 1783. *Quaternary Research* **70**,
404 158-172.

405 Gou, X., Zhou F., Zhang Y., Chen Q. & Zhang J. (2013) Forward modeling analysis of
406 regional scale tree-ring patterns around the northeastern Tibetan Plateau, Northwest
407 China. *Biogeosciences Discuss.* **2013**, 9969-9988.

408 Grace, J., Berninger F. & Nagy L. (2002) Impacts of climate change on the tree line. *Annals*
409 *of Botany* **90**, 537-544.

410 Gričar, J., Zupančič, M., Čufar, K., Koch, G., Schmitt, U. & Oven, P. (2006) Effect of local
411 heating and cooling on cambial activity and cell differentiation in the stem of Norway
412 spruce (*Picea abies*). *Annals of Botany* **97**, 943-951.

413 Gruber, A., Baumgartner, D., Zimmermann, J. & Oberhuber, W. (2009) Temporal dynamic
414 of wood formation in *Pinus cembra* along the alpine treeline ecotone and the effect of
415 climate variables. *Trees - Structure and Function* **23**, 623-635.

416 Hoch, G. & Körner, C. (2003) The carbon charging of pines at the climatic treeline: a global

comparison. *Oecologia* **135**, 10-21.

Holtmeier, F. K. (2003) Mountain timberlines: ecology, patchiness and dynamics. Dordrecht, Kluwe.

Hosoo, Y., Yoshida, M., Imai, T. & Okuyama, T. (2002) Diurnal difference in the amount of immunogold-labeled glucomannans detected with field emission scanning electron microscopy at the innermost surface of developing secondary walls of differentiating conifer tracheids. *Planta* **215**, 1006-1012.

Huang, J.-G., Deslauriers A. & Rossi S. (2014) Xylem formation can be modeled statistically as a function of primary growth and cambium activity. *New Phytologist* **203**, 831-841.

Körner, C. (1998) A re-assessment of high elevation treeline positions and their explanation. *Oecologia* **115**, 445-459.

Körner, C. (2003) Alpine plant life: functional plant ecology of high mountain ecosystems, 2nd edn. Springer, Berlin Heidelberg New York.

Körner, C. (2012) Alpine treelines: functional plant ecology of the global high elevation tree limits. Springer, Berlin Heidelberg New York.

Körner, C. & Paulsen J. (2004) A world-wide study of high altitude treeline temperatures. *Journal of Biogeography* **31**, 713-732.

Kollas, C., Körner, C. & Randin, C.F. (2014). Spring frost and growing season length co-control the cold range limits of broad-leaved trees. *Journal of Biogeography* **41**, 773-783.

Lenz, A., Hoch, G. & Körner, C. (2012) Early season temperature controls cambial activity and total tree ring width at the alpine treeline. *Plant Ecology & Diversity* **6**, 365-375.

Li, M., Xiao, W., Wang, S., Cheng, G., Cherubini, P., Cai, X., Liu, X., Wang, X. & Zhu, W. (2008) Mobile carbohydrates in Himalayan treeline trees I. Evidence for carbon gain limitation but not for growth limitation. *Tree Physiology* **28**, 1287-1296.

Li, X., Liang, E., Gričar, J., Prislan, P., Rossi, S. & Čufar, K. (2013) Age dependence of xylogenesis and its climatic sensitivity in Smith fir on the south-eastern Tibetan Plateau. *Tree Physiology* **33**, 48-56.

Liang, E., Wang Y., Eckstein D. & Luo T. (2011) Little change in the fir tree-line position on the south-eastern Tibetan Plateau after 200 years of warming. *New Phytologist* **190**, 760-769.

Liang, E., Shao, X. & Xu, Y. (2009) Tree-ring evidence of recent abnormal warming on the southeast Tibetan Plateau. *Theoretical and Applied Climatology* **98**, 9-18.

Lupi, C., Morin, H., Deslauriers, A. & Rossi, S. (2012) Xylogenesis in black spruce: does soil temperature matter? *Tree Physiology* **32**, 74-82.

Makinen, H., Nojd P. & Mielikainen K. (2000) Climatic signal in annual growth variation of Norway spruce (*Picea abies*) along a transect from central Finland to the Arctic timberline. *Canadian Journal of Forest Research* **30**, 769-777.

Miehe, G., Miehe, S., Vogel, J., Co, S. & Duo, L. (2007) Highest treeline in the northern hemisphere found in southern Tibet. *Mountain Research and Development* **27**, 169-173.

Moser, L., Fonti, P., Büntgen, U., Franzen, J., Esper, J., Luterbacher, J. & Frank, D. (2010) Timing and duration of European larch growing season along altitudinal gradients in the Swiss Alps. *Tree Physiology* **30**, 225-233.

Oberhuber, W. (2004) Influence of climate on radial growth of *Pinus cembra* within the

alpine timberline ecotone. *Tree Physiology* **24**, 291-301.

Opgenoorth, L., Vendramin, G.G., Mao, K.S., Miehe, G., Miehe, S., Liepelt, S., Liu, J.Q. &

Ziegenhagen, B. (2010) Tree endurance on the Tibetan Plateau marks the world's

highest known tree line of the Last Glacial Maximum. *New Phytologist* **185**, 332-342.

Oribe, Y., Funada, R., Shibagaki, M. & Kubo, T. (2001) Cambial reactivation in locally

heated stems of the evergreen conifer *Abies sachalinensis* (Schmidt) Masters. *Planta*

212, 684-691.

Rathgeber, C.B.K., Rossi, S. & Bontemps, J.-D. (2011) Cambial activity related to tree size in

a mature silver-fir plantation. *Annals of Botany* **108**, 429-438.

Rossi, S., Deslauriers, A., Anfodillo, T. & Carraro, V. (2007) Evidence of threshold

temperatures for xylogenesis in conifers at high altitudes. *Oecologia* **152**, 1-12.

Rossi, S., Deslauriers, A., Gričar, J., Seo, J.-W., Rathgeber, C. B. K., Anfodillo, T., Morin, H.,

Levanic, T., Oven, P. & Jalkanen, R. (2008) Critical temperatures for xylogenesis in

conifers of cold climates. *Global Ecology and Biogeography* **17**, 696-707.

Rossi, S., Morin, H., Deslauriers, A. & Plourde, P.-Y. (2011) Predicting xylem phenology in

black spruce under climate warming. *Global Change Biology* **17**, 614-625.

Seo, J.-W., Eckstein, D., Jalkanen, R., Rickebusch, S. & Schmitt, U. (2008) Estimating the

onset of cambial activity in Scots pine in northern Finland by means of the heat-sum

approach. *Tree Physiology* **28**, 105-112.

Shi, J., Liu Y., Vaganov E., Li J. & Cai Q. (2008a). Statistical and process-based modeling

analyses of tree growth response to climate in semi-arid area of north central China: A

case study of *Pinus tabulaeformis*. *Journal of Geophysical Research-Biogeosciences*

483 **113**, G01026.

484 Shi, P., Körner, C. & Hoch, G. (2008b) A test of the growth-limitation theory for alpine tree
485 line formation in evergreen and deciduous taxa of the eastern Himalayas. *Functional*
486 *Ecology* **22**, 213-220.

487 Smith, W. K., Germino M. J., Hancock T. E. & Johnson D. M. (2003) Another perspective on
488 altitudinal limits of alpine timberlines. *Tree Physiology* **23**, 1101-1112.

489 Steppe K., Sterck F. & Deslauriers A. (2015) Diel growth dynamics in tree stems: linking
490 anatomy and ecophysiology. *Trends in Plant Science* **20**: 335-343

491 Stevens, G.C. & Fox, J.F. (1991) The Causes of Treeline. *Annual Review of Ecology and*
492 *Systematics* **22**, 177-191.

493 Susiluoto, S., Hiltavuori, E. & Berninger, F. (2010) Testing the growth limitation hypothesis
494 for subarctic Scots pine. *Journal of Ecology* **98**, 1186-1195.

495 Tranquillini, W. (1979) Physiological ecology of the alpine timberline. Springer, Berlin
496 Heidelberg New York.

497 Vaganov, E.A., Hughes, M.K., Kirilyanov, A.V., Schweingruber, F.H. & Silkin, P.P. (1999)
498 Influence of snowfall and melt timing on tree growth in subarctic Eurasia. *Nature* **400**,
499 149-151.

500 Vaganov, E. A., M. K. Hughes, and A. V. Shashkin. (2006) Introduction and factors
501 influencing the seasonal growth of trees. Springer, New York.

502 Wang, Y., Li, X., Dawadi, B., Eckstein, D. & Liang, E. (2013) Phenological variation in
503 height growth and needle unfolding of Smith fir along an altitudinal gradient on the
504 southeastern Tibetan Plateau. *Trees - Structure and Function* **27**, 401-407.

Wiley, E. & Helliker B. (2012) A re-evaluation of carbon storage in trees lends greater support for carbon limitation to growth. *New Phytologist* **195**, 285-289.

Zhai, L., Bergeron Y., Huang J. & Berninger F. (2012) Variation in intra-annual wood formation, and foliage and shoot development of three major Canadian boreal tree species. *American Journal of Botany* **99**, 827-837.

Zhang, Y. X., Shao X. M., Xu Y. & Wilmking M. (2011) Process-based modeling analyses of *Sabina przewalskii* growth response to climate factors around the northeastern Qaidam Basin. *Chinese Science Bulletin* **56**, 1518-1525.

Table 1 Mean $\pm$ standard deviation of the threshold daily maximum (MaxT), mean (MeanT) and minimum (MinT) temperatures for the onset and ending of xylogenesis.

Site	Year	Onset of xylogenesis			Ending of xylogenesis		
		MinT(°C)	MeanT(°C)	MaxT(°C)	MinT(°C)	MeanT(°C)	MaxT(°C)
1	2007	0.6 $\pm$ 0.2	4.3 $\pm$ 0.1	9.7 $\pm$ 0.4	0.8 $\pm$ 0.3	3.9 $\pm$ 0.3	9.0 $\pm$ 0.4
	2008	0.8 $\pm$ 0.2	3.7 $\pm$ 0.3	8.9 $\pm$ 0.4	0.6 $\pm$ 0.2	3.5 $\pm$ 0.2	8.7 $\pm$ 0.1
	2009	0.8 $\pm$ 0.3	4.4 $\pm$ 0.1	9.8 $\pm$ 0.2	1.1 $\pm$ 0.1	4.7 $\pm$ 0.1	9.9 $\pm$ 0.1
	2010	0.3 $\pm$ 0.1	3.4 $\pm$ 0.2	9.1 $\pm$ 0.3	0.4 $\pm$ 0.1	2.9 $\pm$ 0.3	8.0 $\pm$ 0.2
2	2007	0.5 $\pm$ 0.3	4.3 $\pm$ 0.2	9.4 $\pm$ 0.5	0.9 $\pm$ 0.4	4.1 $\pm$ 0.5	9.3 $\pm$ 0.6
	2008	1.0 $\pm$ 0.3	3.9 $\pm$ 0.3	8.7 $\pm$ 0.4	0.7 $\pm$ 0.3	3.6 $\pm$ 0.2	8.7 $\pm$ 0.1
	2009	0.8 $\pm$ 0.3	4.4 $\pm$ 0.4	9.8 $\pm$ 0.2	1.1 $\pm$ 0.3	4.5 $\pm$ 0.4	9.8 $\pm$ 0.3

Table 2 The best estimate of parameters of VS model used in this study.

Model parameter	Description(unit)	Value
CT_{min}	Minimum temperature for tree growth ($^{\circ}C$)	0.9
T_{opt1}	Lower end of range of optimal temperatures ($^{\circ}C$)	5.9
T_{opt2}	Upper end of range of optimal temperatures	9.3
T_{max}	Maximum temperature for tree growth ($^{\circ}C$)	19.9
W_{min}	Minimum soil moisture for tree growth (v/v)	0.06
W_{opt1}	Lower end of range of optimal soil moisture (v/v)	0.18
W_{opt2}	Upper end of range of optimal soil moisture (v/v)	0.22
W_{max}	Maximum soil moisture for tree growth (v/v)	0.50
T_{beg}	Temperature sum for initiation of growth ($^{\circ}C$)	30
D_{root}	Depth of root system (mm)	50
P_{max}	Maximum daily precipitation for saturated soil (mm)	20
K_1	Fraction of precipitation penetrating soil (dimensionless)	0.86
K_2	First coefficient for calculation of transpiration (mm/day)	0.12
K_3	Second coefficient for calculation of transpiration ($1/^{\circ}C$)	0.175
K_r	Coefficient for water infiltration from soil (dimensionless)	0.006

Figure legends:

Figure 1. Micrometeorological conditions (2007 – 2010) at the upper treeline in the Sygera Mts., SE Tibetan Plateau, showing (a) daily mean air temperature and daily total precipitation, (b) daily soil temperature (at depths of 10, 20 and 40 cm) and snow depth, and (c) daily mean soil volumetric moisture contents (at depths of 10, 20 and 40 cm).

Figure 2. Onset (a), ending (b) and duration (c) of xylem formation of Smith fir (*Abies georgei* var. *smithii*) based on weekly xylogenesis observations at Site 1 (4360 m a.s.l.) (grey columns) and Site 2 (4250 m a.s.l.) (white columns). Error bars indicate standard deviation among trees.

Figure 3. Lagged (0 - 10 days) Pearson correlation coefficients between xylem cell production (a, b), growth index (c, d) and corresponding climatic variables. P0 represents the weekly climatic mean for the exact period between two sampling dates. P1 to P10 represent the weekly means lagged 1 - 10 days before the sampling date. Dotted horizontal lines show the 95% confidence limits. *Abbreviations:* MaxT = maximum temperature, MeanT= mean temperature, MinT= minimum temperature, P = precipitation and GGD = growing degree days above 5°C.

Figure 4. Critical minimum (black dots), mean (white dots) and maximum (grey dots) air temperatures at Sites 1 and 2, corresponding with the 0.5-probability of the onset and ending

of xylem formation according to xylogenesis observations in Smith fir. Error bars indicate the standard deviation among trees.

Figure 5. Observed (solid line) and simulated (dashed line) tree-ring width indices at Smith fir treeline in the Sygera Mts. on the south-eastern Tibetan Plateau, 1960 - 2006.

Figure 6. Pearson correlation coefficients between the observed and calculated values of tree ring width when the parameter of CT_{min} varied.

Figure 1

Figure 2

Figure 3

Figure 4

577

578

579 Figure 5

580

581

582 Figure 6

583

SUPPORTING INFORMATION

Table S1 Parameters of the Gompertz function (A , β , K), R^2 and day of the inflection point (tp) for Smith fir growing at two treeline sites, 2007 – 2010.

	Year	A	β	$K(10^{-2})$	tp	R^2
Site 1	2007	23.95	5.48	3.25	170.59	0.98
	2008	22.97	5.97	3.32	179.32	0.96
	2009	25.68	7.07	4.08	174.50	0.96
	2010	21.08	5.67	3.25	176.67	0.99
Site 2	2007	24.79	5.03	2.85	176.56	0.98
	2008	21.22	6.97	3.87	178.13	0.98
	2009	23.62	5.70	3.02	188.53	0.98

Figure S1. Minimum air temperatures recorded by the automatic weather station (black line) and temperature data logger at Site 1 (red line) and Site 2 (blue line) from August 5, 2011 to August 5, 2013.

Figure S2. Dynamics of xylem growth (including enlarging, wall thickening and mature xylem cells) at two Smith fir treelines as modeled using a Gompertz function.

Figure S3. Correlations between the simulated tree ring chronology and monthly temperature and precipitation at Smith fir treeline in the Sygera Mountains on the south-eastern Tibetan Plateau. Dotted horizontal lines show the 95% confidence limits. *Abbreviations:* MaxT = maximum temperature, MeanT= mean temperature, MinT= minimum temperature and P = precipitation.