Water availability drives signatures of local adaptation in whitebark pine (*Pinus albicaulis* Englm.) across fine spatial scales of the Lake Tahoe Basin, USA

Brandon M. Lind^{*1}, Christopher J. Friedline[†], Jill L. Wegrzyn[‡], Patricia E. Maloney[§], Detlev R. Vogler^{**}, David B. Neale^{††}, and Andrew J. Eckert[†]

^{*}Integrative Life Sciences program, Virginia Commonwealth University, Richmond, Virginia 23284 USA

[†]Department of Biology, Virginia Commonwealth University, Richmond, Virginia 23284 USA

[‡]Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT 06269 USA

[§]Department of Plant Pathology and Tahoe Environmental Research Center, University of California, Davis, California 95616 USA

^{**}USDA, Forest Service, Pacific Southwest Research Station, Institute of Forest Genetics, 2480 Carson Road, Placerville, California 95667 USA

^{††}Department of Plant Sciences, University of California, Davis California 95616 USA

Data reference numbers (XXXXXXXXXXXX)

¹ Corresponding Author. Current Address: Department of Biology, Virginia Commonwealth University, Richmond, VA 23284. E-mail: lindb@vcu.edu

19 **Running Title:** Local adaptation of *P. albicaulis*

20

21 **Keywords:**

22 Local adaptation, *Pinus albicaulis*, water availability, Sierra Nevada, linkage disequilibrium

23

24 Corresponding Author:

25 Brandon M. Lind

26 1000 West Cary Street

27 Life Science Bldg. - Department of Biology Rm 126

28 Virginia Commonwealth University

29 Richmond, Virginia 23284 USA

30 e-mail: lindb@vcu.edu, fax: 804-828-0820

ABSTRACT

Patterns of local adaptation at fine spatial scales are central to understanding how evolution proceeds, and are essential to the effective management of economically and ecologically important forest tree species. Here, we employ single and multilocus analyses of genetic data ($n = 116,231$ SNPs) to describe signatures of fine-scale adaptation within eight whitebark pine (*Pinus albicaulis* Engelm.) populations across the local extent of the environmentally heterogeneous Lake Tahoe Basin, USA. We show that despite highly shared genetic variation ($F_{ST} = 0.0069$) there is strong evidence for adaptation to the rain shadow experienced across the eastern Sierra Nevada. Specifically, we build upon evidence from a common garden study and find that allele frequencies of loci associated with four phenotypes (mean = 236 SNPs), 18 environmental variables (mean = 99 SNPs), and those detected through genetic differentiation ($n = 110$ SNPs) exhibit significantly higher signals of selection (covariance of allele frequencies) than could be expected to arise, given the data. We also provide evidence that this covariance tracks environmental measures related to soil water availability through subtle allele frequency shifts across populations. Our results replicate empirical support for theoretical expectations of local adaptation for populations exhibiting strong gene flow and high selective pressures, and suggest that ongoing adaptation of many *P. albicaulis* populations within the Lake Tahoe Basin will not be constrained by the lack of genetic variation. Even so, some populations exhibit low levels of heritability for the traits presumed to be related to fitness. These instances could be used to prioritize management to maintain adaptive potential. Overall, we suggest that established practices regarding whitebark pine conservation be maintained, with the additional context of fine-scale adaptation.

INTRODUCTION

The study of local adaptation has been an integral part of evolutionary biology as a whole, as local adaptation influences a wide variety of biological patterns and processes (reviewed in Savolainen *et al.* 2013). Trees in particular have received much attention in this regard because many species are ecologically and economically important, and high outcrossing rates (Neale & Savolainen 2004) result in large effective population sizes (which increase the effectiveness of selection) as well as weak neutral genetic differentiation (which decreases the confounding effects of selection and population structure), which together create ideal circumstances in which to detect selective processes in nature (Savolainen & Pyhäjärvi 2007). Investigators seeking to explain the genetic basis of local adaptation in trees, and plants in general, have been motivated by observations of significant differentiation for quantitative genetic variation across populations (e.g., Q_{ST}). If such phenotypes have a genetic basis, the underlying loci may be differentiated among populations as well (Endler 1977; reviewed in Storz 2005, Haas & Payseur 2016). In these cases, loci contributing to local adaptation could be identified through genetic indices of differentiation, or by targeting trait- or environmentally-associated loci that stand out above background demography. Yet, theoretical (Latta 2003; Le Corre & Kremer 2003) and empirical (Hall *et al.* 2007; Luquez *et al.* 2007) investigations have shown that discordance between Q_{ST} and F_{ST} of causative loci can occur under adaptive evolution. Moreover, as the number of underlying loci increases, the divergence between these indices increases as well, and the contribution of F_{ST} to any individual underlying locus decreases. In cases that exhibit strong diversifying selection and high gene flow, this adaptive divergence results from selection on segregating genetic variation (Hermisson & Pennings 2005; Barret & Schluter 2008) and is attributable to the among-population component of linkage disequilibrium (Ohta 1982, Latta 1998). In the short term, local adaptation will be realized through subtle

coordinated shifts of allele frequencies across populations causing covariance (i.e., LD) among many underlying loci (Latta 1998; Barton 1999; Latta 2003; McKay & Latta 2002; Kremer & Le Corre 2012; Le Corre & Kremer 2012), such that adaptation need not take place through numerous fixation events or sweeping allele frequency changes (MacKay *et al.* 2009; Pritchard & di Rienzo 2010). Over many thousands of generations, these shifts can lead to concentrated architectures of large-effect loci with a reduction of those with small effect (Yeaman & Whitlock 2011). For studies investigating continuous phenotypes such as those often related to fitness, even among populations with highly differentiated phenotypic traits sampled under a robust design (Lotterhos & Whitlock 2015), it may be difficult to identify many of the loci underlying the quantitative trait in question. Thus, for many species, specifically across fine spatial scales, the signal of local adaptation within much of current genetic data may go largely undetected using only single-locus approaches (Latta 1998; 2003; Le Corre & Kremer 2003; Yeaman & Whitlock 2011; Kemper *et al.* 2014), resulting in calls for theory and empiricism that move beyond single-locus perspectives (Pritchard & di Rienzo 2010; Sork *et al.* 2013; Tiffin & Ross-Ibarra 2014; Stephan 2015).

Populations of forest trees, particularly conifers, have a rich history of common garden, provenance tests, and genecological studies that demonstrate abundant evidence for local adaptation among populations, even over short geographic distances (e.g., Mitton 1989; 1999; Budde *et al.* 2014; Csilléry *et al.* 2014; Vizcaíno *et al.* 2014; Eckert *et al.* 2015) providing further support that fine spatial scales are relevant to adaptation (Richardson *et al.* 2014). This extensive history has also revealed the highly polygenic nature of adaptive traits (Langlet 1971; Holland 2007). Even so, the majority of these investigations have been limited to single-locus perspectives using either candidate genes (e.g., González-Martínez *et al.* 2008; Eckert *et al.* 2009) or a large set of molecular markers (e.g., Eckert *et al.* 2010) to explain the genetic basis

of local adaptation. In most cases, a few loci underlying the adaptive trait in question are identified and generally explain a small to moderate proportion of the overall heritability of the trait (Neale & Savolainen 2004; Savolainen *et al.* 2007; Ćalić *et al.* 2016). Yet because of the presumed polygenic nature underlying these adaptive phenotypic traits, and because past investigations have generally applied single-locus perspectives, it is likely that a majority of the genetic architecture of local adaptation in trees remains undescribed (Savolainen 2007; Sork *et al.* 2013; Ćalić *et al.* 2016).

Spurred in part by the advance of theory and availability of genome-wide marker data, attention has been refocused to describe underlying genetic architectures from a polygenic perspective. This transition began in model organisms (e.g., Turchin *et al.* 2012) and has expanded to other taxa such as stick insects (Comeault *et al.* 2014; 2015), salmon (Bourret *et al.* 2014), and trees (Ma *et al.* 2010; Csilléry *et al.* 2014; Hornoy *et al.* 2015). Indeed, species that occupy landscapes with high degrees of environmental heterogeneity offer exemplary cases with which to investigate local adaptation. Near its southern range limit, whitebark pine (*Pinus albicaulis* Engelm.) populations of the Lake Tahoe Basin (LTB) inhabit a diversity of environmental conditions. As exemplified by the strong west to east precipitation gradient (see Figure 1), many of the environmental characteristics of the LTB vary over short physical distances (<1km) and have the potential to shape geographic distributions of *P. albicaulis* at spatial scales below those typically investigated (i.e., range-wide studies) for forest trees. Local spatial scales are of particular interest to resource and conservation agencies as this is the scale at which most management is applied. Here, we build upon past work from a common garden (Maloney *et al.* in review) to investigate the genetic architecture of fine-scale local adaptation across *P. albicaulis* populations of the LTB by exploring the relationships between genotype, 18 environmental variables, and five fitness-related phenotypic traits using both

single and multilocus approaches. Specifically, we use the *P. albicaulis* populations of the LTB to address the following four questions: (i) Is there evidence that a long-lived, outcrossing plant species exhibiting high levels of gene flow can be locally adapted across fine spatial scales? (ii) What is the genetic basis and relationship among loci underlying adaptation in such a species? (iii) How similar are the genetic bases of fitness-related phenotypes to the loci putatively under selection from the environment? Using this information, we will contextualize how instances of fine-scale adaptation have management implications. This study highlights the advantages of a polygenic perspective and investigates signatures of local adaptation using a large set of null markers to determine the extremity of allele covariance among putatively adaptive loci where others have relied on simulation or null candidate genes. Furthermore, this work provides additional empirical evidence for theoretical predictions of covariance among adaptive loci found by other studies in trees.

MATERIALS and METHODS

Focal species, study area, and sampling

A principal component of high elevation forests in California and Nevada, *P. albicaulis* is widespread throughout subalpine and treeline environments and plays a vital role in ecosystem function and services including food resources for wildlife, forest cover, watershed protection, protracting snowmelt, and biodiversity (Hutchins & Lanner 1982; Farnes 1990; Tomback *et al.*, 2001; McKinney *et al.*, 2009; Tomback & Achuff 2010; Tomback *et al.* 2016). Most of the species' distribution is outside of California, extending northward into Oregon, Washington, British Columbia, and Alberta and eastward into northern Nevada, Idaho, Montana, and Wyoming (Critchfield & Little 1966; Tomback & Achuff 2010). Whitebark pine is a foundation species in subalpine ecosystems throughout most of its range in western North America (Ellison *et al.* 2005) and is threatened by fire-suppression, climate change, the non-native pathogen

white pine blister rust, caused by *Cronartium ribicola* J.C. Fisch., and mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Tomback & Achuff 2010; Mahalovich & Stritch 2013).

The LTB lies within California and Nevada in the north-central Sierra Nevada range, varies in elevation from 1900 to 3300m, and is flanked to the west by the Sierra Nevada crest and to the east by the Carson Range. The LTB experiences a Mediterranean climate with warm, dry summers and cool, wet winters. Precipitation falls during the winter months, most often in the form of snow, with a strong west-east gradient. The geology of the region is dominated by igneous intrusive rocks, typically granodiorite, and igneous extrusive rocks, typically andesitic lahar, with small amounts of metamorphic rock (USDA NRCS 2007).

Each of the eight study populations (three subplots per population) were located in a distinct watershed and distributed around the Basin to capture variation in the physical environment (e.g., climate, geology, and topography; Figure 1). Needle tissue was sampled in the summer of 2008 from 244 *P. albicaulis* trees (Table 1). From these eight populations, six populations were chosen to sample cones from 88 of the trees that were sampled for needle tissue. All samples were collected from trees separated by 30 to 1000m, with an average interpopulation distance of 31km. Universal Transverse Mercator coordinates, elevation, slope, and aspect (USDA FS FHTET) were used with the PRISM climatic model (Daly *et al.* 1994) to determine climatic parameters of sampled areas from 1971-2000, while soil survey data (USDA NRCS 2007) were used to describe the edaphic conditions of the LTB (Table 1).

Common gardens and phenotypic measurements

Fitness-related traits related to survival, especially during seedling and juvenile stages, are an important component of total lifetime fitness (e.g., Postma & Ågren 2016), particularly for forest trees, and are likely to be composed of phenotypic traits related to growth, phenology, resource allocation patterns, water-use efficiency, and disease susceptibility. In order to

estimate early-lifetime phenotypes of mother trees, seeds sampled from 11 to 19 maternal trees ($n = 88$) located in six of the eight populations were established in a common garden (Table 1) using a random block design (for further details see Maloney *et al.* in review). Growth (height), phenology (budset), water-use efficiency ($\delta^{13}\text{C}$), and resource allocation [root:shoot biomass, $\text{N}(\mu\text{g})$] were measured when seedlings reached ~2 years in age (see Maloney *et al.* in review for details). Height was recorded in April and October 2011, while 2 seedlings per family per block were harvested, clipped above the root collar, dried, and weighed to determine root and shoot biomass. For $\delta^{13}\text{C}$ and $\text{N}(\mu\text{g})$ analysis, needle tissue from 1 seedling per family per block was harvested, coarsely ground, and dried at 60°C for 96 hours. Between 2-3mg of tissue per sample was sent to the Stable Isotope Facility at UC Davis for isotope analyses (<http://stableisotopefacility.ucdavis.edu/>).

DNA extraction, sequencing, and analysis

Total genomic DNA was isolated from finely ground needle tissue sampled from 244 trees across all eight populations using the Qiagen DNEasy 96 Plant kit according to protocol (Qiagen, Germantown, MD). Restriction site-associated double digests of total genomic DNA using *MseI* and *EcoRI* enzymes (ddRADSeq, Peterson *et al.* 2012) were used to prepare three multiplexed libraries of up to 96 individuals each, as in Parchman *et al.* (2012). Total genomic DNA was digested and ligated with adapters containing amplification and sequencing primers as well as barcodes for multiplexing. These 96 barcodes (Parchman *et al.* 2012) are of 8-10bp sequences that differ by at least four bases. Ligated DNA was then amplified using high-fidelity PCR. Using the QIAquick Gel Extraction Kit (Qiagen), amplified fragments were then isolated near 400bp of pooled PCR product separated in a 1% agarose gel at 100V for one hour. Single-end sequencing of libraries was carried out on the Illumina HiSeq 2500 platform with a single library per flowcell lane. For added coverage, each library was sequenced twice using 50bp

reads and twice for 150bp reads, except Library 3 which was sequenced 4x for 150bp reads to increase optimality of the mapping reference individual. All sequencing was performed at the DNA Sequencing Facility of the University of California at Berkeley (<https://mcb.berkeley.edu/barker/dnaseq/home>). After calling genotypes, SNPs, and further filtering (see Supporting Information), we judged the veracity of our sequence data by mapping the empirical set of SNPs against the sugar pine (*P. lambertiana* Dougl.) reference genome (v1.0) using 85% similarity and 50 % length coverage thresholds (http://dendrome.ucdavis.edu/ftp/Genome_Data/genome/pinerefseq/Pila/v1.0/).

Identifying focal sets of loci

To identify genotype-environmental associations, we implemented *bayenv2* (v2.0; Coop *et al.* 2010; Günther & Coop 2013), a Bayesian single-locus approach that accounts for population history and gene flow before performing association analysis (Coop *et al.* 2010). To ensure convergence, we ran five independent chains of *bayenv2* using the empirical SNPs ($n = 116,231$), with 100,000 iterations for each SNP within each chain. MCMC convergence across chains was inspected using the *coda* library in R. For each SNP, we calculated the harmonic mean across chains for the Bayes factor ($\overline{BF}$) and absolute value of Spearman's ρ (hereafter $\overline{\rho_S}$). When calculating $\overline{BF}$, if a particular SNP returned Bayes factors greater than one for at least 3/5 chains, we would take the harmonic mean from this subset to avoid underestimation of the Bayes factor. However, if this was not the case ($BF > 1$ in $\leq 2/5$ chains) we took the harmonic mean from the values that were less than or equal to one. We identified focal SNPs by the intersection between the upper tail (99.5th percentile) of $\overline{BF}$ and the upper tail (99th percentile) of the absolute value of $\overline{\rho_S}$, as recommended in the *bayenv2* manual (v2.0; page 4).

To associate genotype with phenotype, we implemented a Bayesian sparse linear mixed model (BSLMM) from the *GEMMA* software package (Zhou *et al.* 2013). BSLMM is a hybrid of

LMM and Bayesian variable selection regression (BVSr) that also offers considerable statistical advantages over single-locus GWAS approaches (Guan & Stephens 2011; Ehret *et al.* 2012; Zhou *et al.* 2013; Moser *et al.* 2015). Specifically, to describe the underlying genetic architecture, BSLMM uses priors (described below) and attributes of the genetic data to estimate the number of underlying SNPs (N_{SNP}), the posterior inclusion probability (γ , hereafter PIP) for individual SNPs as well as the proportion of phenotypic variance explained by the polygenic and sparse effects of each SNP (PVE).

Before input to GEMMA, the empirical set of SNPs was reduced to include only those individuals with seedlings in the common garden ($n = 88$), and loci which had MAF below 0.01 due to this reduction were eliminated alongside monomorphic SNPs. For each phenotype, we ran four independent chains for the BSLMM, with 1,000,000 warm-up steps and 50,000,000 steps in the MCMC, sampled every 1000th step. Priors for PVE by the model, h , were set as [0.01,0.9], and the $\log_{10}$ inverse number of SNPs, $\log_{10}(1/p)$, [-3.0,0.0], which equates to between 1 and 300 underlying loci (N_{SNP}). Convergence of the MCMC across chains was inspected using the coda library in R. To summarize the GEMMA output, we report means and 95% credible intervals for PVE and N_{SNP} from the posterior distributions. To assess significance of association of a SNP to a phenotype, we used the PIP from all four independent chains to calculate the harmonic mean ($\overline{PIP}$) and chose SNPs that were greater than or equal to the 99.9th percentile of $\overline{PIP}$ ($n \approx 116$ SNPs/phenotype) for each phenotype. We also explored SNPs with $\overline{PIP} \geq 99.8^{\text{th}}$ percentile ($n \approx 232$ /phenotype).

We implemented the program OutFLANK (Whitlock & Lotterhos 2015) to investigate the opportunity of detecting the environmentally- and phenotypically-associated loci using outlier approaches based on population genetic structure alone (e.g., F_{ST}). Using this approach and excluding loci with expected heterozygosity values below 10% with subsequent trimming of the

lower and upper 5% of empirical F_{ST} values, we inferred a null distribution of F'_{ST} and identified outlier loci with a false discovery rate of 5% from the empirical set of SNPs.

Inferring signatures of local adaptation

To determine if individual sets of focal loci (identified from GEMMA, bayenv2, and OutFLANK analyses) collectively exhibited elevated signatures of selection acting across multiple loci, we investigated the level of allele frequency covariance among all SNP pairs within each focal set. For instance, to calculate the covariance of allele frequencies across populations between two SNPs, SNP_i and SNP_j , within a focal set of SNPs associated with a particular phenotype in GEMMA, we used the global minor allele of each SNP, q , according to the interpopulation component of linkage disequilibrium,

$$\hat{D}_{a(ij)} = \sum_k \frac{n_k}{n} (q_{i,k} q_{j,k} - q_i q_j) \quad \text{Eq. (1)}$$

where n_k is the number of individuals in population k , n is the global population size, $q_{i,k}$ is the allele frequency of the i^{th} SNP in population k , $q_{j,k}$ is the allele frequency of the j^{th} SNP in population k , while q_i and q_j are the respective global allele frequencies of the i^{th} and j^{th} SNP across $k = 6$ populations (Storz & Kelly 2008, their Equation 2; Ma *et al.* 2010, their Equation 3). Because we chose the allele to use in comparisons based on global minor allele frequency, all calculations of $\hat{D}_{a(ij)}$ are therefore referenced to the global minor allele haplotype for a pair of SNPs. For populations that experience high levels of gene flow and divergent phenotypic optima due to selection, $\hat{D}_{a(ij)}$ is expected to be positive between allele frequencies of loci conferring a positive effect on the phenotype, negative between those conferring opposite effect, and zero between (conditionally) neutrally loci (eq. [6] in Latta 1998). Because we were not able to discern the direction of effect for alleles within each population (as in e.g., Gompert *et al.* 2015), and to facilitate comparison among analyses, we identified selective signatures by calculating

the absolute value of $\hat{D}_{a(ij)}$ for each locus pair. We also calculated $\hat{D}_{a(ij)}$ for focal SNPs associated with environmental variables from `bayenv2` and those identified as outliers from `OutFLANK`. In these two cases, we used allele frequencies across all eight populations.

To be able to discern if the level of covariance of allele frequencies among SNPs identified by `GEMMA` (or another method; hereafter focal SNPs) was greater than that from SNPs randomly chosen from our dataset (i.e., than expected given the data), we first separated all SNPs in the dataset by their expected heterozygosity into bins of 0.01 ranging from 0 to 0.50 (e.g., a SNP with H_E of (0.000-0.010] would be binned into the first bin, while an H_E of (0.490-0.500] would be binned into the 50th). We then created a set of SNPs from which to take randomized draws by subtracting the focal SNPs from the full set of SNPs. Next, based on the occupancy of heterozygosity bins for a given focal set, we randomly selected remaining SNPs to create a null set. We chose SNPs randomly in this way, 1000 times, each time calculating the absolute value of $\hat{D}_{a(ij)}$ among SNP pairs within each set. From each of these 1000 distributions, we calculated 1000 median absolute $\hat{D}_{a(ij)}$ values to create a null distribution for use in comparison to the median absolute $\hat{D}_{a(ij)}$ from the focal set of SNPs. If the median $\hat{D}_{a(ij)}$ is greater among our focal SNPs than the 95th percentile of the null distribution of 1000 medians, we will conclude that the signature of selection among loci within our focal sets is greater than could have arisen by chance, given the data.

To infer signatures of allele frequency shifts associated to environment, we implemented an approach similar to Equation 1 but instead of estimating $\hat{D}_{a(ij)}$ across all populations we estimated $\hat{D}_{a(ij)}$ across populations in a pairwise fashion (hereafter $pw\hat{D}_{a(ij)}$) using focal SNPs from a given method. In this case, we calculated global allele frequency (q_i or q_j) based on the frequency of allele q across the $k = 2$ populations (pop_l and pop_m) under consideration (where $n_l + n_m = n$). From these estimates, we created a symmetric matrix of $pw\hat{D}_{a(ij)}$ with columns

and rows for populations, and distances within the diagonal set to zero. We then implemented Mantel tests (Mantel 1967) using $pw\hat{D}_{a(ij)}$ matrices against other population pairwise distance matrices such as geographic distance inferred using great circle distances (km) following Vincenty's method, and Euclidian distance matrices for each of the five phenotypes and 18 environmental variables. Because we chose to take absolute values of $pw\hat{D}_{a(ij)}$ for each locus pair (as with $\hat{D}_{a(ij)}$) we note that the sign of the correlation coefficient, r , from Mantel tests may reflect the opposite directionality for any given SNP pair. Mantel tests were run with 9999 iterations using the `skbio` package (v0.4.2) in `Python`. Each environmental or phenotypic value was centered and standardized across populations before calculating Euclidian distances, but not for $pw\hat{D}_{a(ij)}$ or geographic distance matrices. For each set of focal SNPs associated with phenotype or environment, we also quantified the mean allele frequency differences across populations and compared this to 1000 sets of random SNPs chosen by H_E .

RESULTS

SNP filtering and characterization

After calling genotypes, SNPs, and filtering (see results section of Supporting Information), we retained 116,231 imputed SNPs for use as the empirical set in downstream analyses (Table S1). Of these contigs, 107,354 (92.4%) mapped to the *P. lambertiana* reference genome, thus lending authenticity to our sequence data. However, we avoid further discrimination of loci for (proximity to) genic regions until a future genome update with increased curation and density of annotation.

Overall, populations show little genetic structure with plots accounting for less than 1% of the variance in allele frequencies ($F_{\text{plot,total}} = 0.00687$; 95% credible interval: 0.0067-0.0070). Of this variation, 56.6% was accounted for by populations ($F_{\text{pop,total}} = 0.00389$; 95% CI: 0.0038-0.0040) with the remainder due to plots within populations ($F_{\text{plot,pop}} = 0.00299$; 95% CI: 0.0029-

0.0031). We found similar patterns among the locus-specific estimates of F_{ST} (Figure S1). Moreover, we found no discernable clustering of populations using PCA, respectively accounting for 5.6% and 1.2% of the variance in allele frequencies (Figure S2). To further address applicability of the island model used for calculation of $\hat{D}_{a(ij)}$ and $pw\hat{D}_{a(ij)}$, we analyzed population pairwise F_{ST} according to Weir & Cockerham (1984) using the `hierfstat` package in R. Results show little differentiation among populations (mean = 0.005, max = 0.016) with no evidence of isolation by distance (Mantel's $r = 0.0990$, $p = 0.2310$).

Genotype-environment analyses

To explore the degree of association among environmental variables between populations, we used Mantel tests between Euclidian environmental distance matrices. In most cases, we found significant correlations with many of the edaphic variables measured for this study, as well as between latitude and elevation ($r = 0.3988$, $p = 0.0490$), longitude and annual precipitation ($r = 0.7145$, $p = 0.0030$), and between percent maximum solar radiation and latitude distances ($r = 0.4629$, $p = 0.0370$; Table S2). Additionally, geographic distance among populations was only associated with latitude ($r = 0.9631$, $p = 0.001$), percent maximum solar radiation input ($r = 0.3992$, $p = 0.0468$), and elevation ($r = 0.4062$, $p = 0.0452$), the three of which were correlated environmentally (Table S2), but not to any of the remaining environmental variables (Mantel tests $p > 0.3131$, data not shown).

Through the intersection of the top 0.5% of $\overline{BF}$ and top 1% of $\overline{p_s}$, `bayenv2` analysis revealed between 14 (CEC) and 157 (GDD-Aug) focal SNPs associated with environment (Table S3). However, when calculating the $\overline{BF}$ for each SNP, it was never the case that more than two of the five chains produced $BF > 1$, of which chains with large values were driven primarily by seed number (we used additional seed numbers for a small subset of the data during exploration, data not shown). The range of $\overline{p_s}$ across all focal SNPs across all

environments varied from a minimum of 0.138 to a maximum 0.345 (Table S3). Additionally, the focal SNPs identified by `bayenv2` displayed a bias towards SNPs with low values of H_E (Figures S3-S4, see results section of Supporting Information) when compared to the distribution from the full set of SNPs (Figure S5). As such, our environmental associations should be interpreted with caution, as we did not have any SNPs with $\overline{BF} > 1$ nor do our harmonic mean nonparametric correlations exceed 0.35. Even so, when we compared absolute estimates of $\hat{D}_{a(ij)}$ among focal SNPs against the corresponding 1000 null sets of loci, we found that for all focal sets the median $\hat{D}_{a(ij)}$ was always greater than the 100th percentile of the null distribution (Figure 2, Table S3). The magnitude of this difference varied across environmental variables, being the smallest for percent clay (1.17x) and largest for annual precipitation (5.10x, Table S3). The percentile of the focal $\hat{D}_{a(ij)}$ distribution corresponding to the 100th percentile of the associated null set varied across environmental variables as well, reaching just the 3rd percentile for minimum January temperature and the 43rd percentile for percent clay (Table S3). Upon comparison of focal and null sets in both the distribution of single-locus and multilocus F_{ST} , focal sets were representative of single-locus estimates of the null sets (Figures S6-S7), and generally greater than the distribution of multilocus F_{ST} than the null (Figures S8-S9). Single-locus results suggest that many focal SNPs are unlikely outliers for F_{ST} , while multilocus comparisons exemplify the elevated frequency covariance of SNPs in focal sets. This data demonstrates that for most environmental variables the focal SNPs show higher degrees of $\hat{D}_{a(ij)}$ than could be expected, given the data, despite having low $\overline{BF}$ and $\overline{\rho_S}$.

Through the examination of patterns of allele frequency shifts ($pw\hat{D}_{a(ij)}$) across loci associated with environment we found no significant associations with geographic distance using Mantel tests ($p > 0.1116$). While this suggests the absence of linear allelic clines, it does not necessarily preclude the presence of environmental gradients or correlated patches as

suggested by environmental distance associations (Table S2). When we investigated the association between $pw\hat{D}_{a(ij)}$ matrices against the eponymous environmental distance matrix, we found significant association for annual precipitation ($r = 0.7134$, $p = 0.0027$), GDD-May ($r = 0.8480$, $p = 0.0013$), longitude ($r = 0.6522$, $p = 0.0024$), percent rock coverage ($r = 0.5124$, $p = 0.0145$), percent sand ($r = 0.5574$, $p = 0.0046$), minimum January temperature ($r = 0.5791$, $p = 0.0137$), and WC- $\frac{1}{3}$ bar (field capacity, $r = 0.4806$, $p = 0.0361$; Table 2). Additionally, we examined relationships between a particular $pw\hat{D}_{a(ij)}$ matrix and the 17 remaining environmental distance matrices and found significant associations in an additional 13 comparisons (Table 2), with five of these comparisons having $pw\hat{D}_{a(ij)}$ associated with either annual precipitation or longitudinal Euclidian distance. We also observed shifts of alleles associated with longitude or soil water capacity across six of the remaining eight significant associations (Table 2), with the remaining two significant associations among edaphic conditions of sand, silt, or clay. The magnitude of the mean allele frequency difference across populations of focal SNPs was subtle as expected (range 0.018-0.029) and were generally slightly larger than that predicted from random SNPs of the same heterozygosity (Figures S10-S11). Overall, our results indicate that the vast majority of subtle allele frequency shifts among loci associated with environment ($pw\hat{D}_{a(ij)}$) have significant associations related to interpopulation distances of water availability (Table 2).

Genotype-phenotype analyses

Phenotypic traits were heritable and structured across populations (bud flush $h^2 = 0.3089$, $Q_{ST} = 0.0156$; $\delta^{13}C$ $h^2 = 0.7787$, $Q_{ST} = 0.0427$; height $h^2 = 0.0608$, $Q_{ST} = 0.0418$; N(μ g) $h^2 = 0.3525$, $Q_{ST} = 0.0191$; root:shoot $h^2 = 0.3240$, $Q_{ST} = 0.0110$; Table S4) and were correlated with environmental variables (both climate and soil) in ways unexplainable by neutral evolutionary forces (Maloney *et al.* in review). Additionally, bud flush and $\delta^{13}C$ had significant $Q_{ST} > F_{ST}$

(Maloney *et al.* in review, using the genetic data presented herein). We used a subset of the empirical set of SNPs for use in genotype-phenotype analysis, after filtering we retained 115,632 SNPs. PCA revealed a similar pattern to the empirical set of SNPs (data not shown). Using three significant axes of population structure identified through Tracy-Widom tests, we associated SNPs to phenotypes with BSLMM (Zhou *et al.* 2013) using the top 99.9th and 99.8th percentiles of $\overline{PIP}$ (Figure 4, Table S5, Figure S12). From observations of density and trace plots, we concluded that the posterior distributions across chains were converging (not shown). The H_E of focal loci were generally representative of the empirical set (Figures S13-S14).

Overall, the genetic variance of SNPs included in the polygenic model explained between 14.4% [N(μ g)] and 37.6% (root:shoot) of the variance in the phenotypes measured in our study (PVE , Figure 3, Table S4). For many of the measured phenotypes, a considerable proportion of the narrow sense heritability estimated previously was therefore accounted for in the estimates of PVE (Table S4), as should be the case with sufficient genetic sampling (Gompert *et al.* 2016). Interestingly, in the case of height, PVE exceeded the upper confidence interval of the estimated h^2 (Table S4).

To acquire estimates of PVE from the identification of loci with large effects on phenotype, we conducted single-locus association using univariate linear mixed models implemented in GEMMA (see Supporting Information, Table S6). Across all phenotypes, there were no loci that exceeded the adjusted threshold for inclusion calculated from q -values with an FDR of 0.05 (Storey *et al.* 2015; $\sqrt{2 \cdot \ln(2)}$), with the minimum q -value across SNPs within phenotypes ranging between 0.2046 ($\delta^{13}\text{C}$) and 0.9999 [N(μ g)] (Table S6). Except for root:shoot biomass, the maximum likelihood estimates of PVE differed drastically from the estimates from BSLMM, with PVE never exceeding 1.08e-06 suggesting that a larger proportion of the heritable genetic variation for the traits measured here is explained by multiple SNPs than by individual SNPs

alone. Finally, to determine if LMM loci near the threshold were captured by the BSLMM for a particular phenotype, we isolated the loci from univariate LMM above a reduced threshold of $-\ln(p_{wald}) \geq 10$ (see Figure S15, Supporting Information). By this reduced threshold we identified one unique locus for both bud flush and $N(\mu g)$, four unique loci for both height and root:shoot biomass, and five unique loci for $\delta^{13}C$ (15 unique loci overall). We examined the focal loci sets identified from the 99.9th percentile of $\overline{PTP}$ in BSLMM for these LMM reduced-threshold loci and found 1 of the 4 LMM loci for both root:shoot biomass and height, and 2 of the 5 loci for $\delta^{13}C$. When we assessed the set of loci in the 99.8th percentile of BSLMM $\overline{PTP}$, we recovered all LMM reduced-threshold loci for bud flush and $N(\mu g)$ ($n = 1$), 1 of 4 loci for root:shoot biomass, 3 of 4 loci for height, and 3 of 5 loci for $\delta^{13}C$.

To determine if focal loci associated with phenotype by BSLMM exhibited evidence of selection, we estimated allele frequency covariance ($\hat{D}_{a(ij)}$) among focal SNPs and compared these estimates to 1000 null sets of SNPs. We found evidence for elevated covariance among the 99.9th percentile of $\overline{PTP}$ loci associated with bud flush and root:shoot biomass (Figure 4a, Table S5), with the latter exceeding the 100th percentile of the random distribution. To consider larger numbers of loci representative of the number of underlying loci estimated by BSLMM, we also isolated SNPs from the top 99.8th percentile of $\overline{PTP}$. In these sets, we found evidence for elevated signatures of selection acting across multiple loci for all phenotypes except for height, which did not produce a focal median $\hat{D}_{a(ij)}$ greater than the 95th percentile of null distribution of $\hat{D}_{a(ij)}$ (Figure 4b, Table S5). When focal and null sets of SNPs were compared, focal sets were representative of single-locus (Figure S16) and multilocus (Figure S17) F_{ST} estimates of the null sets.

To identify signatures of allele frequency shifts among focal loci associated with phenotype ($p w \hat{D}_{a(ij)}$), we ran Mantel tests of $p w \hat{D}_{a(ij)}$ matrices against geographic distance and

environmental Euclidian distance matrices. When considering SNPs identified by the 99.9th percentile of $\overline{PIP}$, we see substantial evidence for allele frequency shifts of loci associated with bud flush to Euclidian distances of GDD-May, GDD-Aug, percent maximum radiation input, and minimum January temperature (Table 3). Additionally, when we consider the 99.8th percentile of $\overline{PIP}$, we show evidence for allele frequency shifts among loci associated with bud flush, height, and $\delta^{13}\text{C}$ to Euclidian distances of annual precipitation, as well as for $N(\mu\text{g})$ loci to elevation, and bud flush loci to both longitude (a correlate of annual precipitation) and to percent maximum radiation input (as in the 99.9th $\overline{PIP}$ set). The strong signal from bud flush and water-related variable in Table 3 is intriguing, as bud flush and $\delta^{13}\text{C}$ were the only two phenotypic traits to have significantly larger Q_{ST} than F_{ST} (Maloney *et al.* in review). The magnitude of the mean focal allele frequency differences across populations were subtle as expected (range: 0.054-0.087) and representative of unassociated SNPs of similar H_E (Figure S18).

F_{ST} outlier analysis

From the empirical set of SNPs, OutFLANK analysis revealed 110 focal loci as outliers for F'_{ST} (range: 0.069-0.118). Expected heterozygosity values among the outlier SNPs (Figure S19) varied across the distribution from the full set of SNPs (Figure S5). Upon analysis of patterns of covariance ($\hat{D}_{a(ij)}$) among the OutFLANK focal SNPs, we found that the median focal $\hat{D}_{a(ij)}$ (6.08e-03) was 10.6x greater than the 100th percentile of the null distribution of $\hat{D}_{a(ij)}$ (5.74e-04). Moreover, the maximum median $\hat{D}_{a(ij)}$ from null sets corresponded to just the 12th percentile of the focal $\hat{D}_{a(ij)}$ values, suggesting that the majority of SNPs within the focal set showed higher levels of covariance among other outliers than expected by chance. However, when we analyzed these outlier SNPs for signatures of allele frequency shifts ($pw\hat{D}_{a(ij)}$) we found no significant associations with geographic or environmental distances.

Intersection of SNPs within and across methods

We examined overlap of focal SNPs among the various methods employed in this study (Table S7). While there was considerable overlap of loci found between methods, overall there was more overlap of loci associated with multiple phenotypes or with multiple environments than found across methods. For sets of loci associated with environmental variables, there was a considerable number of loci that were found to overlap among comparisons. This seemed to be driven by the correlations among soil properties, for when ordered by the number of loci within the intersection, 12 of the top 15 comparisons were among edaphic conditions. Additionally, climate-related variables relating to maximum radiation input, degree growing days, minimum and maximum temperature generally shared loci among soil variables relating to water availability while annual precipitation shared 18 loci with longitude, among other variables. Very few of the loci identified by *bayenv2* would have been detected through conventional F_{ST} outlier approaches (Figures S20-S21). Even so, *OutFLANK* captured between 1-3 ($n = 18$) of the loci identified across 10/18 environmental associations from *bayenv2* (many of which were water-related, including annual precipitation), and captured 4 of the loci identified in the 99.8th percentile of $\overline{PTP}$, but not for any of the loci identified from the reduced threshold of LMM. Among loci associated with phenotype (99.8th $\overline{PTP}$), there were between one and three loci which were found in the intersection among pairwise phenotypic comparisons, yet none of these overlap loci were those identified from LMM. Finally, 15 loci associated with environment overlapped with the 99.8th percentile of $\overline{PTP}$ (including two between $\delta^{13}C$ and longitude, a correlate of annual precipitation) while environmental associations did not capture any of the reduced-threshold loci from univariate LMM (Table S7).

DISCUSSION

The spatial extent of local adaptation, particularly in conifers, has generally been investigated at regional scales (Neale & Savolainen 2004; Savolainen *et al.* 2007; Čalić *et al.*

2016). While informative for range-wide inference, management and conservation agencies are often limited to local scales spanning only tens to several hundreds of square kilometers. While there is an expectation that high gene flow (i.e., migration load) exhibited by many conifers can lead to swamping of adaptive alleles, there is mounting empirical evidence that adaptation to the environment can still occur at relatively fine spatial scales (Mitton 1989; 1999; Budde *et al.* 2014; Csilléry *et al.* 2014; Vizcaíno *et al.* 2014; Eckert *et al.* 2015). Thus, studies which investigate adaptation at scales amenable to management may be of relatively greater importance (especially for endangered and threatened species) to reforestation applications such as those carried out through seed sourcing (*sensu* McLane & Aitken 2012) and replanting efforts. Previously, we provided evidence that measured fitness-related phenotypes are heritable, that population explains a significant proportion of phenotypic variation, and Q_{ST} was significantly greater than F_{ST} for bud flush and $\delta^{13}C$ (Maloney *et al.* in review). Here, our genetic analyses indicate selective pressures of *P. albicaulis* are likely driven by water availability (e.g., precipitation gradients) as well as interactions and correlates of soil properties, which lends replicative support to both theoretical and empirical predictions for the patterns of loci underlying quantitative traits undergoing selection with gene flow. For instance, Ma *et al.* (2010) assessed evidence for diversifying selection within European aspen (*Populus tremula* L.) across 23 candidate genes of the photoperiodic pathway using the covariance of allelic effects among loci, albeit across a geographic region of Sweden spanning 10 latitudinal degrees. From this candidate set, they identified high degrees of covariance among phenotypic effects as predicted from theory (Latta 1998), despite minimal allele frequency differentiation among sampled populations. More recently, Csilléry *et al.* (2014) assessed 53 climate-related candidate genes within European beech (*Fagus sylvatica* L.) providing evidence that covariance among loci is attributable to epistatic selection (*sensu* Ohta 1982) across fine spatial scales of less than

100km². While varying across spatial scales, these studies replicate evidence for a signal of local adaptation in trees through elevated among-population linkage disequilibrium between adaptive loci.

Standing genetic variation for fitness-related traits

The populations under study appear to have extensive gene flow, recent divergence, or both. Variation of allele frequencies among populations accounts for less than 1% of the variance observed, which was less than that found for *P. lambertiana* populations within the LTB (Eckert *et al.* 2015), or among isozymes sampled from populations across the Northern *P. albicaulis* range (Krakowski *et al.* 2003). Inspection of PCs showed no distinctive clustering of populations (Figure S2) while population pairwise F_{ST} did not exceed 0.016 and a test for isolation by distance was not significant. Consequently, the mean allele frequency differences between populations for focal SNPs were also subtle. Importantly, the majority of the populations under study exhibited significant, heritable genetic variation underlying the measured phenotypic traits. Biologically, such a pattern of extensive sharing of alleles across populations has likely resulted from a combination of long-distance pollen movement, and seed dispersal by Clark's nutcracker (*Nucifraga columbiana* Wilson) which is known to disperse seeds at distances similar to those between our sampled populations (Tomback 1982; Richardson *et al.* 2002 and references therein). Given this pattern of structure, the island model with symmetric migration used to describe the interpopulation component of linkage disequilibrium among loci ($\hat{D}_{a(ij)}$) and allele frequency shifts ($p_w \hat{D}_{a(ij)}$) is likely suitable to investigate our dataset for signatures of selection across multiple loci.

While bayenv2 did not identify any loci strongly associated with environment, as given from small values of Bayes factors (all $\overline{BF} < 1.0$, Table S3), there is a strong biological signal for adaptation to soil water availability in our dataset (discussed below), evidence that other white

pinus within the LTB are also being structured by precipitation differences among populations (Eckert *et al.* 2015), and elevated signals of selection among focal loci which would be unlikely to arise, given the data. Thus, it seems unlikely that the focal sets of SNPs are driven solely by false positives. However, if the majority of loci across `bayenv2` are not an artifact of the method (i.e., are within or in linkage with causative sites), one possible explanation for elevated covariance among focal loci is that the structure of environmental variables across populations captured variation for unmeasured phenotypic traits which were largely representative of total lifetime fitness (Schoville *et al.* 2012). Structure of unmeasured fitness-related traits may also explain the high covariance of `OutFLANK` loci. Future work could provide validation through functional analyses of loci or from similar patterns found in other systems.

Water availability as a driver of local adaptation

The strongest signal for local adaptation among *P. albicaulis* populations of the LTB came from evidence of adaptation to soil water availability (Figures 2 and 4; Tables 2-3). Indeed, water availability is a critical component shaping standing variation across plant taxa (Vicente-Serrano *et al.* 2013), including the distributions of tree species in general (van Mantgem *et al.* 2009; Allen 2010), and southern populations of *P. albicaulis* specifically (Bower & Aitken 2008; Chang *et al.* 2014). During the Pleistocene-Holocene transition (10,000-12,000yr BP), shifts from mesic to xeric conditions caused proximal *P. albicaulis* populations of the Western Great Basin (~50km distant) to shift from 1380m in elevation to their current position about 1100m down-slope (Nowak *et al.* 1994; *cf.* Table 1). Such shifts in climate and local edaphic conditions in the last 10,000yr may in part explain recent (relative to $4N_e$ generations), and ongoing, selective pressures on *P. albicaulis* populations of the LTB. Because of climatic constraints imposed on the southern range of *P. albicaulis*, phenotypic traits affected by precipitation, soil water availability, or soil water capacity likely have fitness-related consequences for this

species. Additionally, with climatic models predicting warmer temperatures, reduced snow accumulation, and earlier spring melt across the western USA, it is likely that *P. albicaulis* populations of the Sierra Nevada will continue to face selective pressures of this kind. Even so, many of *P. albicaulis* populations of the LTB exhibit substantial genetic variation for the fitness-related traits measured, suggesting that the majority of ongoing adaptation within the LTB will likely be unconstrained by the lack of genetic variation. Instead, other biotic factors (e.g., white pine-blister rust infection) that can lead to negative population growth rates may be of more immediate concern (see Maloney *et al.* 2012).

Implications to whitebark management

As pointed out by McLane & Aitken (2012), distribution models of many species predict climate niches to shift in the upcoming century, leaving great uncertainty that tree species in particular will be able to track suitable environments through natural migration and establishment, as the rate of many of these geographic shifts would far exceed observed post-glacial rates of migration (Davis & Shaw 2001; McLachlan *et al.* 2005). Exacerbating this issue, the presence of *C. ribicola*, and climate-driven outbreaks of mountain pine beetle (*D. ponderosae*) create further challenges to the conservation of *P. albicaulis* (Tomback & Achuff 2010; Mahalovich & Stritch 2013). Without intervention, such cases could lead to population collapse, extirpation, or extinction. As such, assisted gene flow, replanting, or restoration efforts will need to continue to take current and future selective pressures into account (e.g., genetic variation, resistance to *C. ribicola*, etc.), as has generally been the standard of practice (Keane *et al.* 2012; Maloney *et al.* 2012; McLane & Aitken 2012).

At the same time, the choice of seed source will also need to take into account local adaptation at fine spatial scales. While a small proportion of the neutral genetic variation (F_{ST}) is found among most tree populations (often less than 5%, Neale & Savolainen 2004), this does

not necessarily mean that seeds from within an established seed zone will be optimal for any given constituent environment, particularly if the seed zone exhibits high environmental heterogeneity (such as in montane regions), or if the seed zone is relatively broad compared to these environmental gradients, as is the case in California (see Buck *et al.* 1970). Weak neutral genetic differentiation can be misleading in this way, as polygenic traits influenced by selective processes in the face of gene flow may lead to divergent local adaptation through the covariance of alleles among populations without the buildup of substantial genetic differentiation at any given locus. Particularly in cases where there is evidence of local adaptation, and when ethical (McLachlan *et al.* 2007), appropriate (considering e.g., ecological or demographic factors), or plausible, seed source should come from local sources (i.e., relative to scales of geography and environmental gradients) to maximize adaptive potential (McKay & Latta 2002). This is particularly important when maximal fitness (e.g., reproductive output) is a priority for established trees, as while many genotypes may survive, realized phenotypes related to fitness may be suboptimal for a given environment. In cases where local seed source is not plausible, perhaps due to (inbreeding through) isolation, or with prohibitively small population sizes, sources likely to perform well are also likely to come from highly correlated environments, particularly if the populations are not highly diverged. With this taken into consideration, management may be able to prioritize populations for restoration through estimates of trait heritability (the prerequisite for adaptation), as estimates of genetic variation alone may be misleading. For instance, heritability for $\delta^{13}\text{C}$ was near zero for the Rifle Peak population, as was the case for root:shoot ratio in Freel Peak and Little Roundtop, $\delta^{15}\text{N}$ in Rifle Peak, and height for four of six populations. While this could mean the presence of recent, strong natural selection, or, conversely, that these traits do not convey an overwhelming adaptive advantage in these populations, candidate traits with substantial evidence for contributing to total lifetime

fitness should be monitored nonetheless. In appropriate cases, introducing compatible variation would stand to improve adaptive potential in such populations. While there is no one specific solution to conserve whitebark populations across its range, taking into consideration fine-scale local adaptation in addition to established strategies will likely aid in such endeavors.

Concluding remarks, limitations, and future work

The results reported here suggest that focal loci collectively show elevated allele frequency covariance (a signal expected between loci undergoing selection with gene flow) across multiple loci than could have been expected to arise from the data by chance, and that interpopulation levels of allele frequency covariance are often associated with interpopulation distances of soil water availability. Our results further explain a considerable proportion (*PVE*) of the additive genetic variation (h^2) of the quantitative traits under study from a polygenic perspective, as should be the case with sufficient genetic sampling. Thus, we can posit that the general mode of adaptation for *P. albicaulis* across the LTB is facilitated by selection on standing levels of genetic variation that is extensively shared throughout the basin and likely improves performance in early life stages. Finally, if soil and climatic variables continue to influence the extant populations within the LTB as evidenced from our analyses, it is likely that these variables will continue to be important to the long-term success of this threatened keystone species.

While we described associations among genotype, phenotype, and environment that reflect strong evidence for adaptive responses of *P. albicaulis* populations to the environment, we acknowledge several shortcomings. First, our study design was limited in statistical power which could have been improved by increasing the number of individuals sampled, the total number of populations, or both, given an ideal sampling regime (Lotterhos & Whitlock 2015) which together would have further facilitated other methodologies for uncovering evidence of polygenic local

adaptation (e.g., Berg & Coop 2014). Second, while we measured fitness-related traits among seedlings of a species whose lifespan differs by several orders of magnitude, establishment success is one of the primary factors influencing dynamics of forest populations, and early life stages of plants have been shown to be a major component of total lifetime fitness (Postma & Ågren 2016). Third, much of the statistical signal for the association of allele frequency shifts to environment would be lost with correction for multiple tests yet we leverage the fact that, of the few significant $pw\hat{D}_{a(ij)}$ associations, the majority were related to $\delta^{13}C$, annual precipitation, its correlate of longitude, or measures of soil water availability, an outcome highly unlikely by chance alone. Fourth, while we provide evidence for statistical signals predicted by theory, our methodology limited us from making conclusions regarding local adaptation *sensu stricto* as we utilized just a single common garden without reciprocal transplants and were unable to quantify functional differences of putative loci among populations. Reciprocal transplants would have allowed us to differentiate pleiotropic effects and facilitate direct measures of fitness through survival and growth across environments. Finally, a more fully curated, well-annotated genome assembly and accompanying linkage map would have aided in the detection of physical linkage among SNPs, proximity to genomic regions of estimated effect, and detection of false positives. For instance, the *P. lambertiana* genome used to judge authenticity of sequence data does not yet have the density of annotation needed to draw inferences on the causative sites likely within or linked to the loci described here, as its assembly and curation are still ongoing. Because of this, we cannot confidently estimate the proportion of the polymorphism due to coding and non-coding sites nor conclude that the linkage inferred for focal loci are not an artifact due to distant linkage with causative sites of larger effect. For this artifact to be true, however, our results would indicate that many of the loci in focal sets of SNPs would all have to be linked to the same smaller number of larger-effect loci, or that many large-effect loci underlie the trait, both

652 unlikely outcomes given the expectations of quantitative traits and the coverage from ddRADseq
 653 methods found in other white pines (e.g., Friedline *et al.* 2015). Even so, while we may not have
 654 identified all causative sites, linked sites can still maintain signals of evolutionary processes
 655 (McVean 2007). Moreover, signatures of selection presented here are unlikely to have arisen
 656 from our dataset by chance alone (as evidenced from null models) and our inferences were
 657 synthesized from prevailing signals across multiple phenotypic, environmental, and taxonomic
 658 lines of evidence, as opposed to inferences based on single variables or tests. Future work
 659 could address these shortcomings and lead to the corroboration of our results, particularly in
 660 describing patterns exhibited by underlying loci in similar systems.

Acknowledgements

We thank Annette Delfino Mix, Camille Jensen, Tom Burt, and Randi Famula for field, common garden, and lab assistance. Additionally, we thank David Fournier, Joey Keely, Kurt Teuber (USDA Forest Service - LTBMU), Roland Shaw (Nevada Division of Forestry), Bill Champion (Nevada State Parks), Woody Loftis (USDA NRCS) for site information and permission to work on Federal and State lands, the VCU CHiPC for computational resources, Jennifer Ciminelli for GIS tips, and Lindsay Miles who helped improve this manuscript. This work was supported by the Southern Nevada Public Lands Management Act – Rounds 7 and 10, sponsored by the USDA, Forest Service, Pacific Southwest Research Station, Albany, CA. We also thank anonymous reviewers whose suggestions greatly contributed to this manuscript.

REFERENCES

- Aitken SN, Yeaman S, Holliday JA, Wang T, Curtis-McLane S (2008) Adaptation, migration, or extirpation: climate change outcomes for tree populations. *Evolutionary Applications* **1**: 95–111.
- Allen CD, Macalady AK, Chenchouni H, Bachelet D, McDowell N, Vennetier M, Kitzberger ., Rigling A, Breshears DD, Hogg EHT, Gonzalez P, Fensham R, Zhang Z, Castro J, Demidova N, Lim J-H, Allard G, Running SW, Semerci A, Cobb N (2010) A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, **259**, 660–684.
- Barrett R, Schluter D (2008) Adaptation from standing genetic variation. *Trends in Ecology & Evolution*, **23**, 38–44.
- Barton NH (1999) Clines in polygenic traits. *Genetical Research*, **74**, 223–236.
- Berg JJ, Coop G (2014) A population genetic signal of polygenic adaptation. *PLoS Genetics*, **10**, e1004412.
- Bourret V, Dionne M, Bernatchez L (2014) Detecting genotypic changes associated with selective mortality at sea in Atlantic salmon: polygenic multilocus analysis surpasses genome scan. *Molecular Ecology*, **23**, 4444–4457.
- Bower AD, Aitken SN (2008) Ecological genetics and seed transfer guidelines for *Pinus albicaulis* (Pinaceae). *American Journal of Botany*, **95**, 66–76.
- Browning BL, Browning SR (2016) Genotype imputation with millions of reference samples. *The American Journal of Human Genetics*, **98**, 116–126.
- Buck JM, Adams RS, Cone J, Conkle MT, Libby WJ, Eden CJ, Knight MJ (2007) California seed zones. San Francisco: California Region, Forest Service, United States Department of Agriculture, 5p.
- Budde KB, Heuertz M, Hernández-Serrano A, Pausas JG, Vendramin GG, Verdú M, González-Martínez SC (2014) In situ genetic association for serotiny, a fire-related trait, in Mediterranean maritime pine (*Pinus pinaster*). *New Phytologist*, **201**, 230–241.
- Ćalić I, Bussotti F, Martínez-García PJ, Neale DB (2015) Recent landscape genomics studies in forest trees. *Tree Genetics & Genomes*, **12**, 1–7.
- Chang T, Hansen AJ, Piekielek N (2014) Patterns and Variability of Projected Bioclimatic Habitat for *Pinus albicaulis* in the Greater Yellowstone Area (B Bond-Lamberty, Ed.). *PLoS ONE*, **9**, e111669.
- Comeault AA, Flaxman SM, Riesch R, Curran E, Soria-Carrasco V, Gompert Z, Farkas TE, Muschick M, Parchman TL, Schwander T, Slate J, Nosil P (2015) Selection on a genetic polymorphism counteracts ecological speciation in a stick insect. *Current Biology*, **25**, 1975–1981.

- 707 Comeault AA, Soria-Carrasco V, Gompert Z, Farkas TE, Buerkle CA, Parchman TL, Nosil P
708 (2014) Genome-wide association mapping of phenotypic traits subject to a range of
709 intensities of natural selection in *Timema cristinae*. *The American Naturalist*, **183**, 711–727.
- 710 Coop G, Witonsky D, Di Rienzo A, Pritchard JK (2010) Using environmental correlations to
711 identify loci underlying local adaptation. *Genetics*, **185**, 1411–1423.
- 712 Critchfield WB, Little EL (1966) Geographic distribution of the pines of the world (No. 991). US
713 Department of Agriculture, Forest Service.
- 714 Csilléry K, Lalagüe H, Vendramin GG, González-Martínez SC, Fady B, Oddou-Muratorio S
715 (2014) Detecting short spatial scale local adaptation and epistatic selection in climate-
716 related candidate genes in European beech (*Fagus sylvatica*) populations. *Molecular*
717 *Ecology*, **23**, 4696–4708.
- 718 Daly C, Neilson RP, Phillips DL (1994) A statistical-topographic model for mapping
719 climatological precipitation over mountainous terrain. *Journal of Applied Meteorology*, **33**,
720 140–158.
- 721 Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
722 Marth GT, Sherry ST, *et al.* (2011) The variant call format and VCFtools. *Bioinformatics*, **27**,
723 2156–2158.
- 724 Davis MB, Shaw RG (2001) Range shifts and adaptive responses to Quaternary climate
725 change. *Science*, **292**, 673–679.
- 726 Eckert AJ, Maloney PE, Vogler DR, Jensen CE, Mix AD, Neale DB (2015) Local adaptation at
727 fine spatial scales: an example from sugar pine (*Pinus lambertiana*, *Pinaceae*). *Tree*
728 *Genetics & Genomes*, **11**, 42.
- 729 Eckert AJ, van Heerwaarden J, Wegrzyn JL, Nelson CD, Ross-Ibarra J, González-Martínez SC,
730 Neale DB (2010) Patterns of population structure and environmental associations to aridity
731 across the range of loblolly pine (*Pinus taeda* L., *Pinaceae*). *Genetics*, **185**, 969–982.
- 732 Eckert AJ, Wegrzyn JL, Pande B, Jermstad KD, Lee JM, Liechty JD, Tearse BR, Krutovsky KV,
733 Neale DB (2009) Multilocus patterns of nucleotide diversity and divergence reveal positive
734 selection at candidate genes related to cold hardiness in coastal douglas fir (*Pseudotsuga*
735 *menziesii* var. *menziesii*). *Genetics*, **183**: 289–298.
- 736 Ehret GB, Lamparter D, Hoggart CJ, Whittaker JC, Beckmann JS, Kutalik Z, Consortium G. I. O.
737 A. T. (2012) A multi-SNP locus-association method reveals a substantial fraction of the
738 missing heritability. *The American Journal of Human Genetics*, **91**, 863–871.
- 739 Ellison AM, Bank MS, Barton DC, Colburn EA, Elliot K, Ford CR, Foster DR, Kloeppel BD,
740 Knoepp JD, Lovett GM, Mohan J (2005) Loss of foundation species: consequences for the
741 structure and dynamics of forested ecosystems. *Frontiers in Ecology and the Environment*,
742 **3**, 479–486.
- 743 Endler JA (1977) Geographic variation, speciation, and clines. No. 10, Princeton University

- 744 Press.
- 745 Farnes PE (1990) SNOTEL and snow course data: describing the hydrology of whitebark pine
746 ecosystems. General Technical Report INT (USA).
- 747 Friedline CJ, Lind BM, Hobson EM, Harwood DE, Mix AD, Maloney PE, Eckert AJ (2015) The
748 genetic architecture of local adaptation I: The genomic landscape of foxtail pine (*Pinus*
749 *balfouriana* Grev. & Balf.) as revealed from a high-density linkage map. *Tree Genetics &*
750 *Genomes*, **11**, 1-15
- 751 Gladman S, Seeman T (2016) Velvet Optimiser v2.5.5, available online
752 <http://bioinformatics.net.au/software/velvetoptimiser.shtml>.
- 753 Gompert Z, Egan SP, Barret RDH, Feder JL, Nosil P (2016) Multilocus approaches for the
754 measurement of selection on correlated genetic loci. *Molecular Ecology*, 1-18.
- 755 Gompert Z, Jahner JP, Scholl CF, Wilson JS, Lucas LK, Soria-Carrasco V, Fordyce JA, Nice
756 CC, Buerkle CA, Forister ML (2015) The evolution of novel host use is unlikely to be
757 constrained by trade-offs or a lack of genetic variation. *Molecular Ecology*, **24**, 2777–2793.
- 758 González-Martínez SC, Huber D, Ersoz E, Davis JM, Neale DB (2008) Association genetics in
759 *Pinus taeda* L. II. Carbon isotope discrimination. *Heredity*, **101**, 19–26.
- 760 Groover A (2015) Genomic science provides new insights into the biology of forest trees. *New*
761 *Phytologist*, **208**, 302–305.
- 762 Guan Y, Stephens M (2011) Bayesian variable selection regression for genome-wide
763 association studies and other large-scale problems. *Annals of Applied Statistics*, **5**, 1780–
764 1815.
- 765 Günther T, Coop G (2013) Robust identification of local adaptation from allele frequencies.
766 *Genetics*, **195**, 205–220.
- 767 Haasl RJ, Payseur BA (2016) Fifteen years of genomewide scans for selection: trends, lessons
768 and unaddressed genetic sources of complication. *Molecular Ecology*, **25**, 5–23.
- 769 Hall D, Luquez V, Garcia VM, St Onge KR, Jansson S, Ingvarsson PK (2007) Adaptive
770 population differentiation in phenology across a latitudinal gradient in European aspen
771 (*Populus tremula*, L.): A comparison of neutral markers, candidate genes, and phenotypic
772 traits. *Evolution*, **61**, 2849–2860.
- 773 Hermisson J (2005) Soft sweeps: molecular population genetics of adaptation from standing
774 genetic variation. *Genetics*, **169**, 2335–2352.
- 775 Holland J (2007) Genetic architecture of complex traits in plants. *Current Opinion in Plant*
776 *Biology*, **10**, 156–161.
- 777 Hornoy B, Pavy N, Gérardi S, Beaulieu J, Bousquet J (2015) Genetic Adaptation to Climate in
778 White Spruce Involves Small to Moderate Allele Frequency Shifts in Functionally Diverse

- 779 Genes. *Genome Biology and Evolution*, **7**, 3269–3285.
- 780 Hutchins HE, Lanner RM (1982) The central role of Clark's nutcracker in the dispersal and
781 establishment of whitebark pine. *Oecologia*, **55**, 797–807.
- 782 Kemper KE, Saxton SJ, Bolormaa S, Hayes BJ, Goddard ME (2014) Selection for complex traits
783 leaves little or no classic signatures of selection. *BMC Genomics*, **15**, 246–260.
- 784 Krakowski J, Aitken SN, El-Kassaby YA (2003) Inbreeding and conservation genetics in
785 whitebark pine. *Conservation Genetics* **4**, 581–593.
- 786 Kremer A, Le Corre V (2012) Decoupling of differentiation between traits and their underlying
787 genes in response to divergent selection. *Heredity*, **108**, 375–385.
- 788 Langlet O (1971) Two hundred years of genecology. *Taxon*, 653–721.
- 789 Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nature Methods*,
790 **9**, 357–359.
- 791 Latta RG (1998) Differentiation of allelic frequencies at quantitative trait loci affecting locally
792 adaptive traits. *The American Naturalist*, **151**, 283–292.
- 793 Latta RG (2003) Gene flow, adaptive population divergence and comparative population
794 structure across loci. *New Phytologist*, **161**, 51–58.
- 795 Le Corre V, Kremer A (2003) Genetic variability at neutral markers, quantitative trait loci and
796 trait in a subdivided population under selection. *Genetics*, **164**, 1205–1219.
- 797 Le Corre V, Kremer A (2012) The genetic differentiation at quantitative trait loci under local
798 adaptation. *Molecular Ecology*, **21**, 1548–1566.
- 799 Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
800 1000 Genome Project Data Processing Subgroup (2009) The sequence alignment/map
801 format and SAMtools. *Bioinformatics*, **25**, 2078–2079.
- 802 Lotterhos KE, Whitlock MC (2015) The relative power of genome scans to detect local
803 adaptation depends on sampling design and statistical method. *Molecular Ecology*, **24**,
804 1031–1046.
- 805 Luquez V, Hall D, Albrechtsen BR, Karlsson J, Ingvarsson P, Jansson S (2007) Natural
806 phenological variation in aspen (*Populus tremula*): the SwAsp collection. *Tree Genetics &*
807 *Genomes*, **4**, 279–292.
- 808 Ma XF, Hall D, Onge KRS, Jansson S, Ingvarsson PK (2010) Genetic differentiation, clinal
809 variation and phenotypic associations with growth cessation across the *Populus tremula*
810 photoperiodic pathway. *Genetics*, **186**, 1033–1044.
- 811 Mackay TFC, Stone EA, Ayroles JF (2009) The genetics of quantitative traits: challenges and
812 prospects. *Nature Reviews Genetics*, **10**, 565–577.

- 813 Mahalovich MF, Stritch L (2013) *Pinus albicaulis*. The IUCN Red List of Threatened Species.
814 e.T39049A2885918. <http://dx.doi.org/10.2305/IUCN.UK.2013-1.RLTS.T39049A2885918.en>.
- 815 Maloney PE, Vogler DR, Eckert AJ, Jensen CE, Delfino-Mix A (in review) Ecological genetics of
816 three white pine species from the Lake Tahoe Basin, USA: Implications for conservation
817 and evolutionary potential. 1–67.
- 818 Maloney PE, Vogler DR, Eckert AJ, Jensen CE, Delfino-Mix A (2012) Ecology of whitebark pine
819 in relation to white pine-blister rust infection in subalpine forests of the Lake Tahoe Basin,
820 USA: Implications for restoration. *Forest Ecology and Management*, **280**, 166-175.
- 821 Mantel N (1967) The detection of disease clustering and a generalized regression approach.
822 *Cancer Research*, **27**, 209–220.
- 823 McKay JK, Latta RG (2002) Adaptive population divergence: markers, QTL and traits. *Trends in*
824 *Ecology & Evolution*, **17**, 285–291.
- 825 McKinney ST, Fiedler CE, Tomback DF (2009) Invasive pathogen threatens bird-pine mutualism:
826 implications for sustaining a high-elevation ecosystem. *Ecological Applications*, **19**, 597-
827 607.
- 828 McLachlan JS, Clark JS, Manos PS (2005) Molecular indicators of tree migration capacity under
829 rapid climate change. *Ecology*, **86**, 2088-2098.
- 830 McLachlan JS, Hellman JJ, Schwartz MW (2007) A framework for debate of assisted migration
831 in an era of climate change. *Conservation Biology*, **21**, 297-302.
- 832 McLane SC, Aitken SN (2012) Whitebark pine (*Pinus albicaulis*) assisted migration potential:
833 testing establishment north of the species range. *Ecological Applications*, **22**, 142–153.
- 834 McVean, G (2007) The structure of linkage disequilibrium around a selective sweep. *Genetics*,
835 **175**, 1395-1406.
- 836 Mitton JB, Grant MC, Yoshino AM (1998) Variation in allozymes and stomatal size in pinyon
837 (*Pinus edulis*, Pinaceae), associated with soil moisture. *American Journal of Botany*, **85**,
838 1262–1265.
- 839 Mitton JB, Stutz HP, Schuster WS (1989) Genotypic differentiation at PGM in Engelmann
840 spruce from wet and dry sites. *Silvae Genetica*, **38**, 217–221.
- 841 Moser G, Lee SH, Hayes BJ, Goddard ME, Wray N. R., Visscher P. M., 2015 Simultaneous
842 Discovery, Estimation and Prediction Analysis of Complex Traits Using a Bayesian Mixture
843 Model. *PLoS Genetics* **11**: e1004969.
- 844 Neale DB, Savolainen O (2004) Association genetics of complex traits in conifers. *Trends in*
845 *Plant Science*, **9**, 325–330.
- 846 Nowak CL, Nowak RS, Tausch RJ, Wigand PE (1994) A 30000 year record of vegetation
847 dynamics at a semi-arid locale in the Great Basin. *Journal of Vegetation Science*, **5**, 579-

- 848 590.
- 849 Ohta T (1982) Linkage disequilibrium with the island model. *Genetics*, **101**, 139–155.
- 850 Parchman TL, Gompert Z, Mudge J, Schilkey FD, Benkman CW, Buerkle CA (2012) Genome-
851 wide association genetics of an adaptive trait in lodgepole pine. *Molecular Ecology*, **21**,
852 2991–3005.
- 853 Pérez F, Granger BE (2007) IPython: A system for interactive scientific computing. *Computing*
854 *in Science and Engineering*, **9**, 21–29.
- 855 Patterson N, Price AL, Reich D (2006) Population structure and eigenanalysis. *PLoS Genetics*,
856 **2**, e190.
- 857 Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE (2012) Double digest RADseq: An
858 inexpensive method for de novo SNP discovery and genotyping in model and non-model
859 species. *PLoS ONE*, **7**, e37135.
- 860 Postma FM, Ågren J (2016) Early life stages contribute strongly to local adaptation in
861 *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences*, **113**, 7590–7595.
- 862 Pritchard JK, Di Rienzo A (2010) Adaptation – not by sweeps alone. *Nature Reviews Genetics*,
863 **11**, 665–667.
- 864 R Core Team (2015) R: A language and environment for statistical computing R Foundation for
865 Statistical Computing, Vienna, Austria URL <http://www.R-project.org/>
- 866 Richardson BA, Brunfeldt SJ, Klopfenstein NB (2002) DNA from bird-dispersed seed and wind-
867 disseminated pollen provides insights into postglacial colonization and population genetic
868 structure of whitebark pine (*Pinus albicaulis*). *Molecular Ecology*, **11**, 215–227.
- 869 Richardson JL, Urban MC, Bolnick DI, Skelly DK (2014) Microgeographic adaptation and the
870 spatial scale of evolution. *Trends in Ecology & Evolution*, **29**, 165–176.
- 871 Savolainen O, Lascoux M, Merilä J (2013) Ecological genomics of local adaptation. *Nature*
872 *Reviews Genetics*, **14**, 807–820.
- 873 Savolainen O, Pyhäjärvi T, Knürr T (2007) Gene Flow and Local Adaptation in Trees. *Annual*
874 *Review of Ecology, Evolution, and Systematics*, **38**, 595–619.
- 875 Savolainen O, Pyhäjärvi T (2009) Genomic Diversity in Forest Trees. *Current Opinion in Plant*
876 *Biology*, **10**, 162–167.
- 877 Schoville SD, Bonin A, Francois O, Lobreaux S, Melodelima C, Manel S (2012) Adaptive
878 Genetic Variation on the Landscape: Methods and Cases. *Annual Review of Ecology,*
879 *Evolution, and Systematics*, **43**, 23–43.
- 880 Sork VL, Aitken SN, Dyer RJ, Eckert AJ, Legendre P, Neale DB (2013) Putting the landscape
881 into the genomics of trees: approaches for understanding local adaptation and population

- 882 responses to changing climate. *Tree Genetics & Genomes*, **9**, 901–911.
- 883 Stephan W (2015) Signatures of positive selection: from selective sweeps at individual loci to
884 subtle allele frequency changes in polygenic adaptation. *Molecular Ecology*, **25**, 79–88.
- 885 Storey, JD, Bass AJ, Dabney A, Robinson D (2015) qvalue: Q-value estimation for false
886 discovery rate control R package version 2.4.2.
- 887 Storz JF (2005) Using genome scans of DNA polymorphism to infer adaptive population
888 divergence. *Molecular Ecology*, **14**, 671–688.
- 889 Storz JF, Kelly JK (2008) Effects of spatially varying selection on nucleotide diversity and
890 linkage disequilibrium: Insights from deer mouse globin genes. *Genetics*, **180**, 367–379.
- 891 Tiffin P, Ross-Ibarra J (2014) Advances and limits of using population genetics to understand
892 local adaptation. *Trends in Ecology & Evolution*, **29**, 673–680.
- 893 Tigano A, Friesen VL (2016) Genomics of local adaptation with gene flow *Molecular Ecology*
894 *AOP*: 1-21.
- 895 Tomback DF (1982) Dispersal of whitebark pine seeds by Clark's nutcracker: a mutualism
896 hypothesis. *The Journal of Animal Ecology*, **51**, 451–467.
- 897 Tomback DF, Achuff P (2010) Blister rust and western forest biodiversity: ecology, values and
898 outlook for white pines. *Forest Pathology*, **40**, 186-225.
- 899 Tomback DF, Arno, SF, Keane RE (2001) Whitebark pine communities: ecology and
900 restoration. Island Press, Washington, DC.
- 901 Tomback DF, Resler LM, Keane RE, Pansing ER, Andrade AJ, Wagner AC (2016) Community
902 structure, biodiversity, and ecosystem services in treeline whitebark pine communities:
903 potential impacts from a non-native pathogen. *Forests*, **7**, 1-8.
- 904 Turchin MC, Chiang CW, Palmer CD, Sankararaman S, Reich D, Hirschhorn JN (2012)
905 Evidence of widespread selection on standing variation in Europe at height-associated
906 SNPs *Nature Genetics* **44**: 1015–1019.
- 907 USDA Forest Service, *Forest Health Technology Enterprise Team* Ft Collins, CO.
- 908 USDA NRCS, 2007 *Soil survey of the Tahoe Basin area, California and Nevada*.
- 909 van Mantgem PJ, Stephenson NL, Byrne JC, Daniels LD, Franklin JF, Fulé PZ, Harmon ME,
910 Larson AJ, Smith JM, Taylor AH, Veblen TT (2009) Widespread increase of tree mortality in
911 the western United States. *Science*, **323**, 521-524.
- 912 Vicente-Serrano SM, Gouveia C, Camarero JJ, Beguería S, Trigo R, López-Moreno JI, Azorín-
913 Molina C, Pasho E, Lorenzo-Lacruz J, Revuelto J, Morán-Tejeda E, Sanchez-Lorenzo A
914 (2013) Response of vegetation to drought time-scales across global land biomes.
915 *Proceedings of the National Academy of Sciences*, **110**, 52–57.

916 Vizcaíno-Palomar N, Revuelta-Eugercios B, Zavala MA, Alia R, González-Martínez SC (2014)
917 The Role of Population Origin and Microenvironment in Seedling Emergence and Early
918 Survival in Mediterranean Maritime Pine (*Pinus pinaster* Aiton). *PLoS ONE*, **9**, e109132.

919 Weir BC, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure.
920 *Evolution*, 1358-1370.

921 Whitlock MC, Lotterhos KE (2015) Reliable detection of loci responsible for local adaptation:
922 Inference of a null model through trimming the distribution of FST. *The American Naturalist*,
923 **186**, S24–S36.

924 Yeaman S (2015) Local adaptation by alleles of small effect. *The American Naturalist*, **186**, S74-
925 S89.

926 Yeaman S, Whitlock MC (2011) The genetic architecture of adaptation under migration-
927 selection balance. *Evolution*, **65**, 1897–1911.

928 Zerbino DR, Birney E (2008) Velvet: Algorithms for de novo short read assembly using de Bruijn
929 graphs. *Genome Research*, **18**, 821–829.

930 Zhou X, Carbonetto P, Stephens M (2013) Polygenic modeling with bayesian sparse linear
931 mixed models. *PLoS Genetics*, **9**, e1003264.

932 **Data Accessibility**

933 Sequence data is deposited in the short read archive of the National Center for Bio-
934 technology Information (project number: TBD). Scripts used in analyses can be found in IPython
935 notebook format (Pérez and Granger 2007) at https://github.com/brandonlind/whitebark_pine.

936 **Author Contributions**

937 PEM, AJE, DBN, DRV, and JLW conceived the study. PEM oversaw sample collection and
938 common garden maintenance. BML wrote the manuscript and carried out analysis of data, with
939 contributions from CJF and AJE. All authors contributed to the editing of the manuscript.

TABLES

Table 1

	Dick's Pass*	Freel Peak*	Heavenly	Little Round Top*	Mt. Rose Ophir*	Rifle Peak*	Snow Valley Peak*	West Shore Peaks
Population size	25 (15)	48 (19)	25 (0)	25 (14)	49 (11)	24 (15)	24 (14)	24 (0)
Ann. precipitation (mm)	1686	1019	782	1221	1186	1281	869	1585
AWC-25cm (kPa)	1.66	1.57	1.12	1.97	1.95	1.89	2.66	1.20
AWC-50cm (kPa)	2.75	2.38	2.00	2.93	2.75	3.11	4.22	2.02
CEC (cmol _c · kg ⁻¹)	0.00	1.45	0.00	12.50	2.90	0.00	0.00	0.00
Clay (%)	6.50	4.50	6.70	14.60	3.00	6.75	6.80	6.00
Elevation (m)	2806	2865	2851	2875	2717	2819	2740	2780
GDD Aug (days)	295	190	276	211	296	235	289	279.5
GDD May (days)	0	0	6	0	11	0	2	1
Max solar rad input (%)	83.59	79.03	78.40	80.09	90.61	93.28	71.70	76.43
Max Temp – July (°C)	21.1	21.6	23.2	21.5	22.9	22.7	23.4	21.8
Min. Temp – Jan (°C)	-6.5	-8.8	-7.5	-8.0	-7.4	-7.4	-7.7	-6.6
Rock coverage (%)	31.00	18.67	25.00	14.67	7.00	30.00	26.67	42.67
Sand (%)	77.67	87.80	83.50	66.20	90.60	74.00	64.50	85.00
Silt (%)	15.8	7.7	9.7	19.1	6.4	19.2	28.7	9.0
WC-15 bar (kPa)	6.6	4.0	3.3	3.6	5.5	8.7	14.0	2.5
WC-1/3 bar (kPa)	9.7	8.0	8.4	7.3	9.8	11.4	14.4	6.2

TABLE 1 Population location and associated attributes. Population size – total (maternal trees with seedlings in common garden). Climatic values were ascertained from data spanning 1971-2000. Ann. precipitation – annual precipitation; AWC – available water capacity at 25cm or 50cm soil depth; CEC – cation exchange capacity; GDD – growing degree days above 5°C; Max solar rad input – maximum solar radiation input; WC-15bar – water capacity at -15bar (wilting point); WC-1/3bar – water capacity at -1/3bar (field capacity). Asterisks indicates populations from which seeds sampled from cones were planted in a common garden. Environmental variables are averaged across subplots.

947 **Table 2**

$pw\hat{D}_{a(ij)}$	Environmental Euclidian Distance	Mantel's r	p -value
†Ann. precipitation	Ann. precipitation	0.7135	0.0027
†Longitude	Longitude	0.6522	0.0024
†Longitude	Ann. precipitation	0.7716	0.0016
†Rock coverage	Ann. precipitation	0.5542	0.0221
†Tmin-Jan	Ann. precipitation	0.5765	0.0132
†Longitude	Latitude	-0.4257	0.0347
†Rock coverage	Longitude	0.5566	0.0284
†Tmin-Jan	Longitude	0.4822	0.0273
†WC-15bar	Tmax-July	0.3490	0.0309
†WC-1/8bar	WC-1/8bar	0.4806	0.0361
†WC-1/8bar	Tmax-July	0.4539	0.0037
†WC-1/8bar	AWS0-25	0.4329	0.0384
†WC-1/8bar	AWS0-50	0.4538	0.0464
†WC-1/8bar	WC-15bar	0.5126	0.0335
GDD-May	GDD-May	0.8480	0.0013
Rock coverage	Rock coverage	0.5124	0.0145
Sand	Sand	0.5574	0.0046
Tmin-Jan	Tmin-Jan	0.5791	0.0137
Sand	Clay	0.5345	0.0232
Silt	Sand	0.4408	0.0238

948 **TABLE 2** Signatures of allele frequency shifts associated with environmental distance. *Significant Mantel tests (9999 permutations) from*
949 *comparisons among $pw\hat{D}_{a(ij)}$ matrices from SNPs associated with environment (first column) against environmental Euclidian distance (second*
950 *column). Environmental variables as in Table 1. † indicates a comparison in which at least one variable is water-related, or was associated with*
951 *annual precipitation.*

Table 3

Selection criterion	$pw\hat{D}_{a(ij)}$	Environmental Euclidian Distance	Mantel's r	p -value
99.9 th $\overline{PTP}$	Bud flush	GDD August	0.5804	0.0181
	Bud flush	GDD May	-0.5190	0.0458
	Bud flush	Max. radiation input	-0.5486	0.0482
	Bud flush	$T_{\min}$ January	0.6984	0.0191
99.8 th $\overline{PTP}$	[†] Bud flush	Annual precipitation	0.4309	0.0140
	[†] Bud flush	Longitude	0.5532	0.0405
	Bud flush	Max. radiation input	-0.6312	0.0127
	N(μ g)	Elevation	0.5334	0.0246
	[†] Height	Annual precipitation	0.7210	0.0320
	[†] $\delta^{13}\text{C}$	Annual precipitation	0.5952	0.0195

TABLE 3 Signatures of allele frequency shifts associated with environmental distance. *Significant Mantel tests (9999 permutations) from comparisons among allele frequency shifts ($pw\hat{D}_{a(ij)}$) of SNPs associated with phenotype against environmental Euclidian distance. Selection Criterion refers to the process used to identify SNPs associated with phenotype. † indicates a comparison in which at least one variable is water-related, or was associated to annual precipitation.*

957 FIGURES

958

959 **Figure 1** Populations used for sampling *P. albicaulis* within the Lake Tahoe Basin (dark outline). Annual
960 precipitation is given for each population to demonstrate the west-east rain shadow experienced across
961 fine spatial scales. Asterisks indicate populations in the common garden study.

Figure 2

Figure 2. Significantly higher allele frequency covariance ($\hat{D}_{a(ij)}$) among loci associated to environment by *bayenv2* than could be expected to arise, given the data. In white are the median values from $\hat{D}_{a(ij)}$ calculated among focal SNPs associated to environment. Black bars display the 95th percentile of the null distribution of median $\hat{D}_{a(ij)}$. Environmental variables are grouped by those related to soil (CEC through WC-1/3bar) and those related to either climate or geography (Annual precipitation through T-min Jan), with variables related to water availability grouped together in the center of the figure (AWC-25cm through Annual precipitation). All sets of focal loci had median $\hat{D}_{a(ij)}$ greater than the 100th percentile of the null distribution, as indicated by two stars (**). Environmental variables as in Table 1.

Figure 3

A

B

Figure 3 Violin plots for the kernel density estimator of the posterior distributions (light grey) taken from Bayesian sparse linear mixed models (BSLMM) executed in GEMMA for (A) the proportion of variance explained by SNPs included in the model (PVE) and (B) the number of SNPs underlying the phenotypic trait (N_{SNP}). Priors for N_{SNP} and PVE were [1,300] and [0.01,0.9], respectively. Dark grey vertical bars display the first through third interquartile range with the median represented by the white dot.

979 **Figure 4**

Figure 4. Allele frequency covariance ($\hat{D}_{a(ij)}$) among loci associated to phenotype by GEMMA. In white are the median values from $\hat{D}_{a(ij)}$ calculated among focal SNPs associated to phenotype. Black bars display the 95th percentile of the null distribution of median $\hat{D}_{a(ij)}$. (A) SNPs identified in the top 99.9th percentile of $\bar{P}TP$, (B) SNPs identified in the top 99.8th percentile of $\bar{P}TP$. One star (*) indicates that the median focal $\hat{D}_{a(ij)}$ was greater than the 95th percentile of the null distribution, whereas two stars (**) indicate that the focal median $\hat{D}_{a(ij)}$ was greater than the 100th percentile of the null distribution.