Bounds to Parapatric Speciation: A Dobzhansky-Muller incompatibility model involving autosomes, X chromosomes and mitochondria

Ilse Höllinger^{1,2} and Joachim Hermisson¹

¹Mathematics and BioSciences Group, Faculty of Mathematics and
Max F. Perutz Laboratories, University of Vienna, 1090 Vienna,
Austria

²Vienna Graduate School of Population Genetics, Vienna Austria

February 3, 2017

Postal adress:

Joachim Hermisson and Ilse Höllinger
Faculty of Mathematics, University of Vienna
Oskar-Morgenstern-Platz 1
1090 Wien, Austria
Europe

Telephone number:

+43/1/4277 50648 (J. Hermisson)
+43/1/4277 50771 (I. Höllinger)

Email-adress:

joachim.hermisson@univie.ac.at
ilse.hoellinger@univie.ac.at

Key words:

hybrid incompatibility, two-locus DMI, speciation-with-gene-flow, large X-effect,
introgression on X and autosomes

Word count

excluding tables, figures and their captions, and literature cited:

Title: 15 words

—

Abstract: 172 words

Introduction: 650 words

Model and Methods: 1389 words

Results: 2466 words

Discussion: 2713 words

—

Sum: 7390 words

—

Abstract

We investigate the conditions for the origin and maintenance of postzygotic isolation barriers, so called (Bateson-)Dobzhansky-Muller incompatibilities or DMIs, among populations that are connected by gene flow. Specifically, we compare the relative stability of pairwise DMIs among autosomes, X chromosomes, and mitochondrial genes. In an analytical approach based on a continent-island framework, we determine how the maximum permissible migration rates depend on the genomic architecture of the DMI, on sex bias in migration rates, and on sex-dependence of allelic and epistatic effects, such as dosage compensation. Our results show that X-linkage of DMIs can enlarge the migration bounds relative to autosomal DMIs or autosome-mitochondrial DMIs, in particular in the presence of dosage compensation. The effect is further strengthened with male-biased migration. This mechanism might contribute to a higher density of DMIs on the X chromosome (large X-effect) that has been observed in several species clades. Furthermore, our results agree with empirical findings of higher introgression rates of autosomal compared to X-linked loci.

1 Introduction

Historically, speciation research has mostly focused on two idealized scenarios: allopatric speciation (complete geographic isolation of incipient species) and sympatric speciation (divergence of subpopulations in a common habitat) (Orr and Turelli, 2001; Coyne and Orr, 2004; Via and West, 2008). Both scenarios are simplifications of biological reality. While strict sympatry of incipient species seems to be an exception, there is abundant evidence for hybridization even among “good species” with viable and not completely sterile hybrid offspring (reviewed *e.g.* in Coyne and Orr, 2004; Mallet, 2008). Population genetic theory shows that even low levels of gene flow can strongly interfere with population differentiation (Felsenstein, 1981; Slatkin, 1987). This makes it inevitable to assess the impact of limited gene flow at various stages of the speciation process, a scenario commonly referred to as parapatric speciation.

The classical model for the evolution of postzygotic isolation barriers in allopatry is the (Bateson-)Dobzhansky-Muller model (DMM) (Bateson, 1909; Dobzhansky, 1936; Muller, 1942). The DMM assumes that new substitutions occur on different genetic backgrounds. When brought into secondary contact, these previously untested alleles might be mutually incompatible and form Dobzhansky-Muller incompatibilities (DMIs), thus reducing hybrid fitness and decreasing gene flow at linked sites. The emergence of species boundaries due to accumulation of DMIs in allopatry is well understood (Coyne and Orr, 1989; Orr and Turelli, 2001; Coyne and Orr, 2004). More recently, several studies have considered this process in parapatry (Gavrilets, 1997; Feder and Nosil, 2009; Agrawal et al., 2011; Bank et al., 2012; Wang, 2013; Lindtke and Buerkle, 2015). All support that the DMM provides a viable mechanism for the evolution of postzygotic isolation even in the presence of gene flow, although the bounds for maximum permissible migration rates can be quite stringent.

Empirically, there is widespread evidence for DMIs not only among recently diverged sister species (Maheshwari and Barbash, 2011; Presgraves, 2010; Sweigart

and Flagel, 2014), but also segregating within species (Corbett-Detig et al., 2013). Hence, these authors argue that the genetic basis of reproductive isolation is continuously present within natural populations, rendering the independent allopatric evolution of newly incompatible substitutions obsolete.

While most theoretical studies focus on autosomal DMIs, empirical evidence points to a major role of sex chromosomes in speciation. Haldane’s rule (Haldane, 1922, reviewed in Coyne and Orr, 2004), states that in species with sex specific reduced hybrid fitness the affected sex is generally heterogametic. The *large X-effect* (Coyne and Orr, 1989, reviewed in Presgraves, 2008) expresses the disproportional density of X-linked incompatibility genes in postzygotic isolation. For example Masly and Presgraves (2007) report a higher density of incompatibilities causing hybrid male sterility on the X chromosome relative to autosomes in *Drosophila*. Equivalent findings exist of a *large Z-effect* in WZ-systems, such as birds, where WZ-females are heterogametic (Ellegren, 2009). Also cytoplasmic incompatibilities have been described (Ellison and Burton, 2008; Lee et al., 2008; Burton and Barreto, 2012; Barnard-Kubow et al., 2016).

A recent study by Bank et al. (2012) determined stability conditions and maximum permissible migration rates of autosomal two-locus DMIs in a continent-island framework. They distinguished two main mechanisms shaping the evolution of DMIs: selection against (maladapted) immigrants and selection against (unfit) hybrids, which lead to different dependence of maximum migration rates on the model parameters.

Prompted by the empirical observations described above, we extend the model by Bank et al. (2012) to general two-locus DMIs in diploids involving X chromosomes, autosomes, or mitochondria. We include sex-specific fitness effects, in particular, to account for the effect of dosage compensation of hemizygous X-linked genes in males. We also allow for sex-specific migration, as many species display differences in migration patterns for males and females Greenwood (1980).

Following Bank et al. (2012) we derive maximum migration bounds where DMIs

can still originate in parapatry, or resist continental gene flow. In contrast to the autosomal case, we find that sex specific fitness- and sex-biased migration cause substantial differences in the maximum permissible rates and hence influence the prevalence of autosomal DMIs relative to X-linked and mitochondrial DMIs. Especially, we find that X-linkage of (nuclear or cytonuclear) DMIs together with dosage compensation and/or male-biased migration boosts migration bounds and thus enhances the evolution of X-linked DMIs, possibly contributing to a *large X-effect* and to reduced introgression probabilities of X-linked DMI loci.

2 Model and Methods

We consider a diploid, dioecious population with separate sexes (at 1:1 ratio) that is divided into two panmictic subpopulations, continent and island. (See Figure 1 and Figure B.1 in the Supporting Information (SI)). Both demes are sufficiently large that drift can be ignored (drift effects are discussed in SI Section A.3). They are connected by unidirectional sex-dependent migration at rate $m^{\varnothing}$ and m^{σ} from the continent to the island. We fix the average migration rate per individual, $m = \frac{m^{\varnothing} + m^{\sigma}}{2}$, and define

$$R := \frac{m^{\varnothing} - m^{\sigma}}{m^{\varnothing} + m^{\sigma}} \in [-1, 1] \quad (1)$$

as a measure of sex-bias in migration. Sex-specific migration gives rise to distinct migration rates per allele for autosomes, X chromosomes, and mitochondria, $m_{\mathcal{A}}$, $m_{\mathcal{X}}$, and $m_{\mathcal{O}}$ (Eqs. (B.13)-(B.15)). For $-1 \leq R < 0$ migration is male-biased and we obtain $m_{\mathcal{A}} > m_{\mathcal{X}} > m_{\mathcal{O}}$. In contrast, for $0 < R \leq 1$ migration is female-biased, resulting in $m_{\mathcal{A}} < m_{\mathcal{X}} < m_{\mathcal{O}}$.

Figure 1: **Schematic model.** The population inhabits a continent (left) and an island (right), which are connected by unidirectional migration at rate m . The figure shows two out of eight genomic architectures investigated: an X-autosome DMI (upper line) and a cytoplasmic DMI between X and mitochondrion (lower line). Genotypes of female residents are depicted by yellow circles and males by blue circles, respectively. The capital letters denote incompatible alleles, which reduce hybrid fitness.

The DMI

The incompatibility is formed by two unlinked biallelic loci, situated on autosomes $\mathcal{A}$, X chromosomes $\mathcal{X}$, or in the mitochondrial genome (cytoplasmic organelle) $\mathcal{O}$, (cf. Table 1). Both sexes are diploid for autosomes and haploid for the mitochondrial locus. Males are hemizygous for the X chromosomes, whereas females are diploid. The continent is monomorphic for the continental (geno-)type and only acts as source of migrants for the island. Our analysis focuses on the evolutionary dynamics on the island. A stable DMI corresponds to a stable equilibrium on the island where all four alleles are maintained (a two-locus polymorphism), including the pair of incompatible alleles (indicated by capital letters in Table 1).

We model genotypic fitness as the sum of direct allelic fitness and epistasis. Hence any given allele contributes directly to genotype fitness, where it can be locally or globally adapted, and additionally via epistasis if it is incompatible with other alleles in the same genotype. We set the (Malthusian) fitness of genotypes containing no incompatible alleles (only lower case letters) in both sexes to 0. For simplicity, we assume no dominance of the single-locus effects, but we allow for dominance or

Different genomic architectures of DMIs.

Model	DMI	continental type (φ, σ)	island type (φ, σ)
$A \rightarrow A$	$\mathcal{A}-\mathcal{A}$	aaBB	AAbb
$X \rightarrow A$	$\mathcal{A}-\mathcal{X}$	aaXX, aaXy	AAxx, AAxy
$A \rightarrow X$	$\mathcal{A}-\mathcal{X}$	AAxx, AAxy	aaXX, aaXy
$X \rightarrow X$	$\mathcal{X}-\mathcal{X}$	$X_1X_1x_2x_2, X_1x_2y$	$x_1x_1X_2X_2, x_1X_2y$
$A \rightarrow O$	$\mathcal{A}-\mathcal{O}$	AAo	aaO
$O \rightarrow A$	$\mathcal{A}-\mathcal{O}$	aaO	AAo
$X \rightarrow O$	$\mathcal{X}-\mathcal{O}$	XXo, Xyo	xxO, xyO
$O \rightarrow X$	$\mathcal{X}-\mathcal{O}$	xxO, xyO	XXo, Xyo

Table 1: Each genomic architecture is defined by a continental (geno-)type (third column) and an island (geno-)type (fourth column). Mutually incompatible pairs of DMI-alleles are denoted by capital letters. We call the immigrating DMI-allele *continental allele* and its resident incompatible partner *island allele*. The name of each model in the first column is constituted by “the continental allele $\rightarrow$ the island allele”. The $A \rightarrow A$ -model corresponds to the model by Bank et al. (2012).

109 recessitivity of the incompatibility.

110 We define the fitness of an arbitrary female genotype as

$$\omega(G^{\varphi}) = \underbrace{n_C \cdot \sigma_C^{\varphi} + n_I \cdot \sigma_I^{\varphi}}_{\text{allelic fitness}} - \underbrace{\Gamma_*(G^{\varphi})}_{\text{epistasis}} \quad (2)$$

111 or for a male genotype as

$$\omega(G^{\sigma}) = n_C \cdot \sigma_C^{\sigma} + n_I \cdot \sigma_I^{\sigma} - \Gamma_*(G^{\sigma}). \quad (3)$$

112 The allelic fitness is captured by the selection coefficient $\sigma^{\varphi}, \sigma^{\sigma}$ (for females and
 113 males) and weighted with the respective number of incompatible alleles, $n_{C,I} \in$
 114 $\{0, 1, 2\}$ in a given genotype. To match the locus effects of haploid mitochondrial
 115 genes to autosomes, we set $n_{C,I} \in \{0, 2\}$ for the absence or presence of the single
 116 incompatible allele in this case.

We assume $\sigma^{\sigma} = \sigma^{\varphi}$ for autosomes and organelles, but for X-linked alleles the fitness effect may be enhanced in males, $\sigma^{\sigma} = (1 + D)\sigma^{\varphi}$, where $D \in \{0, 1\}$ measures dosage compensation (see below). The contribution of epistasis to hybrid genotype fitness can be summarized by an epistasis vector Γ_* , for each model (*), detailed in Table 2

Strength of the incompatibility: The epistasis vector

DMI	hybrids: φ, σ	epistasis vector Γ_*
$\mathcal{A}\text{-}\mathcal{A}$	φ : AaBb, AaBB, AABb, AABB σ : AaBb, AaBB, AABb, AABB	$\Gamma_{\mathcal{A}\mathcal{A}} = (\underbrace{\gamma_1, \gamma, \gamma, 2\gamma}_{\varphi}, \underbrace{\gamma_1, \gamma, \gamma, 2\gamma}_{\sigma})$
$\mathcal{A}\text{-}\mathcal{X}$	φ : AaXx, AaXX, AAXx, AAXX, σ : AaXy, AAXy	$\Gamma_{\mathcal{A}\mathcal{X}} = (\gamma_1, \gamma, \gamma, 2\gamma, (1 + D)\frac{\gamma}{2}, (1 + D)\gamma)$
$\mathcal{X}\text{-}\mathcal{X}$	φ : X ₁ x ₁ X ₂ x ₂ , X ₁ x ₁ X ₂ X ₂ X ₁ X ₁ X ₂ x ₂ , X ₁ X ₁ X ₂ X ₂ , σ : X ₁ X ₂ y	$\Gamma_{\mathcal{X}\mathcal{X}} = (\gamma_1, \gamma, \gamma, 2\gamma, (1 + 3D)\frac{\gamma}{2})$
$\mathcal{A}\text{-}\mathcal{O}$	φ : AaO, AAO, σ : AaO, AAO	$\Gamma_{\mathcal{A}\mathcal{O}} = (\gamma, 2\gamma, \gamma, 2\gamma)$
$\mathcal{X}\text{-}\mathcal{O}$	φ : XxO, XXO, σ : XyO	$\Gamma_{\mathcal{X}\mathcal{O}} = (\gamma, 2\gamma, (1 + D)\gamma)$

Table 2: The table shows all possible hybrid genotypes with DMIs (second column) and corresponding fitness cost, parametrized by the entries of the epistasis vector (third column). The strength of the incompatibility depends on the number of incompatible alleles in the genotype. Plausibly, the strength increases with the number of incompatible pairs, which can be 1, 2, or 4 (Turelli and Orr, 2000). We focus on two particular epistasis schemes, one with a codominant DMI ($\gamma_1 = \frac{\gamma}{2}$) with fitness cost proportional to the number of incompatible pairs and one with a recessive DMI ($\gamma_1 = 0$) where the fitness cost is zero if there is still a pair of compatible alleles in the genotype. The strength of X-linked incompatibilities in males depends on dosage compensation, captured by $D \in \{0, 1\}$.

121

122 Dosage compensation

Dosage compensation can be related to different mechanisms. For example, in the model organism *Drosophila melanogaster* the expression of the X chromosome is doubled in males. An alternative mechanism has evolved in mammals, where one X chromosome is randomly inactivated in females (Payer and Lee, 2008). Finally, in birds dosage compensation seems to be incomplete, as some genes show elevated expression levels in homogametic ZZ-males compared to heterogametic females,

129 whereas other genes are dosage compensated (Ellegren et al., 2007; Graves et al.,
130 2007).

131 Our model allows for arbitrary sex-dependence of allelic and epistatic effects, but
132 we focus on dosage compensation of the hemizygous X chromosome in males as a
133 key biological mechanism. We model fitness for any X-linked allele in hemizygous
134 males in two ways (Charlesworth et al., 1987):

- 135 • *No dosage compensation*, $D = 0$: A single copy of an X-linked allele has the
136 same allelic ($\sigma^{\sigma} = \sigma^{\varphi}$) and epistatic effects in hemizygous males as in females.
- 137 • *Full dosage compensation*, $D = 1$: Hemizygosity of the X chromosome is
138 compensated in males, and a single X-linked allele has the same effect as a
139 homozygous pair of X chromosomes in females (allelic selection coefficient:
140 $\sigma^{\sigma} = 2\sigma^{\varphi}$).

141 With random deactivation of X in females we naturally obtain a codominant
142 DMI in our model since (average) heterozygous fitness is equal to the mean of the
143 homozygous fitnesses in this case.

144 Dynamics of the general model

145 For our analytical treatment, we assume weak evolutionary forces, such that linkage
146 equilibrium among both loci and Hardy-Weinberg-proportions can be assumed. It
147 is then sufficient to track the frequencies of the continental allele p_C and the island
148 allele p_I on the island. We test this approximation for stronger selection by numerical
149 simulations in SI Section A.2.

150 For each genomic architecture (Table 1) we derive a pair of differential equations
151 in continuous time (see **Box 1**). For the case of an X→A DMI, in particular,
152 p_C measures the frequency of the incompatible X allele that immigrates from the
153 continent and p_I the frequency of the incompatible autosomal allele on the island.

154 We obtain:

$$\begin{aligned}\dot{p}_C &= p_C(1 - p_C) \left(\frac{3+D}{3} \sigma_C^{\varnothing} + \frac{2}{3} p_I \left((2p_C(1 - p_I) + p_I)(2\gamma_1 - \gamma) - 2\gamma_1 - \frac{\gamma}{2}(1 + D) \right) \right) \\ &\quad + (1 - p_C) \left(1 + \frac{R}{3} \right) m \\ \dot{p}_I &= p_I(1 - p_I) \left(\sigma_I^{\varnothing} + \frac{1}{2} p_C \left((2p_I(1 - p_C) + p_C)(2\gamma_1 - \gamma) - 2\gamma_1 - \frac{\gamma}{2}(1 + D) \right) \right) \\ &\quad - p_I m\end{aligned}\tag{4}$$

155 We see that with dosage compensation ($D = 1$), the X-linked allelic fitness is
156 increased ($\frac{4}{3}\sigma_C^{\varnothing}$), because a single X-allele in males now acts as strongly as two
157 X-alleles in females. Similarly, dosage compensation increases the term due to
158 epistasis in males ($\frac{\gamma}{2}(1 + D)$) affecting both the X-linked allele and the autosomal
159 allele. Sex-biased migration, quantified by R (see Eq. 1), affects only the X-linked
160 allele, as males are hemizygous X carriers. Parameterizations for all other cases are
161 provided in the SI Section B.1.2.

Box 1:

Dynamics of the continental allele frequencies p_C :

$$\dot{p}_C = \begin{cases} \text{for } \mathcal{A} : & p_C \left(\frac{\omega_C^{*\varnothing} + \omega_C^{*\sigma}}{2} - \frac{\bar{\omega}^{\varnothing} + \bar{\omega}^{\sigma}}{2} \right) & +(1 - p_C)m \\ \text{for } \mathcal{X} : & p_C \left(\frac{2\omega_C^{*\varnothing} + \omega_C^{*\sigma}}{3} - \frac{2\bar{\omega}^{\varnothing} + \bar{\omega}^{\sigma}}{3} \right) & +(1 - p_C)(1 + \frac{R}{3}) \cdot m \\ \text{for } \mathcal{O} : & p_C (\omega_C^{*\varnothing} - \bar{\omega}^{\varnothing}) & +(1 - p_C)(1 + R) \cdot m \end{cases}$$

Dynamics of the island allele p_I :

$$\dot{p}_I = \begin{cases} \text{for } \mathcal{A} : & p_I \left(\frac{\omega_I^{*\varnothing} + \omega_I^{*\sigma}}{2} - \frac{\bar{\omega}^{\varnothing} + \bar{\omega}^{\sigma}}{2} \right) & -p_I m \\ \text{for } \mathcal{X} : & p_I \left(\frac{2\omega_I^{*\varnothing} + \omega_I^{*\sigma}}{3} - \frac{2\bar{\omega}^{\varnothing} + \bar{\omega}^{\sigma}}{3} \right) & -p_I(1 + \frac{R}{3}) \cdot m \\ \text{for } \mathcal{O} : & p_I (\omega_I^{*\varnothing} - \bar{\omega}^{\varnothing}) & -p_I(1 + R) \cdot m \end{cases}$$

Marginal fitness $\omega_{C/I}^{*\varnothing/\sigma}$ and mean fitness $\bar{\omega}^{\varnothing/\sigma}$ for each sex are functions of genotype fitness (consult SI Eqs.(B.9),(B.10) for explicit expressions). Sex-bias in migration m is measured by R (Eq.(1)).

162

163 The codominant model

164 If the effect of the incompatibility is additive, such that it is proportional to the
165 number of incompatible pairs in a genotype ($\gamma_1 = \frac{\gamma}{2}$ in Table 2), the model simplifies
166 greatly. For the X→A model, in particular,

$$\begin{aligned}\dot{p}_C &= (1 - p_C)\left(\frac{3+D}{3}p_C(\sigma_C^Q - \gamma p_I) + \left(1 + \frac{R}{3}\right)m\right) \\ \dot{p}_I &= p_I\left((1 - p_I)\left(\sigma_I^Q - \frac{(3+D)}{4}\gamma p_C\right) - m\right)\end{aligned}\tag{5}$$

see SI Eqs. (B.34) for the other models.

Evolutionary histories

A parapatric DMI can evolve via different routes, depending on the timing and geographic location of the origin of the two mutations. Following Bank et al. (2012), we distinguish five histories: For *secondary contact*, both substitutions occur during an allopatric phase and can originate in any order. In contrast, if the substitutions originate in the presence of gene flow, the timing matters and we obtain four further scenarios: for a *continent-island* DMI we have the first substitution originating on the continent and the second on the island. Analogously, there are *island-continent*, *island-island*, and *continent-continent* scenarios. Note that the first two scenarios lead to *derived-derived DMIs*, with one substitution in each deme, whereas the last two lead to *derived-ancestral DMIs*, where both substitutions occur in the same deme. In all cases we refer to the immigrating incompatible allele as the *continental allele* and to the resident, incompatible allele as the *island allele*. All five evolutionary histories lead to the same dynamics (as given in **Box 1**) upon appropriate relabeling of genotypes, where different histories correspond to different initial conditions (see SI Section B.2.5 and "Mapping of evolutionary histories" below).

3 Results

Our analytical analysis of the dynamical system in **Box 1** is presented in comprehensive form in SI B. It comprises on the following steps. For the general model ($0 \leq \gamma_1 < \gamma$),

we determine all boundary equilibria and conditions for their stability. Instability of all boundary equilibria implies a protected polymorphism at both loci. Excluding cycling behavior, this is a sufficient condition for a globally stable DMI that will be reached from all starting conditions (all evolutionary histories). An internal stable equilibrium (DMI) can also coexist with a stable boundary equilibrium. In this case, the DMI is only locally stable and will only be reached from favorable starting conditions. Necessary and sufficient conditions for the existence of (locally or globally) stable DMIs can be derived for weak migration by means of perturbation analysis: A stable DMI results if the monomorphic boundary equilibrium ($p_I = 1$, $p_C = 0$) is stable for $m = 0$ and is dragged inside the state space for small $m > 0$. For codominant DMIs, also the internal equilibria can be assessed analytically and conditions for stable DMIs follow from a bifurcation analysis. For the recessive DMIs, we complement our analytical results by numerical work to derive stability conditions for locally stable DMIs under stronger migration.

Below, we summarize the key results for the general model. This is followed by a detailed analysis of the codominant model. In the supplement we added continuative results, first for the recessive model in SI Section A.1. Second, SI Section A.2 contains simulation results to assess the effects of linkage disequilibrium (LD), which is relevant for very strong incompatibilities. Third, we present simulations for finite populations and analyze how migration limits are affected by genetic drift in SI Section A.3. Finally, in SI Section A.4 we calculate adaptive substitution rates for autosomes and X chromosomes with gene flow and derive conditions on dominance favoring the *faster X-effect*, described by Charlesworth et al. (1987).

Stable equilibria: global and local stability of DMIs

The model has three boundary equilibria: A monomorphic state, where the continental genotype swamps the island, which is always reached for strong migration. Furthermore, two single locus polymorphisms (SLPs) where one locus is swamped, but the other is

maintained polymorphic. There is at most one stable internal equilibrium, corresponding to a DMI. It can either be globally or locally stable. In the latter case, one of the boundary equilibria is also locally stable and it depends on the evolutionary history which equilibrium is reached. We therefore obtain two migration thresholds $0 \leq m_{\max}^- \leq m_{\max}^+$:

- For migration rates $0 \leq m < m_{\max}^-$, a globally stable DMI, that is reached for all evolutionary histories.
- For migration rates $0 \leq m_{\max}^- \leq m < m_{\max}^+$, the dynamics are bistable and yield a locally stable DMI. Hence, only certain evolutionary histories permit its evolution, but any existing DMI will be maintained.
- For migration rate $m_{\max}^+ \leq m$ no stable DMI exists.

Mapping of evolutionary histories

Every evolutionary history maps to a distinct initial condition (SI Section B.2.5 for results and proofs). As in Bank et al. (2012), we find three permissive histories that always result in the evolution of a stable DMI for $m < m_{\max}^+$: *secondary contact*, *island-continent*, and *continent-continent*. In all these cases, the second substitution occurs in a deme where the incompatible first substitution is not (yet) present. In contrast, the evolution of a stable DMI in parapatry is more difficult for an *island-island* or *continent-island* substitution history. Here, the second substitution needs to invade on the island despite of competition of the incompatible allele. We need $m < m_{\max}^-$ for a DMI to originate under these circumstances.

Necessary conditions for the existence of a stable DMI

Based on previous results for the model without migration (Rutschman, 1994) or without epistasis (Bürger and Akerman, 2011), and in accordance to Bank et al. (2012), we find that with epistasis and increasing migration a stable DMI can only

exist if the island allele is beneficial and its sex-averaged selection coefficient exceeds migration. Furthermore, any averaged selective advantage of the continental allele must be outweighed by averaged epistasis. For example for $X \rightarrow A$, we obtain $m < \sigma_I^{\frac{\sigma}{\tau}}$ and $\frac{\sigma_C^+}{3} < \frac{\gamma}{2}$. Consult Eqs. B.29, B.31 and Table B.3 for different terms for each model and the SI Sections B.1.3, B.1.6 for proofs.

3.1 Nuclear codominant DMIs

We obtain full analytical solutions for the maximum migration bounds $m_{\max}^{\pm}$ (B.2.4). Below, we discuss how these rates depend on the various genetic architectures, sex dependence of fitness and migration, and on dosage compensation. Figures 2 and 3 compare the $m_{\max}^{\pm}$ for the different types incompatibilities among nuclear genes: DMIs among autosomal genes ($A \rightarrow A$), DMIs among X and autosomes, with either the incompatible X allele immigrating from the continent ($X \rightarrow A$) or the autosomal locus ($A \rightarrow X$), and DMIs among two X-linked loci ($X \rightarrow X$). Figure 2 assumes full dosage compensation of X-linked alleles in males, Figure 3 treats the case without dosage compensation.

Selection against hybrids and against immigrants

Following Bank et al. (2012), we can distinguish two main selective forces maintaining a DMI in the face of gene flow. If the continental allele is beneficial on the island (first column of Fig. 2 and 3), a polymorphism at the respective locus can only be maintained by hybrid formation and selection against the immigrating allele is due to hybrid inferiority (“selection against hybrids”). This type of selection will only be effective as long as the immigrating allele is rare. Once the migration pressure is so high that the immigrating continental allele is in a majority, incompatibility selection rather works against the resident allele on the island. Consequently, we expect a large bistable regime with $m_{\max}^+ \gg m_{\max}^-$ and a small region with global stability, as can indeed be seen for all types of DMIs with a beneficial continental allele. Note

also that $m_{\max}^+$ increases with γ , as should be expected if hybrid incompatibility, *i.e.* epistasis, is the sole cause of (local) stability.

In contrast, with a deleterious immigrating allele (third column of Fig. 2 and 3), a DMI can also be maintained by “selection against immigrants” for small values of epistasis, or via a combination of the two selective forces (selection against hybrids and immigrants) with stronger epistasis. If selection against immigrants predominates, maintenance of the DMI is driven by local adaptation. The fitness advantage of the resident allele depends on its direct effect and the dynamics will usually be frequency independent. Therefore we obtain no or only a small bistable regime, with $m_{\max}^+ \approx m_{\max}^-$. For stronger epistasis, selection against hybrids becomes more important, leading to a relative increase of the bistable regime. The main effect of epistasis now is that it promotes swamping of the island allele: $m_{\max}^{+/-}$ decreases with epistasis. In the case of a neutral immigrating allele, the observed migration bounds exhibit an intermediate pattern.

Sex-biased migration

To understand the differences among the DMI architectures, we take the case of full dosage compensation and strict female-biased migration ($R = 1$) as a starting point (Fig. 2(a)-(c)). In this case, all curves for $m_{\max}^{\pm}$ for the different models collapse onto a single one. Indeed, if only females migrate, the number of migrating X chromosomes and autosomes is equal. Full dosage compensation balances any direct and epistatic effects of loci with different ploidy levels. Consequently, the corresponding Eqs. (B.34) differ only by a constant factor.

If also males migrate (Fig. 2(d)-(i)) genomic architectures involving an X chromosomes experience effectively lower migration rates of the X and hence increasing $m_{\max}^{\pm}$. Male-biased migration boosts $m_{\max}^{\pm}$ most effectively for X→X, as both loci experience reduced migration pressure. For unbiased migration, $m_{\max}^+$ of X→X relative to A→A DMIs increases by $\frac{4}{3}$ (the autosome-X ratio), and doubles for pure male migration

Figure 2: **Codominant nuclear DMIs with dosage compensation, $D = 1$.** The columns show $m_{\max}^{\pm}$ as a function of the strength of epistasis γ for beneficial ($\sigma_C^O = 0.5$), neutral ($\sigma_C^O = 0$), and deleterious ($\sigma_C^O = -0.5$) effect of the immigrating allele. All quantities in the figure ($\sigma_C^O, \gamma, m_{\max}^{\pm}$) are measured relative to the fitness effect of the island allele, which is normalized to $\sigma_I^O = 1$. Note the different scaling of the y-axis in the third column. Strong differences between $m_{\max}^{\pm}$ in the various models occur if migration rates are sex-biased. For female-biased migration $m_{\max}^{\pm}$ coincide for all four models. With increasing proportion of male migrants (top to bottom), migration pressure on the X chromosome is reduced and differences among the models appear. All bounds $m_{\max}^{\pm}$ are derived analytically, see Eqs.(B.40),(B.42).

(corresponding to the 1:2 X-autosome ratio among migrants in this case).

The migration bounds $m_{\max}^{\pm}$ for the A→X and X→A DMIs are intermediate between the A→A and X→X DMIs. Our analytical results (see B.2) show that the upper limit of the bistable regime (*i.e.*, the value of $m_{\max}^+$) is identical for the A→X and the X→A models with dosage compensation. However, the limits for global stability, $m_{\max}^-$, can differ, which can be understood as follows:

For pure selection against migrants (no epistasis $\gamma \rightarrow 0$, and $\sigma_C < 0$, right column in Fig. 2), increased male migration reduces the effective migration pressure on the X chromosome. This leads to a corresponding increase in the migration bound $m_{\max}^-$ ($= m_{\max}^+$ in this case) for all DMIs that are lost for $m > m_{\max}^{\pm}$ because of swamping at an X-locus. This is clearly always the case for X→X DMIs, but also for the X→A model, as long as $|\sigma_C| < |\sigma_I|$ (as in our example: X fixes before A is lost). In contrast, for the A→A and the A→X model (if $|\sigma_C| < |\sigma_I|$) the DMI is lost due to swamping at the autosomal locus (continental locus in these cases). Increased male bias in migration therefore does not change the migration bound $m_{\max}^{\pm}$ in these cases.

For strong epistasis (with a deleterious immigrating allele), where the direct locus effects are less important, it is always the incompatible island allele that cannot invade on the island for migration rates $m > m_{\max}^-$. Here, any incompatible island allele that interacts with an X allele has an advantage from male-biased migration since it feels less gene flow from the competing X. This can be seen for the $m_{\max}^-$ lines in Figure 2(f),(i): While the migration bound is increased for the X→A model (and the X→X model) over the whole range of epistasis, it converges to the value of the autosomal DMI for the A→X model.

No dosage compensation

In Figure 3 we investigate migration bounds without dosage compensation, such that the differences in ploidy between autosomes and X chromosomes are no longer

masked. Relative to the model with dosage compensation, we have weaker allelic
and epistatic effects of the X chromosome. Hence, incompatible island X alleles
are easier go get swamped and also have a more difficult time to keep incompatible
continental (A or X) alleles from swamping.

Codominant nuclear DMIs without dosage compensation ($D = 0$)

Only females migrate ($R = 1$):

Unbiased migration rate ($R = 0$):

Only males migrate ($R = -1$):

----- local stability $m_{\max}^+$ — global stability $m_{\max}^-$
— $X \rightarrow A$ — $A \rightarrow X$ — $X \rightarrow X$ — $A \rightarrow A$

Figure 3: **Codominant nuclear DMIs without dosage compensation, $D = 0$.** Without dosage compensation the ploidy differences between the autosomes and the X chromosome are unmasked, inducing strong asymmetry between the $\mathcal{A}$ - $\mathcal{X}$ -models. this leads to a larger effect per allele. All bounds $m_{\max}^{\pm}$ are derived analytically, see Eqs.(B.40),(B.42). See also Figure 2 for further explanations. Note the different scaling of the y-axis in the third column.

324 The consequences can most easily be seen in the first row of Figure 3(a)-(c)
 325 with pure female migration, where, in contrast to dosage compensation, differences
 326 between the various genomic architectures are not compensated anymore. We
 327 observe a strong asymmetry between $m_{\max}^{\pm}$ of $X \rightarrow A$ and $A \rightarrow X$ -models for all levels
 328 of male migration relative to the corresponding results with dosage compensation.
 329 Here migration bounds for $X \rightarrow A$ always exceed those obtained for $A \rightarrow X$ -models.
 330 Intuitively, one can understand this as follows: In the $X \rightarrow A$ model, three immigrating
 331 X chromosomes “fight” against four resident autosomes, whereas in $A \rightarrow X$ the odds
 332 are in favor of the immigrating autosomes. Thus the island is swamped more easily
 333 in the latter case.

334 As seen for dosage compensation before, for weak epistasis ($\gamma \approx 0$, pure “selection
 335 against immigrants”), it is always the locus with weaker direct effect that is swamped
 336 first. In our example this is always the “continental” locus, because we have stronger
 337 selection on the island locus. For unbiased migration (Fig. 3(f)) all models converge
 338 to the same bound. However, introducing sex-biased migration leads to relative
 339 higher gene flow on the X for a female bias (and therefore lower bounds for models
 340 with immigrating X), as can be seen in Figure 3(c). Similarly, male-biased migration
 341 leads to weaker X-linked gene flow and a higher bounds in these models, *i.e.* $X \rightarrow A$
 342 and $X \rightarrow X$, (Fig. 3(i)).

343 If we compare migration bounds of Figure 2 and 3, we can see that dosage
 344 compensation outbalances most of the differences in $m_{\max}^{\pm}$ between $A \rightarrow X$ and $X \rightarrow A$,
 345 especially for local stability. While dosage compensation strengthens the fitness
 346 effect of the island allele in $A \rightarrow X$, the increased epistatic pressure on the continental
 347 allele in $X \rightarrow A$ is outbalanced by its increased fitness effect.

348 3.2 Cytonuclear (mitochondrial) codominant DMIs

349 Finally, we investigate cytonuclear DMIs in Figure 4, where a gene in the haploid
 350 mitochondrial genome (termed o/O for organelle) is incompatible with a nuclear

locus. Dosage compensation of the X chromosome again means that the male XyO-hybrids suffer as much as the female XXO-hybrids while they suffer only as much as XxO hybrids without dosage compensation. Relative to nuclear DMIs, three main effects lead to changes in $m_{\max}^{\pm}$:

First, the cytoplasmic locus experiences effectively stronger direct and epistatic selection (factor two in Eqs. (B.34c)), because we maintain the per locus effect identical to nuclear loci. Since a single allele already accounts for the full mitochondrial locus effect this leads to a larger effect per allele. As a consequence, $m_{\max}^{\pm}$ without sex-bias in migration is elevated relative to A→A model (gray lines in Fig. 4(a)-(c)).

Second, sex-biased migration has an even stronger effect in cytonuclear DMIs than in the X-linked nuclear DMIs: Since mitochondria are maternally inherited, the effective gene flow for mitochondrial loci is reduced to zero with pure male migration. Consequently, all migration bounds with immigrating incompatible mitochondrial genes diverge to infinity. In Figure 4 (last two rows), we study the case of strong, but not complete male bias ($R = -0.9$). Since the migration pressure on the mitochondrial locus and the X chromosome is reduced, migration bounds $m_{\max}^{\pm}$ increase for all cytonuclear DMIs, especially for those also involving X chromosomes. This increase in $m_{\max}^{\pm}$ is even further promoted by dosage compensation, strengthening the effect of X.

Finally, because of strict maternal inheritance, the dynamics of the mitochondrial locus is not influenced by any fitness effects in males. In $\mathcal{X}\text{-}\mathcal{O}$ models this also entails that dosage compensation only affects the dynamics of the X-locus - in contrast to nuclear DMIs, where also autosomal loci are affected if they interact with a hemizygous X locus. As a consequence, the boosting effect of dosage compensation on $m_{\max}^{\pm}$ is symmetric for O→X and X→O, in stark contrast to nuclear DMIs, where dosage compensation does not change much for X→A while it strongly increases A→X.

Figure 4: **Codominant cytonuclear DMIs.** Maximum permissible migration rates for local stability either coincide for all models (a)-(c), or just for $X \rightarrow O$ and $O \rightarrow X$ as well as for $O \rightarrow A$ and $A \rightarrow O$ in all other cases. Migration bounds for global stability only coincide without dosage compensation or sex-biased migration between $O \rightarrow X$ and $O \rightarrow A$, as well as for $X \rightarrow O$ and $A \rightarrow O$. The $A \rightarrow A$ model is given in panel (a)-(c) in gray as a reference. All bounds $m_{\max}^{\pm}$ are derived analytically, see Eqs.(B.40),(B.42). See Figure 2 for further explanations. Note the different scaling of the y-axis in third column.

4 Discussion

If barriers to gene flow build up among populations in primary or secondary contact, this can have important consequences for their genetic architecture. A lot of recent interest has focused on *islands of speciation (or divergence)* (Wu, 2001; Turner et al., 2005; Butlin et al., 2012; Nosil, 2012; Nosil and Feder, 2012; Via, 2012), yet corresponding empirical findings are equivocal on that matter (Cruickshank and Hahn, 2014; Pennisi, 2014). There are, however, several clear and undisputed genomic patterns of speciation, on which we concentrate here. The most widely known ones are Haldane’s rule, (Haldane, 1922), which has motivated much previous speciation research (see reviews and examples in Coyne and Orr (2004); Presgraves (2008); Lachance and True (2010); Presgraves (2010); Oka and Shiroishi (2013) and the *large X-effect* (reviewed in Presgraves, 2008), which both highlight an important role of the X chromosome (or the Z chromosome in birds) in speciation. In addition, hybrid incompatibilities are frequently observed also between nuclear and cytoplasmic markers. Plants show incompatibilities with plastid genomes (Greiner et al., 2011; Snijder et al., 2007) and mitochondria have been reported to be incompatible with nuclear genes across a wide range of species (Ellison and Burton, 2008; Lee et al., 2008; Burton and Barreto, 2012). In insects, cytoplasmic incompatibilities can also be caused by infections with the intracellular bacterium *Wolbachia* (O’Neill et al., 1992; Werren, 1997; Coyne and Orr, 2004).

In the current study we investigate how the genetic architecture of an initial hybrid incompatibility between incipient sister species can maintain divergence in the presence of ongoing gene flow. Can (primary or secondary) gene flow favor X-linked or cytonuclear DMIs over autosomal ones, and if so under which conditions? We studied this question about a possible first step towards speciation using a minimal model of a two-locus DMI in a continent-island population that allows for analytical treatment. We derive maximum permissible migration bounds which

still permit maintenance of a DMI in the face of gene flow. Conditions that yield increased migration limits facilitate speciation, as they are lost less easily and can subsequently provide more persistent seeds for further ongoing differentiation.

Conditions for parapatric DMIs

Like in the autosomal case (Bank et al., 2012), the origin and maintenance of a two-locus X-linked or cytonuclear DMI requires that at least one of the DMI substitutions (namely: the incompatible variant on the island) is adaptive. If multi-locus barriers to gene flow build up gradually from initial two-locus incompatibilities, this confirms that postzygotic parapatric speciation requires at least some degree of ecological differentiation and local adaptation. Empirically, there is widespread evidence for positive selection on genes involved in DMIs (Macnair and Christie, 1983; Ting et al., 1998; Presgraves et al., 2003; Barbash et al., 2004; Dettman et al., 2007).

For all types of DMIs, we observe two basic selective forces driving their evolution. Selection against immigrants implies that the new migrants have a fitness deficit relative to island residents, resulting in *ecological speciation* scenarios (Schluter and Conte, 2009; Nosil, 2012). A characteristic of this regime is that evolution of a stable DMI is independent of its evolutionary history.

Alternatively, a stable DMI is caused by selection against hybrids, where migrants can even have a positive fitness. If hybrids are unfit, immigrants still suffer an indirect disadvantage as long as they are rare and their genotypes are readily broken down by sex and recombination. This scenario typically leads to a bistable dynamics, where a stable DMI will only evolve from favorable starting conditions or permissive evolutionary histories (such as secondary contact). The scenario has also been referred to as *mutation-order-speciation* (Mani and Clarke, 1990).

We measure the strength of a parapatric DMI by means of two migration bounds. The higher one, $m_{\max}^+$, is the limit beyond which a DMI can neither evolve nor an

existing one can be maintained. The lower bound, $m_{\max}^-$, is the limit up to which a DMI will always evolve in the face of gene flow, irrespective of the evolutionary history (globally stable DMI). For migration rates between both bounds, a DMI is maintained, but will evolve only under favorable histories, such as secondary contact, or if the second incompatible substitution occurs on the continent.

Contrasting different DMI architectures

We find that the genetic architecture of a DMI (with incompatible genes on autosomes, X chromosomes, or in the mitochondrial genome) can have a strong effect on its stability. However, this effect also crucially depends on other factors, such as, in particular, the level of dosage compensation and the sex-bias in the migration rates.

First, without dosage compensation and without sex-biased gene flow, the hemizyosity of the X chromosome in males leads to shifts of $m_{\max}^\pm$ in the presence of epistasis compared to autosome-autosome DMIs. This is due to ploidy differences: “3 X chromosomes fight 4 autosomes”. Therefore, the A→X scenario (where a resident X-linked allele competes with an immigrating incompatible autosomal gene) constitutes a weaker barrier to gene flow than the X→A model. Note that this effect depends crucially on the (negative) epistasis of the DMI and is not observed in a single-locus model of local adaptation. Second, dosage compensation strengthens the X alleles, which leads to higher migration bounds in all X-linked DMIs. In particular, it increases stability of DMIs with an incompatible X locus on the island, compensating the A→X versus X→A asymmetry. Third, sex-biased migration leads to lower/higher limits for DMIs with immigrating X for female/male bias. Fourth, our results in the SI Section A.1 show no large difference between codominant and recessive nuclear DMIs (which lead to Haldane’s rule) concerning the migration bounds. In fact, the difference for X-linked DMIs are even smaller than for autosome-autosome DMIs. Fifth, for cytonuclear DMIs we often observe stronger barriers to gene flow since the haploid cytoplasmic alleles experience the full locus effect. Furthermore, sex-bias

in migration yields an especially strong effect, as for pure male migration effective gene flow at the mitochondrial locus ceases completely.

Our numerical simulations for the effect of LD in the SI Section A.2 agree with the approximate analytical results for weak and moderately strong DMIs. For very strong DMIs, stronger deviations occur for codominant $A \rightarrow A$ and $X \rightarrow X$ DMIs, which maintain very strong LD once all (male and female) hybrids with incompatible alleles are almost inviable/infertile. As a consequence, gene flow among the continent and island haplotypes is blocked and we obtain higher migration bounds relative to $X \rightarrow A$ and $A \rightarrow X$ DMIs. For the latter two, F1 hybrid males carrying the compatible x allele (genotype $Aaxy$) are viable and can produce ax gametes for the F2 generation. This effect of extreme LD and blocked gene flow does not exist for recessive DMIs (see SI Section A.1 for details). Our numerical simulations also show that the effect of drift is usually small and does not lead to qualitative changes (SI Section A.3). Since DMI alleles can be lost by drift, stochastic migration bounds $m_{\max}^N$ are generally smaller than their deterministic counterparts. In SI Section A.3, we present an analytical approximation to estimate this reduction due to drift.

The large X -effect

Summarizing all different cases described above, we find that the most stable DMIs are almost always X-linked, where migration bounds are typically enhanced by a factor of $4/3$ to 2 relative to autosomal DMIs (unless migration is strongly female biased). Although this is not a very strong effect, it is very general and applies whenever gene flow plays a role at any stage of the speciation process. This includes, in particular, scenarios of secondary contact and also later stages of the speciation process where additional barriers to gene flow exist in the genomic background. In this latter case, the gene flow at the focal DMI loci needs to be replaced by appropriate effective migration rates (Barton and Bengtsson, 1986). The pattern that follows from a more stable X barrier is consistent with a higher density of

486 X-linked hybrid incompatibilities, the *large X-effect*.

487 Our results show a clear boost of X migration bounds, in particular, if there is
488 dosage compensation and if migration is male biased. Empirical studies show that
489 sex-biased migration is common in nature and report a prevalence for migration
490 of the heterogametic sex in both mammals, where dispersal is on average male
491 biased, (Lawson Handley and Perrin, 2007) and in birds, where female dispersal
492 dominates (Greenwood, 1980). In the context of our results, these trends strengthen
493 the predicted pattern of a *large X-effect* or *large Z-effect*, respectively.

494 One example stems from the house mouse, *Mus musculus*. There is strong
495 empirical evidence for a *large X-effect* in this species (Tucker et al., 1992; Good
496 et al., 2008; White et al., 2012), such as the major involvement of the X chromosome
497 in hybrid sterility (Oka et al., 2004; Storchova et al., 2004). Mice exhibit rather
498 complete dosage compensation due to X-inactivation in females (Payer and Lee,
499 2008). Furthermore, the house mouse displays male-biased dispersal at breeding
500 age (Greenwood, 1980; Gerlach, 1990).

501 Several alternative mechanisms as potential underlying causes for a *large X-effect*
502 have been discussed in the literature, such as sex ratio meiotic drive, regulation of
503 the X chromosome in the male germ line (Coyne and Orr, 2004; Presgraves, 2008), or
504 faster evolution of the X chromosome (termed *faster-X-effect* Charlesworth et al.,
505 1987). In the panmictic population model by Charlesworth et al. (1987), faster
506 evolution on the X chromosome results if adaptations are, on average, recessive
507 and are thus exposed to stronger selection on the hemizygous X. We note that our
508 model with gene flow predicts an advantage of X-linked genes for island adaptations
509 even if they are not recessive, but codominant (or even slightly dominant, see SI
510 Section A.4 for details and proofs). Since the *faster X-effect* (more adaptations on
511 the X) also favors a *larger X-effect* (more incompatibilities involving the X), this
512 is another way how speciation with gene-flow can contribute to this pattern. In
513 summary a mono-causal explanation for the *large X-effect* seems unlikely, and it

514 remains an open question, to which extent each factor contributes. Our study adds
515 differentiation under gene flow as another element to this mix.

516 Our results relate to Haldane’s rule only in so far as this pattern partially overlaps
517 with the *large X-effect*. Beyond that, we do not obtain a prediction. In particular,
518 the migration bounds for codominant and recessive DMIs are similar (while only the
519 latter lead to Haldane’s rule).

520 Introgression patterns

521 A second conclusion from our results that can be related to data is that X-linked
522 alleles in an incompatibility face stronger barriers to introgression than the corresponding
523 autosomal alleles. This effect rests on two basic observations: the tendency for
524 higher migration bounds of all X-linked DMIs with dosage compensation (which
525 also contributes to a large X-effect), and the asymmetry promoting $A \rightarrow X$ over $X \rightarrow A$
526 introgression that we observe for the incompatible allele if dosage compensation is
527 incomplete or absent (the 3 versus 4 chromosomes effect). Our findings agree with
528 the result of a recent simulation study for DMIs on a cline by Wang (2013), who
529 showed that, for an X-autosome DMIs, the incompatible X allele flows less easily
530 across a cline than the autosomal allele.

531 A pattern of reduced X-introgression relative to autosomal introgression has been
532 recognized in many sister-species in nature. In the complex of *Anopheles gambiae*
533 sister clades Fontaine et al. (2015) found “pervasive autosomal introgression” between
534 different species, in contrast to the X chromosome, which contains disproportionately
535 more factors in reproductive isolation.

536 Liu et al. (2015) report three interspecies hybridization events in mice (*Mus*
537 *musculus domesticus* and *M. spretus*), leading to exclusively autosomal, partially
538 adaptive introgression. Similarly, Macholán et al. (2007) showed weaker introgression
539 patterns and lower selection pressure on the X chromosomes compared to the autosomes
540 in the central European mouse hybrid zone of *Mus musculus musculus* and *M.*

541 *m. domesticus*. The authors suppose that the X is shielded more effectively from
542 introgression due to the *large X-effect*.

543 Further examples exist for birds. Sætre et al. (2003) report “rather extensive
544 hybridization and backcrossing in sympatry” between two populations of flycatchers
545 hybridizing in secondary contact. Nevertheless, gene flow was again predominantly
546 found on the autosome. Hooper and Price (2015) report that derived cross-species
547 inversions among sister species of Estrilid finches are strongly enriched on the Z
548 chromosome. The pattern is strongest in continental clades with high level of
549 sympatry and (plausibly) higher levels of gene-flow during the speciation process. If
550 inversions harbor DMIs, this is consistent with our finding that derived incompatibilities
551 on the Z chromosome are more stable to gene flow than autosomal incompatibilities.

552

553 Also other factors, such as recombination, can influence differential introgression
554 on X chromosomes and autosomes. Indeed, there is empirical evidence that recombination
555 can structure autocorrelation patterns among introgressed loci. However, available
556 data also show that recombination cannot be the the sole explanation for differential
557 introgression among genomic regions, e.g. in mice (Payseur et al., 2004) or finches (Hooper
558 and Price, 2015). As for the *large X-effect* our mechanism is one of several possible
559 ones.

560 **Biological assumptions and limitations of the model**

561 Our study has been intended as a minimal model approach that allows for analytical
562 treatment. As such, it rests on several simplifying assumptions concerning the
563 genetics of the DMI and the ecological setting. These limitations suggest possible
564 model extensions for future work.

565 All our results assume a simple DMI between just two loci. This is in line with
566 most previous theoretical work and known empirical cases (Coyne and Orr, 2004;
567 Maheshwari and Barbash, 2011). Nevertheless, complex DMIs involving multiple

568 loci are clearly relevant at later stages of a speciation process and could lead to new
569 effects that are not captured here (*e.g.* Lindtke and Buerkle, 2015).

570 Our fitness scheme for two-locus DMIs comprises codominant and recessive
571 cases. Empirically, the functional form depends on the underlying mechanisms
572 causing hybrid fitness loss, which is still debated. Hybrid incompatibilities can
573 be due to loss-of-function or gain-of-function mutations (reviewed by Maheshwari
574 and Barbash, 2011). While the former tend to act recessively, the latter will likely
575 affect heterozygotes, and may be better captured by a partially dominant DMI.

576 Recessive DMIs, in turn, occur in a number of different types, (*e.g.* Presgraves,
577 2010; Cattani and Presgraves, 2012; Matsubara et al., 2015), which lead to slightly
578 different models. We have briefly studied some of these alternatives analytically, such
579 as a *recessive-A codominant-X*-DMI or a *codominant-A recessive-X*-DMI (data not
580 shown). We did not detect any noteworthy difference in their evolutionary dynamics
581 or for the migration bounds relative to the results reported here. Still, more relevant
582 changes are clearly possible, for example if the single locus effects can lead to over-
583 or underdominance.

584 For the results presented, we assume that dosage compensation enhances not only
585 the single-locus effect, but also the incompatibility. Empirically, hybrid incompatibilities
586 are frequently dosage-sensitive, *e.g.* in a *Arabidopsis thaliana* / *A. arenosa* cross,
587 where a DMI results due to failure in gene silencing Josefsson et al. (2006), or in
588 a *Mus musculus musculus* / *M. m. domesticus* cross, where X-linked hybrid male
589 sterility results from over-expression of X-linked genes in spermatogenesis (Good
590 et al., 2010). Furthermore, in haplo-diploid Nasonian wasps genetically engineered
591 diploid males were less affected by hybrid sterility than haploid male hybrids, pointing
592 to a strong effect of ploidy on hybrid fertility Beukeboom et al. (2015).

593 Nevertheless, we also investigated the effect of dosage compensation only on the
594 single locus effect or only on the incompatibility (results not shown). As expected,
595 we obtain intermediate patterns between no and full dosage compensation.

Concerning the ecological assumptions, our model assumes unidirectional gene flow between two panmictic demes. While our results readily extend to weak back migration (which leads only to slight shifts of the equilibria), strong bidirectional migration can lead to qualitatively new effects that are not captured by our framework. For example, polymorphisms at single loci can be maintained for arbitrarily strong gene flow if heterogeneous soft selection leads to a rare-type advantage (Levene, 1953). Furthermore, generalist genotypes that are inferior in both demes, but do well on average, can be maintained if (and only if) bidirectional migration is sufficiently strong (see Akerman and Bürger, 2014, for results in a two-locus model without epistasis).

Alternative models for the population structure can also lead to substantial differences. In particular, our two-deme model ignores isolation by distance, which can be captured either in a discrete cline model with a chain of demes, or in a continuous-space framework. It is expected that polymorphisms (and DMIs) can be maintained with much larger gene flow (or weaker selection) in these settings (Barton, 2013). Still, several of our key results, such as reduced introgression of X-linked incompatibility alleles, should still hold under these conditions (see Wang, 2013, for a discrete cline model).

5 Acknowledgements

We thank Andrea Betancourt, Alexandre Blanckaert, Reinhard Bürger, Brian Charlesworth, Andy Clark, Sebastian Matuszewsky, Sylvain Mousset, Mohamed Noor, Sally Otto, Christian Schlötterer, Maria Servedio, Derek Setter, Claus Vogl and three referees for helpful discussions, suggestions and comments on the manuscript. This work was made possible with financial support by the Austrian Science Fund (FWF) via funding for the Vienna Graduate School for Population Genetics.

References

- Agrawal, A., Feder, J., and Nosil, P. (2011). Ecological divergence and the origins of intrinsic postmating isolation with gene flow. *International Journal of Ecology*, 2011.
- Akerman, A. and Bürger, R. (2014). The consequences of gene flow for local adaptation and differentiation: a two-locus two-deme model. *Journal of mathematical biology*, 68(5):1135–1198.
- Bank, C., Bürger, R., and Hermisson, J. (2012). The limits to parapatric speciation: Dobzhansky–Muller incompatibilities in a continent–island model. *Genetics*, 191(3):845–863.
- Barbash, D., Awadalla, P., and Tarone, A. (2004). Functional divergence caused by ancient positive selection of a *Drosophila* hybrid incompatibility locus. *PLoS biology*, 2:839–848.
- Barnard-Kubow, K. B., So, N., and Galloway, L. F. (2016). Cytonuclear incompatibility contributes to the early stages of speciation. *Evolution*, 70(12):2752–2766.
- Barton, N. (2013). Does hybridization influence speciation? *Journal of evolutionary biology*, 26(2):267–269.
- Barton, N. and Bengtsson, B. (1986). The barrier to genetic exchange between hybridising populations. *Heredity*, 57(3):357–376.
- Bateson, W. (1909). Heredity and variation in modern lights. *Darwin and modern science*, 85:101.
- Beukeboom, L., Koevoets, T., Morales, H. E., Ferber, S., and van de Zande, L. (2015). Hybrid incompatibilities are affected by dominance and dosage in the haplodiploid wasp *Nasonia*. *Frontiers in Genetics*, 6:14.
- Bürger, R. and Akerman, A. (2011). The effects of linkage and gene flow on local adaptation: A two-locus continent–island model. *Theoretical population biology*, 80(4):272–288.

- 646 Burton, R. and Barreto, F. (2012). A disproportionate role for mtDNA in
647 Dobzhansky–Muller incompatibilities? *Molecular Ecology*, 21(20):4942–4957.
- 648 Butlin, R., Debelle, A., Kerth, C., Snook, R., Beukeboom, L., Castillo, C., Diao, W.,
649 Maan, M., Paolucci, S., Weissing, F., et al. (2012). What do we need to know about
650 speciation? *Trends in Ecology & Evolution*, 27(1):27–39.
- 651 Cattani, M. and Presgraves, D. (2012). Incompatibility Between X Chromosome Factor and
652 Pericentric Heterochromatic Region Causes Lethality in Hybrids Between *Drosophila*
653 *melanogaster* and Its Sibling Species. *Genetics*, 191(2):549–559.
- 654 Charlesworth, B., Coyne, J., and Barton, N. (1987). The relative rates of evolution of sex
655 chromosomes and autosomes. *American Naturalist*, pages 113–146.
- 656 Corbett-Detig, R., Zhou, J., Clark, A., Hartl, D., and Ayroles, J. (2013). Genetic
657 incompatibilities are widespread within species. *Nature*, 504(7478):135–137.
- 658 Coyne, J. and Orr, H. (1989). Two rules of speciation. *Speciation and its Consequences*,
659 pages 180–207.
- 660 Coyne, J. and Orr, H. (2004). *Speciation*. Sinauer Associates Sunderland, MA.
- 661 Cruickshank, T. and Hahn, M. (2014). Reanalysis suggests that genomic islands of
662 speciation are due to reduced diversity, not reduced gene flow. *Molecular Ecology*,
663 23(13):3133–3157.
- 664 Dettman, J., Sirjusingh, C., Kohn, L., and Anderson, J. (2007). Incipient speciation by
665 divergent adaptation and antagonistic epistasis in yeast. *Nature*, 447(7144):585–588.
- 666 Dobzhansky, T. (1936). Studies on hybrid sterility. II. Localization of sterility factors in
667 *Drosophila pseudoobscura* hybrids. *Genetics*, 21(2):113.
- 668 Ellegren, H. (2009). Genomic evidence for a large-Z effect. *Proceedings of the Royal Society*
669 *B: Biological Sciences*, 276(1655):361–366.

Ellegren, H., Hultin-Rosenberg, L., Brunström, B., Dencker, L., Kultima, K., and Scholz, B. (2007). Faced with inequality: chicken do not have a general dosage compensation of sex-linked genes. *BMC biology*, 5(1):40.

Ellison, C. and Burton, R. (2008). Interpopulation hybrid breakdown maps to the mitochondrial genome. *Evolution*, 62(3):631–638.

Feder, J. and Nosil, P. (2009). Chromosomal inversions and species differences: when are genes affecting adaptive divergence and reproductive isolation expected to reside within inversions? *Evolution*, 63(12):3061–3075.

Felsenstein, J. (1981). Skepticism towards Santa Rosalia, or why are there so few kinds of animals? *Evolution*, pages 124–138.

Fontaine, M., Pease, J., Steele, A., Waterhouse, R. M., Neafsey, D., Sharakhov, I., Jiang, X., Hall, A., Catteruccia, F., Kakani, E., et al. (2015). Extensive introgression in a malaria vector species complex revealed by phylogenomics. *Science*, 347(6217):1258524.

Gavrilets, S. (1997). Hybrid zones with Dobzhansky-type epistatic selection. *Evolution*, pages 1027–1035.

Gerlach, G. (1990). Dispersal mechanisms in a captive wild house mouse population (*Mus domesticus* Ratty). *Biological Journal of the Linnean Society*, 41(1-3):271–277.

Good, J., Dean, M., and Nachman, M. (2008). A complex genetic basis to X-linked hybrid male sterility between two species of house mice. *Genetics*, 179(4):2213–2228.

Good, J., Giger, T., Dean, M., and Nachman, M. (2010). Widespread over-expression of the X chromosome in sterile F1 hybrid mice. *PLoS genetics*, 6(9):e1001148.

Graves, J., Disteche, C., et al. (2007). Does gene dosage really matter. *J Biol*, 6(1):1.

Greenwood, P. (1980). Mating systems, philopatry and dispersal in birds and mammals. *Animal behaviour*, 28(4):1140–1162.

Greiner, S., Rauwolf, U., Meurer, J., and Herrmann, R. (2011). The role of plastids in plant speciation. *Molecular ecology*, 20(4):671–691.

- 696 Haldane, J. (1922). Sex ratio and unisexual sterility in hybrid animals. *Journal of genetics*,
697 12(2):101–109.
- 698 Hooper, D. and Price, T. (2015). Rates of karyotypic evolution in Estrildid finches differ
699 between island and continental clades. *bioRxiv*, page 013987.
- 700 Josefsson, C., Dilkes, B., and Comai, L. (2006). Parent-dependent loss of gene silencing
701 during interspecies hybridization. *Current Biology*, 16(13):1322–1328.
- 702 Kimura, M. (1957). Some problems of stochastic processes in genetics. *The Annals of*
703 *Mathematical Statistics*, pages 882–901.
- 704 Lachance, J. and True, J. (2010). X-autosome incompatibilities in *Drosophila*
705 *melanogaster*: tests of Haldane’s rule and geographic patterns within species. *Evolution*,
706 64(10):3035–3046.
- 707 Lawson Handley, L. and Perrin, N. (2007). Advances in our understanding of mammalian
708 sex-biased dispersal. *Molecular Ecology*, 16(8):1559–1578.
- 709 Lee, H., Chou, J., Cheong, L., Chang, N., Yang, S., and Leu, J. (2008). Incompatibility
710 of nuclear and mitochondrial genomes causes hybrid sterility between two yeast species.
711 *Cell*, 135(6):1065–1073.
- 712 Levene, H. (1953). Genetic equilibrium when more than one ecological niche is available.
713 *American Naturalist*, pages 331–333.
- 714 Lindtke, D. and Buerkle, C. (2015). The genetic architecture of hybrid incompatibilities
715 and their effect on barriers to introgression in secondary contact. *Evolution*.
- 716 Liu, K., Steinberg, E., Yozzo, A., Song, Y., Kohn, M., and Nakhleh, L. (2015). Interspecific
717 introgressive origin of genomic diversity in the house mouse. *Proceedings of the National*
718 *Academy of Sciences*, 112(1):196–201.
- 719 Macholán, M., Munclinger, P., Šugerková, M., Dufková, P., Bímová, B., Božíková, E.,
720 Zima, J., and Piálek, J. (2007). Genetic analysis of autosomal and X-linked markers
721 across a mouse hybrid zone. *Evolution*, 61(4):746–771.

Macnair, M. and Christie, P. (1983). Reproductive isolation as a pleiotropic effect of copper tolerance in *Mimulus guttatus*. *Heredity*, 50(3):295–302.

Maheshwari, S. and Barbash, D. (2011). The genetics of hybrid incompatibilities. *Annual review of genetics*, 45:331–355.

Mallet, J. (2008). Hybridization, ecological races and the nature of species: empirical evidence for the ease of speciation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1506):2971–2986.

Mani, G. and Clarke, B. (1990). Mutational order: a major stochastic process in evolution. *Proceedings of the Royal Society of London B: Biological Sciences*, 240(1297):29–37.

Masly, J. and Presgraves, D. (2007). High-resolution genome-wide dissection of the two rules of speciation in *Drosophila*. *PLoS Biol*, 5(9):e243.

Matsubara, K., Yamamoto, E., Mizobuchi, R., Yonemaru, J., Yamamoto, T., Kato, H., and Yano, M. (2015). Hybrid Breakdown Caused by Epistasis-Based Recessive Incompatibility in a Cross of Rice (*Oryza sativa* L.). *Journal of Heredity*, 106(1):113–122.

Muller, H. (1942). Isolating mechanisms, evolution and temperature. In *Biological Symposia*, volume 6, pages 71–125.

Nosil, P. (2012). *Ecological speciation*. Oxford University Press.

Nosil, P. and Feder, J. (2012). Genomic divergence during speciation: causes and consequences. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1587):332–342.

Oka, A., Mita, A., Sakurai-Yamatani, N., Yamamoto, H., Takagi, N., Takano-Shimizu, T., Toshimori, K., Moriwaki, K., and Shiroishi, T. (2004). Hybrid breakdown caused by substitution of the X chromosome between two mouse subspecies. *Genetics*, 166(2):913–924.

Oka, A. and Shiroishi, T. (2013). Regulatory divergence of X-linked genes and hybrid male sterility in mice. *Genes & genetic systems*, 89(3):99–108.

O'Neill, S., Giordano, R., Colbert, A., Karr, T., and Robertson, H. (1992). 16S
rRNA phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic
incompatibility in insects. *Proceedings of the National Academy of Sciences*,
89(7):2699–2702.

Orr, H. and Turelli, M. (2001). The evolution of postzygotic isolation: accumulating
Dobzhansky-Muller incompatibilities. *Evolution*, 55(6):1085–1094.

Payer, B. and Lee, J. (2008). X chromosome dosage compensation: how mammals keep
the balance. *Annual review of genetics*, 42:733–772.

Payseur, B. A., Krenz, J. G., and Nachman, M. W. (2004). Differential patterns of
introgression across the x chromosome in a hybrid zone between two species of house
mice. *Evolution*, 58(9):2064–2078.

Pennisi, E. (2014). Disputed islands. *Science*, 345(6197):611–613.

Presgraves, D. (2008). Sex chromosomes and speciation in *Drosophila*. *Trends in Genetics*,
24(7):336–343.

Presgraves, D. (2010). The molecular evolutionary basis of species formation. *Nature
Reviews Genetics*, 11(3):175–180.

Presgraves, D., Balagopalan, L., Abmayr, S., and Orr, H. (2003). Adaptive evolution
drives divergence of a hybrid inviability gene between two species of *Drosophila*. *Nature*,
423(6941):715–719.

Rutschman, D. H. (1994). Dynamics of the two-locus haploid model. *Theoretical population
biology*, 45(2):167–176.

Sætre, G., Borge, T., Lindroos, K., Haavie, J., Sheldon, B., Primmer, C., and Syvänen,
A.-C. (2003). Sex chromosome evolution and speciation in *Ficedula* flycatchers.
Proceedings of the Royal Society of London B: Biological Sciences, 270(1510):53–59.

Schluter, D. and Conte, G. (2009). Genetics and ecological speciation. *Proceedings of the
National Academy of Sciences*, 106(Supplement 1):9955–9962.

774 Slatkin, M. (1987). Gene flow and the geographic structure of natural populations. *Science*,
775 236:787–792.

776 Snijder, R., Brown, F., and van Tuyl, J. (2007). The role of plastome-genome
777 incompatibility and biparental plastid inheritance in interspecific hybridization in
778 the genus *Zantedeschia* (Araceae). *Floriculture and Ornamental Biotechnology*,
779 1(2):150–157.

780 Storchova, R., Gregorová, S., Buckiova, D., Kyselova, V., Divina, P., and Forejt, J. (2004).
781 Genetic analysis of X-linked hybrid sterility in the house mouse. *Mammalian Genome*,
782 15(7):515–524.

783 Sweigart, A. and Flagel, L. (2014). Evidence of Natural Selection Acting on a Polymorphic
784 Hybrid Incompatibility Locus in *Mimulus*. *Genetics*, pages genetics–114.

785 Ting, C., Tsaur, S., Wu, M., and Wu, C. (1998). A rapidly evolving homeobox at the site
786 of a hybrid sterility gene. *Science*, 282(5393):1501–1504.

787 Tucker, P., Sage, R., Warner, J., Wilson, A., and Eicher, E. (1992). Abrupt cline for sex
788 chromosomes in a hybrid zone between two species of mice. *Evolution*, pages 1146–1163.

789 Turelli, M. and Orr, H. (2000). Dominance, epistasis and the genetics of postzygotic
790 isolation. *Genetics*, 154(4):1663–1679.

791 Turner, T., Hahn, M., and S., N. (2005). Genomic islands of speciation in *Anopheles*
792 *gambiae*. *PLoS biology*, 3(9):e285.

793 Via, S. (2012). Divergence hitchhiking and the spread of genomic isolation during ecological
794 speciation-with-gene-flow. *Philosophical Transactions of the Royal Society B: Biological*
795 *Sciences*, 367(1587):451–460.

796 Via, S. and West, J. (2008). The genetic mosaic suggests a new role for hitchhiking in
797 ecological speciation. *Molecular Ecology*, 17(19):4334–4345.

- 798 Wang, R. (2013). Gene flow across a hybrid zone maintained by a weak heterogametic
799 incompatibility and positive selection of incompatible alleles. *Journal of Evolutionary*
800 *Biology*, pages 386–396.
- 801 Werren, J. (1997). Biology of wolbachia. *Annual review of entomology*, 42(1):587–609.
- 802 White, M., Stubbings, M., Dumont, B. L., and Payseur, B. (2012). Genetics and evolution
803 of hybrid male sterility in house mice. *Genetics*, 191(3):917–934.
- 804 Wu, C. (2001). The genic view of the process of speciation. *Journal of Evolutionary*
805 *Biology*, 14(6):851–865.
- 806 Yeaman, S. and Otto, S. P. (2011). Establishment and maintenance of adaptive genetic
807 divergence under migration, selection, and drift. *Evolution*, 65(7):2123–2129.