Hybridization affects life-history traits and host specificity in *Diorhabda* spp.

Bitume, E.V., Bean, D. Stahlke, A.R., Hufbauer, R. A.

Addresses

¹ Bioagricultural Science and Pest Management, Plant Sciences C129, Colorado State University, Fort Collins, CO

² Colorado Department of Agriculture, Palisade, CO

Statement of authorship: EB, DB, and RA designed the experiments, EB performed the experiments and analysed the data. EB wrote the first draft of the manuscript, and DB, AS, and RH contributed substantially to revisions.

Running title: Hybridization in *Diorhabda*

Corresponding author:

Ellyn Bitume

BSPM

Plant Sciences C129

Colorado State University

Fort Collins, CO 80523-1177

ebitume@gmail.com

Abstract

Hybridization is an influential evolutionary process that has been viewed alternatively as an evolutionary dead-end or as an important creative evolutionary force. In colonizing species, such as introduced biological control agents, hybridization can negate the effects of bottlenecks and genetic drift through increasing genetic variation. Such changes could be beneficial to a biological control program by increasing the chances of establishment success. However, hybridization can also lead to the emergence of transgressive phenotypes that could alter host specificity; an important consideration when assessing potential non-target impacts of planned agents. In a series of lab experiments, we investigated the effects of hybridization between three species of *Diorhabda* released to control invasive *Tamarix* (saltcedar) on life history traits through two generations, and through the third generation for one cross. Depending on the cross, hybridization had either a positive or neutral impact on development time, adult mass, and fecundity. We evaluated preference for the target (saltcedar) relative to a non-target host *Tamarix aphylla* (athel), and found host specificity patterns varied in two of the three hybrids, demonstrating the possibility for hybridization to alter host preference. Importantly, the overall effects of hybridization were inconsistent by cross, leading to unpredictability in the outcome of using hybrids in biological control.

Keywords:

Biological control
Life-history traits
Host specificity
Hybrid vigor
Inbreeding
Invasive species

1. Introduction

Hybridization is an influential evolutionary process that has been viewed alternatively as an evolutionary dead-end, because hybrids are often less fit than the parental species (Mayr 1963; Dobzhanski 1970) or as an important creative evolutionary force (Anderson & Stebbins 1954; Ellstrand & Schierenbeck 2000). On the detrimental side, hybrid breakdown, or outbreeding depression, can decrease performance of hybrid individuals across a suite of traits linked to fitness, such as development time, mortality, and fecundity (Burton *et al.* 1999; Edmands 2002). On the positive side, hybridization can increase fitness relative to parents directly through heterozygote advantage (overdominance of beneficial traits) (Edmands 2002; Hedrick & Garcia-Dorado 2016; Lee *et al.* 2016) or by alleviating high mutational load (heterosis) and reducing inbreeding depression, and indirectly through restoring genetic variation lost through genetic drift or bottlenecks in population size (genetic rescue). Even in populations that have not experienced strong drift or bottlenecks, hybridization can increase overall population genetic variation, resulting in increased ability to respond to selection pressures (Fisher 1930). Additionally, hybridization can facilitate the formation of novel genotypes, potentially producing ‘transgressive’ phenotypes that fall outside the range of either parent (Rieseberg *et al.* 1999). Alternatively, hybridization can in some cases have minimal effects if the genetic distance between parents is small (Mallet 2005).

Intraspecific hybridization between recently diverged species may be particularly beneficial in colonizing populations that pass through strong bottlenecks in population size, in turn losing genetic variation, and potentially becoming inbred (Ellstrand & Schierenbeck 2000; Dlugosch & Parker 2008; Rius & Darling 2014; Laugier *et al.* 2016). In the planned release of specialized biological control agents, the goal is for the intentionally released population to

establish and propagate (Seastedt 2015), to feed on the target host (typically an invasive weed or insect), and not shift to use other, non-target hosts. Biological control programs have a fairly low success rate (<50%), mostly due to lack of establishment of agents in their new environment (Van Driesche *et al.* 2010). As an evolutionary mechanism, hybridization might allow these establishing populations to better face adaptive challenges in the novel environment. There is some evidence that releasing different “strains” or ecotypes of biological control agents in an effort to increase genetic variation might improve establishment success (Hopper *et al.* 1993; Henry *et al.* 2010). New evidence suggests that increased genetic variation can be even more important than augmenting population size in promoting population growth (Frankham 2015; Hufbauer *et al.* 2015; Frankham 2016). Colonizing populations also experience novel environments in which transgressive phenotypes may, by chance, have higher fitness than parental phenotypes (Ellstrand & Schierenbeck 2000). Yet, the quantification of genetic variation in populations of biological control agents planned for release is not yet a standard procedure, likely because of the lack of studies investigating the effects of increased variation on long-term establishment.

Releasing genetically distinct ecotypes in the same area can promote hybridization. Only a few studies have looked at hybridization in biological control agents (Hoffmann *et al.* 2002; Mathenge *et al.* 2010; Benvenuto *et al.* 2012; Szucs *et al.* 2012). Szucs *et al.* (2012) found that hybridization improved performance in vital life-history traits, which could improve control of the target pest. However, there is also evidence that hybridization can decrease host specificity or increase host range as a result of changes in phenotype (Hoffmann *et al.* 2002; Mathenge *et al.* 2010). Such a change would increase the risk of biological control to non-target species dramatically. Thus, it is imperative that more experiments are executed to understand the

consequences of hybridization for biological control programs, including evaluating the degree to which it will be possible to draw general conclusions versus research being needed on a case-by-case basis.

We present research in which we quantify the effects of hybridization between biological control agents in the genus *Diorhabda* that were released to control *Tamarix* (saltcedar, or tamarisk) in North America. Saltcedar in North America is comprised of a hybrid swarm of *Tamarix chinensis* and *T. ramosissima* (Gaskin and Schall 2002). It is an invasive weed that has colonized riparian habitats from Montana to Mexico (Gaskin & Schaal 2002). In 2001, the USDA's Animal and Plant Health Inspection Service (USDA APHIS) approved the open field release of the central Asian salt cedar leaf beetle, *D. elongata*, as a biological control agent for saltcedar (DeLoach et al. 2003). *Diorhabda elongata* was classified as a single wide-ranging species that specialized on *Tamarix* spp and comprised different subspecies and ecotypes. To match environmental conditions in North America, the saltcedar biological control program eventually utilized seven *Diorhabda* ecotypes with native ranges stretching from north Africa to central Asia (Tracy & Robbins 2009; Bean et al. 2013a). A recent taxonomic revision of the *Tamarix*-feeding *Diorhabda* has used morphological and biogeographical data to define this group as a complex comprising five species: *D. meridionalis*, *D. carinulata*, *D. carinata*, and *D. sublineata* (Tracy & Robbins 2009). Recent genetic studies using amplified fragment length polymorphisms (AFLPs) revealed four major clades within this group which coincide with the four morphospecies (Bean et al. 2013b). There was also a fifth species, *D. meridionalis*, not currently used in the saltcedar biological control program. Currently, *D. carinulata* is the most widespread of the species in North America and covers large areas in Oregon, Idaho, Wyoming, Colorado, Utah, Nevada, northern Arizona, and northern New Mexico (Bean et al. 2013a). The

other three species have all been released in Texas and have started spreading (Michels *et al.* 2013). Hybridization is possible between all four taxa, but hybrids between *D. carinulata* and each of the other three species produce few viable offspring. In contrast, egg viability of hybrids between *D. elongata*, *D. carinata*, and *D. sublineata* is comparable to that of the parents (Bean *et al.* 2013b). We crossed these three species reciprocally, and tracked performance over three generations to quantify the effects of hybridization. We measured several life history traits crucial to fitness, as well as host preference for saltcedar relative to a non-target plant, *Tamarix aphylla* (athel hereafter) for both hybrid offspring and the parental species.

2. Materials & Methods

2.1 Organism

The beetles used in our experiments stemmed from samples originally collected from saltcedar in Eurasia and North Africa. Descendants of these samples were used to establish laboratory populations maintained at the Palisade Insectary, Biological Pest Control Program, Colorado Department of Agriculture, Palisade, CO (CDA Palisade). Colonies were maintained on cuttings of saltcedar, including *T. ramosissima*, *T. chinensis*, and their hybrids (Gaskin & Schaal 2002), in 7.5 liter capacity plastic containers with mesh siding for ventilation in incubators under a light regime of 16:8 and 27°C/16°C. *Diorhabda carinata* (“C” hereafter) used in this study were originally collected in 2002 near Karshi (Qarshi), Uzbekistan (38.86 N, 65.72 E; elevation 350 m), and *Diorhabda sublineata* (“S” hereafter) originated near the town of Sfax, Tunisia (34.66 N, 10.67 E, elevation 10 m). Both these species were maintained in the lab prior to our experiments. *Diorhabda elongata* (“E” hereafter) were collected from Sfakaki, Crete, Greece (35.83 N, 24.6 E, elevation 7 m) and in 2004 they were first released upstream of Esparto, CA along Cache Creek in the Capay Valley. Unlike the other two species, *D. elongata*

used in our experiments were collected in 2015 from the field in the Capay Valley and used to start a laboratory colony. No other species were released into the Capay Valley nor have any been established within 150 miles, and therefore it is reasonable to assume that there was no chance for hybridization before our experiments.

2.2 Crosses

To produce the first generation of hybrids, seven virgin females and seven males of each species were placed together into a plastic bucket with mesh siding (7.5 liter) with saltcedar. Since male female directionality can affect the fitness of hybrid offspring (Payseur & Rieseberg 2016), we crossed each species reciprocally. We thus made the following hybrids: *D. carinulata* x *D. elongata* ($C_f \times E_m$, $E_f \times C_m$), *D. carinulata* x *D. sublineata* ($C_f \times S_m$, $S_f \times C_m$), *D. elongata* x *D. sublineata* ($E_f \times S_m$, $S_f \times E_m$), plus the parents ($C_f \times C_m$, $E_f \times E_m$, $S_f \times S_m$). To keep inbreeding depression to a minimum, we initiated two separate buckets for each of the parental lines so that density remained the same but so the parental generation had 14 families rather than 7 for the crosses. All adults were allowed to remain in the buckets for five days of egg-laying.

2.3 F_1 adult performance test

We counted the number of eggs produced over 48 hours as an estimate of performance of first generation hybrids. Buckets were checked daily for emergence of F_1 adults. On the day of emergence, adults were sexed and mating pairs were placed into a plastic container (0.4L) with a paper towel lining the bottom and food. The containers were checked daily for eggs. The number of eggs produced was counted for 48 hours after the first eggs were laid. After this time, F_1 adults were removed and killed by freezing.

2.4 F_2 larval performance test

We measured percent hatching of all eggs laid in the first 48 hours, development time (in days), and adult mass (mg) attained by each F_2 larva. Upon emergence, the date was recorded as well as the number of eggs that successfully hatched. Counting eggs is challenging due to the three-dimensional nature of the egg clutches. Following (Bean *et al.* 2013b), to ensure accuracy we also counted the number of larvae and compared this with the number of eggs. If the number of eggs was less than the number of larvae, we used the number of larvae as the total number of eggs produced. If the number of eggs was greater than the number of larvae, we conducted a recount of the clutch. Out of the hatched larvae from each mating pair, up to five were randomly chosen and allowed to develop individually.

Larvae were maintained in small plastic cups (0.4L) and given fresh saltcedar with its stem in a water-filled 1.5mL eppendorf tube each day. A paper-towel lined the bottom of each cup. When the larvae reached their last stage of development, 2 cm of sand was placed in each cup to provide conditions favorable for pupation. All larvae were maintained in incubators under a light regime of 16:8 (L:D) and 27°C/16°C, and rotated every other day to standardize environmental effects.

2.5 F_2 adult preference test

We conducted a host preference test to determine if hybridization affected host preference for the non-target species, athel, presenting beetles with a choice between saltcedar and athel. Athel is an ornamental that is found at more southern latitudes in the US and is considered invasive in the southwestern U.S. (Gaskin & Shafroth 2005). *Tamarix* hybrids of *T. ramosissima* and *T. chinensis* (saltcedar) and are considered the preferred field host of *Diorhabda*. Previous host testing showed that the three *Diorhabda* species used in this study can survive as well on athel as on saltcedar, will oviposit on either saltcedar or athel under laboratory

no-choice conditions, and showed an inconsistent preference for saltcedar under choice conditions (Milbrath & DeLoach 2006a; Milbrath & DeLoach 2006b). However, saltcedar is preferred at field sites (Moran *et al.* 2009), and the intrinsic rate of increase of beetle populations is reduced on athel due to smaller egg mass size and a delayed start to oviposition (Milbrath & DeLoach 2006b).

Between 24 and 48 hours after emergence, we sexed and weighed the F_2 adults. The beetles were placed in a plastic tub (3L) with two eppendorf tubes containing equal amounts of either athel or saltcedar. Each beetle was placed in the middle of the tub, with both plants placed equidistantly at 10 cm from the center. The beetle remained in the plastic tub for 24 hours, at which time the amount of frass under each plant was weighed to the nearest 0.1 g (DeLoach *et al.* 2003).

2.6 F_3 larval performance test on two different hosts

We measured F_3 larval performance on athel and saltcedar. After the host-choice test, mating pairs were formed with F_2 adults from the same cross. All F_2 adults were given saltcedar foliage to feed on regardless of what they chose as their host in the adult preference test. They were placed in the same plastic dish as previously described and allowed to mate and oviposit. The date of first oviposition, the number of eggs laid in 48 hours, and the percent hatching was recorded. Larvae from each mating pair were split and a maximum of five larvae were placed in a plastic dish with either athel or saltcedar. We measured development time to adult and adult mass.

2.7 Statistical analysis

Our interests center on comparing the fitness of hybrids to their parental species. Thus, each analysis was done separately for each of the 7 pairs of parental species and their respective

two hybrid crosses (male/female reciprocal). All statistical analysis was conducted using R version 3.3.2 (R_Core_Team 2016). For the first generation, we analyzed differences in the total number of eggs produced between hybrids and parental species using a standard linear model. The number of eggs was log-transformed to meet the assumption of homogeneity of variance. Percent hatching was analyzed as a proportion of hatched eggs compared to the total number of eggs using a generalized linear model with quasibionomial error distribution. Cross was the only fixed effect for number of eggs produced and percent hatching in the first-generation analysis. For the second and third generations, we quantified the development time from egg to adult (days), adult mass (mg), and host choice. For development time and adult weight, we used linear mixed-effects models through the lme4 package (Bates *et al.* 2015) with cross, sex and their interaction as fixed effects, and family as a random effect. For host choice with a binary response (saltcedar or athel), we used a generalized linear mixed effects model with a binomial error distribution. For the third generation, we also included the random effect of cup nested within family for development time and adult weight.

3. Results

3.1 Egg count, percent hatching

Hybridization did not significantly affect the number of F_1 eggs laid in 48 hours for any of the crosses (Tables 1-3). Cross had a marginally significant influence on percent of eggs that hatched with the $E \times E$ and $E_f \times S_m$ cross producing slightly fewer viable eggs than the other crosses ($F_{3,37} = 2.82$, $P = 0.052$, Table 2). In the F_2 , only for the $S \times C$ cross was there a significant effect of cross on the number of eggs laid in 48 hours, where hybrids produced significantly more eggs than either parental species ($F_{3,39} = 2.97$, $P = 0.044$, Table 1, Figure 1).

Cross did not affect the percentage of eggs hatched for any crosses in the second generation (Table 1-3).

3.2 Development time, adult mass

In the S × C crosses, females were larger (effect of sex: $\chi^2 = 12.98$, df = 1, $P < 0.001$, effect of cross: $\chi^2 = 16.39$, df = 3, $P < 0.001$) and hybrids developed faster (effect of sex: $\chi^2 = 9.93$, df = 1, $P = 0.002$, effect of cross: $\chi^2 = 20.60$, df = 3, $P < 0.001$) (Table 1, 4, Figure 2). For the S × E cross, there was a significant interaction between sex and cross in development time, in that males developed slower than females in the S × S line (interaction cross*sex: $\chi^2 = 8.90$, df = 3, $P = 0.031$, Tables 2, 4). We also found a significant effect of sex and cross on adult mass in the S × E cross, with females being overall larger (effect of sex: $\chi^2 = 7.28$, df = 1, $P = 0.007$, effect of cross: $\chi^2 = 17.50$, df = 3, $P < 0.001$, Tables 2, 4). There was no effect of sex or cross on development time or adult mass for the E × C cross, although overall, females tended to be larger.

For the third generation, we were only able to investigate the effects of hybridization for the S × C cross (*D. sublineata* × *D. carinata*) due to limitations in the availability of our host plants. Development time was significantly affected by host plant and by cross, with both parents and hybrids developing slower on athel (effect of cross: $\chi^2 = 9.74$, df = 4, $P = 0.029$; effect of plant: $\chi^2 = 10.22$, df = 1, $P = 0.001$, Tables 1, 5, Figure 3). While there is a trend for hybrids to develop slower than parents regardless of host plant, there was no significant decrease in development time in hybrids compared to parents (effect of cross: Hybrids develop slower than the parents regardless of host plant, yet contrasts between crosses were not significant. Adult weight was not affected by hybridization, however females were larger regardless of cross (effect of sex: $\chi^2 = 10.124$, df = 1, $P = 0.001$, Tables 1, 5).

3.3 Host choice

We tested the host preference of individuals from all crosses in the second generation.

Due to limitations in our host plant resources and, because of differences seen in the second generation, we also examined host preference for the $S \times C$ cross in the third generation. Sex did not affect host choice for any of the crosses in the second generation (Tables 1-3). Cross significantly affected host choice in the $S \times C$ cross (effect of cross: $\chi^2 = 9.87$, $df = 3$, $P = 0.031$) and $S \times E$ cross (effect of cross ($\chi^2 = 9.23$, $df = 3$, $P = 0.026$), whereas there was no difference in host preference between hybrids and their parents in the $E \times C$ cross (Tables 1-4, Figure 4). There was no effect of hybridization on host preference in the $S \times C$ cross in the third generation (effect of cross: $\chi^2 = 1.163$, $df = 3$, $P = 0.7619$, Tables 1, 5).

4. Discussion

In introduced species, the effects of hybridization can influence local adaption and determine the fate of colonization success and establishment (Rius & Darling 2014). Introduced biological control agents undergo similar pressures as newly invading species, and understanding the mechanisms behind population growth and establishment are crucial to the implementation of successful biological control. In this study, we investigated the effects of hybridization on various life history traits and host preference for three different species of the biological control agent *Diorhabda*. We confirmed that all three species were reproductively compatible (Bean *et al.* 2013b), and that found that reciprocal crosses produced viable offspring through at least two generations. Life history traits beyond the production of viable eggs were either unchanged or improved with hybridization when compared to the parental species. These results support the hypothesis that these species have not experienced reproductive isolation for long enough to allow the evolution of genetic incompatibilities.

Hybridization can have positive, neutral, or negative effects on fitness. These effects depend on the genetic distance between mixing populations and the interactions between genes and environment. Hybrid vigor is commonly seen in the first generation of admixture between genetically distinct populations, and is typically thought to be due to masking of deleterious alleles rather than overdominance (Szulkin *et al.* 2010), whereas hybrid breakdown is commonly seen in the second or later generations due to recombination of the parental genes, allowing for the possibility of deleterious allele combinations (heterozygote disadvantage) (Dobzhansky 1950; Edmands 2002). In our study, there was no difference between parents and their hybrid offspring in fecundity or percent hatching in the first generation in any cross. Previous molecular work done by Bean *et al* (2013b) showed that while all four *Diorhabda* species separated into their own clades, the three species examined here were likely more closely related to each other than to the congeneric *D. carinulata*. It is possible that these species are not genetically distinct enough to be detrimentally affected by hybridization. However, the beetles used in our study had been lab reared for varying amounts of time (at least ten generations), and may have become inbred or lost variation via drift, both of which could reduce fitness. Thus, an alternative explanation is that positive effects of crossing, via masking of deleterious mutations could have balanced out potentially negative effects of hybridization, leading to zero, or close to zero, net change in life history traits. The masking of deleterious mutations can persist for many generations (Frankham 2016; Hedrick & Garcia-Dorado 2016), and thus further study investigating the effects of hybridization for more than three generations would increase our understanding.

Our results show that some of our crosses benefited greatly from hybridization in fecundity and development time in the second generation, and thus we see no evidence of hybrid

breakdown. $S \times C$ crosses produced 67% more eggs and developed approximately 7 days shorter than the parental species. The $E \times C$ cross exhibit the same trend, although this was only marginally significant. Other crosses showed no effect of hybridization, and none of our crosses suffered a fitness cost. In the $S \times C$ cross, where we could examine a third generation, we saw no effect of hybridization on fecundity, but we did see a trend that hybrids were developing slower on both host plants. For this analysis, our sample size was lower than for the previous generations, and so further work is necessary to determine if development time slowed because of hybridization.

Changes in host specificity in a released agent are one of the most concerning issues to scientists studying biological control (Van Klinken & Edwards 2002; Brodeur 2012; McEvoy *et al.* 2012). Our results show that host specificity can indeed be affected by hybridization, and that the phenotype can vary depending on the maternal or paternal species. In the $S \times C$ crosses, host preference of the hybrid followed the preference of the maternal species, whereas in the $S \times E$ cross, hybrids showed no preference for either host plant where the parents both showed a strong preference for the target host. Host specificity depends upon a suite of traits, such as behavior, morphology, and life-history strategies and as such is highly constrained (Zwolfer & Harris 1971; Giebink *et al.* 1984; Chang *et al.* 1987). Even so, in more generalist species than are typically used for biological control, host use has been shown to have a genetic basis, and can thus vary between individuals and populations (Singer & Parmesan 1993; Funk 1998). In our study, the inherited pattern for host use depended not only on the cross, but the preference of the maternal species. A growing body of literature suggests that for herbivorous insect species, mothers have been shown to influence host use (Amarillo-Suarez & Fox 2006; Egan *et al.* 2011; Cahenzli & Erhardt 2013). Egan *et al.* (2011) specifically demonstrated that host-use and

performance are traits with sex-linked maternal influence. Consequently, the pattern of host specificity in hybrid crosses can be hard to predict since it will depend not only on the amount of genetic variation across a suite of traits, but also parental influence.

Using hybridization in biological control presents unique challenges. On one hand, increased genetic variation, potentially from hybridization, can buffer introduced populations against adaptive challenges and thus increase the probability of establishment and effective control (Hopper *et al.* 1993). On the other, the genetic admixture of previously isolated populations might give rise to new phenotypes that are less desirable, such as a change in host specificity (Hoffmann *et al.* 2002; Mathenge *et al.* 2010). Our results demonstrate that while some crosses benefit from hybridization in terms of development time and fecundity, differences in host specificity due to hybridization is of concern.

Figure 1: Number of eggs produced in first 48 hours by *D. carinata* ($C \times C$), *D. sublineata* ($S \times S$), and their hybrids in the second generation. Cross significantly affected egg production, with hybrids producing more than either parental species. Bars represent 95% confidence intervals.

Figure 2: Development time from hatching until adult for each sex of *D. carinata* (C × C), *D. sublineata* (S × S) and their hybrids in the second generation. Cross significantly affected development time, with hybrids developing faster than either parental species. Grey and black lines represent females and males, respectively. Bars represent 95% confidence intervals.

Figure 3: Development time on athel (non-target) and saltcedar (target) for *D. carinata* ($C \times C$), *D. sublineata* ($S \times S$), and their hybrids after three generations of hybridization. $S_f \times C_m$ are long close dashes, $C_f \times S_m$ small dashes, $S \times S$ long spaced dashes, and $C \times C$ solid line. Host plant significantly affected development time with beetles developing slower on the non-target host.

Figure 4: Preference for saltcedar (target) over athel (non-target) across all parental species (*D. carinata* “C”, *D. sublineata* “S”, *D. elongata* “E”) and their hybrids in the second generation of hybridization. Hybridization significantly affected host preference for the S × C and S × E crosses. Error bars represent 95% confidence intervals.

Table 1: Trait means (95% CI) for each generation of the *D. sublineata* by *D. carinata* cross. Letters indicate significant differences between crosses.

Gen	Trait		Cross				F value	P
			C × C	C _r × S _m	S _r × C _m	S × S		
1	48 hour egg count	N	25.53 (20.08, 32.46) ^a	21.54 (17.13, 27.08) ^a	30.14 (22.38, 40.59) ^a	20.25 (15.73, 26.08) ^a	F _{3, 69} = 1.733	0.1682
			20	22	13	18		
	Proportion hatching		91.4% (66.0, 99.0) ^a	90.9% (68.0, 98.6) ^a	91.8% (59.9, 99.6) ^a	94.3% (67.2, 99.8) ^a	F _{3, 60} = 0.8425	0.476
			17	20	12	15		
2	Dev time (days) (male)		43.6 (41.2, 45.9) ^a	38.2 (36.3, 40.1) ^b	39.6 (37.4, 41.9) ^{ab}	43.5 (41.0, 45.9) ^a	See Table 4 for statistical results	
		N	14	31	18	17		
	Dev time (days) (female)		44.1 (41.6, 46.7) ^a	36.2 (34.3, 38.2) ^b	38.7 (36.2, 41.3) ^b	39.8 (37.6, 41.9) ^b		
		N	11	22	12	13		
	Adult mass (mg) (male)		21.0 (19.3, 22.8) ^{ab}	22.6 (21.2, 24.0) ^a	19.4 (18.0, 21.0) ^b	18.3 (16.9, 19.8) ^b		
		N	14	31	18	17		
	Adult mass (mg) (female)		22.5 (18.5, 27.5) ^{ab}	25.8 (22.2, 29.9) ^a	20.6 (16.9, 25.1) ^{ab}	19.2 (16.2, 22.8) ^b		
		N	11	22	12	13		
	Preference for saltcedar		54.2% (34.6, 72.5) ^a	53.8% (40.3, 66.8) ^a	76.9% (57.2, 89.2) ^a	80.0% (62.1, 90.7) ^a		
		N	24	52	26	30		
48 hour egg count		25.0 (16.1, 38.8) ^a	41.7 (32.1, 54.1) ^a	45.6 (33.0, 63.1) ^a	27.0 (18.9, 38.7) ^a	F_{3, 39} = 2.9659	0.0437	
	N	6	17	11	9			
Proportion hatching		93.2% (26.5, 99.1) ^a	94.7% (67.7, 99.8) ^a	94.2% (50.0, 99.9) ^a	79.7% (29.7, 98.9) ^a	F _{3, 26} = 1.3756	0.2722	
		3	15	7	5			
3	Dev time (days) saltcedar		31.9 (29.7, 34.3) ^a	34.6 (33.3, 35.9) ^a	35.3, (33.2, 37.6) ^a	32.4 (30.5, 34.3) ^a	See Table 5 for statistical results	
		N	9	38	9	12		
	Dev time (days) athel		34.1 (31.6, 36.9) ^a	37.0 (35.4, 38.6) ^a	37.7 (35.4, 40.2) ^a	34.6 (32.5, 36.9) ^a		
		N	3	24	11	7		
	Adult mass (mg) (male)		19.5 (15.9, 23.9) ^a	19.7 (17.7, 21.8) ^a	18.6 (15.8, 22.0) ^a	16.7 (14.1, 19.7) ^a		
		N	9	38	9	12		
	Adult mass (mg) (female)		22.5 (18.4, 27.6) ^a	22.8 (20.6, 25.3) ^a	21.6 (18.1, 25.6) ^a	19.3 (16.4, 22.7) ^a		
		N	3	24	11	7		
	Preference for saltcedar		69.0% (37.2, 89.3) ^a	61.8% (47.3, 74.5) ^a	50.2% (30.3, 75.7) ^a	71.4% (45.6, 88.1) ^a		
		12	62	20	19			

484
485

Table 2: Trait means (95% CI) for each generation of the *D. sublineata* by *D. elongata* cross. Letters indicate significant differences between crosses.

Gen	Trait		Cross				F value	P
			S × S	S _f × E _m	E _f × S _m	E × E		
1	48 hour egg count		20.25 (15.45, 26.55) ^a	22.46 (16.12, 31.28) ^a	26.24 (18.84, 36.55) ^a	18.04 (10.16, 32.03) ^a	F _{3, 42} = 0.4807	0.5706
		N	18	12	12	4		
	Proportion hatching		94.3% (67.2, 99.8) ^a	90.6% (58.5, 99.3) ^a	87.62% (51.6, 98.9) ^a	87.8% (30.2, 99.9) ^a	F_{3, 37} = 2.8245	0.05193
		N	15	12	10	4		
2	Dev time (days) (male)		43.4 (40.8, 46.2) ^a	37.5 (34.7, 40.5) ^b	39.5 (35.5, 44.0) ^{ab}	42.2 (38.6, 46.3) ^{ab}	See Table 4 for statistical results	
		N	17	9	5	13		
	Dev time (days) (female)		39.8 (37.8, 42.0) ^a	38.0 (35.2, 41.0) ^a	42.4 (39.7, 45.3) ^a	43.3 (39.2, 47.7) ^a		
		N	13	5	10	3		
	Adult mass (mg) (male)		18.1 (16.2, 20.2) ^a	13.1 (11.5, 15.1) ^b	17.1 (14.1, 20.8) ^{ab}	17.0 (14.6, 19.9) ^{ab}		
		N	17	9	5	13		
	Adult mass (mg) (female)		19.1 (16.7, 21.9) ^a	15.6 (12.8, 19.0) ^a	21.5 (18.1, 25.5) ^a	18.8 (14.5, 24.3) ^a		
		N	13	5	10	3		
	Preference for <i>Tamarix</i> spp		80.0% (62.1, 90.7) ^a	38.5% (17.0, 65.6) ^b	50.0% (26.0, 74.0) ^{ab}	80.0 (53.0, 93.4) ^{ab}		
		N	30	13	14	15		
	48 hour egg count		27.0 (17.7, 41.2) ^a	27.0 (14.4, 50.9) ^a	45.1 (21.8, 93.6) ^a	28.0 (11.4, 68.3) ^a	F _{3, 14} = 0.622	0.6124
		N	9	4	3	2		
	Proportion hatching		79.7% (29.7, 98.9) ^a	70.5% (14.2, 98.9) ^a	96.6% (18.1, 96.7) ^a	94.1% (4.3, 92.5) ^a	F _{3, 7} = 0.9465	0.4682
		N	5	3	2	1		

Table 3: Trait means (95% CI) for each generation of the *D. elongata* by *D. carinata* cross. Letters indicate significant differences between crosses.

Gen	Trait		Cross				F Value	P
			E × E	E _r × C _m	C _r × E _m	C × C		
1	48 hour egg count		18.04 (9.93,32.77) ^a	18.90 (14.37, 24.85) ^a	18.36 (13.96, 24.14) ^a	25.53 (19.55, 33.35) ^a	F _{3, 58} = 1.3855	0.2837
		N	4	19	19	20		
	Proportion hatching		87.8% (35.1, 99.4) ^a	86.8% (58.1, 99.7) ^a	90.7% (61.9, 99.1) ^a	91.4% (66.0, 99.0) ^a	F _{3, 42} = 0.7341	0.5376
		N	4	11	14	17		
2	Dev time (days) (male)		42.4 (38.4, 46.3) ^a	39.5 (36.6, 42.4) ^a	42.5 (39.7, 45.4) ^a	43.6 (41.1, 46.3.4) ^a	See Table 4 for statistical results	
		N	13	12	18	14		
	Dev time (days) (female)		43.33 (38.57, 48.09) ^a	39.13 (36.55, 41.71) ^a	40.0 (36.3, 43.7) ^a	44.0 (41.02, 47.17) ^a		
		N	3	14	5	11		
	Adult mass (mg) (male)		17.1 (14.9, 19.6) ^a	18.8 (16.7, 21.0) ^a	20.5 (18.4, 22.9) ^a	21.0 (19.0, 23.3) ^a		
		N	13	12	18	14		
	Adult mass (mg) (female)		18.8 (13.7, 25.8) ^a	22.1 (18.7, 26.1) ^a	19.7 (15.4, 25.2) ^a	22.7 (18.7, 27.5) ^a		
		N	3	14	5	11		
	Preference for <i>Tamarix</i> spp.		81.5% (50.8, 94.9) ^a	45.6% (26.2, 66.4) ^a	54.3% (31.9, 75.1) ^a	54.1% (32.8, 74.0) ^a		
		N	15	26	22	24		
	48 hour egg count		28.0 (15.0, 52.0) ^a	35.6 (27.9, 45.4) ^a	50.1 (33.8, 74.1) ^a	25.0 (17.5, 35.8) ^a	F _{3, 22} = 2.6441	0.07446
		N	2	13	5	6		
	Proportion hatching		94.1% (2.0, 97.2) ^a	74.6% (58.6, 99.9) ^a	93.0% (35.3, 99.3) ^a	93.2% (26.5, 99.1) ^a	F _{3, 14} = 1.5797	0.2387
		N	1	10	4	3		

516

Table 4: Results from generalized linear mixed-effects models for the second generation of hybridization for all crosses.

Trait	S × E Cross			Random effects			
	Cross	Sex	Cross*sex	Family		Residual	
	χ^2 , (df), <i>P</i>	χ^2 , (df), <i>P</i>	χ^2 , (df), <i>P</i>	Variance	Std dev	Variance	Std dev
Dev time	10.82, (3), 0.013	0.47, (1), 0.493	8.90, (3), 0.031	0.006174	0.07857	0.001771	0.04208
Adult mass	17.50, (3), <0.001	7.28, (1), 0.007	2.56, (3), 0.464	0.01266	0.1125	0.02347	0.1532
Preference for saltcedar	9.23, (3), 0.026	0.98, (1), 0.322	0.99, (3), 0.804	0	0	0	0
C × E Cross							
Dev time	5.99, (3), 0.112	0.08 (1), 0.7837	5.79, (3), 0.122	0.008246	0.09081	0.001845	0.04295
Adult mass	5.68, (3), 0.128	3.47, (1), 0.0626	2.65, (3), 0.449	0.01058	0.1029	0.0348	0.1868
Preference for saltcedar	3.66, (3), 0.300	0.89, (1), 0.3464	3.69, (3), 0.297	0.6548	0.8092	0	0
S × C Cross							
Dev time	20.60, (3), <0.001	9.934, (1) 0.002	5.82, (3), 0.120	9.787	3.128	4.205	2.051
Adult mass	16.39, (3), <0.001	12.982, (1), <0.001	2.73, (3), 0.435	0.02041	0.1429	0.02266	0.1505
Preference for saltcedar	9.87, (3), 0.031	0.3953, (1), 0.5295	1.89, (3), 0.596	0	0	0	0

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

Table 5: Results from generalized linear mixed-effects models for the third generation of the *D. sublineata* by *D. carinata* cross.

Trait	S × C Cross					Random effects				
	Cross	Sex	Plant	Plant * Cross	Cup within family		Family		Residual	
	χ^2 , (df), <i>P</i>	χ^2 , (df), <i>P</i>	χ^2 , (df), <i>P</i>	χ^2 , (df), <i>P</i>	Variance	Std dev	Variance	Std dev	Variance	Std dev
Dev time	9.74, (3), 0.030	0.022, (1), 0.882	10.22, (1), 0.001	3.46, (3), 0.326	0.0027	0.052	0.0009	0.0303	0.00305	0.05531
Adult mass	4.16, (3), 0.244	9.74, (1), 0.021	1.39, (1), 0.239	2.67, (3), 0.449	0.02133	0.1461	0	0	0.04805	0.2192
Preference for saltcedar	1.16, (3), 0.762	0.12, (1), 0.728	1.00, (1), 0.317	0.30, (3), 0.960	0	0	0	0	0	0

538

References

1. Amarillo-Suarez, A.R. & Fox, C.W. (2006). Population differences in host use by a seed-beetle: local adaptation, phenotypic plasticity and maternal effects. *Oecologia*, 150, 247-258.
2. Anderson, E. & Stebbins, G.L. (1954). Hybridization as an Evolutionary Stimulus. *Evolution*, 8, 378-388.
3. Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, 67, 1-48.
4. Bean, D., Dudley, T. & Hultine, K. (2013a). Bring on the Beetles! . In: *Tamarix: A Case Study of Ecological Change in the American West* (eds. Sher, A & Quigley, MF). Oxford University Press Oxford.
5. Bean, D.W., Kazmer, D.J., Gardner, K., Thompson, D.C., Reynolds, B., Keller, J.C. *et al.* (2013b). Molecular genetic and hybridization studies of *Diorhabda* spp. released for biological control of tamarix. *Invasive Plant Science and Management*, 6, 1-15.
6. Benvenuto, C., Cheyppé-Buchmann, S., Bermond, G., Ris, N. & Fauvergue, X. (2012). Intraspecific hybridization, life history strategies and potential invasion success in a parasitoid wasp. *Evolutionary Ecology*, 26, 1311-1329.
7. Brodeur, J. (2012). Host specificity in biological control: Insights from opportunistic pathogens. *Evolutionary Applications*, 5, 470-480.
8. Burton, R.S., Rawson, P.D. & Edmands, S. (1999). Genetic architecture of physiological phenotypes: Empirical evidence for coadapted gene complexes. *Am. Zool.*, 39, 451-462.
9. Cahenzli, F. & Erhardt, A. (2013). Transgenerational acclimatization in an herbivore - host plant relationship. *Proc. R. Soc. B-Biol. Sci.*, 280, 8.
10. Chang, N.T., Lynch, R.E., Slansky, F.A., Wiseman, B.R. & Habeck, D.H. (1987). QUANTITATIVE UTILIZATION OF SELECTED GRASSES BY FALL ARMYWORM LARVAE. *Entomologia Experimentalis Et Applicata*, 45, 29-35.
11. DeLoach, C.J., Lewis, P.A., Herr, J.C., Carruthers, R.I., Tracy, J.L. & Johnson, J. (2003). Host specificity of the leaf beetle, *Diorhabda elongata deserticola* (Coleoptera: Chrysomelidae) from Asia, a biological control agent for saltcedars (*Tamarix*: Tamaricaceae) in the Western United States. *Biological Control*, 27, 117-147.
- 12.

578 Dlugosch, K.M. & Parker, I.M. (2008). Invading populations of an ornamental shrub show rapid life
579 history evolution despite genetic bottlenecks. *Ecology Letters*, 11, 701-709.
580 13.

581 Dobzhansky, T. (1950). GENETICS OF NATURAL POPULATIONS .19. ORIGIN OF HETEROSIS THROUGH
582 NATURAL SELECTION IN POPULATIONS OF DROSOPHILA-PSEUDOBBSCURA. *Genetics*, 35, 288-
583 302.
584 14.

585 Edmands, S. (2002). Does parental divergence predict reproductive compatibility? *Trends Ecol. Evol.*, 17,
586 520-527.
587 15.

588 Egan, S.P., Janson, E.M., Brown, C.G. & Funk, D.J. (2011). Postmating isolation and genetically variable
589 host use in ecologically divergent host forms of *Neochlamisus bebbianae* leaf beetles. *J. Evol.*
590 *Biol.*, 24, 2217-2229.
591 16.

592 Ellstrand, N.C. & Schierenbeck, K.A. (2000). Hybridization as a stimulus for the evolution of invasiveness
593 in plants? *Proceedings of the National Academy of Sciences of the United States of America*, 97,
594 7043-7050.
595 17.

596 Fisher, R.A. (1930). *The genetical theory of natural selection*. Clarendon Press, Oxford, England.
597 18.

598 Frankham, R. (2015). Genetic rescue of small inbred populations: meta-analysis reveals large and
599 consistent benefits of gene flow. *Molecular Ecology*, 24, 2610-2618.
600 19.

601 Frankham, R. (2016). Genetic rescue benefits persist to at least the F3 generation, based on a meta-
602 analysis. *Biol. Conserv.*, 195, 33-36.
603 20.

604 Funk, D.J. (1998). Isolating a role for natural selection in speciation: Host adaptation and sexual isolation
605 in *Neochlamisus bebbianae* leaf beetles. *Evolution*, 52, 1744-1759.
606 21.

607 Gaskin, J.F. & Schaal, B.A. (2002). Hybrid *Tamarix* widespread in U.S. invasion and undetected in native
608 Asian range. *Proceedings of the National Academy of Sciences of the United States of America*,
609 99, 11256-11259.
610 22.

611 Gaskin, J.F. & Shaforth, P.B. (2005). HYBRIDIZATION OF TAMARIX RAMOSISSIMA AND T. CHINENSIS
612 (SALT CEDARS) WITH T. APHYLLA (ATHEL) (TAMARICACEAE) IN THE SOUTHWESTERN USA
613 DETERMINED FROM DNA SEQUENCE DATA. *Madroño*, 52, 1-10.
614 23.

615 Giebink, B.L., Scriber, J.M. & Wedberg, J.L. (1984). BIOLOGY AND PHENOLOGY OF THE HOP-VINE BORER,
616 HYDRAECIA-IMMANIS GUENEE, AND DETECTION OF THE POTATO STEM BORER, HYDRAECIA-

617 MICACEA (ESPER) (LEPIDOPTERA, NOCTUIDAE), IN WISCONSIN. *Environmental Entomology*, 13,
618 1216-1224.
619 24.

620 Hedrick, P.W. & Garcia-Dorado, A. (2016). Understanding Inbreeding Depression, Purging, and Genetic
621 Rescue. *Trends Ecol. Evol.*, 31, 940-952.
622 25.

623 Henry, L.M., May, N., Acheampong, S., Gillespie, D.R. & Roitberg, B.D. (2010). Host-adapted parasitoids
624 in biological control: Does source matter? *Ecological Applications*, 20, 242-250.
625 26.

626 Hoffmann, J.H., Impson, F.A.C. & Volchansky, C.R. (2002). Biological control of cactus weeds:
627 implications of hybridization between control agent biotypes. *Journal of Applied Ecology*, 39,
628 900-908.
629 27.

630 Hopper, K.R., Roush, R.T. & Powell, W. (1993). MANAGEMENT OF GENETICS OF BIOLOGICAL-CONTROL
631 INTRODUCTIONS. *Annu. Rev. Entomol.*, 38, 27-51.
632 28.

633 Hufbauer, R.A., Szucs, M., Kasyon, E., Youngberg, C., Koontz, M.J., Richards, C. *et al.* (2015). Three types
634 of rescue can avert extinction in a changing environment. *Proceedings of the National Academy*
635 *of Sciences of the United States of America*, 112, 10557-10562.
636 29.

637 Laugier, G.J.M., Le Moguedec, G., Su, W., Tayeh, A., Soldati, L., Serrate, B. *et al.* (2016). Reduced
638 population size can induce quick evolution of inbreeding depression in the invasive ladybird
639 *Harmonia axyridis*. *Biological Invasions*, 18, 2871-2881.
640 30.

641 Lee, Y.W., Fishman, L., Kelly, J.K. & Willis, J.H. (2016). A Segregating Inversion Generates Fitness
642 Variation in Yellow Monkeyflower (*Mimulus guttatus*). *Genetics*, 202, 1473-+.
643 31.

644 Mallet, J. (2005). Hybridization as an invasion of the genome. *Trends Ecol. Evol.*, 20, 229-237.
645 32.

646 Mathenge, C.W., Holford, P., Hoffmann, J.H., Zimmermann, H.G., Spooner-Hart, R.N. & Beattie, G.A.C.
647 (2010). Hybridization between *Dactylopius tomentosus* (Hemiptera: Dactylopiidae) biotypes and
648 its effects on host specificity. *Bulletin of Entomological Research*, 100, 331-338.
649 33.

650 McEvoy, P.B., Higgs, K.M., Coombs, E.M., Karaçetin, E. & Ann Starcevich, L. (2012). Evolving while
651 invading: Rapid adaptive evolution in juvenile development time for a biological control
652 organism colonizing a high-elevation environment. *Evolutionary Applications*, 5, 524-536.
653 34.

654 Michels, G.J., Royer, T.A., Jones, E.N., Lange, R.A., Bynum, E.D., Ruthven, D.C. *et al.* (2013). New
655 Establishment and County Records for *Diorhabda* spp. (Coleoptera: Chrysomelidae) and

656 Coniatus splendidulus (Coleoptera: Curculionidae) in the Texas Panhandle and Western
657 Oklahoma. *Southwestern Entomologist*, 38, 173-181.
658 35.

659 Milbrath, L.R. & DeLoach, C.J. (2006a). Acceptability and Suitability of Athel, Tamarix aphylla, to the Leaf
660 Beetle Diorhabda elongata (Coleoptera: Chrysomelidae), a Biological Control Agent of Saltcedar
661 (Tamarix spp.). *Environmental Entomology*, 35, 1379-1389.
662 36.

663 Milbrath, L.R. & DeLoach, C.J. (2006b). Host specificity of different populations of the leaf beetle
664 Diorhabda elongata (Coleoptera: Chrysomelidae), a biological control agent of saltcedar
665 (Tamarix spp.). *Biological Control*, 36, 32-48.
666 37.

667 Moran, P.J., DeLoach, C.J., Dudley, T.L. & Sanabria, J. (2009). Open field host selection and behavior by
668 tamarisk beetles (Diorhabda spp.) (Coleoptera: Chrysomelidae) in biological control of exotic
669 saltcedars (Tamarix spp.) and risks to non-target athel (T. aphylla) and native Frankenia spp.
670 *Biological Control*, 50, 243-261.
671 38.

672 Payseur, B.A. & Rieseberg, L.H. (2016). A genomic perspective on hybridization and speciation.
673 *Molecular Ecology*, 25, 2337-2360.
674 39.

675 R_Core_Team (2016). R: A language and environment for statistical computing. R Foundation for
676 Statistical Computing. Vienna, Austria.
677 40.

678 Rieseberg, L.H., Archer, M.A. & Wayne, R.K. (1999). Transgressive segregation, adaptation and
679 speciation. *Heredity*, 83, 363-372.
680 41.

681 Rius, M. & Darling, J.A. (2014). How important is intraspecific genetic admixture to the success of
682 colonising populations? *Trends Ecol. Evol.*, 29, 233-242.
683 42.

684 Seastedt, T.R. (2015). Biological control of invasive plant species: a reassessment for the Anthropocene.
685 *New Phytologist*, 205, 490-502.
686 43.

687 Singer, M.C. & Parmesan, C. (1993). SOURCES OF VARIATIONS IN PATTERNS OF PLANT INSECT
688 ASSOCIATION. *Nature*, 361, 251-253.
689 44.

690 Szucs, M., Eigenbrode, S.D., Schwarzländer, M. & Schaffner, U. (2012). Hybrid vigor in the biological
691 control agent, Longitarsus jacobaeae. *Evolutionary Applications*, 5, 489-497.
692 45.

693 Szulkin, M., Bierne, N. & David, P. (2010). HETEROZYGOSITY-FITNESS CORRELATIONS: A TIME FOR
694 REAPPRAISAL. *Evolution*, 64, 1202-1217.
695 46.

696 Tracy, J.L. & Robbins, T.O. (2009). Taxonomic revision and biogeography of the Tamarix-feeding
697 *Diorhabda elongata* (Brulle, 1832) species group (Coleoptera: Chrysomelidae: Galerucinae:
698 Galerucini) and analysis of their potential in biological control of Tamarisk. *Zootaxa*, 1-152.
699 47.

700 Van Driesche, R.G., Carruthers, R.I., Center, T., Hoddle, M.S., Hough-Goldstein, J., Morin, L. *et al.* (2010).
701 Classical biological control for the protection of natural ecosystems. *Biological Control*, 54, S2-
702 S33.
703 48.

704 Van Klinken, R.D. & Edwards, O.R. (2002). Is host-specificity of weed biological control agents likely to
705 evolve rapidly following establishment? *Ecology Letters*, 5, 590-596.
706 49.

707 Zwolfer, H. & Harris, P. (1971). HOST SPECIFICITY DETERMINATION OF INSECTS FOR BIOLOGICAL
708 CONTROL OF WEEDS. *Annu. Rev. Entomol.*, 16, 159-&.

709