Genome-wide meta-analysis of cognitive empathy: heritability, and correlates with sex, neuropsychiatric conditions and brain anatomy

Varun Warriar^{1,16}, Katrina Grasby², Florina Uzefovsky¹, Roberto Toro^{3,4,5}, Paula Smith¹, Bhismadev Chakrabarti⁶, Jyoti Khadake⁷, Eleanor Mawbey-Adamson⁷, Nadia Litterman⁸, Jouke-Jan Hottenga^{9,10,11}, Gitta Lubke^{9,12}, Dorret I Boomsma⁹, Nicholas G Martin², Peter K Hatemi¹³, Sarah E Medland², David A Hinds⁸, Thomas Bourgeron^{3,4,5,15}, and Simon Baron-Cohen^{1,14,15,16}

¹Autism Research Centre, Department of Psychiatry, University of Cambridge, Cambridgeshire, United Kingdom

²QIMR Berghofer Medical Research Institute, Brisbane, Australia.

³CNRS UMR 3571: Genes, Synapses and Cognition, Institut Pasteur, Paris, France

⁴Université Paris Diderot, Sorbonne Paris Cité, Human Genetics and Cognitive Functions, Paris, France

⁵Institut Pasteur, 5-28 Rue du Dr Roux, 75015 Paris, France

⁶Centre for Integrative Neuroscience and Neurodynamics, School of Psychology and Clinical Language Sciences, University of Reading, Reading, United Kingdom

⁷NIHR Cambridge BioResource, Cambridge University and Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

⁸23andMe Inc., 899 West Evelyn Ave, Mountain View, California 94041, USA

⁹Department of Biological Psychology, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

¹⁰EMGO+ Institute for Health and Care Research, VU Medical Center, Amsterdam, the Netherlands.

¹¹Neuroscience Campus Amsterdam, Vrije Universiteit, Amsterdam, the Netherlands.

¹²Department of Psychology, University of Notre Dame, United States

¹³Political Science, Microbiology and Biochemistry, Pennsylvania State University

¹⁴CLASS Clinic, Cambridgeshire and Peterborough NHS Foundation Trust (CPFT), Cambridgeshire, United Kingdom

¹⁵Joint senior authors.

¹⁶ **Corresponding authors:** Varun Warriar (vw260@medschl.cam.ac.uk) and Simon Baron-Cohen (sb205@cam.ac.uk). Autism Research Centre, Douglas House, 18B Trumpington Road, Cambridge, CB2 8AH, United Kingdom. Telephone: 0044 (0) 1223 746057.

We conducted a genome-wide meta-analysis of cognitive empathy using the ‘Reading the Mind in the Eyes’ Test (Eyes Test) in 88,056 research volunteers of European Ancestry (44,574 females and 43,482 males) from 23andMe Inc., and an additional 1,497 research volunteers of European Ancestry (891 females and 606 males) from the Brisbane Longitudinal Twin Study (BLTS). We confirmed a female advantage on the Eyes Test (Cohen’s $d = 0.21$, $P < 2.2 \times 10^{-16}$), and identified a locus in 3p26.1 that is associated with scores on the Eyes Test in females ($rs7641347$, $P_{meta} = 1.58 \times 10^{-8}$). Common single nucleotide polymorphisms (SNPs) explained 5.8% (95% CI: 0.45 – 0.72; $P = 1.00 \times 10^{-17}$) of the total trait variance in both sexes, and we identified a twin heritability of 0.28 (95% CI: 0.13–0.42). Finally, we identified significant genetic correlation between the Eyes Test and anorexia nervosa, measures of empathy (the Empathy Quotient), openness (NEO-Five Factor Inventory), and different measures of educational attainment and cognitive aptitude, and show that the genetic determinants of volumes of the dorsal striatum (caudate nucleus and putamen) are positively correlated with the genetic determinants of performance on the Eyes Test.

Introduction

Cognitive empathy, defined as the ability to recognize what another person is thinking or feeling, and to predict their behaviour based on their mental states, is vital for interpersonal relationships, which in turn is a key contributor of one’s wellbeing. Cognitive empathy is distinct from affective empathy, defined as the drive to respond to another’s mental states with an appropriate emotion^{1,2}. Difficulties in cognitive empathy have been found in different psychiatric conditions, particularly autism³. The dissociation between cognitive and affective empathy (the latter often being intact in autism, for example) suggests these have independent biological mechanisms.

Differences in cognitive empathy have been identified in individuals with different psychiatric conditions including autism⁴, schizophrenia^{5,6}, and anorexia nervosa⁷. This includes both higher and lower cognitive empathy in comparison to neurotypical controls i.e. not empathizing enough, or empathizing too much, both of which can contribute to difficulties in social interactions and wellbeing⁸. However, little is known about the genetic correlates of cognitive empathy. It is unclear to what extent, genetically, differences in cognitive empathy are a risk factor for developing various psychiatric conditions. Further, as previous studies have typically been conducted using self-report or performance-based tests,

results from these studies may be influenced by the characteristics of the test and/or factors associated with the psychiatric conditions themselves. In other words, from previous studies, it is difficult to tease apart the genetic and non-genetic contributions to strengths and difficulties in cognitive empathy, and how these relate to various psychiatric conditions.

Here, we investigated the genetic architecture of this aspect of social cognition using a well-validated test, the ‘Reading the Mind in the Eyes’ Test (Eyes Test). The Eyes Test is a brief online test where participants are shown photographs of the eye regions and have to identify the appropriate emotion or mental state they express². It has been widely used to investigate differences in cognitive empathy in several neuropsychiatric conditions including autism⁴, schizophrenia⁹, bipolar disorder¹⁰, anorexia nervosa¹¹, and major depressive disorder¹². We conducted a genome-wide association meta-analysis of cognitive empathy in more than 89,000 individuals of European ancestry, and investigated both SNP-based and twin-based heritabilities. We further conducted bivariate genetic regression analyses for psychiatric conditions, psychological traits, and brain volumes. We finally conducted gene based enrichment analysis and investigate potential genetic sources of sex differences.

Methods

Participants

23andMe : Research participants were customers of 23andMe, Inc., and have been described in detail elsewhere^{13,14}. All participants completed an online version of the ‘Reading the Mind in the Eyes’ test (Eyes Test)² online on the 23andMe research participant website (36 items). In total, 88,056 participants (44,574 females and 43,482 males) of European ancestry completed the Eyes Test and were genotyped. All participants provided informed consent and answered questions online according to 23andMe’s human subjects protocol, which was reviewed and approved by Ethical & Independent Review Services, an AAHRPP-accredited private institutional review board (<http://www.eandireview.com>). Only participants who were primarily of European ancestry (97% European Ancestry) were selected for the analysis using existing methods¹⁵. Unrelated individuals were selected using a segmental identity-by-descent algorithm¹⁶.

Brisbane Longitudinal Twin Study (BLTS): In addition, 1,497 participants (891 females and 606 males) of Caucasian ancestry with genotype data from the BLTS completed the short version (14 questions) of the Eyes Test online as part of a study on genetic and environmental

foundations of political and economic behaviors¹⁷. Participant ages ranged from 18 to 73 ($M = 37$, $SD = 14$). All participants provided online consent and the study was approved by the QIMR Berghofer Human Research Ethics Committee.

Measures

The ‘Reading the Mind in the Eyes’ Test is a brief questionnaire of cognitive empathy. Participants are shown scaled, black and white photographs of eye regions of actors and they have to choose the cognitive state portrayed from the four options provided. The Eyes Test has good test-retest reliability (eg: reliability of 0.833 in the Italian version¹⁸, and 0.63 in the Spanish version¹⁹)⁴, and scores are unimodally and near-normally distributed in the general population. In the BLTS dataset, there was a modest test-retest correlation of 0.47 in 259 participants who retok the test after a gap of nearly two years (Supplementary Note section 1). For each correct answer on the Eyes Test, participants score 1 point, so the scores ranged from 0 – 36 on the full version of the Eyes Test and 0 – 14 on the short version of the Eyes Test. Further details are provided in the Supplementary Note section 1.

Genotyping, imputation and quality control

23andMe cohort: DNA extraction, genotyping, imputation and initial quality control were completed by 23andMe, Inc. Participants provided saliva samples, and DNA extraction and genotyping were performed by the National Genetic Institute. All participants were genotyped using one of four different platforms (V1, V2, V3 and V4). Briefly, the V1 and V2 chips were based on the Illumina Human Hap550+ BeadChip (560,000 SNPs), the V3 on the Illumina OmniExpress+ Beadchip (950,000 SNPs). The V4 had a fully customized array of approximately 570,000 SNPs. Across all platforms, a total of 1,030,430 SNPs were genotyped. For this analysis, we included only participants with a call rate greater than 98.5, and SNPs that passed the Hardy-Weinberg Equilibrium Test at $P < 10^{-20}$ and had a genotype rate $> 90\%$. In addition, SNPs present only on platform V1, Chromosome Y and mitochondrial SNPs were excluded due to small sample sizes and unreliable genotype calling respectively. Using trio-data, where available, SNPs that failed the parent-offspring transmission test were also excluded. Imputation was performed using Minimac2²⁰ using the September 2013 release of the 1000 Genomes Phase 1 reference haplotypes phased using Beagle4²¹ (V3.3.1). Our analyses were restricted to SNPs that had a minor allele frequency of at least 1%, which left 9,955,952 SNPs after quality control. Genotyping, imputation, and preliminary quality control were performed by 23andMe.

BLTS cohort: The BLTS participants were genotyped on Illumina Human610-Quadv1_B or HumanCoreExome-12v1-0_C chips. These samples were genotyped in the context of a larger genome-wide association project. Genotype data was screened for genotyping quality (GenCall < 0.7 from the Human610-Quadv1_B chip), individual and SNP call rates (< 0.95 and < 0.99 for exome markers on the HumanCoreExome-12v1-0_C chip), Hardy-Weinberg Equilibrium ($P < 10^{-6}$), and MAF (< 0.01). The data were checked for non-European ancestry, pedigree, sex, and Mendelian errors. Data from the two different chips were separately phased using SHAPEIT2 and imputed to the 1000 Genomes reference panel (Phase 1 v3) using Minimac3. After imputation SNPs with a MAF < 0.05% were excluded, leaving 11,133,794 SNPs for analyses. We further excluded SNPs with imputation $r^2 < 0.6$ for meta-analysis.

Statistical analyses

Association analyses: Linear regression for the 23andMe cohort was performed for the Eyes Test scores using age, sex, and the first four ancestry principal components as covariates. For the sex-stratified analyses, sex was excluded as a covariate. The same regression model was used for the BLTS after accounting for relatedness using RAREMETALWORKER. Inverse variance weighted meta-analysis was performed using Metal²². Post meta-analysis, we excluded SNPs that were only genotyped in the BLTS cohort due to the small sample size, but included SNPs that were only genotyped in the 23andMe cohort. LD pruning was performed using Plink²³ with an r^2 of 0.1. We calculated the variance explained by each individual SNP²⁴ using the following formula:

$$R^2_{\text{glc}}/(1 - R^2_c) = (t^2/((n-k-1)+t^2)) \times 100$$

Where $R^2_{\text{glc}}/(1 - R^2_c)$ is the proportion of variance explained by the SNP after accounting for the effects of the covariates, t is the t -statistic of the regression co-efficient, k is the number of covariates, and n is the sample size. We corrected for winner's curse using an FDR based approach²⁵.

Heritability and genetic correlation: We used Linkage Disequilibrium Score regression coefficient (LDSR to calculate genomic inflation in the meta-analysis due to population stratification²⁶ (<https://github.com/bulik/ldsc>). Intercepts for the non-stratified GWAS was

1.01 (0.006), for the males-only GWAS was 1.006 (0.006), and for the females-only GWAS was 1.005 (0.006). SNP heritability and genetic correlation were calculated using LDSR. Difference in heritability between males and females was quantified using:

$$Z_{\text{diff}} = (h^2_{\text{males}} - h^2_{\text{females}}) / \text{sqrt}(SE^2_{\text{males}} + SE^2_{\text{females}})$$

Where Z_{diff} is the Z score for the difference in heritability for a trait, $(h^2_{\text{males}} - h^2_{\text{females}})$ is the difference SNP heritability estimate in males and females, and SE is the standard errors for the heritabilities. We calculated two-tailed P-values in R. We performed genetic correlation using summary GWAS data using LDSR. For all genetic correlation analyses, we used LD data from the North West European population as implemented in LDSR. Intercepts were not constrained in the analyses. We used a Benjamini-Hochberg FDR approach and a q-value threshold of 0.05 for determining statistical significance of the genetic correlations. Further details about the samples used are provided in the Supplementary Note section 2.

Twin Heritability: Twin heritability was estimated from 749 twin individuals (including 122 complete monozygotic pairs and 176 complete dizygotic pairs) in the BLTS using full information maximum likelihood in OpenMx²⁷ in R, which makes use of all available data. All twins completed the short version of the Eyes Test, and for those who completed the test twice only their first attempt was included in analyses. ADE, ACE, AE, CE, and E models were fit to the data and fit indices compared to determine the best-fitting model. Standardised variance components are reported from the best-fitting model, the AE model. Further details are given in the Supplementary Note section 4.

Gene-based analyses and sex difference analyses: We used MetaXcan²⁸ using tissue weights from the GTEx to perform gene-based analysis (<https://github.com/hakyimlab/MetaXcan>). MetaXcan uses summary statistics to perform gene based association analyses. It incorporates eQTL data from the GTEx consortium to infer gene level expression based on the summary GWAS statistics provided. This can be used to identify tissue-specific gene expression for the trait of interest. Here, we performed gene based analysis for the non-stratified GWAS meta-analysis for nine neural tissues: anterior cingulate cortex (BA24), caudate basal ganglia, cerebellar hemisphere, cerebellum, cortex, frontal cortex (BA9), hippocampus, hypothalamus, nucleus accumbens basal ganglia, and putamen basal ganglia, using gene-expression regression co-efficients for these tissues from the GTEx project. This is based on

tissues from 73 – 103 individuals. We chose neural tissues as we hypothesized that cognitive empathy is largely a neural phenotype. As MetaXcan predicts expression level from SNP information, we filtered out genes whose correlation with predicted models of expression was < 0.01 , as incorporated in MetaXcan. This step helps guard against false positives, by removing genes whose expressions are poorly predicted by the model. We used an FDR based correction to correct for all the tests run across all the tissues. Details of sex-difference analyses are provided in Supplementary Note section 3.

Data Availability

Summary level data may be requested from 23andMe, Inc. and received subject to 23andMe's standard data transfer agreement. We have also provided summary statistics for the first 10,000 LD pruned SNPs for the three GWAS analyses (males-only, females-only, and non-stratified) as supplementary data.

Results

Heritability

In collaboration with 23andMe, Inc. and the Brisbane Longitudinal Twin Study (BLTS) cohort, we conducted three separate genome-wide association study meta-analyses (GWASs) of the Eyes Test: a males-only GWAS ($n = 43,482$), a females-only GWAS ($n = 44,574$), and a non-stratified GWAS ($n = 88,056$). The study protocol is provided in Figure 1. All participants from the 23andMe cohort completed the full version of the Eyes Test online, comprising 36 questions (mean score = 27.47 ± 3.67), while participants from the BLTS cohort completed the short version of the Eyes Test (14 questions, mean = 8.85 ± 2.34) (Supplementary Note Section 1). Scores on the Eyes Test were significantly associated with age and sex in the 23andMe cohort (age: -0.026 ± 0.0007 ; $P < 2.2 \times 10^{-16}$, sex (females): 0.77 ± 0.02 ; $P < 2.2 \times 10^{-16}$). We used LDSC to calculate the heritability explained by all the SNPs in the HapMap3 with minor allele frequency $> 5\%$. We identified a significant narrow sense heritability of 5.8% (95% CI: 4.5% – 7.2%; $P = 1.00 \times 10^{-17}$) in the non-stratified GWAS. We calculated the twin heritability from 749 twin individuals (including

122 complete monozygotic pairs and 176 complete dizygotic pairs) in the BLTS. Heritability, from the best-fitting additive genes/unique environment (AE) model, was 0.28 (95% CI: 0.13–0.42) (Supplementary Note Section 4).

Insert Figure 1 here

Genetic correlation

We next investigated how the non-stratified Eyes Test is genetically correlated to psychiatric conditions and specific psychological and cognitive traits for which summary GWAS data were available (Supplementary Table 5). After correcting for multiple testing, we identified significant positive genetic correlations between Eyes Test scores and two cognitive traits: self-reported empathy measured using the Empathy Quotient ($r_g = 0.18 \pm 0.07$; $P_{\text{FDRadjusted}} = 0.031$)²⁹, and the NEO-Five Factor Inventory measure of openness ($r_g = 0.54 \pm 0.14$; $P_{\text{FDRadjusted}} = 9.73 \times 10^{-4}$)³⁰. We also identified significant positive correlations with different measures of cognition and education: college years ($r_g = 0.40 \pm 0.6$; $P_{\text{FDRadjusted}} = 3.1 \times 10^{-10}$)³¹, educational attainment (0.34 ± 0.04 ; $P_{\text{FDRadjusted}} = 3.7 \times 10^{-16}$)³², and childhood cognitive aptitude (calculated as Spearman's ρ and is, hence, independent of word knowledge)³³ ($r_g = 0.34 \pm 0.10$; $P_{\text{FDRadjusted}} = 0.0075$). In addition, we identified a significant positive genetic correlation between the Eyes Test scores and anorexia nervosa (Anorexia-GCAN $r_g = 0.14 \pm 0.06$; $P_{\text{FDRadjusted}} = 0.047$), which we validated using a non-independent data-freeze with more cases for anorexia nervosa from the Psychiatric Genomics Consortium (Anorexia-PGC $r_g = 0.25 \pm 0.08$; $P_{\text{FDRadjusted}} = 0.009$) (Figure 2). We did not identify a significant genetic correlation between autism and scores on the Eyes Test.

We also investigated if subcortical brain volumes are correlated with performance on the Eyes Test. We used data from the ENIGMA consortium for six subcortical regions and intracranial volume²⁴. We excluded the amygdala, even though it is relevant for social cognition, as the low heritability of the amygdala could not be accurately quantified using LDSR²³. After correcting for multiple testing, we identified a significant positive correlation between the Eyes Test scores and the volumes of the caudate nucleus²⁴ ($r_g = 0.24 \pm 0.09$; $P_{\text{FDRadjusted}} = 0.033$) and volume of the putamen ($r_g = 0.21 \pm 0.08$; $P_{\text{FDRadjusted}} = 0.041$), which together form the dorsal striatum. All genetic correlations are provided in Supplementary Table 5.

We also investigated sex-stratified genetic correlations between the Eyes Test and educational attainment, the only relevant phenotype where we had access to sex-stratified data. We identified a modest, significant genetic correlation between educational attainment and the Eyes Test in the males-only dataset: $r_g = 0.23 \pm 0.05$; $P = 2.6 \times 10^{-5}$. We identified a higher, significant genetic correlation between educational attainment and Eyes Test in the females-only dataset: $r_g = 0.39 \pm 0.06$; $P = 5.88 \times 10^{-11}$. These results suggest that females share greater pleiotropy between general cognition and cognitive empathy than males, indicating different genetic mechanisms for the development of cognitive empathy.

Insert Figure 2 here

Genome-wide association meta-analyses

GWAMA of the non-stratified and the males-only datasets did not identify any significant loci. In the females-only analysis, we identified one locus at 3p26.2 that was significant at a threshold of $P < 5 \times 10^{-8}$. This locus contains 21 significant SNPs in high LD with the leading SNP rs7641347 (Figure 2), with concordant effect direction for 19 SNPs in the 23andMe and BLTS datasets. The leading SNP rs7641347 ($P_{\text{meta}} = 1.58 \times 10^{-8}$) explained 0.067% of the total variance, or 0.013% of the total variance after correcting for winner's curse²⁵. Of the two SNPs with discordant effects in the two datasets, rs114076548 was the most-significant SNP in the 23andMe dataset and had $P = 6.49 \times 10^{-9}$. We did not identify any inflation in the P-values of the GWAMA due to population stratification using LDSR (intercept = 1.01 ± 0.007).

Insert Figure 3 here

The leading SNP (rs7641347) is located in an intron of *SUMF1* and was nominally significant in the non-stratified analysis ($P_{\text{meta}} = 1.1 \times 10^{-5}$), but non-significant in the males-only analysis ($P_{\text{meta}} = 0.4954$). In addition, SNPs in high LD ($r^2 > 0.8$) were also not nominally significant in the males-only analysis. Together, all 21 SNPs span a region of approximately 77kb 3p26.2 (Supplementary Table 1). At this locus, in addition to *SUMF1*, 2 other genes are present: Leucine Rich Neuronal 1 (*LRRN1*) and SET Domain And Mariner Transposase Fusion Gene (*SETMAR*). *LRRN1* is highly expressed in brain tissues³⁴, with median expression the highest in the putamen, nucleus accumbens and the caudate nucleus,

all three of which are part of the striatum. Deletion of 3p26.1 and 3p26.2 can cause developmental delay, hypotonia and epileptic seizures and has been implicated in autism³⁵.

The most significant SNP in the males-only GWAS meta-analysis (rs4300633 in 16p12.3, $P = 9.11 \times 10^{-8}$) explained 0.062% of the variance, and the most significant SNP in the non-stratified GWAS meta-analysis (rs149662397 in 17q21.32 $P = 1.58 \times 10^{-7}$) explained only 0.029% of the variance. All LD pruned SNPs with $P < 1 \times 10^{-6}$ are provided in Supplementary Table 2. The QQ-plot and locus-zoom plot for the females-only meta-analysis, and the Manhattan and QQ-plots for the males-only and non-stratified analyses are provided in the Supplementary Note section 5. Gene-based analyses MetaXcan²⁸ for ten neural tissues (Methods) and functional enrichment analyses for the non-stratified GWAS did not identify any significant results (Supplementary Tables 3 and 4).

Sex differences

We also investigated sex-differences in the Eyes Test. There was a significant female advantage on the scores of the full Eyes Test (males = 27.08 ± 3.75 ; females = 27.85 ± 3.55 ; cohen's $d = 0.21$, $P < 2 \times 10^{-16}$), replicating previous results³⁶ (Figure 4). There was no significant difference in males-only or females-only SNP heritability estimates (males = 0.071 ± 0.011 , females = 0.067 ± 0.011 ; $P = 0.79$). There was a reasonably high but incomplete genetic correlation between males and females ($r_g = 0.68 \pm 0.12$; $P = 2.70 \times 10^{-8}$). Binomial sign test of LD-pruned nominally significant SNPs in the sex-stratified analyses identified that 61% (95% CI: 59% - 62%) of the SNPs had a concordant effect direction ($P < 2.2 \times 10^{-16}$). We further investigated the effect direction and statistical significance of all independent SNPs with $P < 1 \times 10^{-6}$. SNPs that were of suggestive significance in one sex were not nominally significant in the other (Supplementary Table 2 and Supplementary Note section 5). Using MetaXcan²⁸ we identified the top cortically expressed genes ($P < 0.05$) for both sexes and calculated the overlap in the genes. We did not find any enrichment in gene overlap (Fold difference = 1.2, $P = 0.264$). We also investigated if there was an enrichment of female-overexpressed or male-overexpressed cortical genes for the Eyes Test (Methods) and did not find any significant enrichment (Supplementary Note section 3, and Supplementary Tables 6 - 8).

Insert Figure 4 here

Discussion

This is the first large-scale genetic study investigating the genetic architecture of cognitive empathy. We investigated heritability estimates of the Eye Test. In our sample of 749 twin individuals, heritability was approximately 0.28 (95% CI: 0.13–0.42). This is in keeping with previous studies that have investigated heritability of other facets of empathy in twins. A meta-analysis of empathy in twins identified that approximately a third of the variance is heritable³⁷. In our sample of 88,056 unrelated research volunteers from 23andMe Inc, SNP-based heritability was estimated using LDSR, and approximately 5% of the trait was additively heritable. It is likely that heritability of cognitive empathy changes with age (which was significantly correlated with scores on the Eyes Test in this dataset), as is observed in prosocial behaviour³⁷. In our analyses, age was included as a covariate, and thus our SNP heritability is likely to represent the lower bound of the SNP heritability.

We identified significant genetic correlation between volume of the caudate nucleus, putamen and scores on the Eyes Test. Imaging studies have reported activation in both the putamen³⁸ and caudate nucleus³⁹ on tasks of social cognition. In humans, the ventral striatum is composed of the nucleus accumbens and olfactory tubercle, whereas the dorsal striatum is composed of the caudate nucleus and putamen. There is some evidence to support the role of the striatum in theory of mind⁴⁰, though there is no clear consensus that cognitive and affective empathy utilize different neural circuits. Our results suggest that variants that contribute to volumes of regions in the striatum, the dorsal striatum in particular, also contribute to cognitive empathy. It is difficult to delineate the mechanism behind this observed pleiotropy, but we can hypothesize that volumes of striatal structures influence scores on the Eyes Test.

In addition, we identified significant positive genetic correlations with different measures of cognitive ability including educational attainment. This reflects phenotypic correlation between measures of cognitive empathy and cognitive ability. A meta-analysis has identified a significant positive correlation between scores on the Eyes Test and IQ ($n = 3583$; $r = 0.24$; 95% CI: 0.16 – 0.32)⁴¹. Other tests of theory of mind are also positively correlated with cognitive aptitude and measures of intelligence^{42–44}. We also found a significant positive genetic correlation with the NEO-Openness to experience, and self-reported empathy measured using the Empathy Quotient. Again, both the genetic correlations reflect previous correlation at a phenotypic level between measures of empathy and

personality⁴⁵. With psychiatric conditions, there was a significant positive correlation with anorexia which we identified using two non-independent dataset. One study has identified that individuals with anorexia report higher personal distress⁴⁶, though other studies have reported that deficits in social cognition may be attributable to comorbid alexithymia⁴⁷. Our research suggests that higher genetic contribution to cognitive empathy increases the genetic risk for anorexia. However, it is likely that the development of anorexia can contribute to poorer performance on tests of social cognition and cognitive empathy through other factors including alexithymia.

We did not identify a significant genetic correlation between the Eyes Test and autism. Recent research suggests that this may be due to heterogeneity in performance in the Eyes Test, with only a subset of individuals with autism showing impaired performance on the Eyes Test⁴. In addition, the cognitive phenotype of autism involves non-social aspects (such as attention to detail), not just social ones. A meta-analysis has reported global or selective deficits in performance on the Eyes Test in individuals with schizophrenia, anorexia, bipolar disorder, and clinical depression but preserved or even enhanced performance for individuals with non-clinical depression and borderline personality disorder⁴⁸. However, these studies are typically conducted in small sample sizes and presence of psychiatric conditions may impair performance on the Eyes Test. Further, performance on the Eyes Test may be influenced by multiple factors related to psychiatric conditions, and need not necessarily measure the causal relationship between psychiatric conditions and cognitive empathy.

We also identified one locus that is significantly associated with empathy in females. The top SNP (rs7641347) had a P-value = 1.58×10^{-8} . One of the closest gene, *LRRN1*, is highly expressed in striatum according to the GTEx database. However, we were unable to identify any eQTL that specifically linked this locus to the gene. *LRRN1* is a gene that is not well characterized. In chicks, *Lrrn1* is necessary for the formation of the mid-brain hind-brain boundary⁴⁹. The locus was significant in females, nominally significant in the non-stratified analyses, and non-significant in the males-only analyses, suggesting a sex specific involvement of this locus in cognitive empathy measured using the Eyes Test. We note that even with approximately 90,000 individuals this GWAMA was underpowered to detect loci of significant effect, owing to the relatively low variance explained per SNP.

It was also interesting to note that while twin and SNP-based heritability did not vary between the sexes in our study, we replicated the female-advantage on the Eyes Test in the largest sample thus far. Sex-stratified analyses also allowed us to investigate the genetic correlation between males and females, and subsequently, sex-specific imputed gene expression in cortical tissues. Male-female genetic correlation was only modest, which was supported by binomial sign test. In comparison, other traits for which we had sex-stratified data, genetic correlation was considerably higher (eg: self-reported empathy²⁹: $r_g = 0.82$, $sd = 0.16$, systemizing²⁹: $r_g = 1.0$, $sd = 0.16$; educational attainment³²: $r_g = 0.91$, $sd = 0.02$). Further, examining the top SNPs indicated a sex-specific architecture for the most significant set of SNPs ($P < 1 \times 10^{-6}$). We also did not identify a significant overlap between the genes identified for the sex-stratified GWAS. All of this suggests that there is some sex specificity in the genetic architecture of cognitive empathy. How this sex-specific architecture is expressed and interacts with steroid hormones will help shed further light on the biological contributions to the female superiority on the Eyes Test.

In conclusion, we identify a genetic locus that is associated with scores on the Eyes Test in females. We identify significant positive genetic correlations between scores on the Eyes Test and empathy, cognitive aptitude and educational attainment, and openness to experience. We also identify positive correlation between striatal volumes and scores on the Eyes Test. Phenotypic sex-differences for the Eyes Test may be partly due to different genetic architectures in males and females, interacting with postnatal social experience.

Acknowledgements

This study was funded by grants from the Templeton World Charity Foundation, Inc., the Medical Research Council, the Wellcome Trust, the Autism Research Trust, the Institut Pasteur, the CNRS and the University Paris Diderot. VW is funded by St. John's College, Cambridge, and Cambridge Commonwealth Trust. The research was carried out in association with the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care East of England at Cambridgeshire and Peterborough NHS Foundation Trust. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. The research was supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care East of England at Cambridgeshire and Peterborough NHS Foundation Trust and the National Human Genome Research Institute of the National Institutes of Health (grant number R44HG006981). The National Science Foundation (grant numbers 0729493 and 0721707) supported the research on the Brisbane Longitudinal Twin Study. FU was supported by the British Friends of Haifa University, the Israel Science Foundation (grant no. 449/14), the British Friends of Hebrew University, and the Joseph Levy Charitable Foundation. TB was supported by the Institut Pasteur, the Bettencourt-Schueller Foundation, University. We thank the research participants and employees of 23andMe for making this work possible. We also thank the volunteers of the Brisbane Longitudinal Twin Study and The NIHR Cambridge BioResource. Finally, we thank Carrie Allison, Michael Lombardo, Owen Parsons, and Richard Bethlehem for helpful discussions.

Conflict of Interest

David A Hinds and Nadia Litterman are employees of 23andMe, Inc.

References

- 1 *Understanding Other Minds: Perspectives from developmental social neuroscience*. OUP Oxford, 2013 <https://books.google.com/books?id=3nloAgAAQBAJ&pgis=1> (accessed 25 Jul2015).
- 2 Baron-Cohen S, Wheelwright S, Hill J, Raste Y, Plumb I. The 'Reading the Mind in the Eyes' Test revised version: a study with normal adults, and adults with Asperger syndrome or high-functioning autism. *J Child Psychol Psychiatry* 2001; **42**: 241–51.

- 3 Decety J, Moriguchi Y. The empathic brain and its dysfunction in psychiatric populations: implications for intervention across different clinical conditions. *Biopsychosoc Med* 2007; **1**: 22.
- 4 Baron-Cohen S, Bowen DC, Holt RJ, Allison C, Auyeung B, Lombardo M V *et al.* The ‘Reading the Mind in the Eyes’ Test: Complete Absence of Typical Sex Difference in ~400 Men and Women with Autism. *PLoS One* 2015; **10**: e0136521.
- 5 Bora E, Gökçen S, Veznedaroglu B. Empathic abilities in people with schizophrenia. *Psychiatry Res* 2008; **160**: 23–9.
- 6 Michaels TM, Horan WP, Ginger EJ, Martinovich Z, Pinkham AE, Smith MJ. Cognitive empathy contributes to poor social functioning in schizophrenia: Evidence from a new self-report measure of cognitive and affective empathy. *Psychiatry Res* 2014; **220**: 803–10.
- 7 Calderoni S, Fantozzi P, Maestro S, Brunori E, Narzisi A, Balboni G *et al.* Selective cognitive empathy deficit in adolescents with restrictive anorexia nervosa. *Neuropsychiatr Dis Treat* 2013; **9**: 1583–9.
- 8 Tone EB, Tully EC. Empathy as a ‘risky strength’: a multilevel examination of empathy and risk for internalizing disorders. *Dev Psychopathol* 2014; **26**: 1547–65.
- 9 Lam BYH, Raine A, Lee TMC. The relationship between neurocognition and symptomatology in people with schizophrenia: social cognition as the mediator. *BMC Psychiatry* 2014; **14**: 138.
- 10 Cusi AM, Macqueen GM, McKinnon MC. Patients with bipolar disorder show impaired performance on complex tests of social cognition. *Psychiatry Res* 2012; **200**: 258–64.
- 11 Tapajóz Pereira de Sampaio F, Soneira S, Aulicino A, Allegri RF. Theory of mind in eating disorders and their relationship to clinical profile. *Eur Eat Disord Rev* 2013; **21**: 479–87.
- 12 Berlim MT, McGirr A, Beaulieu M-M, Turecki G. Theory of mind in subjects with major depressive disorder: is it influenced by repetitive transcranial magnetic stimulation? *World J Biol Psychiatry* 2012; **13**: 474–9.
- 13 Tung JY, Do CB, Hinds DA, Kiefer AK, Macpherson JM, Chowdry AB *et al.* Efficient replication of over 180 genetic associations with self-reported medical data. *PLoS One* 2011; **6**: e23473.
- 14 Do CB, Tung JY, Dorfman E, Kiefer AK, Drabant EM, Francke U *et al.* Web-based genome-wide association study identifies two novel loci and a substantial genetic

- component for Parkinson's disease. *PLoS Genet* 2011; **7**: e1002141.
- 15 Eriksson N, Tung JY, Kiefer AK, Hinds DA, Francke U, Mountain JL *et al.* Novel associations for hypothyroidism include known autoimmune risk loci. *PLoS One* 2012; **7**: e34442.
- 16 Henn BM, Hon L, Macpherson JM, Eriksson N, Saxonov S, Pe'er I *et al.* Cryptic distant relatives are common in both isolated and cosmopolitan genetic samples. *PLoS One* 2012; **7**: e34267.
- 17 Hatemi PK, Smith K, Alford JR, Martin NG, Hibbing JR. The genetic and environmental foundations of political, psychological, social, and economic behaviors: a panel study of twins and families. *Twin Res Hum Genet* 2015; **18**: 243–55.
- 18 Vellante M, Baron-Cohen S, Melis M, Marrone M, Petretto DR, Masala C *et al.* The 'Reading the Mind in the Eyes' test: systematic review of psychometric properties and a validation study in Italy. *Cogn Neuropsychiatry* 2013; **18**: 326–54.
- 19 Fernández-Abascal EG, Cabello R, Fernández-Berrocal P, Baron-Cohen S. Test-retest reliability of the 'Reading the Mind in the Eyes' test: a one-year follow-up study. *Mol Autism* 2013; **4**: 33.
- 20 Howie B, Fuchsberger C, Stephens M, Marchini J, Abecasis GR. Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nat Genet* 2012; **44**: 955–9.
- 21 Browning SR, Browning BL. Rapid and Accurate Haplotype Phasing and Missing-Data Inference for Whole-Genome Association Studies By Use of Localized Haplotype Clustering. *Am J Hum Genet* 2007; **81**: 1084–1097.
- 22 Willer CJ, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 2010; **26**: 2190–1.
- 23 Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007; **81**: 559–75.
- 24 Hibar DP, Stein JL, Renteria ME, Arias-Vasquez A, Desrivieres S, Jahanshad N *et al.* Common genetic variants influence human subcortical brain structures. *Nature* 2015; **520**: 224–9.
- 25 Bigdeli TB, Lee D, Webb BT, Riley BP, Vladimirov VI, Fanous AH *et al.* A simple yet accurate correction for winner's curse can predict signals discovered in much larger genome scans. *Bioinformatics* 2016; **32**: 2598–2603.
- 26 Bulik-Sullivan BK, Loh P-R, Finucane HK, Ripke S, Yang J, Patterson N *et al.* LD

- Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet* 2015; **47**: 291–295.
- 27 Boker S, Neale M, Maes H, Wilde M, Spiegel M, Brick T *et al*. OpenMx: An Open Source Extended Structural Equation Modeling Framework. *Psychometrika* 2011; **76**: 306–317.
 - 28 Barbeira A, Shah KP, Torres JM, Wheeler HE, Torstenson ES, Edwards T *et al*. MetaXcan: Summary Statistics Based Gene-Level Association Method Infers Accurate PrediXcan Results. Cold Spring Harbor Labs Journals, 2016 doi:10.1101/045260.
 - 29 Warrier V, Toro R, Chakrabarti B, Litterman N, Hinds D, Bourgeron T *et al*. Genome-wide analyses of empathy and systemizing: heritability and correlates with sex, education, and psychiatric risk. Cold Spring Harbor Labs Journals, 2016 doi:10.1101/050682.
 - 30 de Moor MHM, Costa PT, Terracciano A, Krueger RF, de Geus EJC, Toshiko T *et al*. Meta-analysis of genome-wide association studies for personality. *Mol Psychiatry* 2012; **17**: 337–49.
 - 31 Rietveld CA, Medland SE, Derringer J, Yang J, Esko T, Martin NW *et al*. GWAS of 126,559 individuals identifies genetic variants associated with educational attainment. *Science* 2013; **340**: 1467–71.
 - 32 Okbay A, Beauchamp JP, Fontana MA, Lee JJ, Pers TH, Rietveld CA *et al*. Genome-wide association study identifies 74 loci associated with educational attainment. *Nature* 2016; **533**: 539–542.
 - 33 Benjamin B, Pourcain B, Davis OS, Davies G, Hansell NK, Brion M-JA *et al*. Childhood intelligence is heritable, highly polygenic and associated with FBNP1L. *Mol Psychiatry* 2014; **19**: 253–8.
 - 34 The Genotype-Tissue Expression (GTEx) project. *Nat Genet* 2013; **45**: 580–5.
 - 35 Pinto D, Delaby E, Merico D, Barbosa M, Merikangas A, Klei L *et al*. Convergence of genes and cellular pathways dysregulated in autism spectrum disorders. *Am J Hum Genet* 2014; **94**: 677–94.
 - 36 Kirkland RA, Peterson E, Baker CA, Miller S, Pulos S. Meta-analysis reveals adult female superiority in ‘Reading the mind in the eyes test’. *N Am J Psychol* 2013; **15**: 121–146.
 - 37 Knafo A, Uzefovsky F. Variation in empathy: The interplay of genetic and environmental factors. In: Legerstee M, Haley DW, Bornstein MH (eds). *The infant mind: Origins of the social Brain*. The Guilford Press, 2013, pp 97 – 121.

- 38 Campanella F, Shallice T, Ius T, Fabbro F, Skrap M, Adolphs R *et al.* Impact of brain tumour location on emotion and personality: a voxel-based lesion-symptom mapping study on mentalization processes. *Brain* 2014; **137**: 2532–45.
- 39 Kemp J, Berthel M-C, Dufour A, Després O, Henry A, Namer IJ *et al.* Caudate nucleus and social cognition: neuropsychological and SPECT evidence from a patient with focal caudate lesion. *Cortex* 2013; **49**: 559–71.
- 40 Abu-Akel A, Shamay-Tsoory S. Neuroanatomical and neurochemical bases of theory of mind. *Neuropsychologia* 2011; **49**: 2971–2984.
- 41 Baker CA, Peterson E, Pulos S, Kirkland RA. Eyes and IQ: A meta-analysis of the relationship between intelligence and ‘Reading the Mind in the Eyes’. *Intelligence* 2014; **44**: 78–92.
- 42 Ibanez A, Huepe D, Gempp R, Gutiérrez V, Rivera-Rei A, Toledo MI. Empathy, sex and fluid intelligence as predictors of theory of mind. *Pers Individ Dif* 2013; **54**: 616–621.
- 43 Charlton RA, Barrick TR, Markus HS, Morris RG. Theory of mind associations with other cognitive functions and brain imaging in normal aging. *Psychol Aging* 2009; **24**: 338–48.
- 44 Buitelaar JK, van der Wees M, Swaab-Barneveld H, van der Gaag RJ. Verbal memory and Performance IQ predict theory of mind and emotion recognition ability in children with autistic spectrum disorders and in psychiatric control children. *J Child Psychol Psychiatry* 1999; **40**: 869–81.
- 45 Magalhães E, Costa P, Costa MJ. Empathy of medical students and personality: Evidence from the Five-Factor Model. *Med Teach* 2012; **34**: 807–812.
- 46 Beadle JN, Paradiso S, Salerno A, McCormick LM. Alexithymia, emotional empathy, and self-regulation in anorexia nervosa. *Ann Clin Psychiatry* 2013; **25**: 107–20.
- 47 Brewer R, Cook R, Cardi V, Treasure J, Bird G. Emotion recognition deficits in eating disorders are explained by co-occurring alexithymia. *R Soc open Sci* 2015; **2**: 140382.
- 48 Dinsdale N, Morkkonen M, Crespi B, Abu-Akel A, Abu-Akel A, Bailey AL *et al.* The ‘extreme female brain’: increased cognitive empathy as a dimension of psychopathology. *Evol Hum Behav* 2016; **37**: 323–336.
- 49 Tossell K, Andrae LC, Cudmore C, Lang E, Muthukrishnan U, Lumsden A *et al.* *Lrrn1* is required for formation of the midbrain-hindbrain boundary and organiser through regulation of affinity differences between midbrain and hindbrain cells in chick. *Dev Biol* 2011; **352**: 341–52.

Figure 1: Schematic diagram of the study protocol

88,056 Caucasian participants from 23andMe, Inc. completed the full version of the Eyes Test and were genotyped. An additional 1,497 Caucasian participants from the Brisbane Longitudinal Twin Study completed the short version (14 questions) of the Eyes Test and genotyped. Genome-wide association meta-analysis was performed on the combined cohort of 89553 participants. Three separate meta-analyses were performed: males-only, females-only, and non-stratified. Subsequently, functional enrichment and gene-based analysis was performed for the non-stratified meta-analysis GWAS using the 23andMe dataset. SNP heritability and genetic correlation using LDSR was performed for the 23andMe GWAS dataset. Sex differences were also investigated using the same dataset. In parallel, twin heritability was calculated from 749 twin individuals from the Brisbane Longitudinal Twin Study who had completed the short version of the Eyes Test.

Figure 2: Genetic correlations between the Eyes Test and psychiatric conditions, psychological traits and subcortical brain volumes

Genetic correlations and standard errors for the Eyes Test in the 23andMe cohort. Figures above the bars represent FDR adjusted P-values. All P-values with $P_{FDRcorrected} < 0.05$ provided. * represents significant P-values after FDR correction. Point estimate represents the genetic correlation, and the error bars represent the standard errors. AN (GCAN) is the Anorexia Nervosa-GCAN dataset, AN (PGC) is the Anorexia Nervosa PGC dataset, BIP is bipolar disorder, BPD features is borderline personality disorder features, ICV is intracranial volume. We have removed the genetic correlation for agreeableness from this figure due to the high standard errors. The genetic correlations, standard errors, and P-values for all traits including agreeableness are provided in Supplementary Table 5.

Figure 3: Manhattan plot and regional association plot for the Eyes Test (females) meta-analysis GWAS

A. Manhattan plot of the Eyes Test meta-analysis (female). X axis is the chromosomal position of the SNP, and Y axis is the negative logarithm of the P-value. The red line indicates genome-wide significant threshold of 5×10^{-8} . Lead SNP for all loci with $P < 1 \times 10^{-6}$ is provided. $n = 44,574$, and $\lambda_{gc} = 1.05$. LDSR intercept = 1.05. Regional association plot of the significant locus for the Eyes Test (females) meta-analysis.

Figure 4: Mean scores and SNP heritability

4A. Mean phenotypic scores and standard deviations for the Eyes Test in the 23andMe cohort. Point estimate provides the mean score, and the error bars represent standard deviations. Difference in mean scores between males and females was highly significant ($P < 2.2E-16$; Cohen's $d = 0.21$). Numbers in brackets indicate the number of participants in each GWAS. All: non-stratified GWAS; Females: Females-only GWAS; Males: Males-only GWAS. 4B. Mean SNP heritability estimates and standard errors for the Eyes Test in the GWAMA. Point estimate provides mean SNP heritability, and error bar represents standard errors. There was no significant difference in SNP heritability estimates between males and females ($P = 0.79$). Numbers in brackets indicate the number of participants in each GWAMA. All: non-stratified GWAMA; Females: Females-only GWAMA; Males: Males-only GWAMA