Title page

Morphological Alternations of Intraepithelial and Stromal Telocytes in Response to Salinity Challenges

The full names of all authors.

Soha Mohamed Abdel-latif Soliman (corresponding author)

lecturer of Histology

Department of Histology

Faculty of Veterinary Medicine, South Valley university, Qena, Egypt.

Postal code. 83523. Tel. +201006500848. Fax. +2 09652112231627. Email.

Soha.soliman@yahoo.com

soha_soliman@vet.svu.edu.eg

Walaa Fathy Ali Emeish

lecturer of fish Diseases

Department of fish Diseases

Faculty of Veterinary Medicine, South Valley university, Qena 83523, Egypt.

walaavet2002@yahoo.com

The current study was carried out in South Valley University

Short title **telocytes responding to salinity**

Keywords intraepithelial telocytes, stromal telocytes, salinity, chloride cells, stem cells,

Rodlet cells.

31 ***Summary statement***

32 The article represent an experimental study in which we investigated the effect of the
33 ssalinty on the communicating cells (telocytes) and their target cells including chloride,
34 stem, Rodlet cells, myoblasts
35

Morphological Alternations of Intraepithelial and Stromal Telocytes in Response to Salinity Challenges

Soha A. Soliman¹, Walaa F.A. Emeish²

¹ Department of Histology, Faculty of Veterinary Medicine, South Valley University, Qena 83523, Egypt. (e-mail, soha.soliman@yahoo.com)

² Department of fish Diseases Faculty of Veterinary Medicine, South Valley university, Qena 83523, Egypt.(e-mail, walaavet2002@yahoo.com).

Abstract

Telocyte is a communicating cell established relations to various types of cells. Few experimental studies are performed on telocytes. The current study investigated response of telocytes to salinity stress in relations to osmoregulatory, immune and stem cells. We exposed Common carp to salinity level 0.2, 6, 10, 14 ppt. Gill samples were fixed and processed for microscopic and TEM. Two types of telocytes were identified: intraepithelial and stromal telocytes. Intraepithelial telocytes comprised the cellular lining of the lymph spaces where they shed the secretory vesicles. Stromal telocytes shed their secretory vesicles in the secondary circulatory vessels. Telocyte enlarged and exhibited high secretory activities. They exert their effect either by direct contact or by paracrine mode. In salinity treated samples, chloride cells enlarged and the mitochondria became cigar-shaped. pavement cells enlarged and micro-ridges elongated. Stromal telocytes established contact with stem cell and skeletal myoblast. Macrophages and Rodlet cells increased in number. In conclusion, intraepithelial and stromal responded to salinity stress by activation of cellular signaling. They play a major role in osmoregulation, immunity, and regeneration.

63

64 **Keywords** intraepithelial telocytes, stromal telocytes, salinity, chloride cells, stem cells,
65 Rodlet cells.

66

67 **introduction**

68 Telocyte is a distinctive type of interstitial cells, which have a wide range of
69 biological functions in different tissues and organs. Functional diversity of telocytes is
70 regarding affecting different types of cells and structures (**Varga, Danisovic et al. 2016**).
71 Telocytes have unique morphological characteristics. Multiple cell prolongations;
72 telopodes emerge from the cell body and may extend to hundreds of microns. Telopodes
73 may give rise dichotomous branches and establish cellular connections to form a complex
74 labyrinthine system. Telopodes composed of thin segments; podomers and interval
75 expansions; podoms which are rich in calcium release units; mitochondria, endoplasmic
76 reticulum, and caveolae. Telocytes establish a synaptic junction connecting to
77 immunoreactive cells (**Popescu and Faussone-Pellegrini 2010**).

78

79 Telocytes exert their effect on cells either by establishing cellular contact or
80 through paracrine mode. Two types of cellular contact are documented for telocytes;
81 homocellular and heterocellular contact. Homocellular contact is formed between two
82 telopodes or telocytes and telopodes or between the cell body of two adjacent telocytes.
83 Heterocellular type contact between telocytes and stromal cells either fixed or the free
84 cells. Various types of cellular contacts and communication are mentioned in telocytes
85 including direct apposition of the cell membrane of adjacent telocytes, adherence, and

gap junction. Gap junction have a significant role in intercellular signaling pathway (Mirancea 2016). The secretory function of telocytes influence the target cells. Telocytes deliver microvesicles to the cell and providing macromolecules such as proteins or RNAs, microRNA. Telocytes also shed exosomes, ectosomes and multivesicular vesicles (Popescu, Gherghiceanu et al. 2005; Popescu and Faussone-Pellegrini 2010; Cantarero Carmona, Luesma Bartolome et al. 2011)

Telocytes are multifunctional cell functions. They contribute to generation and transmission of nerve impulses to involuntary muscles (Takaki 2003; Hutchings, Williams et al. 2009; Gandahi, Chen et al. 2012; Drumm, Koh et al. 2014). They serve in mechanoreception and may involve in atrial fibrillation (Gherghiceanu, Hinescu et al. 2008). Telocytes exhibit receptors for excitatory and inhibitory neurotransmitters (Iino and Horiguchi 2006). They establish contact with immunoreactive cells such as eosinophil (Cantarero Carmona, Luesma Bartolome et al. 2011), mast cell, and macrophage (Gherghiceanu and Popescu 2012). Telocytes play a role in regeneration of heart, lung, skeletal muscle, skin, meninges and choroid plexus, eye, liver, uterus, urinary system (Bei, Wang et al. 2015).

Several studies are conducted to study telocytes in human and other mammals but few studies are performed in aquatic species. the current study was conducted using the common carp. The carp belong to *Cyprinidae* family which is commonly known in North America as minnow, while in Eurasia is termed as carp. *Cyprinidae* family are freshwater

and are uncommon in brackish water; North America, Africa, and Eurasia (**Nelson 2006**).

Aquatic species regulate ionic exchange to maintain osmotic balance according to environmental salinity. Several organs are involved in osmoregulation including gills, intestine, kidney, skin, operculum (**Marshall and Grosell 2005**). Marine inhabitants face great challenges to establish ionic balance. Therefore, the ion transporting cells; chloride cells or ionocytes; participate in the elimination of excess ions in seawater fish, while in freshwater fish, chloride cells contribute in ion absorption (**Florkin 2014**). IN MARINE FISH, Chloride cells are structurally modified to adopt high salinity levels. Fish exposed to high salinity environment acquire a high proportion of mitochondrial-rich chloride cells (**Fielder, Allan et al. 2007**). Pervious researches studied the salinity in relation to changes of ionocytes cells. Ionocytes serve in osmoregulation via different types of membranous channels; cystic fibrosis transmembrane regulator (CFTR) anion channel, Na,K,2Cl cotransporter (NKCC) and sodium pump (Na,K-ATPase). CFTR is considered as a membrane protein located on the apical surface of many types of epithelial cells. CFTR is a cyclic AMP-dependent chloride channel, a bicarbonate channel and as a modulator of other ion channels (**Derichs 2013**). The Na-K-Cl cotransporter (NKCC) is a membrane transport proteins that involved in the active transport of sodium, potassium, and chloride ions across the cell membrane (**Russell 2000**). Na-K-ATPase is an electrogenic transmembrane enzyme located predominantly on the basolateral surface of the chloride cell and actively transport chloride, rather than sodium across the plasma membrane (**Suhail 2010**). Pavement cells mostly cover the surface the of the filament and

lamellar epithelium. They considered as ion transporting cells; their cell membrane is rich in hydrogen ion channels (Laurent, Goss et al. 1994; Perry and Fryer 1997). In the current study, we focused on the communicating cells; telocytes which influence the large population of stromal, muscular, and epithelial cells. The aim of the present study was to investigate morphological alternations of telocytes subjected to salinity stress and their effect on different types of cells with special reference to the osmoregulatory and immune and stem cells in the common carp.

Materials and methods

I- Fish source and transportation

The common carp, *Cyprinus carpio*. was obtained from a private fish farm at El-Dakahlea Government and transported in large water tanks. During transportation, the oxygen level was maintained at 5 mg/l and water tank temperature was 23°C ±3 and pH value at 7.2 – 7.5.

II- Fish acclimation

Apparently, healthy fingerlings fish measured about the length of 7±2 cm and aged 1 month old. The body weight was 10±2 g, and. Fish were collected and transported to the wet laboratory at Faculty of Veterinary Medicine, South Valley University, Qena, Egypt. Fish were maintained under laboratory conditions during adaptation in running water (salinity = 0.2 ppt) for 3 weeks before conducting the experiments and fed twice daily to ad libitum feed on a commercial floating powdered feed containing 45% protein with a feeding rate of 3% of their body weight.

154

155 **III- Aquaria**

156 Fish were originally kept in a re-circulating system in porcelain aquaria (260 ×65×70cm)
 157 according to the protocol of maintaining bioassay fish as was previously described
 158 (Elsaesser and Clem, 1986). Experiments were conducted in fiberglass aquaria with
 159 dimensions of 60×30×40 cm. Dissolved oxygen level was maintained above 5 mg/l while
 160 water temperature was kept at 23°C ±3 and pH value at 7.2 – 7.5.

161

162 **IV- Salinity exposure**

163 36 acclimated, apparently healthy Common carp, *C. carpio* were selected with a body
 164 weight range of 9 – 11 g to serve as the experimental groups. Fish were divided into 12
 165 fiberglass aquaria (60×30×40 Cm) to serve as 4 experimental groups, each group contains
 166 9 fish, and there were 3 replicates for each salinity group. Three groups were gradually
 167 subjected to three different salinities until concentrations of 6, 10 and 14 ppt with 2 g/L
 168 NaCl increase every two days. The fourth group was reared in freshwater; a dechlorinated
 169 tap water of 0.2 ppt salinity level and considered as control group. Water was changed
 170 every two days with water that had desired salinities, and aquariums were also cleaned at
 171 this time. Salinity was checked and adjusted regularly every two days during a water
 172 change. When common carp reached the final desired salinity, fish were allowed to
 173 acclimate to the new salinities for a minimum of two weeks before sample collections.

174

175 **V- Clinical examination of fish**

Fish were observed daily during the course of an experiment for any apparent clinical signs, lesions or mortality. Mortality rate was calculated from the number of dead fish between each sampling period.

VI- Fish sampling

At the end of the period, nine fish were decapitated in each salinity level. Gill filaments and gill arches of both sides were dissected and fixed in glutaraldehyde (10 mL of 2.5% glutaraldehyde and 90 mL 0.1 M Na-phosphate buffered formalin).

VII- preparation of resin embedding specimens for semi-thin and ultra-thin sectioning

Fixed samples of gill filaments and arches were cut into small pieces. They were washed 4 times for 15 minutes in 0.1 M sodium phosphate buffer (pH 7.2) then were post-fixed in 1% osmic acid in 0.1 M Na-phosphate buffer at 4°C for 2 hours. The osmicated samples were washed 3 times for 20 minutes in 0.1 M phosphate buffer (pH 7.2). Dehydration was performed through graded acetone (70, 80, 90, 100%), 10 minutes for each concentration. The dehydrated samples were immersed in a mixture of acetone/resin (1/1 for 1 day, 1/2 for another day) and pure resin for three days. The resin was prepared by using 10gm ERL, 6gm DER, 26gm NSA and 0.3gm DMAE and thoroughly mixed by a shaker. The specimens were embedded in the resin at 60 °C for 3 days. Polymerized samples were cut to semi-thin sections by using an ultramicrotome Ultracut E (Reichert-Leica, Germany) and stained with toluidine blue (**Bancroft, Layton et al. 2013**).

Semi-thin sections were also used in histochemical studies. The sections were treated with a saturated alcoholic solution of sodium hydroxide for 15 minutes to dissolve the resin (Lloyd 2001). The semi-thin sections were stained by Heidenhain's Iron-Hx (Heidenhai 1896) and methylene blue used for staining of paraffin sections and prepared as a stain for semi-thin sections (Bancroft, Layton et al. 2013).

Ultrathin sections were obtained by a Reichert ultra-microtome. The sections (70 nm) were stained with uranyle acetate and lead citrate (Reynolds, 1963) and examined by JEOL100CX II transmission electron microscope (TEM) at the CENTREAL LABARTORY UNIT of South Valley University.

VIII- Coloring images

Transmission electron microscopy images were colored using photo filter 6.3.2 program. Coloring images required to change the color balance, using the stamp tool to color the objective cells.

Results

The current study was performed to evaluate ranges of acclimation of the common carp to the hypertonic conditions in relation to responses of the communicating cells; telocytes and the related effector cells including immune, chloride and stem cells in gill filaments and arches using semi-thin and ultrathin sections.

Control and 6ppt salinity exposed groups exhibited normal morphology and behavior and had no noticeable signs of stress and no mortality. However, marked reduction of swimming speed and fish were easily caught in 10 and 14 ppt salinity treated fish.

By semi-thin sections, intraepithelial telocytes had a small cell body and well-defenind telopodes in control samples (Fig. 1A, E, I). They were gradually enlarged in size during exposure to different levels of salinity. In 6 ppt, intraepithelial telocytes were satellite in shape (Fig. 1B, F, J). In 10 ppt and 14 ppt they were large satellite cells with multiple telopodes Fig. 1C, D, G, H, K, L). stromal telocytes were small and had spindle-shaped cell body form which extended fine telopodes in control samples (Fig. 2A, E, I). In 6 ppt salinity concentration, some stromal telocytes were enlarged and telopodes formed a network (Fig. 2 B, F, J). Telopodes formed an extensive network; secretory vesicles were large and could be easily recognized in samples treated with 10 and 14 ppt salinity levels (Fig 2 C, D, G, H, K, L).

Telocytes were identified for the first time by TEM in the epithelium of the gill arches. In control samples, telocytes represented the cellular lining of the intraepithelial lymphatic space in which immuno-reactive cells migrate. Intraepithelial telocytes were small, had spindle or satellite-shaped, their telopodes were thin and formed a labyrinth network separating between the compartments of the lymphatic space. Intraepithelial telocytes rest on the basement membrane. Telopodes extended between epithelial and immune cells. Intraepithelial telocytes could establish contact with epithelial cells. The secretory vesicles of the telopodes were excreted in the intraepithelial lymphatic space (Fig 4 A,

B). In samples treated with 60 ppt salinity level, intraepithelial telocytes undergo hypertrophy and telopodes were thickened (Fig. 4 C-H).

In 10 salinity concentration, intraepithelial telocytes was hypertrophy associated with enlargement of the podoms. They also acquired high secretory activity. Intraepithelial telocytes established planar contact with chloride cell (Fig 5A-E). In 14 ppt salinity level, telocytes shed secretory vesicles, exosomes, and multivesicular vesicle into the intra-epithelial lymphatic spaces. Intra-epithelial telocytes established planar contact with chloride cells (Fig 6A- D).

Stromal telocytes appeared small spindle, satellite, rounded, triangular shaped cell body with thin cellular prolongations (telopodes) in control samples. Telopodes consisted of podoms and podomers. Macrophages were small and contained vesicles (Fig. 7 A, B, Fig. 8A, B). Telocytes established homocellular junction (Fig 7 C, D). Stromal telocytes undergo morphologically modifications during increasing the concentration of the salinity. The cell body of some populations of telocytes enlarged in salinity levels of 6 ppt (Fig 4 C-H), and 10 ppt (Fig. 5 A, F). In 10 ppt salinity level, telopodes were thickened and became slightly wavy (Fig. 7E, F). Telopodes frequently formed an extensive network (Fig. 8 F). The prominent features of high salinity changes were irregular surface telocyte, waviness, and thickening of telopodes (Fig 7 G, H). They exhibited higher secretory activities in salinity levels reached 6, 10, and 14 ppt (Fig 8 A-D). Stromal telocytes established contact with the endothelial lining of the secondary circulatory system or lymph vessels and liberated the secretory vesicles in proximity to

the vessels. Trans-endothelial transportation of the secretory vesicles was observed in the secondary circulatory vessels. The transferring vesicles were shed in the lumen of the secondary circulatory pathway (Fig. 4G, Fig. 6E, Fig. 8E, F).

Telocytes established contact with different types of stromal, epithelial, stem cells and skeletal muscles. Telopodes were connected to nerve fiber (Fig 12A-C) and formed point contact (fig. 7B) or multipoint contact (Fig 12A-C) with skeletal muscles. Large amount of secretory vesicles were excreted closed to the muscular fibers in samples treated with 6 ppt concentration of salinity (Fig 12A, C). An extensive network of telopodes was observed and more secretory vesicles were shed from telocytes. Skeletal muscle fiber undergoes hypertrophy after reaching 10 ppt salinity concentration (Fig. 12D, E).

Both intraepithelial and stromal telocytes established contact with macrophages either via telopodes or the cell body (Fig. 4H, Fig. 5 A, F, Fig. 11C, D). In control samples, few macrophages were detected in the gill arches stroma (Fig. 7 A, B). Macrophages became more active and were rich in lysosomes and vesicles in 6 ppt treated samples (Fig. 4 F-H). In 10 ppt salinity level, large number of macrophages in gill arches stroma (Fig. 5 A-F) and epithelial lymphatic spaces (Fig. 5C, Fig. 7E, F). In 14 ppt treated samples, massive lysosomal-rich macrophages were observed in the stroma and epithelial lymphatic spaces (Fig 6B, Fig. 7G).

Telopodes of several telocytes wrapped around stem cells and partially enclosed the stem cell. Telopodes formed a planar contact along the cell membrane of the stem cell, telopodes gave rise small branches extended into the cytoplasm of stem cell (Fig. 11 A,

B). Telocytes and their telopodes surrounded and established direct contact with the skeletal myoblast (Fig. 11 C, E, F). Telopodes also formed contact with Schwann cells (Fig. 11 C, D).

Intraepithelial and stromal telocytes formed contact with immature rodlet cells (granular rodlet cells) (Fig. 4 F-H, Fig. 5A-D, Fig. 9A, B). Telocytes shed secretory vesicles and multi-vesicular body in the vicinity of immature rodlet cells (Fig. 9A, B). The secretory vesicles of the telocytes were observed in the surface epithelium (Fig. 9 C, D). Intraepithelial telocytes established planar contact with pavement cell (Fig. 9E). In control samples, Pavement cells were flattened with short microvilli (Fig 4A). Pavement cells undergo modifications in salinity treated samples. They enlarged in 6 ppt treated samples (Fig. 9C). In 10 ppt salinity concentration, they were cuboidal in shape (Fig. 9E). Pavement cells became elongated and appeared columnar-shaped in 14 ppt level of salinity (Fig. 6A). The micro-ridges became thin, elongated and extended beyond the epithelial surface. Micro-ridges could be seen attached to or enclosing the secretory vesicles of the telocytes. The surface of pavement cells formed pit-like invaginations (Fig. 6 C, D).

Intraepithelial telocytes established planer contact with Chloride cells (Fig. 5D, 6C). Chloride cells undergo structural modifications during elevation the level of the senility. By TEM, they enlarged, and increase in number gradually depending on salinity concentration. the mitochondrial number increased and changed their morphology from rounded or oval in control group to elongated cigar-shaped in treated samples. Chloride

delivered the secretory vesicles of telocytes which were transferred through the intra-epithelial lymphatic space (Fig 10 A-D, Fig. 5C, Fig. 6A). Amount of mitochondria in the chloride cells was also evaluated by using Heidenhain's Iron-Hx. Mitochondria appeared as black granules which increased with the level of the salinity (Fig. 3E-H).

Discussion

The current investigation was carried out to evaluate telocyte response to salinity stress and their relation to osmoregulatory and immune cells. We detected telocytes in semi-thin sections using toluidine blue, methylene blue and Heidenhain's Iron-Hx, and ultra-thin sections to examine ultrastructural modifications in telocytes in relation to epithelial and stromal cells.

In the current study, telocytes undergo morphological alternations during salinity stress. They were spindle-shaped with fine telopodes in control samples. No significant changes occurred in telocytes in 6 ppt salinity levels, while some telocytes exhibited higher secretory activities. Telocytes shed large secretory vesicles, some populations of telocytes had enlarged cell body, and telopodes became thicker and formed an extensive network, in samples treated with 10 and 14 ppt salinity levels. Hormonal administration could affect the secretory activities of telocytes. Exaggerated secretory activities of telocytes are documented in melatonin treatment of ram seminal vesicles (Abd-Elhafeez, Mokhtar et al. 2016).

In the current study, two types of telocytes were detected in the gills of common carp according to location; intraepithelial telocytes and stromal telocytes. Telopodes formed homocellular and heterocellular contacts. Heterocellular contact was established with a wide range of cells and structures.

Intraepithelial telocytes communicated to from labyrinth network which comprised the wall of the lymph spaces. These spaces represented interconnected channels interspersed between epithelial cells. Intraepithelial telocytes shed their secretory vesicles and multivesicular bodies in the intraepithelial lymph space which deliver them to other epithelial and immune cells including chloride, pavement, mucous, rodlet cells and macrophages. Intraepithelial telocytes may also establish contact with epithelial and immune cells either via point or planar contact. Intraepithelial telocyte is previously detected by scanning electron microscopy in the bovine uterine tube. telocyte is located in the basal layer of the epithelial cells and their telopodes extended between epithelial cells (**Abd-Elhafeez and Soliman 2016**)

In the current study, intercellular communication between telocytes chloride cells either by direct contact or paracrine mode revealed that telocyte may have a potential role in osmoregulation. Changing salinity level affect telocytes morphology which in turn influence chloride cells. They undergo hypertrophy, change morphology of mitochondria and increase their number upon elevation of the salinity level. similar results are documented in the Hawaiian goby (*Stenogobius hawaiiensis*). Salinity caused a slight increase in chloride cell number and size (**McCormick, Sundell et al. 2003**). Gill

chloride cells regulate ionic transportation via transport proteins which have a polarized distribution. Three types of transport protein are described in chloride cells; cystic fibrosis transmembrane regulator (CFTR) anion channel, Na,K,2Cl cotransporter (NKCC) and sodium pump (Na,K-ATPase). Expression of Na⁺/K⁺-ATPase, Na⁺/K⁺/2Cl⁻ cotransporter (NKCC) and cystic fibrosis transmembrane conductance regulator (CFTR) in gill chloride cells of the Hawaiian goby (*Stenogobius hawaiiensis*) is variant in freshwater and 20 per thousand and 30 per thousand salinity concentration for 10 days. Na⁺/K⁺-ATPase and NKCC have a basolateral/tubular localization whereas, CFTR expressed in the apical surface of chloride cells. Gill Na⁺/K⁺-ATPase expression is not affected by salinity, while CFTR immunoreactivity increase in salinity (McCormick, Sundell et al. 2003).

In the present study, pavement cells were connected with telocytes. The secretory vesicles could reach the surface epithelium and attached or partially enclosed by the micro-ridges of the pavement cells. These cells were flattened in the control samples and gradually enlarged depending on salinity level till became columnar-shaped in 14 ppt salinity concentration. Micro-ridges became thinner and elongated in 10 and 14 salinity levels. Pavement cells developed surface invaginations at 14 ppt salinity. A similar result is described in the gill epithelia of the Adriatic sturgeon *Acipenser naccarii*. Pavement cells acquired a complex system of microridges on their apical surface during exposure to the hypertonic environment (salinity 35) (CARMONA, GARCIA- GALLEGO et al. 2004). Pavement cells play an important role in gas exchange (Evans, Claiborne et al.

2013). They were rich in proton pumps which regulate acid-based balance (Laurent, Goss et al. 1994; Perry and Fryer 1997).

In the current study, common carp couldn't sustain salinity levels more than 10 ppt. High mortality rate in fish aquarium began in 12 ppt and was markedly increased in 14 ppt. Mangat and Hundal investigated salinity effect on *Cyprinus carp* survival in different seasons. They used 0, 1.5, 3, 6, and 12 ppt for 60 days. All fish are viable and survive at 0 ppt to 6 ppt salinity during all seasons. Only 50% survive at 12 ppt salinity during winter (14.50C-19.00C), while Fish mortality percentage reached 100% during summer (28.00C-37.00C) and autumn (22.50C-30.50C) (Mangat and Hundal 2014).

In the current study, both intraepithelial and stromal telocytes maintained relation with immune cells; particularly macrophages and rodlet cells; either by cellular contact and paracrine signaling. Thus, telocytes could maintain and enhance immune response in different salinity levels. Macrophages increased in size, number and acquired exaggerated phagocytic activities in samples treated with 14 ppt salinity level. They enlarged and became rich in lysosomes and vesicles. Connection of telocytes and macrophage is documented in mouse eye and rat urinary tract (Zheng, Zhu et al. 2012; Luesma, Gherghiceanu et al. 2013). The contact point is identified by an electron-dense nanostructure in the human heart (Gherghiceanu and Popescu 2012). Mouse peritoneal macrophages are activated and secrete macrophage cytokines and enzymes when co-cultured in telocytes conditioned media (Chi, Jiang et al. 2015). Macrophage activity is investigated in relation to salinity level and ration. Phagocytic activities of the

macrophages of black sea bream; *Mylio macrocephalus* Basilewsky juveniles are primarily affected by ration size rather than salinity (**Narnaware, Kelly et al. 2001**).

In the current study, telocytes formed a planar contact with the stem cell, some telopodes extended into the cytoplasm of stem cell. Moreover, telocytes and their telopodes enclosed and established direct contact with the skeletal progenitor cell which began organization of the intracellular myofilament proteins. Relations between telocytes and stem cells is mentioned in the heart (**Popescu, Curici et al. 2015**), lung, skeletal muscle, meninges, and choroid plexus (**Popescu and Nicolescu 2013**). Cardiac telocytes secrete cytokines and growth factors which promote stem cell proliferation and differentiation such as interleukin (IL)-6, IL-2, IL-10, IL-13, VEGF, macrophage inflammatory protein 1 α (MIP-1 α), MIP-2 and MCP-1 and some chemokines like, GRO-KC (**Albulescu, Tanase et al. 2015**).

The present study provided evidence for relations of intraepithelial and stromal telocytes with immature rodlet cells (granular stage). Telocytes exert their effect on rodlet cells either through direct contact or paracrine signaling. Telocytes modified responding to high salinity and subsequently affect rodlet cells. They were increased in number in samples exposed 6, 10 and 14 ppt salinity. Thus, we suggested that telocytes may have a role regulation of the biological activities and in maturation of rodlet cells. Many researches are conducted to investigate the nature and function of rodlet cells. Rodlet cells are thought to act as ion transporting cells and involve in osmoregulation (**Ostrander 2000**). The widely acceptable hypothesis is that rodlet cells participate in

immune response. They are common in helminthic infestations and other noxious agents and considered as a type of eosinophilic granulocyte **(Reite and Evensen 2006; Matisz, Goater et al. 2010)**. Rodlet cells undergo significant changes depending on salinity level. They are increased during reduction of salinity in European sea bass *Dicentrarchus labrax* **(Giari, Manera et al. 2006)**.

In the present study, stromal telocytes established a direct contact with secondary vascular vessels or lymph vessels. The secretory vesicles of the stromal telocytes were secreted in lymph vessel. Lymphatic vessels deliver the inflow from arterial vessels via arterio-arterial anastomoses and drain into the venous circulation **(Kapoor and Bhavna 2004)**. Thus, we suggested that the secondary vascular vessels represented a principal pathway for trafficking of telocytes sections in the blood circulation. Thus, telocytes may exert their paracrine effect on remote tissues and organs. The secondary vascular vessels are implicated in gaseous exchange and ion transportation **(Steffensen and Lomholt 1992)**.

In conclusion, fish body accommodated changing level of the salinity by activation of an adaptive response through cellular communications. Telocytes represented a major component in the communicating system. They regulated the function of a wide variety of cells either by direct contact or paracrine mode. Morphological modification of telocytes in different levels of salinity reflected increase their activities which influence epithelial, immune and stromal effector cells. Intraepithelial telocytes affected chloride, pavement cells, immature rodlet cells and macrophages while stromal telocytes influence

stem cells, skeletal myoblasts and also macrophages and rodlet cells. Thus, telocytes enhanced immunity, osmoregulation in gill lamellar and filament epithelium and regeneration of stromal cells. Thus, the fish could sustain hyperosmotic environments reaching 10 ppt and maintain internal homeostasis.

References

- Abd-Elhafeez, H. H., Mokhtar, D. M., Hassan, A. H. (2016). Effect of Melatonin on Telocytes in the Seminal Vesicle of the Soay Ram: An Immunohistochemical, Ultrastructural and Morphometrical Study. *Cells Tissues Organs*.
- Abd-Elhafeez, H. H., Soliman, S. A. (2016). New Description of Telocyte Sheaths in the Bovine Uterine Tube: An Immunohistochemical and Scanning Microscopic Study. *Cells Tissues Organs*.
- Albulescu, R., Tanase, C., Codrici, E., Popescu, D. I., Cretoiu, S. M., Popescu, L. M. (2015). The secretome of myocardial telocytes modulates the activity of cardiac stem cells. *J Cell Mol Med* 19(8):1783-1794.
- Bancroft, J. D., Layton, C., Suvana, S. K. (2013). *Bancroft's Theory and Practice of Histological Techniques*. Churchill Livingstone; 7 edition.
- Bei, Y., Wang, F., Yang, C., Xiao, J. (2015). Telocytes in regenerative medicine. *J Cell Mol Med* 19(7):1441-1454.
- Cantarero Carmona, I., Luesma Bartolome, M. J., Junquera Escribano, C. (2011). Identification of telocytes in the lamina propria of rat duodenum: transmission electron microscopy. *J Cell Mol Med* 15(1):26-30.
- CARMONA, R., GARCIA- GALLEGU, M., SANZ, A., DOMEZAIN, A., OSTOS-GARRIDO, M. V. (2004). Chloride cells and pavement cells in gill epithelia of *Acipenser naccarii*: ultrastructural modifications in seawater-acclimated specimens. *Journal of Fish Biology* 64:553–566.
- Chi, C., Jiang, X. J., Su, L., Shen, Z. J., Yang, X. J. (2015). In vitro morphology, viability and cytokine secretion of uterine telocyte-activated mouse peritoneal macrophages. *J Cell Mol Med* 19(12):2741-2750.
- Derichs, N. (2013). Targeting a genetic defect: cystic fibrosis transmembrane conductance regulator modulators in cystic fibrosis. *Eur Respir Rev* 22(127):58-65.
- Drumm, B. T., Koh, S. D., Andersson, K. E., Ward, S. M. (2014). Calcium signalling in Cajal-like interstitial cells of the lower urinary tract. *Nat Rev Urol* 11(10):555-564.
- Evans, D. H., Claiborne, J. B., Currie, S. (2013). *The Physiology of Fishes*, Fourth Edition. CRC Press:152.
- Fielder, D. S., Allan, G. L., Pepperall, D., Pankhurst, P. M. (2007). The effects of changes in salinity on osmoregulation and chloride cell morphology of juvenile Australian snapper, *Pagrus auratus*. *Aquaculture* 272:656-666.
- Florkin, M. (2014). *Deuterostomians, Cyclostomes, and Fishes*. Elsevier:166.

492 Gandahi, J. A., Chen, S. F., Yang, P., Bian, X. G., Chen, Q. S. (2012). Ultrastructural
493 identification of interstitial cells of Cajal in hen oviduct. *Poult Sci* 91(6):1410-1417.
494 Gherghiceanu, M., Hinescu, M. E., Andrei, F., Mandache, E., Macarie, C. E., Faussone-
495 Pellegrini, M. S., Popescu, L. M. (2008). Interstitial Cajal-like cells (ICLC) in
496 myocardial sleeves of human pulmonary veins. *J Cell Mol Med* 12(5A):1777-1781.
497 Gherghiceanu, M., Popescu, L. M. (2012). Cardiac telocytes - their junctions and
498 functional implications. *Cell Tissue Res* 348(2):265-279.
499 Giari, L., Manera, M., Simoni, E., Dezfali, B. S. (2006). Changes to chloride and rodlet
500 cells in gills, kidney and intestine of *Dicentrarchus labrax* (L.) exposed to reduced
501 salinities. *Journal of Fish Biology* 69(2):590-600.
502 Heidenhai, N. M. (1896). Noch einmal uber die Darstellung der centralkorper durch
503 Eisenhamatoxylin nebst einigen allgemeinen Bemerkungen uber die Hamatoxylin
504 fabren. *Zeitschrift fur wissenschaftliche Mikroskopie und fur Mikroskopische Technik*
505 13:180.
506 Hutchings, G., Williams, O., Cretoiu, D., Ciontea, S. M. (2009). Myometrial interstitial
507 cells and the coordination of myometrial contractility. *J Cell Mol Med* 13(10):4268-
508 4282.
509 Iino, S., Horiguchi, K. (2006). Interstitial cells of Cajal are involved in neurotransmission
510 in the gastrointestinal tract. *Acta Histochem Cytochem* 39(6):145-153.
511 Kapoor, B. G., Bhavna, K. (2004). *Ichthyology Handbook*. Springer Science & Business
512 Media:289.
513 Laurent, P., Goss, G. G., Perry, S. F. (1994). Proton pumps in fish gill pavement cells?
514 *Arch Int Physiol Biochim Biophys* 102(1):77-79.
515 Lloyd, R. V. (2001). *Morphology Methods: Cell and Molecular Biology Techniques*.
516 Springer Science & Business Media.
517 Luesma, M. J., Gherghiceanu, M., Popescu, L. M. (2013). Telocytes and stem cells in
518 limbus and uvea of mouse eye. *J Cell Mol Med* 17(8):1016-1024.
519 Mangat, H. K., Hundal, S. S. (2014). Salinity tolerance of laboratory reared fingerlings
520 of common carp, *Cyprinus carpio* (Linn.) during different seasons. *International Journal*
521 *of Advanced Research* 2(11):491-496.
522 Marshall, W. S., Grosell, M. (2005). Ion transport, osmoregulation and
523 acid-base balance.
524 . In *Physiology of Fishes 3*, (ed. D. Evans and J. B. Claiborne):pp. 177-230. Boca Raton:
525 CRC Press.
526 Matisz, C. E., Goater, C. P., Bray, D. (2010). Density and maturation of rodlet cells in
527 brain tissue of fathead minnows (*Pimephales promelas*) exposed to trematode cercariae.
528 *Int J Parasitol* 40(3):307-312.
529 McCormick, S. D., Sundell, K., Bjornsson, B. T., Brown, C. L., Hiroi, J. (2003).
530 Influence of salinity on the localization of Na⁺/K⁺-ATPase, Na⁺/K⁺/2Cl⁻ cotransporter
531 (NKCC) and CFTR anion channel in chloride cells of the Hawaiian goby (*Stenogobius*
532 *hawaiiensis*). *J Exp Biol* 206(Pt 24):4575-4583.
533 Mirancea, N. (2016). Telocyte - a particular cell phenotype. Infrastructure, relationships
534 and putative functions. *Rom J Morphol Embryol* 57(1):7-21.
535 Narnaware, Y. K., Kelly, S. P., Woo, N. Y. S. (2001). Effect of salinity and ration size
536 on macrophage phagocytosis in juvenile black sea bream (*Mylio macrocephalus*).
537 *Journal of Applied Ichthyology* 16(2):86 - 88

Nelson, J. S. (2006). *Fishes of the World*. John Wiley & Sons, Inc 4th ed:140.

Ostrander, G. K. (2000). *The Laboratory Fish*. Elsevier:288.

Perry, S. F., Fryer, J. N. (1997). Proton pumps in the fish gill and kidney. *Fish Physiology and Biochemistry* 17(1-6):363-369.

Popescu, L. M., Curici, A., Wang, E., Zhang, H., Hu, S., Gherghiceanu, M. (2015). Telocytes and putative stem cells in ageing human heart. *J Cell Mol Med* 19(1):31-45.

Popescu, L. M., Faussone-Pellegrini, M. S. (2010). TELOCYTES - a case of serendipity: the winding way from Interstitial Cells of Cajal (ICC), via Interstitial Cajal-Like Cells (ICLC) to TELOCYTES. *J Cell Mol Med* 14(4):729-740.

Popescu, L. M., Gherghiceanu, M., Cretoiu, D., Radu, E. (2005). The connective connection: interstitial cells of Cajal (ICC) and ICC-like cells establish synapses with immunoreactive cells. Electron microscope study in situ. *J Cell Mol Med* 9(3):714-730.

Popescu, L. M., Nicolescu, M. I. (2013). Resident stem cells and regenerative therapy, Chapter: Telocytes and stem cells, . Publisher: Oxford: Academic Press, Editors: dos Santos Goldenberg RC, Campos de 7 Carvalho AC:205-231.

Reite, O. B., Evensen, O. (2006). Inflammatory cells of teleostean fish: a review focusing on mast cells/eosinophilic granule cells and rodlet cells. *Fish Shellfish Immunol* 20(2):192-208.

Russell, J. M. (2000). Sodium-potassium-chloride cotransport. *Physiol Rev* 80(1):211-276.

Steffensen, J. F., Lomholt, J. P. (1992). *Fish Physiology XIIA - Circulation*, Chapter: The secondary vascular system. . Publisher: Academic Press, Editors: Hoar, W. S. Randall, D. J. & Farrell, A. P:185-217.

Suhail, M. (2010). Na, K-ATPase: Ubiquitous Multifunctional Transmembrane Protein and its Relevance to Various Pathophysiological Conditions. *J Clin Med Res* 2(1):1-17.

Takaki, M. (2003). Gut pacemaker cells: the interstitial cells of Cajal (ICC). *J Smooth Muscle Res* 39(5):137-161.

Varga, I., Danisovic, L., Kyselovic, J., Gazova, A., Musil, P., Miko, M., Polak, S. (2016). The functional morphology and role of cardiac telocytes in myocardium regeneration. *Can J Physiol Pharmacol*:1-5.

Zheng, Y., Zhu, T., Lin, M., Wu, D., Wang, X. (2012). Telocytes in the urinary system. *J Transl Med* 10:188.

Figure 1: Morphological changes of the intraepithelial telocytes responding to salinity

Semi-thin sections of gill arches and filaments stained with toluidine blue (A-D), methylene blue (E-H) and Heidenhain's Iron-Hx (I-L). A, E, I: showed intraepithelial

580 telocytes in control samples. They had small cell body (arrows) and prominent telopodes
581 (arrowheads). Note the red arrows refer to the basal lamina. B, F, J: showed
582 intraepithelial telocytes during exposure to 6 ppt salinity level. The cell body enlarged
583 and were satellite in shape (arrows). Note telopodes (arrowheads). C, G, K: cell body
584 undergo hypertrophy and became large satellite (arrows) of the intraepithelial telocytes
585 exposed to 10 ppt salinity level. Note telopodes (arrowheads). D, H, L: the cell body
586 (arrows) of the intraepithelial telocytes increased in size in 14 ppt salinity level. Note
587 telopodes (arrowheads).
588

Figure 2: Morphological changes of the stromal telocytes responding to salinity

Semi-thin sections of gill arches and filaments stained with toluidine blue (A-D), methylene blue (E-H) and Heidenhain's Iron-Hx (I-L). A, E, I: showed stromal telocytes in control samples. They had small cell body (arrows) associated with long telopodes (double arrowheads) . Note the secretory vesicles (arrowheads). B, F, J: Some stromal

595 telocytes had enlarged cell body (arrows) during exposure to 6 ppt salinity level. Note
 596 telopodes formed a network (double arrowheads). C, G, K: stromal telocytes (arrows)
 597 exposed to 10 ppt salinity level had an extensive network of telopodes (double
 598 arrowheads) Note the secretory vesicles (arrowheads). D, H, L: stromal telocytes
 599 (arrows), telopodes (double arrowheads). Note the secretory vesicles (arrowheads).
 600

Figure 3: Changes of rodlet and chloride cells responding to salinity stress

Semi-thin sections of gill arches and filaments stained with toluidine blue (A-D), and Heidenhain's Iron-Hx (E-H). A: few number of immature rodlet cells (granular stage) (arrowheads) in control samples. B: Number of immature rodlet cells (arrowheads) increased in branchial epithelium in 6 ppt treated samples. C: Enormous number of rodlet

607 cells (arrowheads) in branchial epithelium exposed to 10 ppt salinity level. D: Massive
 608 number of rodlet cells (arrowheads) in branchial epithelium in 14 ppt treated samples. E:
 609 chloride cell (arrow) in control samples was small contained few mitochondria which
 610 appeared as dark dots by iron Hx stain. F: large chloride cells (arrow) in 6 ppt treated
 611 samples had larger number of mitochondria. G: Chloride cell (arrow) enlarged and
 612 contained abundant mitochondria in 10 ppt treated samples. H: hypertrophy of the
 613 chloride cells with massive mitochondrial content in 14 ppt salinity level.

614

Figure 4: effect of low level (6 ppt) of salinity on intraepithelial telocytes.

Colored ultra-thin sections in gill arches (A-F, H) and filaments (G) of control (A, B) and 6 ppt treated samples (C-H). A, B: intraepithelial telocytes were small, had spindle or satellite-shaped (arrows) and thin telopodes (arrowheads). They rest on the basal lamina which directly opposed on the stratum compactum (st). They organized in a labyrinth

network which comprised the wall of the epithelial lymph spaces. The secretory vesicles (V) were liberated in the lymph spaces. Note podoms (double arrowheads), pavement cells (P). stromal telocytes had longer and thinner telopodes. C-H: Enlargement of the both intraepithelial and stromal telocytes (arrows), thickening of the telopodes (arrowheads). intraepithelial telocytes liberated the secretory vesicles in the lymph spaces and stromal telocytes shed their vesicles in the secondary circulatory vessels (SCV) or lymphatic vessels. Note stratum compactum (st), pavement cells (p). branchial blood vessels contained red blood cells (RBC). Rodlet cells (r) and lysosome-rich macrophages (m) in the lymph spaces. red circles refer to point of contact between telopodes and rodlet cells, green circle refer to contact between telocytes and the macrophage in branchial epithelium.

Figure 5: effect of 10 ppt salinity level on intraepithelial telocytes.

Colored ultra-thin sections in gill arches of 10 ppt treated samples. The prominent feature in 10 ppt level treated samples was increase in size of telocytes either intraepithelial or stromal. A: Basal telocytes were telocytes in the basal layer of the branchial epithelium established a communicating network between the lymph spaces (LS) in which rodlet

cells (R) migrated. Note telopodes (arrowheads), podom (double arrowheads). The sub-epithelial telocytes connected with blood capillaries (bc), secondary circulatory vessels (SCV) and lysosome-rich macrophages (m). Both Basal intraepithelial and sub-epithelial telocytes undergo hypertrophy. B: high magnification of the podom. C: Superficial intraepithelial telocytes established contact with different types of epithelial cells and formed the lymph spaces (LS) where they shed the secretory vesicles (V). Note telopodes (arrowheads), podoms (double arrowheads). chloride cell (c), rodlet cells (R), pavement cell (P), Mucous cell (mu), macrophages (m). D, E: intraepithelial telocytes established planar contact with chloride cell (dashed line). Telocytes shed the numerous secretory vesicles (V) in the lymph space (LS). Note telopodes (arrowheads). F: Stromal telocytes formed a network connected with different types of stromal cells including rodlet cells (R) and macrophages (m). note telopodes (arrowheads), blood capillary (bc).

Figure 6: effect of high salinity level (14 ppt) on intraepithelial telocytes

Colored ultra-thin sections in gill arches of 14 ppt treated samples. A: Superficial intraepithelial telocytes communicated forming a labyrinth network between epithelial cells and established the lymph spaces (LS). Note telopodes (arrowheads), Chloride cells (C), rodlet cells (R), pavement cell (P) with short microvilli. B: Basal telocytes organized

658 a network which enclose the lymph spaces (LS). massive number of macrophages (M) in
659 the lymph spaces. Telopodes (arrowheads). Stromal telocytes established contact with
660 secondary circulatory vessels (SCV), stromal macrophages (m). note podom (double
661 arrowhead). C, D: Intraepithelial telocytes formed a network between the epithelial cells
662 and construct the wall of the lymph spaces (LS). Note Intraepithelial telocytes formed a
663 planar contact (dashed line). The secretory vesicles and multivesicular body (arrow),
664 exosomes (double arrowhead) of the intraepithelial telocytes were shed in the lymph
665 spaces. Note telopodes (arrowheads).
666

Figure 7: Effect of saintly on stromal telocytes

Colored ultra-thin sections in gill arches of control (A, B), 6 ppt (C, D), 10 ppt (E, F), 14 ppt (G, H) treated samples. A, B: telocytes appeared small spindle or satellite shaped (arrows). Telocyte established direct contact with skeletal muscle (double arrowhead). C, D: small spindle-shaped telocytes established homocellular junction (red circle). E: large

673 population of telocytes surrounding the secondary circulatory vessel (SCV). The
674 secretory vesicles (red arrowhead) of telocytes transferred through the endothelial lining
675 of the secondary circulatory vessel (red double arrowheads) and shed in the lumen of the
676 vessel (red arrows). note thick telopodes (black double arrowhead) and slight waving
677 (black arrowhead) of the telopodes shape, macrophages (m). F: subepithelial telocytes (black
678 black arrows), telopodes became enormous (black arrowheads) telopodes undergo
679 thickening in certain areas (double arrowheads). The basal lamina (red arrowheads). Note
680 the basal intraepithelial telocytes gave rise short basal telopodes (red arrows). Note
681 podom of the basal intraepithelial telocyte (double arrowhead), lymph space (LS), rodlet
682 cells (r), macrophages (m). G, H: The most prominent features of high salinity changes
683 were the cell body had an irregular surface (arrows) and wavy telopodes (arrowheads)
684 and thickening of telopodes (double arrowhead) note macrophage (m) enlargement and
685 was filled with lysosomes (m), blood vessel (bv).
686

Figure 8: Increase the secretory activities of the stromal telocytes responding to salinity and liberation of the secretory vesicles in the secondary circulatory vessels.

Colored ultra-thin sections in gill arches treated with 6 ppt (A-D) and 10 ppt (E, F) level of salinity. A, B: enlarged telocytes acquired rounded or triangular shape (arrows). Note large secretory vesicle (V), cell body of stromal telocyte established contact with

693 macrophage (m). C, D: spindle-shaped telocytes (arrows) shed large secretory vesicle
694 (V). note enormous telopodes (arrowheads). E: large number of enlarged subepithelial
695 telocytes (arrows). Podom (double arrow). Some telocytes established contact with the
696 endothelial lining of the secondary circulatory vessel (SCV). Note transferring of the
697 secretory vesicles (red arrow) secreted by telocytes to the cytoplasm of endothelial cells
698 (e). Basal lamina (red arrowheads). intraepithelial telocytes (IT) and their telopodes
699 (arrowheads). F: spindle shaped stromal telocytes (arrows) and their telopodes formed an
700 extensive network (arrows), podom (double arrowhead), the secretory vesicles (V). note
701 the secretory vesicles (red arrow) were observed in the lumen of the secondary
702 circulatory vessel (SCV). endothelial cells (e).
703

Figure 9: Intraepithelial telocytes in relations to rodlet, chloride and pavement cells.

Colored ultra-thin sections in gill arches treated with 6 ppt (B), 10 ppt (A, C, E, F) and 14 ppt (D) level of salinity. A: stromal telocytes (arrow) in direct contact with rodlet cells (arrowheads). Note the telocytes shed the secretory vesicles (V) in vicinity to rodlet cells (R). Macrophage (m). B: rodlet cell (granular stage) had immature rodlet granules

710 (double arrows) which contained an electron dense central core. Telopodes (arrowheads)
 711 in contact with rodlet cell (R). note multi-vesicular body (arrow). C, D: some pavement
 712 cells (P) in 10 and 14 ppt salinity samples had long microvilli (arrows) which deliver the
 713 secretory vesicles (arrowheads) of telocytes at the surface of the branchial epithelium.
 714 surface invaginations or pits (asterisk) in the pavement cells Note secretory vesicles in
 715 lymph space (double arrow), chloride cells (C), rodlet cell (R). E: Intraepithelial telocyte
 716 in planer contact (arrows) with the pavement cells (P) and chloride cell (C). note
 717 Telopodes (arrows), macrophages (m), secretory vesicles (V), Lymph space (LS). F: two
 718 types of mitochondrial rich chloride cells; dark (d) and light (L) were connected to
 719 telocytes. Note telopodes (arrows), secretory vesicles (V).
 720

Figure 10: changes of chloride cells in different salinity levels

Colored ultra-thin sections in gill arches control (A) and treated samples with 6 ppt (B), 10 ppt (C) and 14 ppt (D) level of salinity. A: chloride cell (C) appeared elongated and had few oval mitochondria (arrows). B: chloride cell (C) increase in size and changed the morphology and appeared more elongated. The mitochondria increased in number

and became elongated cigar-shaped (arrows). D: chloride cells (C) enlarged and appeared cuboidal in shape. They had a large number of mitochondria; some of which were elongated cigar-shaped (arrows). Note the secretory vesicles of telocytes in vicinity to chloride cell. E: Chloride cell (C) hypertrophied and appeared oval-shaped. They had a massive mitochondrial content. Some mitochondria were cigar-shaped (arrows). Note telocytes in closed relation to chloride cell. Telopodes (double arrows). The secretory vesicles (V).

Figure 11: stromal telocytes relation with stem cell and skeletal myoblast.

Colored ultra-thin sections in gill arches treated samples with 6 ppt (C-F), 10 ppt (A, B) level of salinity. A, B: several stromal telocytes wrapped around stem cell (S) which contained mitochondria (m). telopodes established a planar contact with stem cells (dashed line). telopodes formed an extensive network (double arrowheads). They secreted

741 vesicles (V), multi-vesicular body (arrowhead). Several telopodes interdigitated with the
 742 stem cells (arrows). note secretory vesicles attached to stem cell (double arrow). Some
 743 telopodes were thickened (asterisk). C-F: telocytes connected with Schwann cell (s) and
 744 macrophage (m). note nerve fiber (arrow), secretory vesicles (V), multi-vesicular body
 745 (arrowhead). Telocytes established direct contact (red circle) with myoblast which
 746 contained ill-organized myofibrils (My).
 747

Figure 12: skeletal muscle fibers undergo hypertrophy in response to salinity

Colored ultra-thin sections in gill arches control (A) and treated samples with 6 ppt (A-C), 10 ppt (D, E) level of salinity. A, B, C: telocytes established multi-point contact (double arrows) with skeletal muscle fiber (m). telopode formed a direct contact with the nerve fiber (n). note epithelium (ep). D: telocytes established direct contact (double

754 arrowheads) with skeletal muscle fibers increased in diameter (m). note telopodes
755 organized an extensive network (arrows). Note blood vessel (bv), secretory vesicles (V).
756