Comparative transcriptome profiling conferring of resistance to *Fusarium oxysporum* infection between resistant and susceptible tomato

Min zhao,^{*, †} Hui-Min Ji,^{*, †} Yin Gao,[†] Xin-Xin Cao,[†] Hui-Yin Mao,[†] Peng Liu,^{‡, 1}
Shou-Qiang Ouyang^{†, §, **, 1}

[†] College of Horticulture and Plant Protection, [‡]Texting Center, [§]Joint International
Research Laboratory of Agriculture and Agri-Product Safety and ^{**} Key Laboratory
of Plant Functional Genomics of the Ministry of Education, Yangzhou University,
48 East Wenhui Road, Yangzhou, Jiangsu, 225009, China

^{*} Contributing equally. ¹ Corresponding author

Running title:

Comparative transcriptome profiling

Key words:

Tomato, *Fusarium oxysporum* f. sp. *Lycopersici*, comparative transcriptome, pathogen, resistance

Corresponding author:

College of Horticulture and Plant Protection, Yangzhou University, 48 East Wenhui Road, Yangzhou, Jiangsu, 225009, China. E-mail: oysq@yzu.edu.cn; and Texting Center, Yangzhou University, 48 East Wenhui Road, Yangzhou, Jiangsu, 225009, China. E-mail: pengliu@yzu.edu.cn

ABASTRCAT

Tomato Fusarium wilt caused by *Fusarium oxysporum* f. sp. *lycopersici* (FOL) is a destructive disease of tomato worldwide which causes severe yield loss of the crops. As exploring gene expression and function approaches constitute an initial point for investigating pathogen– host interaction, we performed a transcriptional analysis to unravel regulated genes in tomato infected by FOL. Differentially expressed genes (DEG) upon inoculation with FOL were presented at twenty-four hours post-inoculation including four treatments: Moneymaker_H₂O, Moneymaker_FOL, Motelle_H₂O and Motelle_FOL. A total of more than 182.6 million high quality clean reads from the four libraries were obtained. A large overlap was found in DEGs between susceptible tomato cultivar Moneymaker and resistant tomato cultivar Motelle. All Gene Ontology terms were mainly classified into catalytic activity, metabolic process and binding. However, Gene Ontology enrichment analysis evidenced specific categories in infected Motelle. Statistics of pathway enrichment of DEGs resulted that the taurine and hypotaurine metabolism, the stibenoid, diarylheptanoid and gingerol biosynthesis, the starch and sucrose metabolism were the top three pathway affected in both groups. Interestingly, plant-pathogen pathway was greatly regulated in Motelle treated with FOL. Combining with qRT-PCR facilitated the identification of regulated pathogenicity associated genes upon infected resistant or susceptible tomato. Our data showed that a coordinated machinery played a critical role in prompting the response, which could help in generating models of mediated resistance responses with assessment of genomic gene expression patterns.

INTRODUCTION

Fusarium oxysporum f. sp. *lycopersici* (hereafter referred to as FOL) is a biotrophic pathogen which is the causal agent of tomato wilt. Accumulating data indicate that

F. oxysporum is a large species complex, with more than 150 host-specific forms causing disease in vegetables, fruit trees, wheat, corn, cotton and ornamental crops (Di Pietro *et al.* 2003; Leslie and Summerell 2006). *F. oxysporum* infects vascular bundles in the plant host, leading to clogged vessels, yellowing of leaves, wilting and finally death of the whole plant. According to their specific pathogenicity to tomato cultivars, three physiological races (Di Pietro *et al.* 2003; Leslie and Summerell 2006; Takken and Rep 2010) of *F. oxysporum* are distinguished (Kawabe *et al.* 2005).

Tomato (*Solanum lycopersicum*) is a worldwide economic crop, and also has been studied as a crucial model plant for studying the genetics and molecular basis of resistance mechanisms. Four plant resistance (*R*) genes have been discovered in cultivated tomato from wild tomato species including the *I* and *I-2* genes from *S. pimpinellifolium*, and the *I-3* and *I-7* gene from *S. pennellii*. Among these four *R* genes, so far, *I-2*, *I-3* and *I-7* have been cloned, encode an NB-LRR protein like most known *R* genes (Ori *et al.* 1997; Simons *et al.* 1998; Kawabe *et al.* 2005; Catanzariti *et al.* 2015; Gonzalez-Cendales *et al.* 2016). Previous works have demonstrated that the *I-2* and *I-3* gene confers resistance to race 2 and race 3 strains of FOL, respectively (Simons *et al.* 1998; Catanzariti *et al.* 2015). The *I-2* locus encodes an *R* protein that recognizes the *avr2* gene product from *F. oxysporum* (race 2) (Houterman *et al.* 2009). The *I-3* encodes an S-receptor-like kinase (SRLK) genes that confers *Avr3*-dependent resistance to FOL (race 3) (Catanzariti *et al.* 2015). Previously, two near-isogenic tomato cultivars susceptible Moneymaker (*i-2/i-2*) and resistant Motelle (*I-2/I-2*) were recruited to study the interaction between tomato and FOL (Ouyang *et al.* 2014). The genotypes of these two tomato cultivars are for *I-2* and respond to FOL infection (Di Pietro and Roncero 1998; De Ilarduya *et al.* 2001; Yu and Zou 2008). We unveiled the microRNA diversifications responding

to FOL infection in tomato by high-throughput RNA sequencing (RNA-seq) approach (Ouyang *et al.* 2014). Basically, transcriptome analysis is a very important tool to discover the molecular basis of plant-pathogen interaction globally, allowing dissection of the pattern of pathogen activities and molecular repertoires available for defense responses in host plant. By taking advantage of RNA-seq technology, a few of transcriptome profiling studies of plants following inoculation with *Fusarium* fungus have been reported, including studies in banana (Guo *et al.* 2014), cabbage (Xing *et al.* 2016), watermelon (Liu *et al.* 2015), mango (Liu *et al.* 2016), and *Arabidopsis* (Chen *et al.* 2014; Gupta *et al.* 2014). Upon to pathogens infection, plants activate a few of defense responses to resistant diseases caused by according pathogens. Resistance response may associated with hypersensitive reaction (HR), structural alterations, reactive oxygen species (ROS) accumulation, synthesis of secondary metabolites and defense molecules (Park *et al.* 2003; Shah 2003; Ros *et al.* 2004).

The objects of this study were to determine the transcript profile between susceptible Moneymaker and resistant Motelle tomato plants in response to FOL infection and to reveal genes underlying the innate immune response against the fungal pathogen. To achieve these goals, we performed transcriptome analysis using RNA-seq approach. In addition to genes known to response to pathogen infection, our results also uncovered a bunch of novel fungal pathogen-responsive genes for further functional characterization, and provided a broader view of the dynamics of tomato defense transcriptome triggered by FOL infection.

MATERIALS and METHODS

Tomato materials and fungal culture

Two tomato near-isogenic cultivars (cv.) Motelle (*I-2/I-2*) and Moneymaker (*i-2/i-2*) that exhibit different susceptibilities to the root pathogen FOL were used for plant infection and libraries construction. Profiling experiments were performed on two-week-old tomato seedlings grown at 25°C with a 16/8-h light/dark cycle. The wild-type *Fusarium oxysporum* f. sp. *lycopersici* strain used for all experiments is FGSC 9935 (also referred to as FOL 4287 or NRRL 34936). Two-week-old tomato seedlings were removed from soil and roots incubated in a solution of FOL conidia at a concentration of 1×10^8 /ml for 30 min. Control tomato plants were treated with water. Plants were then replanted in soil and maintained in a growth chamber at 25°C for 24 h with constant light. Plants were removed from soil, and roots were rinsed and excised, then immediately frozen in liquid nitrogen and stored at -80°C.

RNA extraction, library preparation, and sequencing

Total RNA was isolated from roots using TRIzol® Reagent (#15596026, Life Technologies, CA, USA) according to the manufacturer's recommendations. After the total RNA extraction and DNase I treatment, magnetic beads with Oligo (dT) were used to isolate mRNA. Mixed with the fragmentation buffer, the mRNA was sheared into short fragments. Then cDNA was synthesized using the mRNA fragments as templates. cDNAs were purified and resolved with EB buffer for end reparation and single nucleotide A (adenine) addition followed by adding adapters to cDNAs. After agarose gel electrophoresis, the suitable cDNAs were selected for the PCR amplification as templates. During the quality control (QC) steps, Agilent 2100 Bioanalyzer and ABI StepOnePlus Real-Time PCR System were used in quantification and qualification of the sample library. The libraries were sequenced using Illumina HiSeq™ 2000.

RNA-seq analysis, normalization of sequence reads and identification of differentially expressed genes (DEGs)

Primary sequencing data that produced by Illumina HiSeqTM 2000, called as raw reads, were subjected to QC. After QC, raw reads were filtered into clean reads which were aligned to the reference sequences as described by previous report. (Trapnell *et al.* 2012). All sequence reads were trimmed to remove the low-quality sequences. The trimmed reads were then aligned to the tomato reference genome downloaded from the Sol Genomics Network using Bowtie v0.12.5 (Langmead *et al.* 2009) and TopHat v2.0.0 (Trapnell *et al.* 2009; Trapnell *et al.* 2012) with default settings. Cufflinks v0.9.3 (Trapnell *et al.* 2010) was used to calculate transcript abundance based on fragments per kilo base of transcript permillion fragments mapped (FPKM) using all parameters on default settings. The transcript was considered as expressed when the FPKM value was greater than 0.1 and the lower boundary for FPKM value was greater than zero at 95% confidence interval. Once the transcript abundance was calculated for individual sample files using Cufflinks, the output files were further merged pairwise for each comparison (in vitro comparison between two populations, in planta comparison between two populations and in planta versus *in vitro* for each population) using Cufflinks utility program-Cuffmerge (Trapnell *et al.* 2012). The pairwise comparisons of gene expression profiles between the two populations were done using the Cuffdiff program of the Cufflinks version 1.3.0 (Trapnell *et al.* 2010). The genes were considered significantly differentially expressed if Log₂ FPKM (fold change) was ≥ 1.0 and false discovery rate (FDR, the adjusted P value) was < 0.01 . The q-value which was a positive FDR analogue of the p-value was set to < 0.01 (Storey and Tibshirani 2003).

Functional categorization of DEGs.

DEGs were functionally categorized online for all pairwise comparisons according to the Munich Information Center for Protein Sequences (MIPS) functional

catalogue (Ruepp *et al.* 2004). The functional categories and subcategories were regarded as enriched in the genome if an enrichment P- and FDR- value was below <0.05. The Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analyses were performed using interface on blast2GO (Blast2GO v2.6.0, <http://www.blast2go.com/b2ghome>) for all DEGs to identify gene enrichment on a specific pathway.

Gene Ontology (GO) and pathway enrichment analysis

Gene Ontology (GO) and pathway enrichment were performed using DAVID software (Smyth 2005). Graphs of the top 20 enriched GO terms for each library were generated using the Cytoscape Enrichment Map plugin (Smoot *et al.* 2011; Merico *et al.* 2010).

Quantitative real time-PCR (qRT-PCR) analysis.

qRT-PCR analysis was performed according to our previous protocol (Ouyang *et al.* 2014). The reverse transcription reaction was done on 1 µg of total RNA using the SMART MMLV Reverse Transcriptase (Takara, Mountain View, CA). cDNA was diluted two times and used as template for quantitative RT-PCR, which was performed with the CFX96 real-time PCR system (Bio-Rad, Hercules, California, USA). Primers used for qRT-PCR were designed from 3-UTR for individual gene. For each cDNA sample, three replications were performed. Each reaction mixture (20 µL) contained 1 µL of cDNA template, 10 µL of SYBR1 Green PCR Master Mix (Applied Biosystems, Foster, CA) and 1 µL of each primer (10 µM). Relative expression levels of genes were normalized using the 18S rRNA as internal control, and were calculated as the fold change by comparison between in water treated and in FOL treated samples.

Statistical analyses

All data in this study were subjected to ANOVA analysis or Student's t-test analysis using SPSS 11.5 (SPSS Company, Chicago, IL).

RESULTS

General features of Moneymaker and Motelle transcriptomes

We investigated transcriptomes in roots of tomato during infection with the tomato wilt disease fungus FOL through construction of transcript libraries and RNA-seq. By taking advantage of two near-isogenic cultivars that show differential interaction with FOL – Moneymaker (susceptible) and Motelle (resistant), we generated four libraries including: Moneymaker treated with water (MM_H₂O), Moneymaker treated with FOL (MM_Foxy), Motelle treated with water (Mot_H₂O) and Motelle treated with FOL (Mot_Foxy). Using Illumina sequencing, we obtained a total of more than 182.6 million high quality clean reads from the four libraries. Of these, 45,616,330 from MM_H₂O, 45,635,428 from MM_Foxy, 45,680,034 from Mot_H₂O, and 45,661,734 from Mot_Foxy. The number of expressed transcripts were 22,796 and 22,639 from MM_H₂O and MM_Foxy library respectively, and 22,825 and 22,725 from Mot_H₂O and Mot_Foxy library respectively (Table 1). Sequence reads presented reasonable correlation between related two populations ($t > 0.95$, $p = 0.29$) (Figure 1). A number of 21,808 and 21,753 genes were co-expressed between MM_H₂O and MM_Foxy library and Mot_H₂O and Mot_Foxy library, respectively. The co-expressed genes increased slightly to 21,887 between MM_Foxy and Mot_Foxy library (Figure 2). The scatter of all expressed genes of each pair were presented in figure 3.

Of the sequence reads from MM_H₂O and MM_Foxy library, 75.49% and 67.89% were mapped to the reference genome of tomato, respectively. For Mot_H₂O and Mot_Foxy library, 75.87% and 70.46% were aligned to the reference genome of

Table 1 Summary of sequence reads (in millions) from four libraries.

Sample name	Clean reads	Genome map rate	Gene map rate	Expressed gene	Novel gene	Alternative splicing	SNP	Indel
MM_H ₂ O	45,616,330	75.49%	76.87%	22,796	775	32,482	14,046	1,232
MM_Foxy	45,635,428	67.89%	68.47%	22,639	761	32,689	13,371	1,228
Mot_H ₂ O	45,680,034	75.87%	75.92%	22,825	795	33,706	17,286	1,506
Mot_Foxy	45,661,734	70.46%	71.26%	22,725	705	32,965	16,409	1,360

Figure 1 Correlations value between each two libraries.

tomato, respectively. Among the reads mapped to the tomato genome, perfect match reads were 63.00% and 55.52% for MM_H₂O and MM_Foxy library respectively, and 62.61% and 57.41% for Mot_H₂O and Mot_Foxy library respectively (Table 2).

Figure 2 Venn Chart of Co-expressed Genes between MM_H₂O and MM_Foxy, Mot_H₂O and Mot_Foxy, and MM_Foxy and MM_Foxy.

Figure 3 Scatter chart of all expressed genes of each pair between MM_H₂O and MM_Foxy, Mot_H₂O and Mot_Foxy, and MM_Foxy and MM_Foxy.

Analysis of differentially expressed genes (DEGs) and functional classification of DEGs by gene ontology (GO) enrichment analysis

After expression levels Fragments Per Kilobase of exon model per Million mapped reads (FPKM) for each gene were calculated. Differentially expressed genes (DEGs) were defined as genes with fold-change > 2 fold and P_{adjust} value < 0.05. A total number of 3,942 and 4,168 genes showed significantly differential expression in MM_H₂O vs. MM_Foxy library and Mot_H₂O vs. Mot_Foxy library, respectively.

247 **Table 2** Mapping statistics of four libraries.

		MM_H ₂ O	MM_Foxy	Mot_H ₂ O	Mot_Foxy
Total mapped reads	Map to genome	75.49%	67.89%	75.87%	70.46%
	Map to gene	76.87%	68.47%	75.92%	71.26%
Perfect match	Map to genome	63.00%	55.52%	62.61%	57.41%
	Map to gene	66.21%	57.68%	64.46%	59.64%
Mismatch	Map to genome	12.49%	12.38%	13.26%	13.05%
	Map to gene	10.66%	10.79%	11.46%	11.63%
Unique match	Map to genome	74.32%	66.85%	74.79%	69.46%
	Map to gene	69.47%	61.75	68.83%	64.43%
Multi-position match	Map to genome	1.17%	1.04%	1.07%	0.99%
	Map to gene	7.40%	6.72%	7.09%	6.83%
Total Unmapped reads	Map to genome	24.51%	32.11%	24.13%	29.54%
	Map to gene	23.13%	31.53%	24.08%	28.74%

248

249 Among these DEGs, 221/219 genes were down-regulated, and 261/415 genes were
250 up-regulated (MM_H₂O vs. MM_Foxy/ Mot_H₂O vs. Mot_Foxy) (Figure 4). A
251 majority of these DEGs were overlapped in both water and FOL treated two tomato
252 cultivars.

253 To explore the distribution of DEGs, gene ontology (GO) enrichment analyses
254 were conducted based on these DEGs. A total of 530 and 769 GO terms were
255 discovered in MM_H₂O vs. MM_Foxy and Mot_H₂O vs. Mot_Foxy library,
256 respectively. For both libraries, all GO terms were assigned to three groups including
257 the biological process, the cellular component and the molecular. All GO terms were
258 mainly classified into catalytic activity (104 out of 530 in MM_H₂O vs. MM_Foxy
259 library, and 141 out of 769 in Mot_H₂O vs. Mot_Foxy library, (the same define in

Figure 4 Statistics of Differentially Expressed Genes (DEGs).

Figure 5 Gene Ontology Analysis of DEGs. The results were basically summarized into three main categories: biological processes, cellular components, and molecular functions. All statistically significant genes from four libraries were assigned to GO terms. A MM_H2O vs MM_Foxy. B Mot_H2O vs Mot_Foxy. C MM_Foxy vs Mot_Foxy.

the following text), metabolic process (81 out of 530, and 118 out of 769), and binding (72 out of 530, and 104 out of 769). For the class of response to stimulus, however, no significant change was presented between these two libraries (31 out of 530, and 36 out of 769) (Figure 5).

To further understand the biological functions, top 20 statistics of pathway enrichment of DEGs were performed to discover the affection of FOL to host plant. Total of 356 and 469 DEGs from MM_H₂O vs. MM_Foxy library and Mot_H₂O vs. Mot_Foxy library respective were annotated for pathway enrichment. The taurine and hypotaurine metabolism, the stilbenoid, diarylheptanoid and gingerol biosynthesis, the starch and sucrose metabolism were the top three pathway affected in both groups, but in different ranking. The metabolic pathway was the most abundant DEGs in both groups with 122 out of 356 and 148 out of 469 DEGs in MM_H₂O vs. MM_Foxy library and Mot_H₂O vs. Mot_Foxy library, respectively (Figure 6). Be worth mentioning, plant-pathogen pathway was ranked in the 24th (24 out of 356 DEGs) in MM_H₂O vs. MM_Foxy library, however, it was presented in the 8th (40 out of 469 DEGs) in Mot_H₂O vs. Mot_Foxy library (Figure 6). When compared with Mot_H₂O vs. Mot_Foxy library, 19 DEGs were presented in MM_H₂O vs. MM_Foxy library.

Figure 6 Top 20 of pathway enrichment analysis of DEGs. A MM_H₂O vs MM_Foxy. B Mot_H₂O vs Mot_Foxy. C MM_Foxy vs Mot_Foxy. Blue circle in B and C highlighted the plant-pathogen interaction.

Expression profiles of DEGs selected in plant-pathogen interaction by qRT-PCR

To verify the DEGs in plant-pathogen interaction pathway, ten disease related DEGs were selected to characterize the gene expression profiles between water and FOL treated Moneymaker and Motelle by qRT-PCR using primers listed in table 3. These DEGs were Solyc00g174330 (Pathogenesis related protein PR-1), Solyc09g007010 (Pathogenesis related protein PR-1), Solyc02g084890 (Cc-nbs-lrr, resistance protein), Solyc07g054120 (LRR receptor-like serine/threonine-protein kinase, RLP), Solyc10g011910 (WRKY transcription factor 23), Solyc03g124110 (Pathogenesis-related transcriptional factor and ERF, DNA-binding), Solyc03g026280 (Pathogenesis-related transcriptional factor and ERF, DNA-binding), Solyc12g009240 (Pathogenesis-related transcriptional factor and ERF, DNA-binding) and Solyc02g080070 (RLK, Receptor like protein, putative resistance protein with an antifungal domain).

The results of qRT-PCR showed the similar pattern with sequencing results with minute difference. Among these DEGs, Solyc00g174330, Solyc10g011910, Solyc03g124110, Solyc02g084890 and Solyc12g009240 were induced greatly in Motelle affected by FOL, however, no significant changes were present in Moneymaker between water and FOL treatment (Figure 7).

DISCUSSION

In this study, we explored the availability of near-isogenic susceptible and resistant cultivars of tomato infected by FOL to uncover a global transcriptomic profile of tomato-FOL interaction using Illumina sequencing. The components of plant responding to pathogen challenging may lead to understand the underlying defense mechanisms. Plants have evolved a complicate defense system against pathogens

including cascade signaling activation, the regulation of gene expression, synthesis of defensive metabolites as well as hormone balancing (Mukhtar *et al.* 2011; Andolfo *et al.* 2014). So far, by taking advantage of high-throughput RNAsequencing (RNA-seq) approach, a few of transcriptome studies discovering the *F. oxysporum*-host interaction have been reported in plants such as banana,

Table 3 Annotation of pathogenesis related genes and primers used for qRT-PCR in this study

Gene ID	Sequence (5' – 3')	Gene annotation
Solyc00g174330	F: AAGTGGATCGGATCGATCTC R: GAACCTAAGCACGATACCATG	Pathogenesis related protein PR-1
Solyc09g007010	F: ATTTACGTAAGGACGGTTC R: GGACTCAAGATCTCTGATCAAG	Pathogenesis related protein PR-1
Solyc02g084890	F: AAGAGCACTATGGACAACGC R: GCTGTGACTTTTCATGCCCA	Cc-nbs-lrr, resistance protein
Solyc07g054120	F: CCGATTAGGAGAAAGGTCTG R: CAGAGAAGATTAGCATGGCC	LRR receptor-like serine/threonine-protein kinase, RLP
Solyc10g011910	F: ACTGATAAAGGACACGTGGC R: TCTTCCAATCTCTAACGTAC	WRKY transcription factor 23
Solyc03g124110	F: TTTTACCCCGTACCCAACCTC R: GGCGGTGATTTGAGTGTTAC	Pathogenesis-related transcriptional factor and ERF, DNA-binding
Solyc03g026280	F: TTATGGGGATTCAATGG R: TGGTGGCACTACTATCTACC	Pathogenesis-related transcriptional factor and ERF, DNA-binding
Solyc12g009240	F: ACACACGGTTTACGCTACTC R: GACGATGCAAAATATTGTTGC	Pathogenesis-related transcriptional factor and ERF, DNA-binding
Solyc02g080070	F: ACGTGTATTCTAGCTAGCAG R: ATGGAATGGAAGGAGGTTCC	RLK, Receptor like protein, putative resistance protein with an antifungal domain

watermelon, mango and Arabidopsis (Chen *et al.* 2014; Guo *et al.* 2014; Gupta *et al.* 2014; Liu *et al.* 2015; Liu *et al.* 2016; Xing *et al.* 2016), shedding light on the cross-talking among different signaling pathways involving in plant-pathogen interaction.

When plant is attacked by pathogen, the host reprograms metabolism balance between development and the resources to support defense to pathogen, involving biological process, cellular components and molecular functions (Mithöfer and Boland 2012). Upon to our results, the tomato–FOL interaction basically followed

Figure 7 Validation of DEGs selected in plant-pathogen interaction pathway by qRT-PCR. Total tomato root RNA was used for qRT-PCR with gene-specific primers. Each column represents an average of three replicates, and error bars represent the standard error of means.

the typical reaction of biotrophic phase pathogens infection. Gene Ontology analysis of DEGs between two tomato cultivars revealed specific enriched categories in both interactions. In resistant tomato cultivar Motelle, cellular component organization or

biogenesis, signaling, molecular transducer activity, and signal transducer activity were evidenced when compared to susceptible tomato cultivar Moneymaker. Among them, cellular component organization or biogenesis was a critical metabolic activities required by plants to survive under fungus-inflicted stresses (Paul *et al.* 2011). Generally, the genes involved in GO analysis presented in Motelle more than in Moneymaker upon FOL infection which was due to different resistant cultivar.

Two main mechanisms, pathogen-associated molecular patterns (PAMPs) or microbe-associated molecular patterns (MAMPs) (Boller and Felix 2009; Cui *et al.* 2014; Yang and Huang 2014) and the adaptive immune system composed of resistance (R) genes (Dangl and Jones 2001; Van Ooijen *et al.* 2007; Marone *et al.* 2013), are involved in plant responses to pathogenic microorganisms in plant. At least five different classes of R genes have been classified based on functional domain (Van Ooijen *et al.* 2007). Among these classes, a nucleotide-binding site (NBS) and leucine-rich repeats (LRRs) (NBS-LRR) is known as the most numerous R-gene class (Dangl and Jones 2001). Previously, we reported that tomato endogenous microRNA slmiR482f and slmiR5300 conferred to tomato wilt disease resistance. Two predicted mRNA targets each of slmiR482f and slmiR5300, encoded protein with full or partial NBS domains respectively, confirmed to exhibit function of resistance to FOL (Ouyang *et al.* 2014). A few of investments have been demonstrated that NB-LRR proteins are required for the recognition of a specific Avr and disease resistance in several plant species, including rice, *N. benthamiana*, Arabidopsis and wheat (Sinapidou *et al.* 2004; Peart *et al.* 2005; Lee *et al.* 2009; Loutre *et al.* 2009; Narusaka *et al.* 2009; Okuyama *et al.* 2011; Ouyang *et al.* 2014). The corresponding R genes were located tightly in physical linkage. However, in spite this physical linkage, not all these R gene pairs were homologous (Sinapidou *et al.* 2004; Lee *et al.* 2009). We found that genes related to plant-pathogen

interaction were activated in resistant cultivar Moltelle once treated with FOL. Our qRT-PCR results demonstrated that some of these genes were up-regulated specifically in Motelle but not in Moneymaker. In particular, most of these genes were NBS-LRR or like genes which may imply that NBS-LRR genes played a critical role in resistance to FOL in tomato. Investigation of differentially regulated pathogen-induced NBS-LRR genes could lead to uncover the specific modulation patterns upon FOL infection in tomato.

To conclude, our abroad genome transcriptome RNA-seq data provided a comprehensive overview of the gene expression profiles between two different tomato cultivars Moneymaker and Motelle treated with FOL. Our results will facilitate further analysis of putative molecular mechanism of resistance in tomato upon to FOL, which eventually lead to improvement of Fusarium wilt disease resistance in tomato. It remains to be determined whether or how these candidate pathogen-related genes confirmed by qRT-PCR are overexpressed/knockouted in Moneymaker/Motelle plant to reveal the Fusarium wilt disease resistance. In this scenario, we would expect that overexpressing of these candidate pathogen-related genes will enhance resistance to *F. oxysporum* and would therefore develop a useful molecular tool to uncover functional roles for the increasing number of discovered genes in tomato.

ACKNOWLEDGMENTS

We gratefully acknowledge support from JSSF: BK20161330, Jiangsu Province, China.

LITERATURE CITED

400 Andolfo, G., F. Ferriello, L. Tardella, A. Ferrarini, L. Sigillo, *et al.*, 2014 Tomato
401 genome-wide transcriptional responses to *Fusarium* wilt and Tomato Mosaic Virus
402 PLoS One 9: e94963.

403 Boller, T., and G. Felix, 2009 A renaissance of elicitors: perception of microbe-
404 associated molecular patterns and danger signals by pattern-recognition receptors.
405 Annu. Rev. Plant Biol. 60: 379-406.

406 Catanzariti, A.M., G.T. Lim, and D.A. Jones, 2015 The tomato I-3 gene: a novel
407 gene for resistance to *Fusarium* wilt disease. New Phytol. 207:106-118.

408 Chen, Y.C., C.L. Wong, F. Muzzi, I. Vlaardingerbroek, B.N. Kidd, *et al.*, 2014 Root
409 defense analysis against *Fusarium oxysporum* reveals new regulators to confer
410 resistance. Sci Rep. 4: 5584.

411 Cui, H., K. Tsuda, and J.E. Parker, 2014 Effector-Triggered Immunity: From
412 Pathogen Perception to Robust Defense. Annu. Rev. Plant Biol. 66: 487-511.

413 Dangl, J.L., and J.D.G. Jones, 2001 Plant pathogens and integrated defence
414 responses to infection. Nature 411: 826–833.

415 Dangl, J.L., J.D.G. and Jones, 2001 Plant pathogens and integrated defense
416 responses to infection. Nature 411: 826–833.

417 De Ilarduya, O.M., A.E. Moore, and I. Kaloshian, 2001 The tomato Rme1 locus is
418 required for Mi-1-mediated resistance to root-knot nematodes and the potato aphid.
419 Plant J 27: 417–425.

420 Di Pietro, A., and M.I. Roncero, 1998 Cloning, expression, and role in pathogenicity
421 of pg1 encoding the major extracellular endopolygalacturonase of the vascular wilt
422 pathogen *Fusarium oxysporum*. Mol Plant Microbe Interact 11: 91–98.

423 Di Pietro, A., M.P. Madrid, Z. Caracuel, J. Delgado-Jarana, and M.I.G. Roncero,
424 2003 *Fusarium oxysporum*: exploring the molecular arsenal of a vascular wilt fungus.
425 *Molecular Plant Pathology*. 4: 315–325.

426 Gonzalez-Cendales, Y., A.M. Catanzariti, B. Baker, D.J. Mcgrath, and D.A. Jones,
427 2016 Identification of I-7 expands the repertoire of genes for resistance to *Fusarium*
428 wilt in tomato to three resistance gene classes. *Mol Plant Pathol*. 17: 448-463.

429 Guo, L., L. Han, L. Yang, H. Zeng, D. Fan, *et al.*, 2014 Genome and transcriptome
430 analysis of the fungal pathogen *Fusarium oxysporum* f. sp. *cubense* causing banana
431 vascular wilt disease. *PLoS One* 9: e95543.

432 Gupta, K.J., L.A. Mur, and Y. Brotman, 2014 *Trichoderma asperelloides* suppresses
433 nitric oxide generation elicited by *Fusarium oxysporum* in *Arabidopsis* roots. *Mol*
434 *Plant Microbe Interact* 27: 307-314.

435 Houterman, P.M., L. Ma, G. van Ooijen, M.J. de Vroomen, B.J. Cornelissen, *et al.*,
436 2009 The effector protein Avr2 of the xylem-colonizing fungus *Fusarium*
437 *oxysporum* activates the tomato resistance protein I-2 intracellularly. *Plant J* 58:
438 970–978.

439 Kawabe, M., Y. Kobayashi, G. Okada, I. Yamaguchi, T. Teraoka *et al.*, 2005 Three
440 evolutionary lineages of tomato wilt pathogen, *Fusarium oxysporum* f. sp.
441 *lycopersici*, based on sequences of IGS, MAT1, and pg1, are each composed of
442 isolates of a single mating type and a single or closely related vegetative
443 compatibility group. *J. Gen. Plant Pathol*. 71: 263–272.

444 Langmead, B., C. Trapnell, M. Pop, and S.L. Salzberg, 2009 Ultrafast and memory-
445 efficient alignment of short DNA sequences to the human genome. *Genome biology*
446 10: R25.

447 Lee, S.K., M.Y. Song, Y.S. Seo, H.K. Kim, S. Ko, *et al.*, 2009 Rice Pi5-mediated
448 resistance to *Magnaporthe oryzae* requires the presence of two coiled-
449 coilnucleotide-binding-leucine-rich repeat genes. *Genetics* 181: 1627–1638.

450 Leslie, J.F. and B.A. Summerell, 2006 *The Fusarium Laboratory Manual*. Ames, IA:
451 Blackwell Publishing. 388 pages p.

452 Liu, F., J.B. Wu, R.L. Zhan, and X.C. Ou, 2016 Transcription Profiling Analysis of
453 Mango-*Fusarium mangiferae* Interaction. *Front Microbiol.* 7: 1443.

454 Liu, N., J. Yang, X. Fu, L. Zhang, K. Tang, *et al.*, 2015 Genome-wide identification
455 and comparative analysis of grafting-responsive mRNA in watermelon grafted onto
456 bottle gourd and squash rootstocks by high-throughput sequencing. *Mol Genet*
457 *Genomics* 291: 621-33.

458 Loutre, C., T. Wicker, S. Travella, P. Galli, S. Scofield, *et al.*, 2009 Two different
459 CC-NBS-LRR genes are required for Lr10-mediated leaf rust resistance in tetraploid
460 and hexaploid wheat. *Plant J* 60: 1043–1054.

461 Marone, D., M.A. Russo, G. Laidò, A.M. De Leonardis, and A.M. Mastrangelo,
462 2013 Plant nucleotide binding site-leucine-rich repeat (NBS-LRR) genes: active
463 guardians in host defense responses. *Int J Mol Sci.* 14: 7302-7326.

464 Merico, D., R. Isserlin, O. Stueker, A. Emili, and G.D. Bader, 2010 Enrichment Map:
465 A Network-Based Method for Gene-Set Enrichment Visualization and Interpretation.
466 *PLoS ONE* 5: e13984.

467 Mithöfer, A., and W. Boland, 2012 Plant defense against herbivores: chemical
468 aspects. *Annual Review of Plant Biology* 63: 431–50.

- 469 Mukhtar, M.S., A.R. Carvunis, M. Dreze, P. Epple, J. Steinbrenner, *et al.*,
 470 2011 Independently evolved virulence effectors converge onto hubs in a plant
 471 immune system network. *Science* 333: 596-601.
- 472 Narusaka, M., K. Shirasu, Y. Noutoshi, Y. Kubo, T. Shiraishi, *et al.*, 2009 RRS1 and
 473 RPS4 provide a dual Resistance-gene system against fungal and bacterial pathogens.
 474 *Plant J* 60: 218–226.
- 475 Okuyama, Y., H. Kanzaki, A. Abe, K. Yoshida, M. Tamiru, *et al.*, 2011 A
 476 multifaceted genomics approach allows the isolation of the rice Pia-blast resistance
 477 gene consisting of two adjacent NBS-LRR protein genes. *Plant J* 66: 467–479.
- 478 Ori, N., Y. Eshed, I. Paran, G. Presting, D. Aviv, *et al.*, 1997 The I2C family from
 479 the wilt disease resistance locus I2 belongs to the nucleotide binding, leucine-rich
 480 repeat superfamily of plant resistance genes. *Plant Cell* 9: 521–532.
- 481 Ouyang, S., G. Park, H. Atamian, C. Han, J. Stajich, *et al.*, 2014 MicroRNAs
 482 suppress NB domain genes in tomato that confer resistance to *Fusarium oxysporum*.
 483 *PLoS Pathogens*. 10: e1004464.
- 484 Park, Y.S., H.J. Min, S.H. Ryang. K.J. Oh, J.S. Cha, *et al.*, 2003 Characterization of
 485 salicylic acid-induced genes in Chinese cabbage. *Plant Cell Reports* 21: 1027-1034.
- 486 Paul, J.Y., D.K. Becker, M.B. Dickman, R.M. Harding, H.K. Khanna, *et al.*, 2011
 487 Apoptosis-related genes confer resistance to *Fusarium* wilt in transgenic ‘Lady
 488 Finger’ bananas. *Journal of Plant Biotechnology* 9: 1141–1148.
- 489 Peart, J.R., P. Mestre, R. Lu, I. Malcuit, and D.C. Baulcombe, 2005 NRG1, a CC-
 490 NBLRR protein, together with N, a TIR-NB-LRR protein, mediates resistance
 491 against tobacco mosaic virus. *Curr Biol*. 15: 968–973.

492 Ros B, Tummler F, Wenzel G (2004) Analysis of differentially expressed genes in
493 a susceptible and moderately resistant potato cultivar upon *Phytophthora infestans*
494 infection. *Molecular Plant Pathology* 5: 191-201.

495 Ruepp, A., A, Zollner, D, Maier, K, Albermann, J, Hani, *et al.*, 2004 The FunCat, a
496 functional annotation scheme for systematic classification of proteins from whole
497 genomes. *Nucleic acids research* 32: 5539-5545.

498 Shah, J, 2003 The salicylic acid loop in plant defense. *Current Opinion in Plant*
499 *Biology* 6: 365-371.

500 Simons, G., J. Groenendijk, J. Wijbrandi, M. Reijans, J. Groenen, *et al.*, 1998
501 Dissection of the *Fusarium I2* gene cluster in tomato reveals six homologs and one
502 active gene copy. *Plant Cell* 10: 1055–1068.

503 Sinapidou, E., K. Williams, L. Nott, S. Bahkt, M. Tor, *et al.*, 2004 Two
504 TIR:NB:LRR genes are required to specify resistance to *Peronospora parasitica*
505 isolate Cala2 in Arabidopsis. *Plant J* 38: 898–909.

506 Smoot, M., K. Ono, J, Ruscheinski, P, Wang, and T. Ideker, 2011 Cytoscape 2.8:
507 new features for data integration and network visualization. *Bioinformatics* 27: 431–
508 432.

509 Smyth, G.K., 2005 Limma: linear models for microarray data. In: Gentleman R,
510 Carey V, editors. *Bioinformatics and Computational Biology Solutions Using R and*
511 *Bioconductor*. New York: Springer 397–420.

512 Storey, J.D., and R. Tibshirani, 2003 Statistical significance for genome wide studies.
513 *Proceedings of the National Academy of Sciences* 100: 9440-9445.

514 Takken, F. and M. Rep, 2010 The arms race between tomato and *Fusarium*
515 *oxysporum*. *Mol Plant Pathol.* 11: 309–314.

516 Trapnell, C., A. Roberts, L. Goff, G. Pertea, D. Kim, *et al.*, 2012 Differential gene
517 and transcript expression analysis of RNA-seq experiments with TopHat and
518 Cufflinks. *Nature protocols* 7: 562-578

519 Trapnell, C., B.A. Williams, G. Pertea, A. Mortazavi, G. Kwan, *et al.*, 2010
520 Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts
521 and isoform switching during cell differentiation. *Nature biotechnology* 28: 511-515.

522 Trapnell, C., L. Pachter, and S.L. Salzberg, 2009 TopHat: discovering splice
523 unctons with RNA-Seq. *Bioinformatics (Oxford, England)* 25: 1105-1111.

524 Van Ooijen, G., H.A. van den Burg, B.J. Cornelissen, and F.L. Takken, 2007
525 Structure and function of resistance proteins in solanaceous plants. *Annu. Rev.*
526 *Phytopathol.* 45: 43–72.

527 Xing, M., H. Lv, J. Ma, D. Xu, H. Li, *et al.*, 2016 Transcriptome profiling of
528 Resistance to *Fusarium oxysporum* f. sp. *conglutinans* in cabbage (*Brassica oleracea*)
529 Roots. *PLoS One* 11: e0148048.

530 Yang, L., and H. Huang, 2014 Roles of small RNAs in plant disease resistance. *J.*
531 *Integr. Plant Biol.* 56: 962–970.

532 Yu, S.C., and Y.M. Zou, 2008 A co-dominant molecular marker of *Fusarium* wilt
533 resistance gene *I-2* derived from gene sequence in tomato. *Yi Chuan* 30: 926-932.