Marine fish intestine responds to ocean acidification producing more carbonate aggregates

Sílvia F. Gregório, Ignacio Ruiz-Jarabo, Edison M Carvalho and Juan Fuentes *

Centre of Marine Sciences (CCMar), Universidade do Algarve, Campus de
Gambelas, 8005-139 Faro, Portugal

Short title: Fish intestinal response to acidification

Keywords: osmoregulation, sea bream, bicarbonate secretion and CO₂

* Author for correspondence:

Juan Fuentes (jfuentes@ualg.pt)

Centre of Marine Sciences (CCMar)

Universidade do Algarve

Campus de Gambelas,

8005-139 Faro, Portugal

Tel: 00351289800900 x 7375

ABSTRACT:

Marine fish contribute to the carbon cycle by producing mineralized intestinal aggregates generated as by-products of their osmoregulation. Here we aimed at characterizing the control of intestinal aggregate production in the gilthead sea bream in response to near future increases of environmental CO₂. Our results demonstrate that hypercapnia (800 and 1200 µatm CO₂) elicits higher intestine epithelial HCO₃⁻ secretion and the subsequent parallel increase of intestinal aggregate production when compared to present values (400 µatm CO₂). Intestinal gene expression analysis revealed the up-regulation of crucial transport mechanisms involved not only in the intestinal secretion cascade (*Slc4a4*, *Slc26a3* and *Slc26a6*) of sea bream, but also in other mechanisms involved in intestinal ion uptake linked to water absorption such as *NKCC2* and the *Aquaporin 1b*. These results highlight the important role of fish in the marine carbon cycle, and their potential growing impact of intestinal biomineralization processes in the scenario of ocean acidification.

INTRODUCTION:

In seawater, fish live in a hyperosmotic environment which leads to a loss of body fluids. The water lost passively through the epithelial surface is compensated by drinking (Fuentes and Eddy, 1997; Smith et al., 1930). Due to the osmotic imbalance between the ingested SW and the internal body fluids, an intense desalinization in the first part of the digestive tract is required i. e. oesophagus and stomach (Hirano and Mayer-Gostan, 1976; Parmelee and Renfro, 1983). This ion capture, mainly Na^+ and Cl^- , results in the decrease of intestinal fluid osmolality from circa 1050 to 400 mOsm kg^{-1} due to the action of active and passive transport processes (Esbaugh and Grosell, 2014; Loretz, 1995; Marquez and Fuentes, 2014). The osmotic pressure of the intestinal fluid is then decreased to a level that matches plasma osmolality (Gregorio et al., 2013), allowing the water to be absorbed passively or by the action of aquaporins (Whittamore, 2012; Wood and Grosell, 2012).

Water absorption in the anterior part of the intestine of marine fish was associated to ion fluxes driven by $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ co-transporters (Musch et al., 1982). However, it was recently established that part of the process is also linked to the action of chloride-dependent movements coupled to bicarbonate secretion, and mediated by apical anion transporters (Grosell et al., 2005; Heuer et al., 2012). Members of the Slc26 family are responsible for this apical mechanism in fish enterocytes (Carvalho et al., 2015; Gregorio et al., 2013; Ruhr et al., 2016). HCO_3^- enters the cell through a basolateral $\text{Na}^+/\text{HCO}_3^-$ co-transporter belonging to the Slc4 family (Chang et al., 2012) or is formed by the intracellular action of carbonic anhydrases that hydrate the CO_2 resulting in the production of HCO_3^- and H^+ ions (Grosell et al., 2009).

The secretion of HCO_3^- into the intestinal luminal fluid promotes an increase of pH (Wilson and Grosell, 2003). In parallel calcium and magnesium are present in high concentrations in SW and accumulate in the intestinal fluid. Under both conditions, precipitation and formation of carbonate aggregates in the intestinal lumen of marine fish is promoted. As a result of this process, an osmolality reduction of the fluid occurs, thus facilitating water absorption (Wilson et al., 2009; Wilson et al., 2002). Apical bicarbonate secretion in the enterocyte is regulated by biotic factors such as hormones, peptides, blood HCO_3^- concentration and abiotic factors e.g. salinity (Genz and Grosell, 2011; Genz et al., 2008; Gregorio et al., 2013). On the other hand, previous studies demonstrated that bicarbonate secretion in the intestine of marine fish is also regulated by an intracellular pH/ CO_2 / HCO_3^- sensor, the soluble

adenylyl cyclase (Carvalho et al., 2012; Tresguerres et al., 2010). Moreover, a challenge with high environmental levels of CO₂ (hypercapnia) elicited a response in the gilthead sea bream (*Sparus aurata*), decreasing its plasma pH during the initial 24 h, which is then buffered by increased levels of plasma bicarbonate (Michaelidis et al., 2007). Such an accumulation of bicarbonate in plasma, but also in other body fluids was also observed in several organisms (Portner et al., 2004). It should be of interest to establish if those physiological modifications alter intestinal carbonate aggregates production in marine fish.

In a global scenario, as a result of anthropogenic emissions in the atmosphere CO₂ increased from 280 up to 380 μ atm in the 20th century. The ocean absorbs part of this CO₂ resulting in a decrease of pH (IPCC, 2007; Kroeker et al., 2013). The ocean surface CO₂ levels are predicted to duplicate by the year 2100 to achieve levels circa 800 μ atm CO₂, thus decreasing 0.3-0.4 pH units (Orr et al., 2005). Previous studies have reported that some marine calcifying biota is affected when submitted to elevated CO₂ levels and the subsequent pH decrease close to values expected by the year 2100 (pH 7.8) and by the end of the next century (pH 7.6) (Mackenzie et al., 2014; Munday et al., 2009; Munday et al., 2010). Moreover, by the year 2300 the seawater CO₂ concentration will be 1900 μ atm that in turn decreased the pH in up to 0.77 units (Caldeira and Wickett, 2003; Kroeker et al., 2013; Zeebe et al., 2008). In contrast to other taxa, marine fish contribute to the carbon cycle as they fix/cycle around 15 % of the oceanic carbon, producing mineralized intestinal aggregates generated as by-products of their osmoregulation (Walsh et al., 1991; Wilson et al., 2009). The nature of those aggregates makes them relevant as they provide neutralizing alkaline buffer in the form of soluble carbonates, preferentially in the uppermost, most productive section of the water column (Perry et al., 2011). This study aimed at understanding the intestinal response and control of aggregate production of marine fish intestine in response to near future concentrations of CO₂ in seawater.

METHODS

Animals

Sea bream (*Sparus aurata*) juveniles were purchased from CUPIMAR SA (Cadiz, Spain) and transported to Ramalhete Marine Station (CCMAR, University of Algarve,

Faro, Portugal). Fish were maintained for 60 days in 1000 L tanks with running seawater at a density 9-10 kg m⁻³ and fed 2 % ration (fish wet weight, Sorgal, S.A., Portugal; Balance 3) twice daily until the start of the experiment. For experimental purposes 5 fish (200 g body weight) were transferred to 100 L tanks. Eight replicates were established per CO₂ concentration in an open circuit where temperature was maintained constant (25 °C), photoperiod was natural (April-June, Algarve, Portugal) and the feeding regime was maintained as above. Food was withheld for 36 h before sacrifice and sampling. No mortality was observed during the experiments.

The experiments comply with the guidelines of the European Union Council (86/609/EU). All fish protocols were performed under a “Group-C” licence from the Direcção-Geral de Veterinária, Ministério da Agricultura, do Desenvolvimento Rural e das Pescas, Portugal.

Experimental conditions and seawater chemistry

The rate of CO₂ injection into the systems was controlled by the pH level of seawater using pH probes connected to CO₂ controllers (EXAxt PH450G; Yokogawa Iberia-Portugal) with an internal controller following guidelines provided by the manufacturer. Each independent circuit was injected with CO₂, thus obtaining three groups constantly maintained at 400, 800 and 1200 µatm CO₂. To calibrate the automated negative feedback system, it was necessary to measure seawater pH, and extrapolate a pH/pCO₂ standard curve to indirectly measure pCO₂. The linearity of the relationship was confirmed through IRGA CO₂ analysis (WMA-4 PP Systems, Amesbury, MA, USA).

Total alkalinity (TA) was measured using an automated, open-cell potentiometric titration unit (SOP 3b) (Dickson et al., 2007), with a combination DL15 titrator and a DG115-SC probe (Mettler-Toledo) using certified acid titrant (0.1 M HCl, Fluka Analytical, Sigma-Aldrich). The results are expressed as µmol kg⁻¹ seawater. The water chemistry parameters were calculated using the CO₂Calc Software (Robbins et al., 2010). Along with temperature and salinity, CO₂Calc algorithms produce a suite of carbonate system parameters using only two measured variables (of five potential inputs); the remaining three possible inputs are also calculated as algorithm outputs. In our samples, pH, TA, and/or TCO₂ were routinely measured, and precedence of use is: 1) measured pH and TA 2) measured pH and TCO₂ (total dissolved inorganic carbon). In situ carbonate system parameters were also

calculated using temperature recorded in the field (“adjusted” outputs), as explained before (Robbins et al., 2010).

General sampling

After 3 months of acclimation to the altered water CO₂ levels, fish were captured and anesthetized in 2-phenoxyethanol (1: 10,000 v/v; Sigma- Aldrich, St Louis, MO, USA). Blood samples were collected by caudal puncture using heparinized syringes. Plasma was obtained by centrifugation (10,000 *rpm*, 5 min, 4 °C) and stored at –20 °C until analysis. Fish were sacrificed by decapitation and the whole intestine was isolated. The intestinal fluid of individual fish was collected from the excised intestinal tract clamped (from pyloric caeca to anal sphincter) with two haemostatic forceps, emptied into pre-weighed vials and centrifuged (12,000 *rpm*, 5 min, RT) to separate fluid from precipitates. The fluid was transferred to pre-weighed vials and the volume was measured to the nearest 0.1 µL (0.1 mg, assuming a density of 1).

Plasma and fluid analysis

Osmolality of plasma and intestinal luminal fluid was measured in 10 µL samples with a Vapro 5520 osmometer (Wescor, South Logan, UT, USA). Sodium concentrations were measured by flame photometry (BWB-XP Performance Plus, BWB Technologies, UK). The results are expressed as mmol L⁻¹. Chloride was determined by Coulomb-metric titration (SAT-500, DKK-TOA, Japan). Calcium and magnesium were measured by colorimetric tests, using commercial kits (Spinreact, Reactivos Spinreact, SA, Girona, Spain), according to the manufacturer instruction in a microplate reader (Biorad Benchmark, Bio-Rad, USA). Intestinal fluid titratable alkalinity (HCO₃⁻ + CO₃²⁻) was manually measured with the double titration method with a pH electrode (HI1330B, Hanna Instruments, Smithfield, RI, USA) attached to a pH meter (PHM84, Radiometer, Copenhagen, Denmark): 50 µL of the intestinal fluid samples was diluted in 10 mL NaCl (40 mmol L⁻¹), gassed with CO₂-free gas for 30 min to remove CO₂ and titrated to pH 3.8 with 10 mmol L⁻¹ HCl; an additional gassing period of 20 min was applied to remove any remaining CO₂. The sample was back titrated to its original pH with 10 mmol L⁻¹ NaOH. The volume difference between added acid and base in both titrations and titrant molarities was used to calculate total HCO₃⁻ equivalents (mEquiv.L⁻¹) as described before (Gregorio et al., 2014).

Intestinal Bicarbonate Secretion

A segment of fish anterior intestine was excised, mounted on tissue holders (P2413, 0.71 cm², Physiologic Instruments, San Diego, CA, USA) and positioned between two half-chambers (P2400, Physiologic Instruments) containing 1.5 mL of basolateral and apical saline. The composition of the basolateral saline was: 160 mmol L⁻¹ NaCl, 1 mmol L⁻¹ MgSO₄, 2 mmol L⁻¹ NaH₂PO₄, 1.5 mmol L⁻¹ CaCl₂, 5 mmol L⁻¹ NaHCO₃, 3 mmol L⁻¹ KCl, 5.5 mmol L⁻¹ glucose and 5 mmol L⁻¹ HEPES, pH 7.800, gassed with 0.3% CO₂+ 99.7% O₂. Apical saline: 88 mmol L⁻¹ NaCl, 9.5 mmol L⁻¹ MgCl₂, 3 mmol L⁻¹ KCl, 7.5 mmol L⁻¹ CaCl₂, 126.5 mmol L⁻¹ MgSO₄ and 1 mmol L⁻¹ Na₂HPO₄, gassed with 100% O₂ and pH maintained at 7.800 throughout the experiments by pH-Stat (see below). The temperature was maintained at 25 °C throughout all experiments. All bioelectrical variables were monitored by means of Ag/AgCl electrodes (with tip asymmetry <1 mV) connected to either side of the Ussing chamber with 3 mm- bore agar bridges (1 mol L⁻¹ KCl in 3% agar). Transepithelial electrical potential (TEP, mV) was monitored by clamping of epithelia to 0 μA cm⁻². Epithelial resistance (R_t, Ω cm²) was manually calculated (Ohm's law) using the voltage deflections induced by a 10 μA cm⁻² bilateral pulse of 2 s every minute. Current injections were performed by means of a VCC 600 amplifiers (Physiologic Instruments). For pH-Stat control, a pH electrode (PHC 4000-8, Radiometer) and a microburette tip were immersed in the luminal saline and connected to a pH-Stat system (TIM 854, Radiometer). To allow pulsing (for R_t calculation) during pH measurements, the amplifier was grounded to the titration unit. The configuration of amplifier/pH-Stat system used in this study is similar to that first established for the characterization of HCO₃⁻ secretion in the intestine of the Gulf toadfish and sea bream (Gregorio et al., 2013; Grosell and Genz, 2006; Guffey et al., 2011; Taylor et al., 2010) and provides rates of intestinal secretion similar to those obtained by the double titration method in the intestine of the sea bream (Fuentes et al., 2010). Measurement of HCO₃⁻ secretion was performed on luminal saline at physiological pH 7.800 during all experiments. The volume of the acid titrant (2.5 mmol L⁻¹ HCl) was recorded and the amount of HCO₃⁻ secreted (nmol h⁻¹ cm⁻²) was calculated from the volume of titrant added, the concentration of the titrant and the surface area (cm²). All experiments comprised 1 h of tissue stable voltage and HCO₃⁻ secretion.

qPCR

After anaesthesia and decapitation, a portion of the anterior intestine was collected from individual fish, and stored in RNA Later at 4 °C (Sigma- Aldrich) until utilized for RNA extraction within 2 weeks. Total RNA was extracted from samples of anterior intestine with the Total RNA Kit I (E.Z.N.A, Omega, US) following the manufacturer's instructions and the quantity and quality of RNA assessed (Nanodrop 1000, Thermo Scientific, US). Prior to cDNA synthesis RNA was treated with DNase using the DNA-free Kit (Ambion, UK) following the supplier's instructions. Reverse transcription of RNA into cDNA was carried out using the RevertAid First Strand cDNA Synthesis Kit (TermoFisher Scientific, UK) with 500 ng of total RNA in a reaction volume of 20 µL. Table 4 shows primer sequences and amplicon sizes.

Real-time qPCR amplifications were performed in duplicate in a final volume of 10 µL with 5 µL SsoFast EvaGreen Supermix (Bio-Rad, UK) as the reporter dye, around 20 ng cDNA, and 0.3 µM of each forward and reverse primers. Amplifications were performed in 96-well plates using the *One-step Plus* sequence detection system (Applied Biosystems, California, USA) with the following protocol: denaturation and enzyme activation step at 95 °C for 2 min, followed by 40 cycles. After the amplification phase, a temperature-determining dissociation step was carried out at 65 °C for 15 s, and 95 °C for 15 s. For normalization of cDNA loading, all samples were run in parallel using 18S ribosomal RNA (18S). To estimate efficiencies, a standard curve was generated for each primer pair from 10-fold serial dilutions (from 0.1 to 1e10⁻⁸ ng) of a pool of first-stranded cDNA template from all samples. Standard curves represented the cycle threshold value as a function of the logarithm of the number of copies generated, defined arbitrarily as one copy for the most diluted standard. All calibration curves exhibited correlation coefficients R²>0.98, and the corresponding real-time PCR efficiencies between 95 and 99%.

Statistics

All results are shown as mean ± standard error of the mean (mean ± s.e.m.). After assessing homogeneity of variance and normality, statistical analysis of the data was carried out by using one-way analysis of variance using CO₂ concentration as a factor of variance, followed by the *post hoc* Bonferroni t-test (Prism 5.0, GraphPad Software for McIntosh). The level of significance was set at *p*< 0.05.

RESULTS:

No mortality was observed in any of the experimental groups.

Experimental conditions - seawater chemistry

The environmental condition values were controlled twice a day in this study and the results are shown in Table 1. Temperature and salinity levels were maintained within a small range of variation, and no differences were described between CO₂ treatments (Table 1). The concentration of CO₂ and pH in the control treatment averaged $408 \pm 20 \mu\text{atm}$ and pH 8.01 ± 0.01 , in the medium CO₂-concentration group $797 \pm 29 \mu\text{atm}$ and pH 7.80 ± 0.01 , whereas in the enriched-CO₂ treatment it averaged $1259 \pm 46 \mu\text{atm}$ and pH 7.64 ± 0.01 . Total alkalinity was not significantly different between treatments ranging between 2582 and 2605 $\mu\text{mol kg}^{-1}$ SW (Table 1).

Plasma and intestinal fluid analysis

The osmolality and the ion composition of plasma and intestinal fluid of sea bream are shown in Table 2 and 3, respectively. No differences were detected in plasma osmolality which ranged between 343 to 346 mOsm kg^{-1} in controls and fish exposed to elevated CO₂ concentrations (Table 2). Additionally, no changes in Cl⁻ (163.5 to 165.3 mmol L⁻¹) and Na⁺ (178.7 to 186.2 mmol L⁻¹) concentration (Table 2) were observed in the plasma between controls and fish exposed to elevated water CO₂. The content of Ca²⁺ ($9.5 \pm 0.6 \text{ mmol L}^{-1}$) in intestinal fluid did not change with the different exposure to elevated water CO₂ (Table 3) and the concentration of Na⁺ (72.1 to 81.8 mmol L⁻¹) in the intestinal fluid was not different between controls and fish exposed to elevated CO₂ (Table 3). The content of Cl⁻ in the intestinal fluid of fish exposed to both levels of elevated water CO₂ was significantly lower $95.0 \pm 5.3 \text{ mmol L}^{-1}$ ($p < 0.05$, one-way ANOVA) when compared to controls $116 \pm 5.5 \text{ mmol L}^{-1}$ (Figure 1A). However, the content of Mg²⁺ in the intestinal fluid of fish exposed to elevated water CO₂ was statistically higher than in control fish ($p < 0.05$, one-way ANOVA, Table 3).

Bicarbonate and carbonate aggregates in the intestinal lumen

HCO₃⁻ concentration in the intestinal fluid of control fish was in the range of 44 mEq L⁻¹ (Figure 1B) and the exposure to both levels of elevated CO₂ resulted in a significant ($p < 0.05$, one-way ANOVA) increase of HCO₃⁻ concentration in the fish

intestinal fluid to 66 and 65 mEq L⁻¹ respectively (Figure 1B). When bicarbonate content was normalized to fluid volume plus the weight of carbonate aggregates (Ca[Mg]CO₃) and fish body mass, we still observed that fish exposed to both levels of elevated CO₂ responded with a 3-fold increase in carbonate aggregate formation ($p < 0.05$, one-way ANOVA, Figure 1C).

Bicarbonate secretion

HCO₃⁻ secretion in the anterior intestine is shown in Figure 2A. In control fish (400 µatm CO₂) bicarbonate secretion was 511 ± 7.3 nmol.h⁻¹.cm⁻², while it averaged 631 ± 7.2 and 842 ± 6.9 nmol.h⁻¹.cm⁻² for fish acclimated to 800 and 1200 µatm CO₂, respectively. No significant effects were observed in bicarbonate secretion between the 400 and 800 µatm CO₂ groups, while the 1200 µatm CO₂-group was significantly higher than the control fish. In terms of epithelial resistance, no changes were observed between groups (Figure 2B).

qPCR

The *Slc26a3* anion exchanger paralleled the accumulation of carbonate aggregates with a significant gradual increase of *Slc26a3* expression in response to increased water CO₂ (Figure 3A). The apical exchanger *Slc26a6* presents a significantly ($p < 0.05$, one-way ANOVA) higher expression in fish exposed to high water CO₂ levels compared to controls (Figure 3B).

Changes in *SLC4a4* expression are shown in Fig. 3C. Expression of the basolateral co-transporter *SLC4a4* was significantly ($p < 0.05$, one-way ANOVA) higher in fish exposed to medium and high CO₂ levels than in control fish (Figure 3E).

The expression of *NKCC2*, the absorptive form of the co-transporter in the anterior intestine of sea bream was significantly increased in response to hypercapnia (Figure 4A)

As expected, the water channel *Aquaporin 1b* also responded positively to the CO₂ challenge, with significantly higher expression of *Aquaporin 1b* gene in fish exposed to medium and high CO₂ concentrations than in controls (Figure 4B).

DISCUSSION:

The sea bream, acclimated from 400 up to 1200 $\mu\text{atm CO}_2$ for three months, is able to maintain within a narrow range its plasma levels of osmolality, Na^+ and Cl^- without suffering any casualties. This homeostatic balance is partially achieved by changes in intestinal epithelia related to ion transport. In this regard, elevated water CO_2 induces increased rates of epithelial HCO_3^- secretion and a molecular response that is in keeping with enhanced water absorption processes.

Gilthead sea bream acclimated to elevated water CO_2 for 3 months increased the amount of HCO_3^- and carbonate aggregates in the intestinal fluid (Figure 1). More precisely the groups of fish acclimated to hypercapnia (800 and 1200 $\mu\text{atm CO}_2$), when compared to the control group (400 $\mu\text{atm CO}_2$), showed a 1.5-fold and 3.0-fold increase in the bicarbonate and carbonate aggregates present in the intestinal lumen, respectively. These results and the fold-change of variation are in good agreement, with those previously described by us in seawater-acclimated sea bream challenged with 55ppt salinity (Gregorio et al., 2013). These effects in relation to salinity challenge, were related to an increase in water processing at the level of enterocytes. An enrichment of the intestinal fluid with Mg^{2+} and other divalent ions was described reflecting high levels of water ingestion and processing in the intestine (Kurita et al., 2008; Whittamore, 2012). Here, a noticeable increase of Mg^{2+} the intestinal fluid in response to hypercapnia is exposed, which suggests different water processing and/or drinking rates in response to higher water CO_2 .

In the intestinal fluid, both levels of hypercapnia elicited significant decreases in the intestinal fluid Cl^- content of about 21 mM. Interestingly, this decrease was paralleled by a ~22 mM increase in HCO_3^- content of the same fluid, pointing to an activation of $\text{Cl}^-/\text{HCO}_3^-$ exchangers at the epithelial level (Grosell, 2006). To test this possibility, we measured bicarbonate secretion in the anterior intestine, since this portion of the intestine shows the highest secretion rates along the intestinal tract in the sea bream (Carvalho et al., 2012; Gregorio et al., 2013). Accordingly, intestinal bicarbonate secretion increased in parallel with increased environmental CO_2 levels (Figure 2). The measured bicarbonate secretion fold-change between the control group and fish challenged with 1200 $\mu\text{atm CO}_2$ from 511 to 842 $\text{nmol HCO}_3^- \text{ cm}^{-2} \text{ h}^{-1}$, coincided well with the alterations reported in the intestine of the sea bream challenged long term with 55 ppt seawater, from 495 to 783 $\text{nmol HCO}_3^- \text{ cm}^{-2} \text{ h}^{-1}$ (Gregorio et al., 2013). In this sense, the effects of hypercapnia could be carefully compared to a hyperosmotic

challenge in this species, as they both enhanced bicarbonate secretion and formation of carbonate aggregates in the intestine.

It was previously reported that the seabream responds to hypercapnia with a plasma pH drop, which is buffered within the first 5 days of exposure by increasing plasma bicarbonate levels (Michaelidis et al., 2007). We hereby postulated that the accumulation of plasma bicarbonate generated in response to high environmental CO₂, could be levelled off by activation of intestinal bicarbonate secretion at the level of the enterocytes. Plasma bicarbonate enters the enterocyte through an electrogenic Na⁺-coupled HCO₃⁻ transporter located at the basolateral membrane, Slc4a4, which is important for a transepithelial HCO₃⁻ transport (Kurita et al., 2008; Taylor et al., 2010). Our results show that the expression of *Slc4a4* increased in the groups challenged with high CO₂ levels, supporting a hypothesis for enhanced bicarbonate secretion capacity in the intestine, in response to water hypercapnia and the consequent plasma HCO₃⁻ accumulation.

In the apical membrane of the fish enterocytes, both bicarbonate secretion and chloride uptake are mediated through Slc26 family transporters, such as the Slc26a3 and Slc26a6 (Grosell et al., 2005; Kurita et al., 2008; Wilson et al., 1996). Likewise those exchangers are expressed in the intestine of sea bream and are modulated by salinity and endocrine regulation (Gregorio et al., 2014; Gregorio et al., 2013). In addition, the expression of this transporter is modulated in response to a CO₂ challenge in the gill of toadfish (Esbaugh et al., 2012). Here, we confirmed that the expression of *Slc26a6* and *Slc26a3* are also modulated positively in response to elevated water CO₂ levels in the intestine of the sea bream (Figure 3). This increase is in good agreement with the parallel expression increase of the *Slc4a4* co-transporter, which would substantiate a boosted secretory flow of bicarbonate from the plasma to the intestinal lumen, subsequently enhancing chloride absorption. It is challenging to establish transporters stoichiometry from ion concentration alone. However, here we have established a 1:1 relationship between the variation of Cl⁻ and HCO₃⁻ in the intestinal fluid after 3 months in hypercapnia (Figures 1A and B). It appears, that as a consequence of the increased bicarbonate levels on the intestinal fluid, carbonate aggregate formation is increased, sustaining a decrease of fluid osmolality and that enables water absorption (Wilson et al., 2002).

In marine fish, water absorption in the intestinal tract is mediated through the activity of ion absorption, especially Cl⁻. One of the most important ion transporters in this

species is the $\text{Na}^+\text{-K}^+\text{-2Cl}^-$ co-transporter (NKCC) (Musch et al., 1982), which mediates the electroneutral movement of 1 Na^+ , 1 K^+ and 2 Cl^- across cell membranes. Two isoforms, *NKCC1* and *NKCC2*, are currently known and are encoded by different genes (Flatman, 2002). In the intestine of marine fish the apical co-transporter NKCC2 activity seems essential for ion regulation and, in parallel with the *Slc26a6*, it is believed to regulate water homeostasis in salinity-challenged sea bream (Gregorio et al., 2013; Grosell, 2006). Here, we show that elevated water CO_2 causes an upregulation of *NKCC2* expression in the anterior intestine (Figure 4A) which adds up to increased expression of the apical anion transporters, *Slc26a6* and *Slc26a3*. This combination has the potential to enhance intestinal water absorption linked to chloride movements. Remarkably, sea bream exposed to elevated water CO_2 showed no significant changes in plasma ions concentration, indicating that the intestinal response to elevated water CO_2 is part of the allostatic control of plasma homeostasis. In order to understand these changes further, we analysed the expression of *Aquaporin 1b*, a functional water channel highly expressed in intestinal epithelial cells of the sea bream anterior intestine (Martos-Sitcha et al., 2015; Raldua et al., 2008). We observed solid increases of *Aquaporin 1b* expression in the anterior intestine of sea bream exposed to elevated water CO_2 . Providing further evidence of the functional and molecular changes of the intestine in response to elevated water CO_2 , that could be the foundation of increased energetic expenditure recently reported in the toadfish intestine in response to elevated water CO_2 (Heuer and Grosell, 2016).

ACKNOWLEDGEMENTS

We appreciate the technical input of Dr. João Eugenio Reis (Ramalhete Marine Station, CCMar, University of Algarve, Portugal) for fish care and assistance during experimental periods.

FUNDING

This work was partially supported by the Ministry of Science and Higher Education and European Social Funds through the Portuguese National Science Foundation (FCT) by studentship SFRH/BD/113363/2015 to SG and by Project PTDC/MAR-BIO/3034/2014 to JF. CCMar is supported by national funds from the Portuguese Foundation for Science and Technology (FCT) through project UID/Multi/04326/2013.

REFERENCES

- Caldeira, K. and Wickett, M. E. (2003). Oceanography: anthropogenic carbon and ocean pH. *Nature* **425**, 365.
- Carvalho, E. S., Gregorio, S. F., Canario, A. V., Power, D. M. and Fuentes, J. (2015). PTHrP regulates water absorption and aquaporin expression in the intestine of the marine sea bream (*Sparus aurata*, L.). *Gen Comp Endocrinol* **213**, 24-31.
- Carvalho, E. S., Gregorio, S. F., Power, D. M., Canario, A. V. and Fuentes, J. (2012). Water absorption and bicarbonate secretion in the intestine of the sea bream are regulated by transmembrane and soluble adenylyl cyclase stimulation. *J Comp Physiol B* **182**, 1069-80.
- Chang, M. H., Plata, C., Kurita, Y., Kato, A., Hirose, S. and Romero, M. F. (2012). Euryhaline pufferfish NBCe1 differs from nonmarine species NBCe1 physiology. *Am J Physiol Cell Physiol* **302**, C1083-95.
- Dickson, A. G., Sabine, C. L. and Christen, J. R. (2007). Guide to best practice for ocean CO₂ measurements. Sidney, Canada: North Pacific Marine Science Organization.
- Esbaugh, A. J. and Grosell, M. (2014). Esophageal desalination is mediated by Na(+), H(+) exchanger-2 in the gulf toadfish (*Opsanus beta*). *Comp Biochem Physiol A Mol Integr Physiol* **171**, 57-63.
- Esbaugh, A. J., Heuer, R. and Grosell, M. (2012). Impacts of ocean acidification on respiratory gas exchange and acid-base balance in a marine teleost, *Opsanus beta*. *J Comp Physiol B* **182**, 921-34.
- Flatman, P. W. (2002). Regulation of Na-K-2Cl cotransport by phosphorylation and protein-protein interactions. *Biochim Biophys Acta* **1566**, 140-51.
- Fuentes, J. and Eddy, F. B. (1997). Drinking in freshwater, euryhaline and marine teleosts. In *Ionic regulation in animals*, eds. N. Hazon F. B. Eddy and G. Flik), pp. 135-149. New York: Heidelberg: Springer-Verlag.
- Fuentes, J., Power, D. M. and Canario, A. V. (2010). Parathyroid hormone-related protein-stanniocalcin antagonism in regulation of bicarbonate secretion and calcium precipitation in a marine fish intestine. *Am J Physiol Regul Integr Comp Physiol* **299**, R150-8.

- Genz, J. and Grosell, M.** (2011). *Fundulus heteroclitus* acutely transferred from seawater to high salinity require few adjustments to intestinal transport associated with osmoregulation. *Comp Biochem Physiol A Mol Integr Physiol* **160**, 156-65.
- Genz, J., Taylor, J. R. and Grosell, M.** (2008). Effects of salinity on intestinal bicarbonate secretion and compensatory regulation of acid-base balance in *Opsanus beta*. *J Exp Biol* **211**, 2327-35.
- Gregorio, S. F., Carvalho, E. S., Campinho, M. A., Power, D. M., Canario, A. V. and Fuentes, J.** (2014). Endocrine regulation of carbonate precipitate formation in marine fish intestine by stanniocalcin and PTHrP. *J Exp Biol* **217**, 1555-62.
- Gregorio, S. F., Carvalho, E. S., Encarnacao, S., Wilson, J. M., Power, D. M., Canario, A. V. and Fuentes, J.** (2013). Adaptation to different salinities exposes functional specialization in the intestine of the sea bream (*Sparus aurata* L.). *J Exp Biol* **216**, 470-479.
- Grosell, M.** (2006). Intestinal anion exchange in marine fish osmoregulation. *J Exp Biol* **209**, 2813-27.
- Grosell, M. and Genz, J.** (2006). Ouabain-sensitive bicarbonate secretion and acid absorption by the marine teleost fish intestine play a role in osmoregulation. *Am J Physiol Regul Integr Comp Physiol* **291**, R1145-56.
- Grosell, M., Genz, J., Taylor, J. R., Perry, S. F. and Gilmour, K. M.** (2009). The involvement of H⁺-ATPase and carbonic anhydrase in intestinal HCO₃⁻ secretion in seawater-acclimated rainbow trout. *J Exp Biol* **212**, 1940-8.
- Grosell, M., Wood, C. M., Wilson, R. W., Bury, N. R., Hogstrand, C., Rankin, C. and Jensen, F. B.** (2005). Bicarbonate secretion plays a role in chloride and water absorption of the European flounder intestine. *Am J Physiol Regul Integr Comp Physiol* **288**, R936-46.
- Guffey, S., Esbaugh, A. and Grosell, M.** (2011). Regulation of apical H⁽⁺⁾-ATPase activity and intestinal HCO₃⁽⁻⁾ secretion in marine fish osmoregulation. *Am J Physiol Regul Integr Comp Physiol* **301**, R1682-91.
- Heuer, R. M., Esbaugh, A. J. and Grosell, M.** (2012). Ocean acidification leads to counterproductive intestinal base loss in the gulf toadfish (*Opsanus beta*). *Physiol Biochem Zool* **85**, 450-9.
- Heuer, R. M. and Grosell, M.** (2016). Elevated CO₂ increases energetic cost and ion movement in the marine fish intestine. *Sci Rep* **6**, 34480.
- Hirano, T. and Mayer-Gostan, N.** (1976). Eel esophagus as an osmoregulatory organ. *Proc Natl Acad Sci U S A* **73**, 1348-50.
- IPCC.** (2007). Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Geneva, Switzerland.
- Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte, C. M. and Gattuso, J. P.** (2013). Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Global change biology* **19**, 1884-96.
- Kurita, Y., Nakada, T., Kato, A., Doi, H., Mistry, A. C., Chang, M. H., Romero, M. F. and Hirose, S.** (2008). Identification of intestinal bicarbonate transporters involved in formation of carbonate precipitates to stimulate water absorption in marine teleost fish. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology* **294**, R1402-R1412.

- 497 **Loretz, C. A.** (1995). Electrophysiology of ion transport in teleost intestinal cells.
498 In *Cellular and molecular approaches to fish ionic regulation*, vol. 14 eds. C. M. Wood and
499 T. J. Shuttleworth), pp. 25-56. Amsterdam: Academic Press.
- 500 **Mackenzie, C. L., Lynch, S. A., Culloty, S. C. and Malham, S. K.** (2014). Future
501 oceanic warming and acidification alter immune response and disease status in a
502 commercial shellfish species, *Mytilus edulis* L. *PLoS One* **9**, e99712.
- 503 **Marquez, L. and Fuentes, J.** (2014). In vitro characterization of acid secretion in
504 the gilthead sea bream (*Sparus aurata*) stomach. *Comp Biochem Physiol A Mol Integr*
505 *Physiol* **167**, 52-8.
- 506 **Martos-Sitcha, J. A., Campinho, M. A., Mancera, J. M., Martinez-Rodriguez, G.**
507 **and Fuentes, J.** (2015). Vasotocin and isotocin regulate aquaporin 1 function in the sea
508 bream. *J Exp Biol* **218**, 684-93.
- 509 **Michaelidis, B., Spring, A. and Portner, H. O.** (2007). Effects of long-term
510 acclimation to environmental hypercapnia on extracellular acid-base status and
511 metabolic capacity in Mediterranean fish *Sparus aurata*. *Marine Biology* **150**, 1417-1429.
- 512 **Munday, P. L., Dixon, D. L., Donelson, J. M., Jones, G. P., Pratchett, M. S.,**
513 **Devitsina, G. V. and Doving, K. B.** (2009). Ocean acidification impairs olfactory
514 discrimination and homing ability of a marine fish. *Proc Natl Acad Sci U S A* **106**, 1848-
515 52.
- 516 **Munday, P. L., Dixon, D. L., McCormick, M. I., Meekan, M., Ferrari, M. C. and**
517 **Chivers, D. P.** (2010). Replenishment of fish populations is threatened by ocean
518 acidification. *Proc Natl Acad Sci U S A* **107**, 12930-4.
- 519 **Musch, M. W., Orellana, S. A., Kimberg, L. S., Field, M., Halm, D. R., Krasny, E.**
520 **J., Jr. and Frizzell, R. A.** (1982). Na⁺-K⁺-Cl⁻ co-transport in the intestine of a marine
521 teleost. *Nature* **300**, 351-3.
- 522 **Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A.,**
523 **Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F. et al.** (2005). Anthropogenic ocean
524 acidification over the twenty-first century and its impact on calcifying organisms.
525 *Nature* **437**, 681-6.
- 526 **Parmelee, J. T. and Renfro, J. L.** (1983). Esophageal desalination of seawater in
527 flounder: role of active sodium transport. *Am J Physiol* **245**, R888-93.
- 528 **Perry, C. T., Salter, M. A., Harborne, A. R., Crowley, S. F., Jelks, H. L. and**
529 **Wilson, R. W.** (2011). Fish as major carbonate mud producers and missing components
530 of the tropical carbonate factory. *Proceedings of the National Academy of Sciences of the*
531 *United States of America* **108**, 3865-3869.
- 532 **Portner, H. O., Langenbuch, M. and Reipschlager, A.** (2004). Biological impact
533 of elevated ocean CO₂ concentrations: Lessons from animal physiology and earth
534 history. *Journal of Oceanography* **60**, 705-718.
- 535 **Raldua, D., Otero, D., Fabra, M. and Cerda, J.** (2008). Differential localization
536 and regulation of two aquaporin-1 homologs in the intestinal epithelia of the marine
537 teleost *Sparus aurata*. *American journal of physiology. Regulatory, integrative and*
538 *comparative physiology* **294**, R993-1003.
- 539 **Robbins, L. L., Hansen, M. E., Kleypas, A. and Meylan, S. C.** (2010). CO₂calc: a
540 user-friendly seawater carbon calculator for Windows, Mac OS X, and iOS (iPhone), pp.
541 17: U.S. Geological Survey Open-File Report.
- 542 **Ruhr, I. M., Takei, Y. and Grosell, M.** (2016). The role of the rectum in
543 osmoregulation and the potential effect of renoguanlylin on SLC26a6 transport activity
544 in the Gulf toadfish (*Opsanus beta*). *American journal of physiology. Regulatory,*
545 *Integrative and Comparative Physiology* **311**, R179-91.

Smith, H. W., Farinacci, N. and Breitweiser, A. (1930). The absorption and excretion of water and salts by marine teleosts. *American Journal of Physiology* **93**, 480-505.

Taylor, J. R., Mager, E. M. and Grosell, M. (2010). Basolateral NBCe1 plays a rate-limiting role in transepithelial intestinal HCO₃⁻ secretion, contributing to marine fish osmoregulation. *J Exp Biol* **213**, 459-68.

Tresguerres, M., Levin, L. R., Buck, J. and Grosell, M. (2010). Modulation of NaCl absorption by [HCO₃⁻] in the marine teleost intestine is mediated by soluble adenylyl cyclase. *Am J Physiol Regul Integr Comp Physiol* **299**, R62-71.

Walsh, P. J., Blackwelder, P., Gill, K. A., Danulat, E. and Mommsen, T. P. (1991). Carbonate Deposits in Marine Fish Intestines - a New Source of Biom mineralization. *Limnology and Oceanography* **36**, 1227-1232.

Whittamore, J. M. (2012). Osmoregulation and epithelial water transport: lessons from the intestine of marine teleost fish. *J Comp Physiol B* **182**, 1-39.

Wilson, R., Gilmour, K., Henry, R. and Wood, C. (1996). Intestinal base excretion in the seawater-adapted rainbow trout: a role in acid-base balance? *J Exp Biol* **199**, 2331-43.

Wilson, R. W. and Grosell, M. (2003). Intestinal bicarbonate secretion in marine teleost fish-source of bicarbonate, pH sensitivity, and consequences for whole animal acid-base and calcium homeostasis. *Biochim Biophys Acta* **1618**, 163-74.

Wilson, R. W., Millero, F. J., Taylor, J. R., Walsh, P. J., Christensen, V., Jennings, S. and Grosell, M. (2009). Contribution of fish to the marine inorganic carbon cycle. *Science* **323**, 359-62.

Wilson, R. W., Wilson, J. M. and Grosell, M. (2002). Intestinal bicarbonate secretion by marine teleost fish--why and how? *Biochim Biophys Acta* **1566**, 182-93.

Wood, C. M. and Grosell, M. (2012). Independence of net water flux from paracellular permeability in the intestine of *Fundulus heteroclitus*, a euryhaline teleost. *J Exp Biol* **215**, 508-17.

Zeebe, R. E., Zachos, J. C., Caldeira, K. and Tyrrell, T. (2008). Oceans. Carbon emissions and acidification. *Science* **321**, 51-2.

Table 1 - Chemical conditions of seawater where sea bream were kept under different CO₂ concentrations over a three month period.

	400 μ atm CO ₂	800 μ atm CO ₂	1200 μ atm CO ₂
pH	8.01 $\pm$ 0.01	7.80 $\pm$ 0.01	7.64 $\pm$ 0.01
pCO ₂ (μ atm)	408 $\pm$ 20	797 $\pm$ 29	1259 $\pm$ 46
Salinity (ppt)	37.5 $\pm$ 0.0	37.5 $\pm$ 0.0	37.5 $\pm$ 0.0
Alkalinity (μ Mol kg ⁻¹ SW)	2605 $\pm$ 10	2582 $\pm$ 25	2585 $\pm$ 14
T (°C)	25.4 $\pm$ 0.2	25.5 $\pm$ 0.2	25.6 $\pm$ 0.2

Table 2 – Osmolality, Na⁺ and Cl⁻ levels in plasma of sea bream acclimated for 3 months to different CO₂ concentration (400, 800 and 1200 μ atm) at 37 ppt. Values are means $\pm$ s.e.m. (N=11-19). No differences were described between any of the experimental groups ($p > 0.05$, one-way ANOVA).

	400 μ atm CO ₂	800 μ atm CO ₂	1200 μ atm CO ₂
Osmolality (mOsm kg ⁻¹)	342 $\pm$ 2	343 $\pm$ 2	346 $\pm$ 2
Na ⁺ (mmol L ⁻¹)	178.7 $\pm$ 2.3	183.4 $\pm$ 2.6	186.2 $\pm$ 2.1
Cl ⁻ (mmol L ⁻¹)	163.5 $\pm$ 1.3	165.2 $\pm$ 1.1	164.6 $\pm$ 1.3

Table 3 - Osmolality, Na⁺, Ca²⁺ and Mg²⁺ levels in intestinal fluid of sea bream acclimated for 3 months to different CO₂ concentration (400, 800 and 1200 μ atm) at 37 ppt. Values are means $\pm$ s.e.m. (N=8-18). Different letters indicate significant differences among groups ($p < 0.05$, one-way ANOVA).

	400 μ atm CO ₂	800 μ atm CO ₂	1200 μ atm CO ₂
Osmolality (mOsm kg ⁻¹)	353 $\pm$ 8	355 $\pm$ 7	359 $\pm$ 9
Na ⁺ (mmol L ⁻¹)	77.0 $\pm$ 2.4	72.1 $\pm$ 4.3	81.8 $\pm$ 5.0
Ca ²⁺ (mmol L ⁻¹)	9.5 $\pm$ 0.6	10.3 $\pm$ 0.6	8.6 $\pm$ 0.5
Mg ²⁺ (mmol L ⁻¹)	123.2 $\pm$ 4.3 ^a	143.9 $\pm$ 7.8 ^b	143.8 $\pm$ 6.5 ^b

597 **Table 4** - Details of primers used for qPCR.

Gene	Primer	Sequence (5' to 3')	T _m (°C)	Product size (bp)	NCBI accession no.
<i>Slc26a3</i>	Forward	ATCTCGGCTCTGAAGGGACT	60	107	AM973894
	Reverse	AGCGAGCA TTTCTGTCCCTGCTC			
<i>Slc26a6</i>	Forward	GCGGGACTGTTCAGCGGAGG	60	153	FM155691.1
	Reverse	TGCGAACACGCCTGAACGGCA			
<i>Slc4a4</i>	Forward	ACCTTCA TGCCACCGCAGGG	60	101	FM157528.1
	Reverse	CGCCGCCGCCGATAACTCTT			
<i>NKCC2</i>	Forward	ACGGAGTCCAAGAAAACCACGGG	60	128	FP335045
	Reverse	CCAGCCAGGATTCCGGTCGC			
<i>Aquaporin1a</i>	Forward	GGCTCTCACGTACGATTTCC	60	158	AY626939
	Reverse	TCTGTGTGGGACTATTTTGACG			
<i>18S</i>	Forward	AACCAGACAAATCGCTCCAC	60	139	AY993930
	Reverse	CCTGCGGCTTAATTTGACTC			

598

FIGURE LEGENDS

Figure 1. Characterization of Cl^- , HCO_3^- content in intestinal fluid and $\text{Ca}(\text{Mg})\text{CO}_3$ aggregates and total carbonate in intestinal lumen of sea bream in response to different environmental CO_2 levels (400, 800 and 1200 $\mu\text{atm CO}_2$) for up to three months. Each bar represents the mean $\pm$ s.e.m. (N= 7 - 9). Different superscript letters indicate significant differences ($p<0.05$, one-way ANOVA followed by Bonferroni *post hoc* test).

Figure 2. Bicarbonate secretion (BCS) and tissue resistance (R_t) measured in Ussing chambers by pH-Stat from anterior intestinal regions of sea bream in response to different environmental CO_2 concentrations (400, 800 and 1200 $\mu\text{atm CO}_2$) for up to three months. Each bar represents the mean $\pm$ s.e.m (N= 5 - 7). Different superscript letters indicate significant differences ($p<0.05$, one-way ANOVA followed by Bonferroni *post hoc* test).

Figure 3. Relative expression (fold change of gene expression using *18S* as the housekeeping gene) of *Slc4a4*, *Slc26a3*, *Slc26a6* in the anterior intestine of sea bream in response to different environmental CO_2 concentrations (400, 800 and 1200 $\mu\text{atm CO}_2$) for up to three months. Each bar represents the mean $\pm$ s.e.m (N= 9 - 10). Different superscript letters indicate significant differences ($p<0.05$, one-way ANOVA followed by Bonferroni *post hoc* test).

Figure 4. Comparative analysis of fold change of gene expression in sea bream juveniles using qPCR of *NKCC2* and *Aquaporin 1b* in the anterior intestine of sea bream in response to different CO_2 concentrations (400, 800 and 1200 $\mu\text{atm CO}_2$) for up to three months. Each bar represents the mean $\pm$ s.e.m (N= 8 - 11). Different superscript letters indicate significant differences ($p<0.05$, one-way ANOVA followed by Bonferroni *post hoc* test).

629 Figure 1 - Gregório et al.

630

631

632 Figure 2 - Gregório et al.

633

634 Figure 3 - Gregório et al.

635

636 Figure 4 - Gregório et al.

637