1 **Morphine selectively promotes glutamate release from glutamatergic terminals**
 2 **of projection neurons from medial prefrontal cortex to dopamine neurons of**
 3 **ventral tegmental area**

4

5 Li Yang, Ming Chen, Ping Zheng

6 State Key Laboratory of Medical Neurobiology, Collaborative Innovation Center for
 7 Brain Science, School of Basic Medical Sciences and Institutes of Brain Science,
 8 Fudan University, Shanghai, China

9

10 **Corresponding author:**

11 Ping Zheng, PhD

12 State Key Laboratory of Medical Neurobiology, Shanghai Medical College, Fudan
 13 University, 138 Yixueyuan Road, Shanghai 200032, People's Republic of China.

14 Phone: (86)(21) 54237437

15 Fax: (86)(21) 64174579

16 E-mail: pzheng@shmu.edu.cn

17

18 **Manuscript Information:**

19 1. The number of words in the abstract: 149

20 2. The number of words in the main text: 1708

21 3. The number of figure (s): 4

22

Abstract

Recently, we found that morphine promoted presynaptic glutamate release of dopamine (DA) neurons in the ventral tegmental area (VTA), which constituted the main mechanism for morphine-induced increase in VTA-DA neuron firing and related behaviors (Chen et al., 2015). However, what source of presynaptic glutamate release of DA neurons in the VTA is promoted by morphine remains unknown. To address this question, we used optogenetic strategy to selectively activate glutamatergic inputs from different projection neurons and then observed the effect of morphine on them. The result shows that morphine promotes glutamate release from glutamatergic terminals of projection neurons from the medial prefrontal cortex (mPFC) to VTA DA neurons, but has no effect on that from the basolateral amygdala (BLA) or the lateral hypothalamus (LH) to VTA DA neurons, and the inhibition of glutamatergic projection neurons from the mPFC to the VTA significantly reduces morphine-induced increase in locomotor activity of mice.

Key words:

morphine; dopamine neurons; ventral tegmental area; optogenetic method; glutamate release; medial prefrontal cortex; basolateral amygdala; lateral hypothalamus

Introduction

Morphine-induced addictive behaviors are strongly dependent on the activation of dopamine (DA) neurons of the ventral tegmental area (VTA) (Wise, 1989, Gardner, 2011, Luscher and Malenka, 2011). One previously reported mechanism for morphine to activate VTA-DA neurons is the disinhibition model of VTA-DA neurons (Johnson and North, 1992, Kalivas, 1993, White, 1996). Recently, we found that morphine could promote presynaptic glutamate release of VTA-DA neurons, which constituted the main mechanism for morphine-induced increase in VTA-DA neurons firing and related behaviors (Chen et al., 2015). However, what source of presynaptic glutamate release of DA neurons in the VTA is promoted by morphine remains unknown.

It has been known that DA neurons of the VTA receive glutamatergic inputs from the medial prefrontal cortex (mPFC), the bilateral amygdala (BLA) and the lateral hypothalamus (LH) (Stuber et al., 2012, Li et al., 2013). To study which glutamatergic inputs onto DA neurons of the VTA are modulated by morphine, we used optogenetic strategy to selectively activate glutamatergic inputs from these different projection neurons and then observed the effect of morphine on them. We also studied the contribution of morphine-induced increase in presynaptic glutamate release of DA neurons in the VTA from specific brain region to morphine-induced increase in locomotor activity of mice, which is an index of enhanced DA function in the VTA (Jalabert et al., 2011).

Results

Morphine promotes glutamate release from glutamatergic terminals of projection neurons from the mPFC to VTA DA neurons, but has no effect on that from the BLA or the LH to VTA DA neurons

To study whether morphine promoted glutamate release from glutamatergic terminals of projection neurons from the mPFC, the BLA or the LH to VTA DA neurons, we used a pair of optical pulses at intervals of 50 ms at 0.1 Hz to evoke excitatory postsynaptic currents (EPSC) in VTA DA neurons from these different brain areas and then observed the effect of morphine on paired-pulse ratio (PPR) of blue light (470 nm)-evoked EPSCs pulses (intervals of 50 ms at 0.1 Hz). PPR, measured as the ratio of EPSC amplitude in response to two successive stimulation pulses, reflects presynaptic release probability; a lower PPR correlates with higher release probability (Zucker and Regehr, 2002, Li et al., 2013). Left panel of **Fig. 1a** showed the expression of ChR2-mCherry in the mPFC at 6 weeks after the injection of the virus into the mPFC and right panel of **Fig. 1a** showed the expression of ChR2-mCherry in the VTA at 6 weeks after the injection of the virus into the mPFC. Left panel of **Fig. 1b** showed typical original traces of the first and second EPSCs before and after morphine (10 μ M) as well as superimposed traces of the first and second EPSCs normalized to the amplitude of the first EPSC in DA neurons of the VTA. The average first EPSC was increased by 50.33 ± 10.39 % ($n = 7$, from 5 animals), whereas the average second EPSC was increased by 8.03 ± 13.71 % ($n = 7$, from 5 animals) at 15 min after morphine. So the superimposition of the two traces normalized to the first EPSC showed that PPR significantly decreased after morphine (left panel of **Fig. 1b**).

The average PPR decreased from 1.01 ± 0.07 before morphine to 0.72 ± 0.1 at 15 min after morphine (right panel of **Fig. 1b**, $n = 7$, from 5 animals, paired t test, $P = 0.0278$, compared to control before morphine). However, morphine ($10 \mu\text{M}$) had no significant effect on PPR evoked by optical stimulation of ChR2-mCherry-positive excitatory terminals from the BLA or the LH to VTA DA neurons. The average PPR was 0.77 ± 0.1 before morphine and 0.8 ± 0.06 at 15 min after morphine in BLA-to-VTA group (right panel of **Fig. 2b**, $n = 11$, from 8 animals, paired t test, $P = 0.8239$, compared to control before morphine) and the average PPR was 0.84 ± 0.07 before morphine and 0.87 ± 0.08 at 15 min after morphine in LH-to-VTA group (right panel of **Fig. 3b**, $n = 7$, from 5 animals, paired t test, $P = 0.8082$, compared to control before morphine). These results suggest that morphine can promote glutamate release from glutamatergic terminals of projection neurons from the mPFC to VTA DA neurons, but has no effect on that from the BLA or the LH to VTA DA neurons.

Inhibition of glutamatergic projection neurons from the mPFC to the VTA significantly reduces morphine-induced increase in locomotor activity of mice

To evaluate the contribution of morphine-induced increase in glutamate release from mPFC-VTA projection neurons to morphine-induced increase in locomotor activity of mice, we observed the influence of the inhibition of glutamatergic projection neurons from the mPFC to the VTA using chemogenetic method (Armbruster et al., 2007) on morphine-induced increase in locomotor activity of mice. We bilaterally injected a retrograde canine adenovirus expressing the Cre recombinase (CAV-GFP-Cre) into

the VTA (Penzo et al., 2015), followed by the injection of the AAV-DIO-hM4Di-mCherry into the mPFC. Six weeks after the infection, the expression of AAV-DIO-hM4D(Gi)-mCherry was observed in the mPFC (left panel of **Fig. 4a**) due to retrograde transported CAV-GFP-Cre from the VTA (right panel of **Fig. 4a**) to the mPFC. This intersectional strategy can effectively target glutamatergic projection neurons from the mPFC to the VTA, which, subsequently, can be suppressed by treating mice with clozapine-N-oxide (CNO), the agonist of hM4Di. On this basis, we measured morphine-induced increase in locomotor activity of mice with and without CNO. The result showed that in control group where mice with hM4D(Gi) were given saline (i.p), morphine (10 mg/kg) could increase locomotor activity of mice. The average distance traveled in morphine+hM4D(Gi)+saline group was 305.4 ± 66.95 cm before and 4913 ± 392.6 cm during 5 min at 15 min after i.p morphine (10 mg/kg, $n = 12$ mice, two-way ANOVA, $F_{(1, 21)} = 11.63$, $*P < 0.001$, compared with control before morphine, left panel of **Fig. 4b**). However, in the CNO group where mice with hM4D(Gi) were given CNO (i.p), morphine-induced increase in locomotor activity of mice significantly decreased. The average distance traveled by mice in morphine+hM4D(Gi)+CNO group was 372.6 ± 51.67 cm before and 2973 ± 356.9 cm during 5 min at 15 min after i.p morphine ($n = 11$ mice, two-way ANOVA, $F_{(1, 21)} = 11.63$, $*P < 0.0001$, compared with control before morphine; $F_{(1, 21)} = 14.22$, $\#P = 0.0011$, compared with morphine+hM4D(Gi)+saline group, left panel of **Fig. 4b**). The average change percentage of distance traveled by mice in morphine+hM4D(Gi)+CNO was 964.1 ± 251.3 %, which was significantly lower than

that (2551 ± 615.9 %) without CNO (independent t test, $P = 0.0313$, compared with morphine+hM4D(Gi)+saline group, right panel of **Fig. 4b**). This result suggests that morphine-induced increase in glutamate release from mPFC-VTA projection neurons contributes to morphine-induced increase in locomotor activity of mice.

Discussion

It has been known that the activation of DA neurons of the VTA plays an important role in both natural stimulus and drug-induced reward response. Under natural state without drugs, direct optogenetic stimulation of VTA DA neurons has been demonstrated to be sufficient to modulate reward-related behaviors (Stuber et al., 2012). In further researches that involved how specific glutamatergic afferents modulate the activity of DA neurons in the VTA, it was found that the LH sent the largest subcortical glutamatergic projection to the VTA (Stuber et al., 2012) and electrical stimulation of the LH predominately increases the firing rates of VTA neurons (Stuber et al., 2012). This evidence suggests that LH-VTA glutamatergic projection neurons may be an important circuit in natural stimulus-induced reward. Another major source of glutamatergic input to the VTA comes from a long-range projection from the mPFC, which can target DA neurons of the VTA (Stuber et al., 2012). Stimulation of the mPFC led to the increase in extracellular glutamate in the VTA (Stuber et al., 2012), the activation of DA neurons of the VTA (Stuber et al., 2012) and the elevation of DA release in the forebrain (Stuber et al., 2012). These evidences suggest that glutamatergic projection neurons from the mPFC to VTA DA neurons

may be another important circuit in natural stimulus-induced reward. In addition, DA neurons of the VTA also receive glutamatergic projections from the BLA (Stuber et al., 2012). However, for the drug-induced reward response, it is still unknown that morphine acts at what glutamatergic pathways to activate DA neurons in the VTA. Here, we find that morphine selectively promotes glutamate release from glutamatergic terminals of projection neurons from the mPFC to the VTA, but has no effect on that from the BLA or the LH to the VTA. This result suggests that morphine may mainly acts at glutamatergic terminals of projection neurons from the mPFC to the VTA to activate DA neurons of the VTA. This result is consistent with the report that the microinjection of morphine into the mPFC cannot produce rewarding effects (Liu et al., 2015) because the site of action of morphine on mPFC-VTA glutamatergic projection neurons is at terminals, rather than at the cell body in the mPFC.

What needs to be explained is that in our previous study in the VTA (Chen et al., 2015), when we use a pair of electrical pulses at intervals of 50 ms at 0.1 Hz to evoke EPSC, it always produces paired pulse facilitation, but here when we use a pair of optical pulses at intervals of 50 ms at 0.1 Hz to evoke EPSC in the VTA, it always produces paired pulse depression. This phenomenon also existed in the study by Vincent Pascoli et al (Pascoli et al., 2011). The reason for this difference remains unknown.

We further evaluate the contribution of morphine-induced increase in glutamate release of mPFC-VTA projection neurons to morphine-induced increase in locomotor activity of mice, which is an index of enhanced DA function in the VTA (Jalabert et al.,

2011). The result shows that the inhibition of glutamatergic projection neurons from the mPFC to the VTA significantly reduces morphine-induced increase in locomotor activity of mice. This result was consistent with the report that the inactivation of the mPFC by the intra-mPFC injection of TTX could eliminate the acute morphine-induced excitation of DA neurons in the VTA (Liu et al., 2015).

Materials and methods

Virus injection

Bilateral injections of purified and concentrated pAAV-CaMKII-hChR2 (H134R)-mCherry, CAV-GFP-cre and AAV-hSyn-DIO-hM4D(Gi)-mCherry virus (2.05×10^{12} vector genomes/ml, Neuron Biotech Company, China) were stereotactically performed in 6-week-old male C57/BL6 mice. Each side of the mPFC (final coordinates: AP, 1.95 mm; ML, $\pm$ 0.33 mm; DV, - 2.4 mm from the skull surface), the BLA (final coordinates: AP, - 1.4mm; ML, $\pm$ 3.0 mm; DV, - 4.8 mm from skull surface) and the LH (final coordinates: AP, - 2.8 mm; ML, $\pm$ 0.4 mm; DV, - 4.2 mm from the skull surface) were respectively injected with 0.5 μ l hChR2 (H134R)-mCherry. Each side of the mPFC (final coordinates: AP, 1.95 mm; ML, $\pm$ 0.33 mm; DV, - 2.4 mm from the skull surface) was injected with 0.5 μ l DIO-hM4D(Gi)-mCherry and each side of the VTA (final coordinates: AP, - 3.08 mm; ML, $\pm$ 0.35 mm; DV, - 4.8 mm from skull surface) was injected with 0.5 μ l CAV-GFP-cre in the same manner (Chen et al., 2015). After the injection of virus, the animals were housed individually and were allowed to recover for over one week.

196

197 **VTA slice preparation**

198 Male mice were anesthetized with chloral hydrate (400 mg/kg, i.p.). All experimental
199 procedures conformed to Fudan University as well as international guidelines on the
200 ethical use of animals. All efforts were made to minimize animal suffering and reduce
201 the number of animals used. VTA slices were prepared according to procedures
202 described previously (Hopf et al., 2007). The brain was removed rapidly from the skull
203 and placed in modified ACSF containing 75 mM sucrose, 88 mM NaCl, 2.5 mM KCl,
204 1.25 mM NaH₂PO₄, 7 mM MgCl₂, 0.5 mM CaCl₂, 25 mM NaHCO₃, and saturated with
205 95% O₂ and 5% CO₂ at ~0°C. Horizontal 250 µm midbrain slices containing VTA were
206 cut on a vibratome (VT-1200, Leica, Wetzlar, Germany) and transferred to normal
207 ACSF containing 126 mM NaCl, 2.5 mM KCl, 1.25 mM NaH₂PO₄, 2 mM MgSO₄, 2.5
208 mM CaCl₂, 25 mM NaHCO₃, and 10 mM glucose at 32°C. Slices were incubated for at
209 least 60 min before patch-clamp recording.

210

211 **In vitro Optogenetics approach for electrophysiology**

212 The medial terminal nucleus of the accessory optic tract (MT) was used as the
213 anatomical structure to define the VTA (Hopf et al., 2007). VTA neurons were
214 visualized on an upright microscope (BX50WI, Olympus, Tokyo, Japan) using infrared
215 differential interference contrast or fluorescent optics. Whole cell current and
216 voltage-clamp recordings were made using an EPC10 amplifier and PatchMaster 2.54
217 software (HEKA, Lambrecht, Germany). Electrodes had a resistance of 3–4 MΩ when

filled with the patch pipette solution. The internal pipette solution contained 130 mM K-gluconate, 8 mM NaCl, 0.1 mM CaCl_2 , 0.6 mM EGTA, 2 mM Mg-ATP, 0.1 mM $\text{Na}_3\text{-GTP}$, and 10 mM HEPES (pH 7.4). Cells were held at -70 mV under a voltage-clamp mode to record evoked EPSC. To observe PPR, two synaptic responses were evoked by flashing 470 nm light (5 ms pulses, 0.1 Hz), which delivered via an optical fiber (core diameter 200 μm , N.A. = 0.39, ThorLabs) coupled to a LED light source (Mightex) 500 μm above the recording cell. The series resistance (R_s) was monitored by measuring the instantaneous current in response to a 5 mV voltage step command. R_s compensation was not used, but cells where R_s changed by $> 15\%$ were discarded.

Identification of VTA-DA neurons

After forming whole-cell recording mode, we identified DA neurons based on electrophysiological characteristics, which included both a spontaneous pacemaker-like firing and expression of a hyperpolarization-induced current (I_h) in the voltage-clamp configuration, by 1 s hyperpolarizing voltage steps (-70 mV to -150 mV) (Grace and Onn, 1989, Margolis et al., 2006, Zhang et al., 2010, Chieng et al., 2011).

Locomotor behavior

The locomotor activity test was conducted as described previously with some modifications (Borgland et al., 2006). The locomotor activity of animals was monitored

with a near infrared video camera within the operant chambers (Med Associates, St. Albans, USA). Distance traveled was measured using Open Field Activity Software (Med Associates) and analyzed locomotion estimates based on movement over a given distance and resting delays (movement in a given period of time). All animals were habituated to the test room for 2 h prior to the start of the experiment. Mice were habituated to the operant chambers for 5 min after 15 min intraperitoneal injection. After 15 minutes drug administration, the mice were placed in the chambers for the 5 min testing session. On days 1, 2, and 3, all mice were only given saline (intraperitoneal injections) to habituate them to the test protocol. On day 4, mice were given drugs according to experimental group. After the behavioral tests, all mice were anesthetized with an overdose of chloral hydrate and perfused with 0.9% saline. The brain was removed and fixed in 4% paraformaldehyde for 24 hr. Coronal sections (80 μ m) were cut by a vibratome and fluorescence was observed under a microscope. Animals where the injection site was outside the VTA were discarded.

Drug

K-gluconate, ATP·K₂, GTP·Na₃, 4-(2-hydroxyethyl) piperazine-1-ethanesulfonic acid (HEPES), ethyleneglycol-bis(b-aminoethyl ether) N,N,N',N'- tetraacetic acid (EGTA), Triton X-100 and 0.01 M PBS were purchased from Sigma. Morphine was from Shenyang No.1 Pharmaceutical Factory, China. Clozapine-N-Oxide (CNO) was from Yihui Biological Technology Co, Shanghai, China. CNO was dissolved in 0.9% saline and administered at 10 mg/kg 30 min before the test (Ray et al., 2011, Li et al., 2013,

Ray et al., 2013, Mahler et al., 2014).

Data analysis

Numerical data were expressed as mean $\pm$ SE. Off-line data analysis was performed using a Mini Analysis Program (Synaptosoft), Clampfit (Axon Instruments), SigmaPlot (Jandel Scientific) and Origin (Microcal Software). Statistical significance was determined using Student's t-test for comparisons between two groups or ANOVA followed by Student-Newman-Keuls test for comparisons among three or more groups, n refers to the number of cells. Every cell was from different slice and a group of cells in each experiment was from at least 5 animals. All post hoc comparisons were made using Tukey's test. Results with $P < 0.05$ were accepted as being statistically significant.

Acknowledgements

This work was supported by the National Program of Basic Research sponsored by the Ministry of Science and Technology of China (2009CB52201 and 2013CB835100), Science and Technology Program of Yunnan Province (2013GA003) and Project of Foundation of National Natural Science of China (31121061, 91332204, 81371466 and 31070932).

Competing interest:

All authors declare no conflict of interest.

284

285 **References**

- 286 Armbruster BN, Li X, Pausch MH, Herlitze S, Roth BL (2007) Evolving the lock to fit the key to create a
287 family of G protein-coupled receptors potentially activated by an inert ligand. *Proceedings of*
288 *the National Academy of Sciences of the United States of America* 104:5163-5168.
- 289 Borgland SL, Taha SA, Sarti F, Fields HL, Bonci A (2006) Orexin A in the VTA is critical for the induction
290 of synaptic plasticity and behavioral sensitization to cocaine. *Neuron* 49:589-601.
- 291 Chen M, Zhao Y, Yang H, Luan W, Song J, Cui D, Dong Y, Lai B, Ma L, Zheng P (2015) Morphine
292 disinhibits glutamatergic input to VTA dopamine neurons and promotes dopamine neuron
293 excitation. *eLife* 4.
- 294 Chieng B, Azriel Y, Mohammadi S, Christie MJ (2011) Distinct cellular properties of identified
295 dopaminergic and GABAergic neurons in the mouse ventral tegmental area. *The Journal of*
296 *physiology* 589:3775-3787.
- 297 Gardner EL (2011) Addiction and brain reward and antireward pathways. *Advances in psychosomatic*
298 *medicine* 30:22-60.
- 299 Grace AA, Onn SP (1989) Morphology and electrophysiological properties of immunocytochemically
300 identified rat dopamine neurons recorded in vitro. *The Journal of neuroscience : the official*
301 *journal of the Society for Neuroscience* 9:3463-3481.
- 302 Hopf FW, Martin M, Chen BT, Bowers MS, Mohamedi MM, Bonci A (2007) Withdrawal from
303 intermittent ethanol exposure increases probability of burst firing in VTA neurons in vitro.
304 *Journal of neurophysiology* 98:2297-2310.
- 305 Jalabert M, Bourdy R, Courtin J, Veinante P, Manzoni OJ, Barrot M, Georges F (2011) Neuronal circuits
306 underlying acute morphine action on dopamine neurons. *Proceedings of the National*
307 *Academy of Sciences of the United States of America* 108:16446-16450.
- 308 Johnson SW, North RA (1992) Opioids excite dopamine neurons by hyperpolarization of local
309 interneurons. *The Journal of neuroscience : the official journal of the Society for*
310 *Neuroscience* 12:483-488.
- 311 Kalivas PW (1993) Neurotransmitter regulation of dopamine neurons in the ventral tegmental area.
312 *Brain research Brain research reviews* 18:75-113.
- 313 Li H, Penzo MA, Taniguchi H, Kopec CD, Huang ZJ, Li B (2013) Experience-dependent modification of a
314 central amygdala fear circuit. *Nature neuroscience* 16:332-339.
- 315 Liu C, Fang X, Wu Q, Jin G, Zhen X (2015) Prefrontal cortex gates acute morphine action on dopamine
316 neurons in the ventral tegmental area. *Neuropharmacology* 95:299-308.
- 317 Luscher C, Malenka RC (2011) Drug-evoked synaptic plasticity in addiction: from molecular changes to
318 circuit remodeling. *Neuron* 69:650-663.
- 319 Mahler SV, Vazey EM, Beckley JT, Keistler CR, McGlinchey EM, Kaufling J, Wilson SP, Deisseroth K,
320 Woodward JJ, Aston-Jones G (2014) Designer receptors show role for ventral pallidum input
321 to ventral tegmental area in cocaine seeking. *Nature neuroscience* 17:577-585.
- 322 Margolis EB, Lock H, Hjelmstad GO, Fields HL (2006) The ventral tegmental area revisited: is there an
323 electrophysiological marker for dopaminergic neurons? *The Journal of physiology*
324 577:907-924.
- 325 Pascoli V, Turiault M, Luscher C (2011) Reversal of cocaine-evoked synaptic potentiation resets

326 drug-induced adaptive behaviour. *Nature* 481:71-75.

327 Penzo MA, Robert V, Tucciarone J, De Bundel D, Wang M, Van Aelst L, Darvas M, Parada LF, Palmiter RD,

328 He M, Huang ZJ, Li B (2015) The paraventricular thalamus controls a central amygdala fear

329 circuit. *Nature* 519:455-459.

330 Ray RS, Corcoran AE, Brust RD, Kim JC, Richerson GB, Nattie E, Dymecki SM (2011) Impaired

331 respiratory and body temperature control upon acute serotonergic neuron inhibition. *Science*

332 333:637-642.

333 Ray RS, Corcoran AE, Brust RD, Soriano LP, Nattie EE, Dymecki SM (2013) Egr2-neurons control the

334 adult respiratory response to hypercapnia. *Brain research* 1511:115-125.

335 Stuber GD, Britt JP, Bonci A (2012) Optogenetic modulation of neural circuits that underlie reward

336 seeking. *Biological psychiatry* 71:1061-1067.

337 White FJ (1996) Synaptic regulation of mesocorticolimbic dopamine neurons. *Annual review of*

338 *neuroscience* 19:405-436.

339 Wise RA (1989) Opiate reward: sites and substrates. *Neuroscience and biobehavioral reviews*

340 13:129-133.

341 Zhang TA, Placzek AN, Dani JA (2010) In vitro identification and electrophysiological characterization of

342 dopamine neurons in the ventral tegmental area. *Neuropharmacology* 59:431-436.

343 Zucker RS, Regehr WG (2002) Short-term synaptic plasticity. *Annual review of physiology* 64:355-405.

344

345 **Legends**

346 **Figure 1.** Effect of morphine on paired-pulse ratio (PPR) of light-evoked EPSC in DA

347 neurons of the VTA from the mPFC. (a) Left panel: image of coronal brain slice

348 showing expression of ChR2-mCherry (red color) in the mPFC at 6 weeks after the

349 injection of the virus into the mPFC. Right panel: image of coronal brain slice showing

350 expression of ChR2-mCherry in the VTA at 6 weeks after the injection of the virus into

351 the mPFC. (b) Left panel: typical original traces of the first and second EPSCs evoked

352 by a pair of blue light pulses (470 nm, intervals of 50 ms at 0.1 Hz) before and after

353 morphine (10 μ M) as well as superimposed traces of the first and second EPSCs

354 normalized to the amplitude of the first EPSC in DA neurons of the VTA. Right panel:

355 average PPR evoked by a pair of blue light pulses (470 nm, intervals of 50 ms at 0.1

356 Hz) before and after morphine (10 μ M) in VTA DA neurons from the mPFC (n = 7 cells

from five mice, $P < 0.05$, compared to control before morphine).

Figure 2. Effect of morphine on paired-pulse ratio (PPR) of light-evoked EPSC in DA neurons of the VTA from the BLA. (a) Left panel: image of coronal brain slice showing expression of ChR2-mCherry (red color) in the BLA at 6 weeks after the injection of the virus into the BLA. Right panel: Image of coronal brain slice showing expression of ChR2-mCherry in the VTA at 6 weeks after the injection of the virus into the BLA. (b) Left panel: typical original traces of the first and second EPSCs evoked by a pair of blue light pulses (470 nm, intervals of 50 ms at 0.1 Hz) before and after morphine (10 μ M) as well as superimposed traces of the first and second EPSCs normalized to the amplitude of the first EPSC in DA neurons of the VTA. Right panel: average PPR evoked by a pair of blue light pulses (470 nm, intervals of 50 ms at 0.1 Hz) before and after morphine (10 μ M) in VTA DA neurons from the BLA ($n = 11$ cells from eight mice, $P > 0.05$, compared to control before morphine).

Figure 3. Effect of morphine on paired-pulse ratio (PPR) of light-evoked EPSC in DA neurons of the VTA from the LH. (a) Left panel: image of coronal brain slice showing expression of ChR2-mCherry (red color) in the LH at 6 weeks after the injection of the virus into the LH. Right panel: image of coronal brain slice showing expression of ChR2-mCherry in the VTA at 6 weeks after the injection of the virus into the LH (b) Left panel: typical original traces of the first and second EPSCs evoked by a pair of blue light pulses (470 nm, intervals of 50 ms at 0.1 Hz) before and after morphine (10 μ M)

as well as superimposed traces of the first and second EPSCs normalized to the amplitude of the first EPSC in DA neurons of the VTA. Right panel: average PPR evoked by a pair of blue light pulses (470 nm, intervals of 50 ms at 0.1 Hz) before and after morphine (10 μ M) in VTA DA neurons from the LH (n = 7 cells from five mice, P > 0.05, compared to control before morphine).

Figure 4. Influence of the inhibition of glutamatergic projection neurons from the mPFC to the VTA using chemogenetic method on morphine-induced increase in locomotor activity of mice. (a) Left panel: image of coronal brain slice showing AAV-hSyn-DIO-hM4D(Gi)-mCherry in the mPFC. Right panel: images of coronal brain slice showing CAV-GFP-cre in the VTA. (b) Left panel: average distance traveled by mice in morphine+hM4D(Gi)+saline (i.p, 0.2 ml/mice) group (n = 12, *P < 0.001, compared with control before morphine) and in morphine+hM4D(Gi)+CNO (i.p, 0.2 ml/mice group (n = 11, *P < 0.001, compared with control before morphine, #P = 0.0016, compared with morphine+hM4D(Gi)+saline) during 5 min before and at 15 min after morphine (i.p, 10mg/kg). Right panel: average change percentage of distance traveled by mice in morphine+hM4D(Gi)+saline group (n = 12) and in morphine+hM4D(Gi)+CNO group (n = 11, P = 0.0313, compared with morphine+hM4D(Gi)+saline). Data are shown as the mean $\pm$ s.e.m.

Figure 1

Figure 2

Figure 3

Figure 4