1 **Running title:** Biocontrol bacteria alleviate replanting problem

2

3 **Inoculation of biocontrol bacteria alleviated *Panax ginseng* replanting problem**

4

5 **Linlin Dong, Jiang Xu, Lianjuan Zhang, Guangfei Wei, He Su, Juan Yang, Jun**
6 **Qian, Ran Xu, Baosheng Liao, Liang Shen, Mingli Wu, Ruiyang Cheng, Shilin**
7 **Chen***

8 **Key Laboratory of Beijing for Identification and Safety Evaluation of Chinese**
9 **Medicine, Institute of Chinese Materia Medica, China Academy of Chinese**
10 **Medical Sciences, Beijing 100700, China.**

11

12 **Highlight:** Changes in rhizospheric microbial communities driven by ginseng plants
13 of different ages and developmental stages could cause microecological degradation.
14 Biocontrol using microbial antagonists effectively alleviated the replanting problem.

15

16 **Linlin Dong E-mail: lldong@icmm.ac.cn; Jiang Xu E-mail: jxu@icmm.ac.cn;**

17 **Lianjuan Zhang E-mail: zlj0970@163.com;**

18 **Guangfei Wei E-mail: 15098855936@163.com; He Su E-mail: suhe.gd@qq.com;**

19 **Juang Yang E-mail: yangjuan6263@163.com;**

20 **Jun Qian E-mail: gianjun@biozeron.com; Ran Xu E-mail: 53509011@qq.com;**

21 **Baosheng Liao E-mail : swjs082lbs@126.com;**

22 **Liang Shen E-mail: lshen@icmm.ac.cn; Mingli Wu E-mail: justuswu@163.com;**

23 **Ruiyang Cheng E-mail: 18612545661@163.com;**

24 **Shilin Chen* *To whom correspondence should be addressed**

25 **Tel: 0086-10-57203877; Fax: 0086-10- 62899776**

26 **E-mail: slchen@icmm.ac.cn**

27 **Linlin Dong and Jiang Xu have equal contributions to this work.**

28 **The date of submission: 5/29/2017**

29 **With 5121 words, 7 figures, and supplementary data (Figures 6 and Tables 5).**

30

31 **Abstract**

32 Replanting problem is a common and serious issue hindering the continuous
33 cultivation of *Panax* plants. Changes in soil microbial community driven by plant
34 species of different ages and developmental stages are speculated to cause this
35 problem. Inoculation of microbial antagonists is proposed to alleviate replanting
36 issues efficiently.

37 High-throughput sequencing revealed that bacterial diversity evidently decreased,
38 and fungal diversity markedly increased in soils of adult ginseng plants in the root
39 growth stage. Relatively few beneficial microbe agents, such as *Luteolibacter*,
40 Cytophagaceae, *Luteibacter*, *Sphingomonas*, Sphingomonadaceae, and Zygomycota,
41 were observed. On the contrary, the relative abundance of harmful microorganism
42 agents, namely, Brevundimonas, Enterobacteriaceae, *Pandoraea*, Cantharellales,
43 *Dendryphon*, *Fusarium*, and Chytridiomycota, increased with plant age. Furthermore,
44 *Bacillus subtilis* 50-1 was isolated and served as microbial antagonists against
45 pathogenic *Fusarium oxysporum* of ginseng root-rot, and its biocontrol efficacy was
46 67.8% using a dual culture assay. The ginseng death rate and relative abundance of
47 *Fusarium* decreased by 63.3% and 46.1%, respectively, after inoculation with 50-1
48 in replanting soils. Data revealed that changes in the diversity and composition of
49 rhizospheric microbial communities driven by ginseng of different ages and
50 developmental stages could cause microecological degradation. Biocontrol using
51 microbial antagonists was an effective method for alleviating the replanting problem.

52

53 **Keywords**

54 *Panax ginseng*; microbial communities; plant age; developmental stage; replanting
55 problem; biocontrol.

56

57 **Introduction**

58 *Panax ginseng* C.A. Meyer demonstrates anti-inflammatory and antitumor effects,
 59 and it is commonly used in traditional Chinese medicine (Choo *et al.*, 2008; Ernst,
 60 2010). The current annual global market value of this species is approximately 3.5
 61 billion dollars (Hong *et al.*, 2006). Wild ginseng resources are becoming scarce
 62 because of excessive and predatory exploitation, and wild ginseng is gradually
 63 substituted with cultivated ginseng in mainstream market (Li *et al.*, 2012; Wu *et al.*,
 64 2013). Ginseng is continuously cultivated in fixed plots for 4–5 years, and
 65 subsequent replanting usually fails because of continuous cropping obstacles (Ying
 66 *et al.*, 2012a). Several years of crop rotation is needed in successful replanting.
 67 Large-scale deforestation increases to satisfy the market demand for ginseng.
 68 Moreover, conflict between the operation of ginseng industry and environmental
 69 conservation has worsened. Replanting issue is a severe drawback hindering the
 70 development of ginseng industry. Hence, this problem must be addressed urgently.

71 Factors such as deterioration of soil physicochemical properties,
 72 allelopathy/autotoxicity, outburst of soil-borne disease, and soil microbial
 73 community changes cause replanting problem (Ogweno & Yu, 2006; Wu *et al.*, 2008;
 74 Huang *et al.*, 2013). Changes in soil microbial community is one of the main factors
 75 hindering crop replantation (Nayyar *et al.*, 2009; Manici *et al.*, 2013). Xiao *et al.*
 76 (2016) reported that the lack of balance in rhizospheric microbial communities is one
 77 of the key factors causing discontinuous ginseng cultivation. Microbial communities
 78 exhibit evident changes during ginseng cultivation, thereby causing an imbalance in
 79 microbial community (Ying *et al.*, 2012b). Increase in the relative abundance of
 80 pathogenic microorganism could lead to the occurrence of soil-borne disease. Taken
 81 together, changes in rhizospheric microbial community are suggested to cause the
 82 replanting issues.

83 Many biotic or abiotic factors lead to changes in rhizospheric microbial
 84 communities. Plants of different ages can alter microbial community (Wu *et al.*,
 85 2015). Differences in diversity and composition of soil microbial community were
 86 observed under continuous cropping of *P. notoginseng*; moreover, fungal diversity

and microbial taxa in soil microbial community are related to seedling mortality (Dong *et al.*, 2016). Li *et al.* (2012) found that differences between rhizospheric and non-rhizospheric soil microbial community in a particular site tend to become increasingly significant with ginseng growth. *Panax* plants with different ages in a site drive changes in microbial community. Plant developmental stage influences microbial community structure and activity, and marked shifts in diversity and relative activity are observed in soil microbial communities of different developmental stages (Houlden *et al.*, 2008). However, the information whether *Panax* plants of different ages and developmental stages mediate microbial community is unclear. We hypothesized that changes in the rhizospheric microbial community driven by ginseng plants of different ages and developmental stages could cause microecological degradation.

Root rot is a severe disease that hinders the replantation of *Panax* plants (Guo *et al.*, 2009). This soil-borne disease is caused by the pathogenic fungus *Fusarium oxysporum*, which is the main agent in *Panax* plants (Miao *et al.*, 2006; Dong *et al.*, 2016). The relative abundance of *F. oxysporum* increased with the notoginseng cultivation and death rate of seedlings was significantly related to its abundance (Dong *et al.*, 2016). Thus decrease in the abundance of pathogenic *F. oxysporum* could alleviate the occurrence of root-rot. Biological control using microbial antagonists has attracted interest as an effective method to control plant pathogens due to its non-toxic nature (Zheng *et al.*, 2011). Many studies reported that biocontrol bacteria, such as *Pseudomonas fluorescens* and *Bacillus amyloliquefaciens* RWL-1, show evident growth inhibition activity against *F. oxysporum* in tomato (Kamilova *et al.*, 2006; Shahzad *et al.*, 2017). Nevertheless, microbial antagonists against ginseng root rot is scarce, and application of biocontrol bacteria could be effective to alleviate the replanting problem.

Recently, herbgonomics has been raised for medicinal plants research using genomic tools including investigating rhizospheric environment using metagenomic sequencing (Chen *et al.*, 2015). In the present study, high-throughput sequencing analysis of 16S and 18S rRNA genes was used to demonstrate changes in the

diversity and composition of soil microbial communities in the rhizosphere of ginseng seedlings of different ages and developmental stages. Furthermore, biocontrol bacteria against *F. oxysporum* were isolated using a dual culture technique, and their inhibition activity in a replanting soil was confirmed. These results increased our understanding of the reasons behind replanting issues caused by rhizospheric microbial community. Data provided an effective method for soil bioremediation to alleviate issues in replanting of Chinese herbal medicine.

124

125 **Materials and Methods**

126 **Field experiment and soil extraction**

This field experiment was performed in a ginseng plantation in Jingyu, Jilin Province (42°20'N, 126°50'E, 775 m a.s.l.), which is the main ginseng-producing region in China. This region receives a northern temperate continental climate and an annual precipitation of approximately 767 mm. The plough layer in the plantation consists of gray brown soil.

Ginseng seedlings are consecutively grown for 4–5 years in a fixed site before their root can be harvested. The disease and mortality of ginseng seedlings generally occur after being consecutively grown for 2 years. Thus, we analyzed the influence of two-, three-, and four-year-old transplanted seedlings on rhizospheric microbial communities. In this experiment, two-, three-, and four-year-old ginseng seedlings were transplanted in each plot in our plantation and denoted as 2-y, 3-y, and 4-y, respectively. In our plantation, field plots were arranged in a completely randomized block design, and three replicate plots (1.7 m × 8.0 m) were used for each plant age. Ginseng was cultivated strictly in accordance with the standard operating procedures of good agricultural practice (Heuberger *et al.*, 2010; Zhang *et al.*, 2010).

The distinct stages of ginseng development are as follows: vegetative, flowering, fruiting, root growth, and annual dormancy (Table S1). During dormancy, the aboveground plant parts of ginseng wither, and the root becomes dormant. In this stage, the underground root activities weaken, and soil samples obtained during this stage are not included in analyses. This experiment included 36 soil samples

147 obtained from two-, three-, and four-year-old ginseng seedlings under four
148 developmental stages, namely, vegetative (2-Ve, 3-Ve, and 4-Ve), flowering (2-Fl,
149 3-Fl, and 4-Fl), fruiting (2-Fr, 3-Fr, and 4-Fr), and root growth (2-Ro, 3-Ro, and
150 4-Ro). Six ginseng seedlings were randomly obtained from each plot (1.7 m × 8.0 m);
151 the soils were removed, and rhizosphere fractions were brushed and pooled into one
152 sample. Soil samples were obtained from three replicates per treatment. Thirty-six
153 soil samples were homogenized by passing through a 2 mm sieve for further
154 processing. The characteristics of the soil samples are described in Table S2.

155

156 **Calculation of emergence and survival rates**

157 The ginseng emergence rate was analyzed in May. The emergence rate in each plot
158 was calculated as follows: number of emerging ginseng seedlings divided by the
159 total number of transplanted seedlings. The mortality of ginseng seedlings was
160 observed from June to August, and the survival rate was determined at the end of
161 August. The survival rate of ginseng in each plot was calculated as follows: number
162 of living seedlings divided by the total number of emerging seedlings in the area.
163 Three plots per treatment served as three replicates.

164

165 **DNA extraction and PCR amplification**

166 Total soil DNA was extracted from 0.1 g of freeze-dried soil using a MoBio
167 Powersoil Kit (MoBio Laboratories Inc., Carlsbad, CA) according to the
168 manufacturer's instructions; the DNA was stored at −20 °C for further processing.
169 For each sample, fragments of 16S and 18S rRNA genes were amplified using the
170 conserved primers 27F/338R (Fierer *et al.*, 2008) and 817F/1196R (Rousk *et al.*,
171 2010). The forward and reverse primers contained an eight-base pair barcode (Table
172 S3). Amplification reactions and purification were performed as previously
173 described (Rodrigues *et al.*, 2013). Purified PCR products were quantified by
174 Qubit®3.0 (Life Invitrogen, Germany), and these amplicons were pooled in
175 equimolar ratios for sequencing.

176

177 **High-throughput sequencing**

178 The pooled DNA product was used to construct Illumina pair-end library according
179 to the Illumina's genomic DNA library preparation procedure. The amplicon library
180 was subsequently paired-end sequenced (2×250) on an Illumina HiSeq platform
181 (Shanghai Biozeron Co., Ltd.) according to standard protocols.

182 Raw FASTQ files were demultiplexed and quality-filtered using QIIME with
183 the following pipeline (Caporaso *et al.*, 2010). Operational taxonomic units were
184 clustered using 97% similarity cutoff by using UPARSE (version 7.1
185 <http://drive5.com/uparse/>), and chimeric sequences were identified and removed
186 using UCHIME. The phylogenetic affiliation of each 16S and 18S rRNA gene
187 sequence was analyzed using Ribosomal Database Project (Wang *et al.*, 2007) and
188 Silva schemes (Quast *et al.*, 2013). Rarefaction analysis based on Mothur v.1.21.1
189 (Schloss *et al.*, 2009) was performed to reveal diversity indexes, including Chao 1
190 and Shannon diversity (H') indexes. We clustered the taxa obtained from the RDP
191 Classifier through complete linkage hierarchical clustering by using R package
192 HCLUST (<http://sekhon.berkeley.edu/stats/html/hclust.html>). PCoA was used to
193 compare groups of samples based on unweighted UniFrac distance metrics in QIIME
194 (Caporaso *et al.*, 2010). Linear discriminant analysis effect size (LEfSe)
195 (<http://huttenhower.sph.harvard.edu/lefse/>) was used to characterize the features
196 differentiating the bacterial and fungal communities in soils according to the
197 methods described by Segata *et al.* (2011). All metagenomic data were submitted to
198 the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov>);
199 the accession numbers of 16S rRNA and 18S rRNA genes are SRP103168 and
200 SRP103176, respectively.

201

202 **Quantitative PCR of *Fusarium***

203 To compare the relative abundance of *Fusarium* in soils of ginseng seedlings, the
204 copy numbers of *Fusarium* were calculated using the ITS-Fu-F/ITS-Fu-R
205 (Abd-Elsalam *et al.*, 2003). Quantitative PCR was performed according to the
206 description of Rousk *et al.* (2010) with minor modification. Standard curves were

generated using 10-fold serial dilution of a plasmid containing a full-length copy of *F. oxysporum* 18S rRNA gene to estimate the copy numbers of *Fusarium*. qPCR reactions (25 µL) were performed using a SYBR Green PCR Master mix (Takara, Toyobo, Japan). *Fusarium* copy numbers were generated using a regression equation for each assay relating the cycle threshold value to the known number of copies in the standards.

213

Isolation and selection of antagonistic bacteria

Biological control is one of the most remarkable potential approaches for plant disease control. Due to the increase in the abundance of pathogenic agents of ginseng root rot,

a dual culture assay was used to screen microbial antagonists against *F. oxysporum*.

Rhizosphere soils of three-year ginseng seedlings under root growth stage were used

to isolate and select the antagonistic bacteria against *F. oxysporum*. The pathogenic *F.*

oxysporum was isolated and confirmed in our previous study (Wang *et al.*, 2016).

Soil (10 g) was homogenized in 100 mL of sterile distilled water, and the bacteria

were isolated through serial dilution technique. The isolated single strain was

screened on the basis of its antagonistic activity against *F. oxysporum* in a dual

culture according to the description of Shahzad *et al.* (2017). The experiment was

replicated three times. The zone of inhibition was measured following the

description of Kaiser *et al.* (2005) to screen the antagonistic bacterium and examine

the antagonistic activity of the candidates.

229

Identification of antagonistic bacterial strain

Antagonistic bacterium (named strain 50-1) was identified by morphological and

molecular methods. The morphology of strain 50-1 was recorded after it was

incubated on Luria–Bertani (yeast extract (5 g), peptone (10 g), NaCl (10 g), and

agar (10 g) in 1 L of water) medium for 24 h later. Strain 50-1 was molecularly

identified by amplifying 16S rRNA according to the description by Cai *et al.* (2012).

The amplified PCR product was analyzed on a 3730 XL sequencer (Applied

237 Biosystems, Foster City, CA, USA), and the generated sequence (accession number
238 KY962803) was submitted to GenBank. Neighbor-joining trees were constructed in
239 MEGA v6.0 to generate Kimura 2-parameter distance matrixes for each sequence
240 following standard parameters. The numbers at the branch knots were bootstrap
241 values based on 1000 resamplings for the maximum likelihood.

242 To further identify the strain 50-1, total genomic DNA from strain 50-1 was
243 extracted (Tiangen, Beijing, China) and purified by RNase-free DNase I (Takara,
244 Kyoto, Japan) to analyze its genome. The complete genome sequence was assembled
245 using a hybrid sequencing strategy combining PacBio RS II and Illumina HiSeq
246 sequencing platforms (Koren *et al.*, 2012). Genome sequencing was performed with
247 Illumina HiSeq2500 using the PE250 strategy following the manufacturer's protocol.
248 The reads obtained with Illumina PCR adapter and the filtered low-quality reads
249 were assembled by SOAPdenovo (Li *et al.*, 2008; Li *et al.*, 2010) to generate
250 scaffolds. Gene prediction genome assembly was performed using Glimmer and
251 gene functions were annotated by BLASTP against NR, COG and KEGG databases.
252 GeneMarkS (Besemer *et al.*, 2001) with an integrated model combining the
253 GeneMarkS-generated parameters and Heuristic model parameters. A genome
254 overview with annotation information was created using Circos (Krzywinski *et al.*,
255 2009). The whole genome was deposited in NCBI (BioProject ID PRJNA383782).

256

257 **Evaluation of the biocontrol efficacy of antagonists in replanting soils**

258 A pot experiment was performed to assess the biocontrol efficacy of bacterium 50-1
259 in replanting soils in our phytotron. Each pot contained 1 kg of soils cultivated with
260 ginseng seedlings for three years and happened root rot. Two-year-old ginseng
261 seedlings (three plants) were transplanted in each pot. The pots were placed in the
262 phytotron under the following conditions: 26 ± 2 °C, 60% humidity, and 14 h of light
263 alternating with 10 h of darkness. The rhizospheres of uniform ginseng seedlings
264 were inoculated with 1 mL (10^6 cfu mL⁻¹) of cultures containing biocontrol strains
265 when the seedlings were cultivated for one week. Pots were inoculated with cultures
266 containing inactivated strains as controls. Five pots served as one replicate, and three

replicates were prepared. After a two months following inoculation, the death rate of ginseng seedlings was calculated as follows: number of dead seedlings divided by the total number of transplanted seedlings in each treatment. The rhizosphere soils of three seedlings were randomly selected from each treatment, and they served as one sample for the analysis of the relative abundance of *Fusarium*. The experiment was replicated three times.

Statistical analyses

SPSS version 16.0 software (SPSS Inc., Chicago, IL, USA) was used for statistical analyses. Variables were considered for all treatment replicates and subjected to ANOVA. Mean values were compared by calculating the least significant difference at 5% level.

Results

Emergence and survival rates of ginseng seedlings

The ginseng plants grown on field showed a healthy appearance with no sign of disease during the growing season (Figs. 1A–1C). The emergence and survival rates exhibited no significant difference among two-, three-, or four-year-old ginseng seedlings (Figs. 1D, 1E). The emergence rate of two-, three-, or four-year-old ginseng seedlings ranged from 82.3% to 86.9% (Fig. 1D), and their survival rate was 94.1%–97.1% (Fig. 1E). The growth of ginseng seedlings of different ages was favorable, which guaranteed the success of the field experiment.

Taxonomic diversity of bacterial and fungal communities

A total of 2,296,684 classifiable 16S rRNA sequence reads were obtained from 36 soil samples for analysis (Table S3). The mean number of classifiable sequences per sample was 63,796 (dominant length: 283–293 bp). The bacterial diversity indexes (H' and Chao 1) markedly changed in the rhizosphere of ginseng seedlings of different ages in late developmental stages (Fig. 2). H' and Chao 1 values were significantly higher in the rhizosphere of 4-y seedlings than those in the rhizosphere

of 2-y seedlings in fruiting stage (Figs. 2A, 2B). During root growth, H' and Chao 1 values evidently decreased in the rhizospheres of 3-y and 4-y seedlings than those in the rhizosphere of 2-y seedlings.

A total of 1,321,521 classifiable fungi sequences with a mean of 36,709 sequences were obtained per soil sample (dominant length: 411–414 bp; Table S3). H' and Chao 1 values markedly decreased in the rhizosphere with increasing ginseng age under vegetative stage (Figs. 2C, 2D). During root growth, H' and Chao 1 values were evidently higher in the soils of 3-y and 4-y seedlings than those in the soil of 2-y seedlings.

Variation in bacterial and fungal community compositions

PCoA ordination and Bray–Curtis distance matrix revealed evident difference in bacterial communities in soils of ginseng plant of different ages and developmental stages (Figs. 3, S1). In vegetative and flowering stages, the second principal components (14.41% and 14.22% contributions, respectively) demonstrated that the bacterial communities in soils of 2-y seedlings differed from those in soils of 4-y seedlings (Figs. 3A, 3B). In fruiting stage, the first principal component axis (22.00% contributions) indicated that the bacterial communities in soils of 2-y seedlings significantly differed from those in soils of 3-y seedlings (Fig. 3C). Additionally, the first principal component axis (22.29% contributions) indicated that the bacterial communities in soils of 3-year-old seedlings significantly differed from those in soils of 2-y and 4-y seedlings; furthermore, the second principal component (13.98% contributions) demonstrated that the bacterial communities in soils of 2-y seedlings differ from those in soils of 4-y seedlings (Fig. 3D). LEfSe revealed differences in the rhizospheric bacterial communities of ginseng seedlings of different ages and developmental stages (Fig. S2).

PCoA ordination and Bray–Curtis distance matrix revealed that fungal communities obviously differed in the soils of ginseng seedlings of different ages and developmental stages (Figs. 3, S1). In vegetative stage, the first principal component axis (33.94% contribution) of the fungal communities in soils of 4-y

seedlings markedly differed from those of the fungal communities in soils of 2-y seedlings. Moreover, the second principal component axis (17.55% contribution) of the fungal communities in soils of 2-year-old seedlings considerably differed from those of the fungal communities in the soils of 3-y seedlings (Fig.3E). At the flowering and fruiting stages, the first principal components (25.71% and 38.96% contributions, respectively) in the fungal communities in soils of 2-y ginseng seedlings significantly differed from those of the fungal communities in the soils of 3-y and 4-y seedlings (Figs. 3F, 3G). During root growth, the first principal component (26.97% contribution) of fungal communities in soils of 3-y seedlings considerably differed from those of fungal communities in soils of 2-y and 4-y seedlings, and the second principal component (14.93% contribution) in fungal communities in soils of 2-y seedlings differed from those of fungal communities in soils of 4-y seedlings (Fig. 3H). According to LEfSe analysis, fungal composition differed in the soils of ginseng seedlings of different ages and developmental stages (Fig. S3).

342

343 **Changes in the relative abundance of bacterial taxa**

344 The relative abundance of bacterial groups changed in soils of ginseng plants of
345 different ages and developmental stages (Fig. 4). The relative abundance of
346 Chthoniobacteraceae, Chthonomonadales, Chthonomonadetes, *Chthoniobacter*,
347 *Granulicella*, and *Blastocatella* declined with plant age in vegetative stage (Fig. 4A).
348 The relative abundance of *Arthrobacter*, *Brevundimonas*, Micrococcaceae,
349 Rhodobiaceae, Intrasporangiaceae, and Micrococcales was significantly higher in
350 soils of 3-y and 4-y seedlings than that in soils of 2-y seedlings. The relative
351 abundance of *Luteolibacter*, Phyllobacteriaceae, *Acidovorax*, *Moheibacter*, and
352 Cytophagaceae in soils of 3-y and 4-y seedlings evidently decreased, and the
353 abundance of Elusimicrobia and Armatimonadetes was markedly increased in the
354 flowering stage (Fig. 4B). The relative abundance of *Bacillus*, Enterobacteriales,
355 Enterobacteriaceae, *Brevundimonas*, and Anaerolineae was significantly higher in
356 soils of 3-y and 4-y seedlings than that in soils of 2-y seedlings; additionally, the

abundance of *Luteibacter*, Clostridia, and Clostridiales declined in soils of 3-y and 4-y seedlings in fruiting stage (Fig. 4C). The relative abundance of *Paralcaligenes*, Sphingomonadaceae, Saccharibacteria, *Sphingomonas*, and Alcaligenaceae significantly decreased with plant age, whereas the relative abundance of *Pandora*, Chlamydiales, and Chlamydiae was obviously higher in soils of 3-y and 4-y seedlings than that in soils of 2-y seedlings in root growth stage (Fig. 4D).

Changes in the relative abundance of fungal taxa

The relative abundance of fungal taxa changed in soils of ginseng plants of different ages and developmental stages (Fig. 5). The relative abundance of Cystofilobasidiales, Ophiostomataceae, *Ophiostoma*, Ophiostomatales, and Cystofilobasidiaceae significantly declined with seedling age, and the abundance of Pezizales, Cantharellales, *Dendryphon*, Pezizomycetes, and Tubeufiaceae was evidently higher in soils of 4-y seedlings than those in soils of 2-y and 3-y seedlings in the vegetative stage (Fig. 5A). The relative abundance of Microascales, *Helicoma*, and Tubeufiaceae was obviously higher in soils of 4-y seedling than those in soils of 2-y and 3-y seedlings in the flowering stage (Fig. 5B). The relative abundance of Tremellales, Acrospermales, *Occultifur*, *Acrospermum*, Cystobasidiales, Cystobasidiaceae, and Cystobasidiomycetes was markedly higher in the soils of 4-y seedlings than that in the soils of 2-y and 3-y seedlings in fruiting stage (Fig. 5C). The relative abundance of Zygomycota was obviously lower in soils of 3-y and 4-y seedlings than that in soils of 2-y seedlings, and the abundance of Tremellomycetes, Chytridiomycota, and Sordariales significantly increased in soils of 4-y seedlings than those in soils of 2-y and 3-y seedlings in root growth stage (Fig. 5D).

The relative abundance of *Fusarium* showed increasing trends in the soils of the adult ginseng plants under later developmental stages (Figs. 5E, S4). According to the high-throughput sequencing analysis, the relative abundance of *Fusarium* increased by 22.5%–25.0%, 35.7%–50.0%, and 18.2%–36.4% in soils of 3-y and 4-y seedlings in the flowering, fruiting, and root growth stages, respectively (Fig. 5E). The copy numbers of *Fusarium* also showed similar trends based on the analysis of

quantitative PCR (Fig. S4). These results revealed that the relative abundance of pathogenic agent increased in soils during ginseng growth.

Bacillus subtilis* 50-1 were responsible for the biocontrol of *F. oxysporum

Due to the relative abundance of pathogenic agent (*Fusarium*) increase in the ginseng soils, dual culture techniques were used to isolate microbial antagonists against *F. oxysporum* to control root rot. Antagonistic bacterium, namely, *B. subtilis* 50-1, was isolated and confirmed using a dual culture technique (Fig. 6). Strain 50-1 exhibited a broad spectrum of growth inhibition activity against *F. oxysporum*, thereby resulting in 67.8% inhibition percentage (Fig.6A). Strain 50-1 is a Gram-positive, oxidase- and catalase-positive, rod-shaped bacterial species (Fig.6B, Table S4). Analysis of the 16S rRNA sequences revealed that strain 50-1 belonged to *B. subtilis* with the bootstrap value of 100% (Fig. 6C). Strain 50-1 was further confirmed according to the genome sequencing. The complete genome was composed of a circular chromosome of 4,040,837 bp with 43.86% GC content (Fig. 6D, Table S5). The total numbers of genes were 4,193, which covered 88.6% of the genome encoding 3,176 proteins. The function of these annotated genes was mainly energy production and conversion, amino acid transport and metabolism, transcription, carbohydrate transport and metabolism, inorganic ion transport and metabolism, and signal transduction mechanisms based on the COG function classification (Fig. S5).

Inoculation of biocontrol bacteria reduced ginseng death rate in a replanting soil

The pot experiment analysis revealed that ginseng death rate and the relative abundance of *Fusarium* significantly decreased by 63.3% and 46.1% in the replanting soils inoculated with strain 50-1, respectively (Fig. 7). Furthermore, the height and leaf area increased by 62.7% and 22.5%, respectively, after inoculation with strain 50-1 (Fig. S6). These results revealed that inoculation of biocontrol bacteria alleviated the ginseng replanting problem.

417

418 **Discussion**

419

420 Replanting problem is a common and serious issue in cultivation of medicinal plants.
 421 Following replantation, *Rehmannia glutinosa* suffers from disease, and its biomass
 422 and tuber quality decline (Qi *et al.*, 2009; Wu *et al.*, 2011; Yang *et al.*, 2011). The
 423 survival rate of ginseng seedlings is lower than 25% after replantation for three years
 424 (Zhao *et al.*, 2001). Multiple factors have caused this replanting problem, and
 425 researchers suggested that changes in microbial communities influence soil health
 426 and crop yield (Bisseling *et al.*, 2009; Bulgarelli *et al.*, 2013). In addition,
 427 rhizospheric microbial community is governed by plant species and growth (Incroğlu
 428 *et al.*, 2011). Thus, we hypothesized that changes in the rhizospheric microbial
 429 community were driven by ginseng seedlings of different ages and development
 430 stages, which caused the replanting problem of *Panax* plants. Reducing the
 431 occurrence of the disease is a useful strategy to alleviate the replanting issues.
 432 Biological control is one of the most remarkable potential approaches for plant
 433 disease control because of its safety and environmental friendliness.

434 Diversity in bacterial and fungal communities changed in the rhizosphere of
 435 ginseng seedlings of different ages and developmental stages. H' and Chao1 values
 436 revealed that bacterial diversity was obviously low, whereas fungal diversity was
 437 significantly higher in the rhizosphere of adult ginseng seedlings in the root growth
 438 stage. A similar study reported that increase in ginseng cultivation ages reduced
 439 bacterial diversity and increased fungal diversity (Xiao *et al.*, 2016). Moreover,
 440 diversity of microbial community in the rhizosphere of *Pseudostellaria heterophylla*
 441 decreases with increased number of cropping years (Zhao *et al.*, 2016). Additionally,
 442 the developmental stage of crops is an important driver of microbial community
 443 structure (Houlden *et al.*, 2008). Plant ages and developmental stages alter the
 444 microbial diversity during growth. The diversity of soil microorganisms is critical in
 445 the maintenance of soil health and quality, and it serves as a sensitive bioindicator
 446 (He *et al.*, 2008). The death rate of notoginseng and fungal diversity are significantly

negatively correlated, which suggest that fungal diversity is a potential bioindicator of soil health (Dong *et al.*, 2016). A relationship exists between microbial diversity and root disease suppression (Nitta, 1991; Workneh & van Bruggen, 1994). Mazzola (2004) reported that reduced soil microbial diversity is responsible for the development of soil-borne diseases. Reduced bacterial diversity in response to adult plants is possibly the key indicator of ecological variations and functional impairment in a root growth stage.

The bacterial and fungal compositions evidently differed in the rhizospheres of ginseng plants of different ages and developmental stages. Variation in bacterial and fungal community composition is observed during continuous cropping of notoginseng (Dong *et al.*, 2016). Dynamics of microbial species is occurred in the rhizosphere during ginseng growth (Li *et al.*, 2012). Sugiyama *et al.* (2014) reported that soybeans affected the bacterial communities under vegetative stage, with further alterations possibly occurring during later growth stages. The most divergent community structures occur in the young plant stage of tomato, and the flowering and senescence stages show increasingly similar community structure (Inceoğlu *et al.*, 2011). The composition of microbial community was affected by plant age and development stage. In our study, the agents of beneficial microbes were relatively few, whereas those of harmful microorganisms increased with plant age. The agents of beneficial microbes include the microorganisms that degrade compounds (*Luteolibacter*, Cytophagaceae, *Luteibacter*, and *Sphingomonas*), and resist disease and improve growth (Sphingomonadaceae and Zygomycota). The agents of harmful microorganisms mainly infect plants (*Brevundimonas*, Enterobacteriaceae, *Pandoraea*, Cantharellales, *Dendryphon*, and Chytridiomycota). Another study has showed similar results, i.e., the population of beneficial microbe decreases, whereas that of harmful microorganisms increases with increasing number of cropping years (Zhao *et al.*, 2016). Soil microbial community is an important bioindicator of soil function (Zuppinger-Dingley *et al.*, 2014). Changes in functional groups revealed that micro-ecological environment was degraded in the rhizosphere with increasing age of ginseng.

477 Rhizospheric microbial community can be influenced by soil characteristics and
 478 plant species (Lauber *et al.*, 2008; Berg and Smalla, 2009). In our study, the pH and
 479 available K and organic matter contents exhibited no significant difference in
 480 ginseng plants of different ages and developmental stages (Table S2). Plant species
 481 could be one of the most important factors that influence rhizobacterial communities
 482 (Micallef *et al.*, 2009). Root exudates offer substrates for microbial metabolism and
 483 act as intermediates for biogeochemical reactions in rhizosphere (Uren, 2000; Berg
 484 and Smalla, 2009). In the early developmental stages of *Arabidopsis*, cumulative
 485 secretion levels of sugars and sugar alcohols are high, which enhanced the richness
 486 of rhizospheric microbial community (Chaparro *et al.*, 2014). By contrast, root
 487 exudates contain allelochemical that disturbs the balance in a microbial community
 488 (Wu *et al.*, 2015). These results imply that root exudates are one of the main drivers
 489 of changes in rhizospheric microbial communities during ginseng growth. Moreover,
 490 our results showed that the diversity of microbial communities markedly changed in
 491 the rhizosphere of ginseng plants with different ages in root growth stage. This
 492 phenomenon is possibly a result of different root types and root exudates. The
 493 composition of *Arabidopsis* root exudates changes throughout developmental
 494 gradient in plants, and tomato root exudates in reproductive stage are more
 495 phytotoxic than those in vegetative stage (Yu and Matsui, 1994; Chaparro *et al.*,
 496 2013). Ginseng root rapidly grows after cropping for three years before harvest,
 497 especially in root growth stage. Ginseng root growth stage is possibly characterized
 498 by a specific but different root exudation pattern that drives different bacterial
 499 communities.

500 Biological control using microbes is an environmentally friendly approach of
 501 controlling disease (Bargabus *et al.*, 2003; Tjamos *et al.*, 2005). In our study, we
 502 isolated *B. subtilis* 50-1 as an effective antagonist against *F. oxysporum*; the results
 503 of antagonist inoculation revealed that biocontrol bacterium reduced the ginseng
 504 morbidity and alleviated the replanting problem. *Bacillus* strains, as potent biological
 505 control agents of plant diseases, have been reported in many studies (Zouari *et al.*,
 506 2016; Shahzad *et al.*, 2017). The biocontrol efficacy of *B. amyloliquefaciens* 54

507 against bacterial fruit blotch is evident in a greenhouse (Jiang *et al.*, 2015). *B.*
508 *megaterium* (B5), *B. cereus* sensu lato (B25), and *Bacillus* sp. (B35) display the
509 highest antagonistic activity against *Fusarium verticillioides*, and antagonistic
510 activity reveals that these strains produce glucanases, proteases, chitinases,
511 siderophores, and auxins (Figueroa-López *et al.*, 2016). Furthermore, data suggest
512 that microbial strains producing bioactive constituents can help the inoculated plant
513 to reduce the negative effects of pathogenesis and abiotic stresses (Shahzad *et al.*,
514 2017). The microbial antagonists could be used for efficient control of plant
515 pathogens and comprise an environmentally friendly approach. Additionally, genome
516 sequence analysis provides insights into the pathways of functional bacteria and
517 facilitates their exploration. The genome sequence analysis of *Ganoderma lucidum*
518 revealed key genes encoding cytochrome P450s in secondary metabolism (Chen *et*
519 *al.*, 2012). In our study, the genome sequence of strain 50-1 helps accelerate the
520 development and application of biological inoculant.

521

522 **Supplementary data**

523 The details were online.

524 **Table S1** Description of developmental stages of ginseng and the sampling times.

525 **Table S2** Soil chemical characteristics in soils of ginseng cropping.

526 **Table S3** Barcodes, numbers of bacterial and fungal sequences and average length in
527 each sample.

528 **Table S4** Physiological and biological characteristics of *Bacillus subtilis* 50-1.

529 **Table S5** Genome features of *Bacillus subtilis* strain 50-1.

530 **Figure S1** Changes in bacterial and fungal community in rhizosphere of ginseng
531 seedlings of different ages and developmental stages.

532 **Figure S2** LEfSe identified the taxa in the soil bacterial communities of plants of
533 different ages in vegetative (A), flowering (B), fruiting (C), and root growth stages
534 (D).

535 **Figure S3** LEfSe identified the taxa in the fungal communities of plants of different
536 ages in vegetative (A), flowering (B), fruiting (C), and root growth stages (D).

537 **Figure S4** Relative abundance of *Fusarium* in the soils based on the analysis of
538 real-time PCR.

539 **Figure S5** COG function classification in the *Bacillus subtilis* strain 50-1.

540 **Figure S6** The height (A) and leaf area (B) of ginseng seedlings after inoculation of
541 strain 50-1 in a replanting soil.

542

543 **Acknowledgements**

544 This study was supported by grants from the National Science Foundation of China
545 (81603238 and 81403053).

546

References

- Abd-El salam KA, Aly IN, Abdel-Satar MA, Khalil MS, Verreet JA.** 2003. PCR identification of *Fusarium* genus based on nuclear ribosomal-DNA sequence data. *African Journal of Biotechnology* **2**, 82–85.
- Bargabus RL, Zidack NK, Sherwood JE, Jacobsen BJ.** 2003. Oxidative burst elicited by *Bacillus mycoides* isolate Bac J: a biological control agent, occurs independently of hypersensitive cell death in sugar beet. *Molecular Plant-Microbe Interactions* **16**, 1145–1153.
- Besemer J, Lomsadze A, Borodovsky M.** 2001. GeneMarkS: a self-training method for prediction of gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. *Nucleic Acids Research* **29**, 2607–2618.
- Berg G, Smalla K.** 2009. Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *FEMS Microbiology Ecology* **68**, 1–13.
- Bisseling T, Dangl JL, Schulze-Lefert P.** 2009. Next-generation communication. *Science* **324**, 691.
- Bulgarelli D, Schlaeppi K, Spaepen S, Ver Loren van Themaat E, Schulze-Lefert P.** 2013. Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology* **64**, 807–838.
- Cai Z, Chen Q, Wang H, He Y, Wang W, Zhao X, Ye Q.** 2012. Degradation of the novel herbicide ZI0273 by *Amycolatopsis* sp. M3-1 isolated from soil. *Applied Microbiology and Biotechnology* **96**, 1371–1379.
- Caporaso JG, Bittinger K, Bushman FD, DeSantis TZ, Andersen GL, Knight R.** 2010. PyNAST: a flexible tool for aligning sequences to a template alignment. *Bioinformatics* **26**: 266–267.
- Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM.** 2013. Root exudation of phytochemicals in *Arabidopsis* follows specific patterns that are developmentally programmed and correlate with soil microbial functions. *PLoS One* **8**, e55731.
- Chaparro JM, Badri DV, Vivanco JM.** 2014. Rhizosphere microbiome assemblage

- is affected by plant development. *The ISME Journal* **8**, 790–803.
- Chen S, Song J, Sun C, et al.** 2015. Herbal genomics: Examining the biology of traditional medicines. *Science* **347(6219)**, S27–S29.
- Chen S, Xu J, Liu C, et al.** 2012. Genome sequence of the model medicinal mushroom *Ganoderma lucidum*. *Nature Communications* **3**, 913.
- Choo MK, Sakurai H, Kim DH, Saiki I.** 2008. A ginseng saponin metabolite suppresses tumor necrosis factor-alpha-promoted metastasis by suppressing nuclear factor-kappa B signaling in murine colon cancer cells. *Oncology Reports* **19**, 595–600.
- Dong L, Xu J, Feng G, Li X, Chen S.** 2016. Soil bacterial and fungal community dynamics in relation to *Panax notoginseng* death rate in a continuous cropping system. *Scientific Reports* **6**, 31802.
- Ernst E.** 2010. *Panax ginseng*: an overview of the clinical evidence. *Journal of Ginseng Research* **34**, 259–263.
- Fierer N, Hamady M, Lauber CL, Knight R.** 2008. The influence of sex, handedness, and washing on the diversity of hand surface bacteria. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 17994–17999.
- Figuerola-López AM, Cordero-Ramírez JD, Martínez-Alvarez JC, López-Meyer M, Lizárraga-Sánchez GJ, Félix-Gastélum R, Castro-Martínez C, Maldonado-Mendoza IE.** 2016. Rhizospheric bacteria of maize with potential for biocontrol of *Fusarium verticillioides*. *SpringerPlus* **5**, 1–12.
- Guo R, Liu X, Li S, Miao Z.** 2009. In vitro inhibition of fungal root-rot pathogens of *Panax notoginseng* by rhizobacteria. *The Plant Pathology Journal* **25**, 70–76.
- He JZ, Zheng Y, Chen CR, He YQ, Zhang LM.** 2008. Microbial composition and diversity of an upland red soil under long-term fertilization treatments as revealed by culture-dependent and culture-independent approaches. *Journal of Soil and Sediments* **8**, 349–358.
- Heuberger H, Bauer R, Friedl F, Heubl G, Hummelsberger J, Nögel R, Seidenberger R, Torres-Londoño P, Planta M.** 2010. Cultivation and

- breeding of Chinese medicinal plants in Germany. *Planta Medica* **76**, 1956–1962.
- Hong SG, Lee KH, Kwak J, Bae KS.** 2006. Diversity of yeasts associated with *Panax ginseng*. *Journal of Microbiology* **44**, 674–679.
- Huang LF, Song LX, Xia XJ, Mao WH, Shi K, Zhou YH, Yu JQ.** 2013. Plant-soil feedbacks and soil sickness: From mechanisms to application in agriculture. *Journal of Chemical Ecology* **39**, 232–242.
- Houlden A, Timms-Wilson TM, Day MJ, Bailey MJ.** 2008. Influence of plant developmental stage on microbial community structure and activity in the rhizosphere of three field crops. *FEMS Microbiology Ecology* **65**, 193–201.
- Inceoglu Ö, Al-Soud WA, Salles JF, Semenov AV, van Elsas JD.** 2011. Comparative analysis of bacterial communities in a potato field as determined by pyrosequencing. *PLoS One* **6**, e23321.
- Jiang CH, Wu F, Yu ZY, Xie P, Ke HJ, Li HW, Yu YY, Guo JH.** 2015. Study on screening and antagonistic mechanisms of *Bacillus amyloliquefaciens* 54 against bacterial fruit blotch (BFB) caused by *Acidovorax avenae* subsp. *citrulli*. *Microbiological Research* **170**, 95.
- Kamilova F, Kravchenko LV, Shaposhnikov, AI, Makarova N, Lugtenberg B.** 2006. Effects of the tomato pathogen *Fusarium oxysporum* f. sp. *radicis-lycopersici* and of the biocontrol bacterium *Pseudomonas fluorescens* WCS365 on the composition of organic acids and sugars in tomato root exudate. *Molecular Plant-Microbe Interactions* **19**, 1121–1126.
- Kaiser C, van der Merwe R, Bekker TF, Labuschagne N.** 2005. *In-vitro* inhibition of mycelial growth of several phytopathogenic fungi, including *Phytophthora cinnamomi* by soluble silicon. *South African Avocado Growers' Association Yearbook* **28**, 70–74.
- Koren S, Schatz MC, Walenz BP, et al.** 2012. Hybrid error correction and *de novo* assembly of single-molecule sequencing reads. *Nature Biotechnology* **30**, 693–700.
- Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ,**

- Marra MA.** 2009. Circos: an information aesthetic for comparative genomics. *Genome Research* **19**, 1639–1645.
- Lauber CL, Strickland MS, Bradford MA, Fierer N.** 2008. The influence of soil properties on the structure of bacterial and fungal communities across land-use types. *Soil Biology Biochemistry* **40**, 2407–2415.
- Li R, Li Y, Kristiansen K, Wang J.** 2008. SOAP: short oligonucleotide alignment program. *Bioinformatics* **24**, 713–714.
- Li R, Zhu H, Ruan J, et al.** 2010. De novo assembly of human genomes with massively parallel short read sequencing. *Genome Research* **20**, 265–272.
- Li Y, Ying Y, Zhao D, Ding W.** 2012. Microbial community diversity analysis of *Panax ginseng* rhizosphere and non-rhizosphere soil using randomly amplified polymorphic DNA method. *Open Journal of Genetics* **2**, 95–102.
- Manici LM, Kelderer M, Franke-Whittle IH, Rühmer T, Baab G, Nicoletti F, Caputo F, Topp A, Insam H, Naef A.** 2013. Relationship between root-endophytic microbial communities and replant disease in specialized apple growing areas in Europe. *Applied Soil Ecology* **72**, 207–214.
- Mazzola M.** 2004. Assessment and management of soil community structure for disease suppression. *Annual Review of Phytopathology* **42**, 35–59.
- Miao Z, Li S, Liu X, Chen Y, Li Y, Wang Y, Guo R, Xia Z, Zhang K.** 2006. The causal microorganisms of *Panax notoginseng* root rot disease. *Scientia Agricultura Sinica* **39**, 1371–1378.
- Micallef SA, Shiaris MP, Colón-Carmona A.** 2009. Influence of *Arabidopsis thaliana* accessions on rhizobacterial communities and natural variation in root exudates. *Journal of Experimental Botany* **60**, 1729–1742.
- Nayyar A, Hamel C, Lafond G, Gossen B, Hanson K, Germida J.** 2009. Soil microbial quality associated with yield reduction in continuous-pea. *Applied Soil Ecology* **43**, 115–121.
- Nitta T.** 1991. Diversity of root fungal floras: Its implications for soil-borne diseases and crop growth. *Jarq-japan Agricultural Research Quarterly* **25**, 6–11.
- Ogwen JO, Yu JQ.** 2006. Autotoxic potential in soil sickness: a Re-examination.

- Allelopathy Journal **18**, 93–101.
- Qi J, Yao H, Ma X, Zhou L, Li X.** 2009. Soil microbial community composition and diversity in the rhizosphere of a Chinese medicinal plant. Communications in Soil Science and Plant Analysis **40**, 1462–1482.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.** 2013. The SILVA ribosomal RNA gene database project: improved data processing and Web-based tools. Nucleic Acids Research **41**, D590–D596.
- Rodrigues JLM, Pellizari VH, Mueller R, et al.** 2013. Conversion of the Amazon rainforest to agriculture results in biotic homogenization of soil bacterial communities. Proceedings of the National Academy of Sciences of the United States of America **110**, 988–993.
- Rousk J, Bååth E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, Knight R, Fierer N.** 2010. Soil bacterial and fungal communities across a pH gradient in an arable soil. The ISME Journal **4**, 1–12.
- Schloss EP, Flanagan DM, Culler CL, Wright AL.** 2009. Some hidden costs of faculty turnover in clinical departments in one academic medical center. Academic Medicine **84**, 32–36.
- Segata N, Abubucker S, Goll J, et al.** 2011. Microbial community function and biomarker discovery in the human microbiome. Genome Biology **12**, P47.
- Shahzad R, Khan AL, Bilal S, Asaf S, Lee IJ.** 2017. Plant growth-promoting endophytic bacteria versus pathogenic infections: an example of *Bacillus amyloliquefaciens* RWL-1 and *Fusarium oxysporum* f. sp. lycopersici in tomato. Peer Journal **5**, e3107.
- Sugiyama A, Ueda Y, Zushi T, Takase H, Yazaki K.** 2014. Changes in the bacterial community of soybean rhizospheres during growth in the field. PLoS One **9**, e100709.
- Tjamos SE, Flemetakis E, Paplomatas EJ, Katinakis P.** 2005. Induction of resistance to *Verticillium dahliae* in *Arabidopsis thaliana* by the biocontrol agent K-165 and pathogenesis-related proteins gene expression. Molecular

- Plant-Microbe Interactions **18**, 555–561.
- Uren NC.** 2000. Types, amounts and possible functions of compounds released into the rhizosphere by soil grown plants. In: Pinton R, Varani Z, Nanniperi P, eds. *The rhizosphere: biochemistry, and organic substances at the soil interface*. New York, NY, USA: Marcel Dekker Inc., 19–40.
- Wang R, Dong L, Xu J, Zhang N, Naoki F, Li X, Shen L, Chen S.** 2016. Identification and prevention of root rot pathogen in model of ginseng cultivated in farmlands. *China Journal of Chinese Materia Medica* **41**, 1787–1791.
- Wang Q, Garrity GM, Tiedje JM, Cole J.** 2007. Naïve Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* **73**, 5261–5267.
- Workneh F, van Bruggen AHC.** 1994. Microbial diversity, composition, and diversity in organically and conventionally managed rhizosphere soil in relation to suppression of corky root of tomatoes. *Applied soil Ecology* **1**, 219–230.
- Wu L, Wang J, Huang W, Wu H, Chen J, Yang Y, Zhang Z, Lin W.** 2015. Plant-microbe rhizosphere interactions mediated by *Rehmannia glutinosa* root exudates under consecutive monoculture. *Scientific Reports* **5**, 15871.
- Wu H, Yang H, You X, Li Y.** 2013. Diversity of endophytic fungi from roots of *Panax ginseng* and their saponin yield capacities. *SpringerPlus* **2**, 107.
- Wu LJ, Zhao YH, Guan YM, Pang SF.** 2008. A review on studies of the reason and control methods of succession cropping obstacle of *Panax ginseng* C.A. Mey. *Special Wild Economic Animal and Plant Research* **2**, 68–72.
- Wu L, Wang H, Zhang Z, Lin R, Zhang Z, Lin W.** 2011. Comparative metaproteomic analysis on consecutively *Rehmannia glutinosa*-monocultured rhizosphere soil. *PLoS One* **6**, e20611.
- Xiao C, Yang L, Zhang L, Liu C, Han M.** 2016. Effects of cultivation ages and modes on microbial diversity in the rhizosphere soil of *Panax ginseng*. *Journal of Ginseng Research* **40**, 28–37.
- Yang Y, Chen X, Chen J, Xu H, Li J, Zhang Z.** 2011. Differential miRNA

- expression in *Rehmannia glutinosa* plants subjected to continuous cropping. BMC Plant Biology **11**, 53.
- Ying YX, Ding WL, Li Y.** 2012. Characterization of soil bacterial communities in rhizospheric and nonrhizospheric soil of *Panax ginseng*. Biochemical Genetics **50**, 848–859.
- Ying Y, Ding W, Zhou Y, Li Y.** 2012. Influence of *Panax ginseng* continuous cropping on metabolic function of soil microbial communities. Chinese Herbal Medicine **4**, 329–334.
- Yu JQ, Matsui Y.** 1994. Phytotoxic substances in root exudates of cucumber (*Cucumis sativus* L.). Journal of Chemical Ecology **20**, 21–31.
- Zhang B, Peng Y, Zhang Z, Liu H, Qi Y, Liu S, Xiao P.** 2010. GAP production of TCM herbs in China. Planta Medica **76**, 1948–1955.
- Zhao RF.** 2001. The mechanism of re-plantation problem in ginseng and American ginseng. Special Wild Economic Animal and Plant Research **1**, 40–45.
- Zhao YP, Lin S, Chu L, Gao J, Azeem S, Lin W.** 2016. Insight into structure dynamics of soil microbiota mediated by the richness of replanted *Pseudostellaria heterophylla*. Scientific Reports **6**, 26175.
- Zheng Y, Xue QY, Xu LL, Xu Q, Lu S, Gu C, Guo JH.** 2011. A screening strategy of fungal biocontrol agents towards Verticillium wilt of cotton. Biocontrol Control **56**, 209–216.
- Zouari I, Jlaiel L, Tounsi S, Trigui M.** 2016. Biocontrol activity of the endophytic of the endophytic *Bacillus amyloliquefaciens* strain CEIZ-11 against *Pythium aphanidermatum* and purification of its bioactive compounds. Biological Control **100**, 54–62.
- Zuppinger-Dingley D, Schmid B, Petermann JS, Yadav V, De Deyn GB, Flynn DF.** 2014. Selection for niche differentiation in plant communities increases biodiversity effects. Nature **515**, 108–111.

548 **Figure captions**

549

550 **Figure 1** Growth status of ginseng seedlings. (A–C) Two-, three-, and four-year-old
551 ginseng seedlings, respectively. (D) Emergence rate of ginseng seedlings. (E)
552 Survival rate of ginseng seedlings. 2-y, 3-y, and 4-y represents the transplanted two-,
553 three-, and four-year-old seedlings, respectively. Data are presented as mean $\pm$ SD (n
554 = 3). Identical letters denote nonsignificant difference at 0.05 significance level.

555

556 **Figure 2** Bacterial and fungal diversities in the rhizosphere of ginseng seedlings of
557 different ages and developmental stages. (A, B) show the Chao 1 and Shannon
558 diversity (H') of bacterial community. (C, D) show the Chao 1 and H' of fungal
559 community. Ve, Fl, Fr, and Ro represent vegetative, flowering, fruiting, and root
560 growth stages, respectively. Data are presented as mean $\pm$ SD ($n = 3$). Non-identical
561 letters denote significant difference in the same developmental stage of ginseng plant
562 with different ages at 0.05 significance level.

563

564 **Figure 3** Changes in bacterial and fungal communities in the rhizosphere of ginseng
565 seedlings of different ages and developmental stages. (A–D) PCoA ordination plots
566 display the relatedness of samples separated using Unweighted UniFrac distance of
567 classified 16S rRNA gene sequences at vegetative, flowering, fruiting, and root
568 growth stages, respectively. (E–H) PCoA ordination plots show the relatedness of
569 samples separated using Unweighted UniFrac distance of classified 18S rRNA gene
570 sequences at the Ve, Fl, Fr, and Ro stages, respectively. Red, green, and blue
571 represent the samples in the rhizosphere of 2-y, 3-y, and 4-y transplanted ginseng
572 seedlings, respectively.

573

574 **Figure 4** Relative abundance of the bacterial taxa detected by the linear discriminant
575 analysis effect size (LEfSe) as biomarker. (A–D) represent the vegetative, flowering,
576 fruiting, and root growth stages, respectively. 2-, 3- and 4- represent the samples in
577 the rhizosphere of 2-y, 3-y, and 4-y transplanted ginseng seedlings. Data represent

578 the mean values of $n = 3$.

579

580 **Figure 5** Relative abundance of the fungal taxa detected by LEfSe as biomarker and
581 *Fusarium*. (A–D) represent the vegetative, flowering, fruiting, and root growth
582 stages, respectively. (E) Relative abundance of *Fusarium*. Data are presented as
583 mean $\pm$ SD ($n = 3$).

584

585 **Figure 6** *Bacillus subtilis* 50-1 antagonized *Fusarium oxysporum*. (A) Colony
586 diameter measured in a dual culture. (B) Morphological features of bacterium 50-1.
587 (C) Relationships of 16S rRNA sequences between *B. subtilis* strain 50-1 (black
588 body) and published 16S rDNA sequences. (D) Genome map of strain 50-1. The six
589 circles (outer to inner) represent the scale line, forward strand CDSs (color by COG
590 categories), reverse strand CDSs (color by COG categories), RNA genes, GC content,
591 and GC skew. From outside to center: genome size, genes on the forward strand
592 (color by COG categories), genes on the reverse strand (color by COG categories),
593 RNA genes (tRNAs, orange; rRNAs, red), GC content (red and blue), and GC skew.
594 Only bootstrap values higher than 70% are shown. Bars represent the mean $\pm$ SE (n
595 = 3). Asterisks denote significant differences between the colony diameters of *F.*
596 *oxysporum* and *F. oxysporum* + strain 50-1 at $P < 0.05$.

597

598 **Figure 7** Inoculation of strain 50-1 in ginseng replanting soils. (A) Death rate of
599 ginseng after inoculation with strain 50-1. (B) Copy numbers of *Fusarium* in soils.
600 No-inoculation and inoculation represent the samples inoculated with cultures
601 containing inactivated and activated strain 50-1, respectively. Data are presented as
602 mean $\pm$ SD ($n = 3$). Asterisks denote significant differences between the
603 No-inoculation and inoculation at $P < 0.05$.

604

605

606 **Figure 1**

607

608

609 **Figure 2**

610

611

612 **Figure 3**

613

614

615 **Figure 4**

616

617

618 **Figure 5**

621 **Figure 6**

622

623

624 **Figure 7**

625

626