Structure of *Allium* lachrymatory factor synthase elucidates catalysis on sulfenic acid substrate

Takatoshi Arakawa ^{a 1 2}, Yuta Sato ^{a 1}, Jumpei Takabe ^{a 1}, Noriya Masamura ^b,
Masahiro Kato ^b, Morihiro Aoyagi ^b, Takahiro Kamoi ^b, Nobuaki Tsuge ^b,
Shinsuke Imai ^b, Shinya Fushinobu ^a

^aGraduate School of Agricultural and Life Sciences, The University of Tokyo,
Tokyo, 113-8657, Japan. ^b Basic Research Division, Central Research &
Development Institute, House Foods Group Inc., Yotsukaido, 284-0033, Japan.

Classification: Agricultural science in Biological Sciences

Keywords: lachrymatory factor, isoalliin, isomerase, SRPBCC, pericyclic
reaction, sulfenic acid, reactive sulfur species, pyrabactin-resistant protein,
polyketide cyclase

Abstract

Natural lachrymatory effects are invoked by small volatile *S*-oxide compounds. They are produced through alkene sulfenic acids by the action of lachrymatory factor synthase (LFS). Here we present the crystal structures of onion LFS (AcLFS) revealed in solute-free and two solute-stabilized forms. Each structure adopts a single seven-stranded helix-grip fold possessing an internal pocket. Mutagenesis analysis localized the active site to a layer near the bottom of the pocket, which is adjacent to the deduced key residues Arg71, Glu88, and Tyr114. Solute molecules visible on the active site have suggested that AcLFS accepts various small alcohol compounds as well as its natural substrate, and they inhibit this substrate according to their chemistry. Structural homologs have been found in the SRPBCC superfamily, and comparison of the active sites has demonstrated that the electrostatic potential unique to AcLFS could work in capturing the substrate in its specific state. Finally, we propose a rational catalytic mechanism based on intramolecular proton shuttling in which the microenvironment of AcLFS can bypass the canonical [1,4]-sigmatropic rearrangement principle established by microwave studies. Beyond revealing

35 how *AcLFS* generates the lachrymatory compound, this study provides insights
36 into the molecular machinery dealing with highly labile organosulfur species.

37

Significance statement

Crushing of onion liberates a volatile compound, *syn*-propanethial *S*-oxide (PTSO), which causes lachrymatory effect on humans. We present the crystal structures of onion LFS (*AcLFS*), the enzyme responsible for natural production of PTSO. *AcLFS* features a barrel-like fold, and mutagenic and inhibitory analyses revealed that the key residues are present in the central pocket, harboring highly concentrated aromatic residues plus a dyad motif. The architecture of *AcLFS* is widespread among proteins with various biological functions, such as abscisic acid receptors and polyketide cyclases, and comparisons with these homologs indicate that unique steric and electronic properties maintain the pocket as a reaction compartment. We propose the molecular mechanism behind PTSO generation and shed light on biological decomposition of short-lived sulfur species.

Introduction

Organosulfur compounds are often accumulated in vegetables, and they have improved the quality of human life in the form of particular sensory experiences as well as nutritional and health benefits (1-5). Pungent smells evoked by the *Allium* plants stem from cytosolic (+)-*S*-alk(en)yl cysteine sulfoxides (CSOs), which are produced abundantly from γ -glutamyl-*S*-alk(en)yl cysteines (6). CSOs are odorless precursors, but disruption stresses such as injury, biting and cutting allow them to contact vacuolar C-S lyase (alliinase; EC 4.4.1.4), after which spontaneous decomposition and polymerization occur (Fig. 1). The population of mostly volatile end products [*e.g.*, dialk(en)yl thiosulfinates (5), cepathiolanes (6), cepaenes (7), polysulfides (8), and zweieberanes (9)] contributes to determining alliaceous smells and flavors (7, 8).

In addition, crushing of bulb onion (*Allium cepa*) and other limited plants induce weeping. *syn*-Propanethial *S*-oxide (PTSO) (3) is a prototype of lachrymatory agents (9) and generated by the conjunct catalysis initiated by the alliinase-mediated cleavage of *trans*-*S*-1-propenyl L-cysteine sulfoxide

(*trans*-1-PRENCISO/isoalliin) (**1**) (7). Lachrymatory factor synthase (LFS; EC 5.3.-.-) serves on the subsequent conversion to PTSO, and thus branches the common spontaneous degradation pathway of CSOs (10) (Fig. 1). Onion LFS (AcLFS, 169 aa) is assumed to be an irreversible isomerase against (*E*)-1-propene-1-sulfenic acid (1-PSA) (**2**), but molecular machinery of AcLFS remains unveiled, only implicating that the two residues (R71 and E88) associate in this catalysis (11). Before the discovery of LFS, Block and colleagues have established a canonical theory of PTSO formation through 1-PSA, termed [1,4]-sigmatropic rearrangement (Fig. 5A), employing flash vacuum pyrolysis (FVP) (12). This non-enzymatic, intramolecular pericyclic reaction-based principle proceeds under vacuum and heat, while AcLFS mediates an equivalent reaction under ambient conditions and without the assistance of any cofactors.

LFS is a unique enzyme because it acts on a the short-lived intermediate. Sulfenic acid (R-SOH) is one of the reactive sulfur species (RSS), which is completely sensitive to thiol and sulfenyl groups as well as oxidants due to its bifacial preference as a nucleophile and an electrophile (13, 14). Sulfenic acids

appear as a protein modification of cysteine residues (15-17) and function in specific regulated activities (18-24). However, they have rarely been identified as substrates due to them being too labile to be detected, with an estimated half-life of several milliseconds (25) and not more than a minute for methane sulfenic acid generated in the vacuum gas phase (26). In organisms lacking LFS genes, for example, garlic (*Allium sativa*), free sulfenic acids are condensed immediately, and thus are just destined to produce dialk(en)yl thiosulfinates (2, 6). Gene silencing of onion LFS derives increases of (5) and (9) isomers, displaying a metabolite composition similar to that of garlic (8, 27).

Here, we present the crystal structures of AcLFS. They enable us to elucidate the natural mechanism by which coincident and spontaneous RSS reactions are selectively modified by this enzyme.

Results

Overall structure and confirmation of the active site

Crystal structures of AcLFS were determined at a resolution of 1.7 to 2.1 Å (Table S1). In the present models, the N-terminal region (1–22), which is not required for the activity (11), was not determined due to conformational disorder. AcLFS (23–169) formed a single globular domain with dimensions of 40×40×43 Å, consisting of a seven-stranded antiparallel β-sheet (S1–S7), a long α-helix (H3), two flanking helices (H1, H2), and connecting loops (Fig. 2A). The molecular architecture in which the β-sheet bends so as to wrap H3 is known as a helix-grip fold (28).

There are five cysteine residues on the molecular periphery of AcLFS (Fig. S1A). Inactivation of the free thiols by an excess amount of iodoacetamide (IAA) caused no significant change in PTSO generation (Fig. 2D). Hence, the catalysis does not involve any cysteine residues, which is also supported by the result that the activity of C107A remained at the same level as that of intact AcLFS.

AcLFS possesses an elongated pocket extending 18 Å from the molecular surface. This pocket can be divided into three layers, termed Gate, Wall, and Floor (Fig. 2B). Wall and Floor are pocket-lining apparatuses provided by side chains of S1, S3–7, and H2–3, and consist mainly of aromatic and aliphatic residues, respectively (Fig. 2C, S4). A series of mutagenesis analyses proved that the active site resides in the Wall region. Specifically, variants of Wall residues (F84A, Y102A, F104A, Y114A, W133A, W155A) exhibited equivalent reductions in activities to less than 10%, while those of Floor (L47A, M51A, L54A, V73A, M118A) and a part of Gate (C107A, T109A, W112Q, M143A) maintained the activities roughly at the level of the wild type (Fig. 2D). Bulky side chains of the Wall residues appear to be necessary for the catalysis since substitutions that lead to maintenance of the aromatic rings (*e.g.*, Y102F and F104Y) partially retained the activities compared with their alanine mutants.

R71 and E88, which are suspected of being involved in catalysis (11), were positioned proximally at the boundary of Wall and Floor. E88A and E88Q comparably caused severe decreases in the activities to less than 1% of that of the wild type. We could not measure the activity of R71A due to protein

instability. Instead, a double mutant, R71A/E88A, was examined and showed that the activity declined more than that of E88A. Although R71 influenced the activity, it was unlikely to contact 1-PSA directly, since it is located slightly away from the pocket surface (Fig. 2C, 3A). It should be noted that E88D showed a low level of activity like E88A, and R71K maintained activity at nearly half that of the wild type (Fig. 2D). These findings indicate that the catalysis is ruled strictly by the geometry of the two residues. The charge groups of these residues were associated tightly with two hydrogen bonds (Fig. 2C). Closely aligned acid and base residues can form a dyad motif in which a proton is localized between the residues, and induce a decrease in pK_a of the acid residue (29).

The hydroxyl group of Y114 is an additionally important factor, as suggested by the activities of Y114A and Y114F being at the same level as those of the alanine mutants of Wall residues. In turn, the activities of Y102F and F104Y were restored compared with those of the alanine mutants (Fig. 2D). In the condensed hydrophobic side chains, Y114, Y102, R71, and E88 were arrayed to form a polar zone, in which Y114 is positioned near the entrance.

Therefore, its hydroxyl group is the first contacting point for solute molecules (Fig. 3A). From these findings, we concluded that Y114 and E88, supported by R71, are the primal key residues for the catalysis of AcLFS, and the surrounding aromatic residues work in auxiliary regulation of this activity.

Alcohols as inhibitors

In this study, three AcLFS structures were determined together with exogenous molecules (Table S1). The first one is glycerol, which originates from cryoprotectant solution (Fig. 3A, top). A glycerol molecule is stabilized near the bottom of the pocket, and two of the three hydroxyl groups are fixed by hydrogen bonds to E88, Y102, and Y114. The carbon moiety orients toward W133 and W155; thus, the conformation is also maintained by hydrophobic interactions. When trehalose was used as a cryoprotectant instead of glycerol, three water molecules were observed at the corresponding site of the glycerol (Fig. 3A, low). Subsequently, 1,2-propanediol was added to the cryoprotectant solution to obtain its complex structure (Fig. 3A, middle). A 1,2-propanediol molecule was observed in the pocket and adopts a conformation like that for

glycerol, except for the lack of a hydrogen bond to Y114. Trehalose molecules were seen only at minor peripheral sites; thus, the pocket is unlikely to accommodate the disaccharide molecules (Fig. 3B).

Next, the inhibition of AcLFS by these compounds was examined. 1,2-Propanediol and crotyl alcohol exhibited concentration-dependent decreases of PTSO production (Fig. 3C). They are deduced to behave as competitive inhibitors, considering that 1,2-propanediol was stabilized at the active site. We further investigated the inhibition patterns of cognate compounds, and demonstrated that the efficiencies are determined by the number of hydroxyl groups (Fig. S2); while linear alcohols ($C_nOH_{2n+2/2n}$; $n \leq 6$) inhibited the reaction with half-inhibition concentration (IC_{50}) values from 20 to 80 mM, diol compounds [ethylene glycol (**19**), 1,2-propanediol (**20**), and 2-methyl-2,4-pentanediol (**21**)] did so at least at 600 mM. Glycerol (**22**) did not affect the activity, even at excess concentrations (Fig. 3C). These findings indicate that the active site of AcLFS prefers alcohols rather than glycerol and diols, and suggest that alcohols stay in the pocket with disordered conformations due to perturbation of their aliphatic portions. Linear alcohols with larger

182 aliphatic chains were prone to having lower IC_{50} values, except for 1-pentanol
 183 (**12**) and *cis*-pentene-1-ol, (**17**) (Fig. S2A). Concerning an exchange of solute
 184 molecules with 1-PSA at the active site, the flexible linear alcohols would gain
 185 less entropy than diols and glycerol, which would gain entropy due to the
 186 release from the two or more hydrogen bonds (Fig. 3A). Assuming that the
 187 differential entropies are the major factor behind the net free-energy changes
 188 upon replacements, the diols and glycerol are feasibly stable, but poor inhibitors
 189 of AcLFS.

190

Discussion

Comparison with structural homologs

The seven-stranded helix-grip fold is an architectural feature shared by members of the recently established superfamily, START/RHO alpha C/PITP/Bet v 1/CoxG/CalC (SRPBCC; cl14643) (30). LFS presents no significant sequence homologies with other proteins, but a structure similarity search highlighted 3 out of 23 SRPBCC subsets (Table S2, Fig. S3A). Pyrabactin-resistant protein-like proteins (PYLs) [maximal score for *Arabidopsis thaliana* PYL10 (31)] are intracellular receptors for a plant hormone, abscisic acid (ABA). Ginseng major latex-like protein 151 (MLP151), which is thought to be associated with lysophosphatidic acid regulation, is a member of the PR-10/Bet v 1-type plant allergens (32, 33). ZhuI of *Streptomyces* sp. R1128 (34) belongs to the aromatase/cyclase components of bacterial type II polyketide synthases (PKCs) and catalyzes the C7–C12-specific first ring-closing reaction toward octaketide–acyl carrier protein conjugate (35).

Each SRPBCC protein has a central cavity that provides a platform for

hydrophobic molecules, as observed in complex structures of PYLs and PKCs (36, 37). Although no potent ligand of MLP151 has been identified, extra non-protein electron densities were found inside the pocket in the MLP151-cognate allergen proteins (38, 39). Compared with PYL10 and ZhuI, *AcLFS* has a small cavity (Fig. 4D). These differences are apparently brought about by displacements of L1, S3, and S4 (Fig. S3B). In PYL proteins, L1 corresponds to the Pro-cap region, which arises from a conformational change upon the binding of ABA and constructs part of the binding site for downstream regulator proteins (36, 40). On the other hand, the C α traces of *AcLFS* were identical among the present solute molecule complexes; thus, no conformational changes were indicated during catalysis (Fig. 3B). The displacement seems not only to ensure the narrow entrance of *AcLFS*, in which bulky hydrophobic residues gather on the Gate layer (Fig. S3C), but also to restrict the dimensions of their original ligands and substrates (Fig. S3A).

Among the four proteins, conserved residues are mapped mainly on their structural frameworks rather than the active sites (Fig. S4). PYL10 harbors the ABA-binding site extending to peripheral L1 and L2 (Fig. 4A). ABA is

224 recognized in the stretched conformation and a Floor residue K56 acts in
 225 orienting the carboxylate end (31). Three key residues of *AcLFS* are fully
 226 conserved among *PYL* proteins, but R75, E90, and the hydroxyl group of Y116
 227 do not directly contribute to the binding of ABA. Studies on mutagenesis and
 228 molecular docking have shown that the active site of *ZhuI* exists at a deeper
 229 position, around which a CAPS molecule is stabilized (34) (Fig. 4D). The
 230 substrate is thought to be placed in a compact conformation, which is ready for
 231 the nucleophilic attack from D146. R66 and H109 (Fig. 4B) are assumed to play
 232 an essential role in binding of the polyketide substrate (Fig. S3A). On the
 233 structures of *WhiE* aro/cyc and *TcmN* aro/cyc (41, 42), which are PKC isoforms
 234 possessing distinct regiospecificities toward dodeca- and decaketides, two of the
 235 deduced key residues needed for the corresponding catalytic reactions do not lie
 236 at the same positions: Y35 and E34 in *WhiE* aro/cyc instead of D146 and H109
 237 in *ZhuI* (Fig. 4B). R66 is located on S3 and conserved among these PKCs (41).
 238 On S3 and S4, *ZhuI* has two other arginine residues, which are structurally not
 239 aligned on the R-E dyad motif of *AcLFS* and *PYL* proteins (Fig. 4C), and the
 240 side chains of R50 and R65 are pointed away from the molecule by one-frame

shifts. These findings indicate that the R-E dyad is a functional motif unique to LFS and not shared with the structural homolog proteins.

Although the key residues of SRPBCC proteins are totally divergent, the electrostatic landscapes in the pockets are comparable to the properties of respective acceptor molecules. In *PYL10* and *ZhuI*, positive patches are distributed on nearly neutral intramolecular surfaces (Fig. 4D). They fit well with the carboxylate of ABA and negatively polarized acyl oxygen atoms of polyketide substrates. On the other hand, *AcLFS* possesses an interior with comprehensively high negative potential, which would be effective at prohibiting the entry of anionic substances. In fact, a sulfate ion is visible only on the outer surface of *AcLFS* (Fig. 3B). Sulfenic acid is regarded as being in protonation equilibrium ($\text{R-SOH} \rightleftharpoons \text{R-SO}^- + \text{H}^+$) under physiological and aqueous conditions (13, 43). Therefore, the electronic filter is likely to have an advantage in isolating single 1-PSA in its protonated form, not 1-propene-1-sulfenate.

257 *Proposed catalytic mechanism*

258 Taking these findings together, a catalytic cycle of AcLFS is proposed, as
 259 shown in Fig. 5B. The Wall apparatus first confines 1-PSA in a similar site of
 260 glycerol and 1,2-propanediol. The hydroxyl group of Y114 forms a hydrogen
 261 bond to the sulfenyl oxygen, and the adjacent hydrogen atom of the sulfenic acid
 262 orients toward the R71-E88 dyad, in which E88 constitutively rests in the
 263 dissociated form. The sulfenic acid is polarized and then deprotonated safely in
 264 the highly negative electrostatic potential inside the molecule, which is
 265 convenient for retention of a cation. The liberated proton forms a bond directly
 266 through the π electron pair of the 1-propenyl group, considering that there is no
 267 possible proton acceptor in the pocket. Indeed, the decrease in activity of E88D
 268 to the same level as its nonpolar variants indicates that AcLFS would not
 269 exchange any electron pairs with the substrate (Fig. 2D). Y114 acts as not only a
 270 trap of 1-PSA but also an anchor of sulfenate anion, which is prone to leaving
 271 the site due to charge repulsion. Upon release of the product, Y114 no longer
 272 keeps PTSO in the pocket because of displacement around the S-linked oxygen,
 273 which arises from a valence change of the sulfur atom.

The reaction scheme based on the intramolecular proton shuttle is consistent with a previous deuterium tracer study, which showed that 1-PSA is the proton source of PTSO (44). This scheme is apparently true in [1,4]-sigmatropic rearrangement, but the retro ene-reaction-like process occurs in a concerted fashion without yielding any intermediates (12) (Fig. 5A). *AcLFS* provides a tailored compartment in order to increase the chances of correctly aligned overlap of binding orbitals, which is needed for the concerted reaction. Further, the compartment can precisely avoid self-condensation of 1-PSA monomers, involving a similar function to protection groups equipped in stable sulfenic acid compounds (45, 46). The roles in the catalysis are distinct from those of regular enzymes since the active site of *AcLFS* is prepared for regulating the inherently labile nature of 1-PSA, rather than activating the substrates into respective transition states.

The present study provides a rationale for an RSS catalysis by revealing the three-dimensional structures of *AcLFS*, which has a simple protein architecture. This work might not only decipher the principle of *Allium*-specific lachrymator generation, but also provide inspirations for multiple lines of work, particularly

291 the synthesis of transient *S*-oxide compounds, which are not abundant and thus

292 are rarely exploited.

293

Materials and Methods

Construction of plasmids

The sequences were amplified from a template plasmid harboring the complete open reading frame of AcLFS (GenBank accession: AB089203). Fragments were digested with NdeI/XhoI and HindIII/BamHI, and then inserted into the corresponding sites of pET26b (Novagen) and pPAL7 (Bio-Rad) vectors to prepare the expression plasmids pET26b-AcLFS and pPAL7-AcLFS, respectively. Expression plasmids for AcLFS mutants were prepared following the instructions of the PrimeStar mutagenesis kit (TaKaRa) using pET26b-AcLFS as the template.

Expression and purification of AcLFS

AcLFS and its variants were expressed in *Escherichia coli* BL21AI cells (Life Technologies). Cells were cultured at 37°C in LB medium containing 50 mg/L kanamycin. After OD₆₀₀ reached 0.6, AcLFS was induced for 2 hours by adding a final concentration of 1 mM IPTG and 0.2% L-arabinose. Harvested

310 cell pellets were suspended in solution A [50 mM Hepes-Na (pH7.5), 0.1 μ M
 311 PMSF] and disrupted by sonication (Sonifier 250D, Branson); then, insoluble
 312 particles were removed by centrifugation (20,000 $\times$ g, 30 min). The supernatants
 313 were loaded onto HisTrap FF crude 5 mL (GE Healthcare) equilibrated with
 314 solution B [50 mM Hepes-Na (pH 7.5), 500 mM NaCl] and the eluates for
 315 solution C (solution B containing 500 mM imidazole) were pooled. The samples
 316 were loaded onto Superdex 200 10/300 GL (GE Healthcare) equilibrated with
 317 solution D [20 mM Hepes-Na (pH 7.5), 150 mM NaCl]. Peak fractions
 318 monitored by absorbance at 280 nm were collected and concentrated to around
 319 20 mg/mL with the buffer exchanged by 20 mM Hepes-Na (pH 7.5) using
 320 Vivaspin turbo 10K MWCO (Sartorius).

321 For the production of selenomethionine (SeMet)-derivatized AcLFS,
 322 pPAL7-AcLFS was introduced into *E. coli* BL21 CodonPlus(DE3) RP-X cells.
 323 These cells were cultured in LeMaster medium containing 100 mg/L ampicillin
 324 and 35 mg/L chloramphenicol, and IPTG was added to the medium at a final
 325 concentration of 1 mM for induction. Extraction and purification were
 326 performed as per the protocols of non-derivatized AcLFS, except for using a

Profinity eXact column (Bio-Rad) instead of a HisTrap FF crude column, in accordance with the manufacturer's instructions. In each step, purity was checked by SDS-PAGE and protein concentration was estimated using BCA protein assay kit (Thermo Fisher Scientific).

Crystallography and structure determination

Crystallization was performed by the sitting-drop vapor diffusion method at 20°C. Crystals appeared within a week from the several batches of the conditions of the JCSG core I–IV and plus kits (Qiagen); then, the reservoir compositions were specifically optimized to the following condition: 2.0 M ammonium sulfate, 0.1 M MES-Na (pH 6.0). Each single crystals on the drops was picked out on CryoLoop (Hampton Research) and directly snapped in liquid nitrogen after cryo-protection, which was performed by transferring the crystal gradually to the reservoir solution containing 20% (v/v) glycerol or 16% D-trehalose. Immersion of the crystals in 1,2-propanediol [final 2% (v/v)] was performed together with the D-trehalose treatment.

Diffraction experiments were carried out at synchrotron facilities of KEK

(Tsukuba, Japan), employing the mounting robot PAM (47) and client software UGUIs. The data acquisition scheme was basically set as follows: exposure of a $100 \times 100 \mu\text{m}^2$ beam for 1 sec per frame and the collection of 180 (360 for the SeMet dataset) frames with 1° oscillation steps. Diffraction images were indexed, integrated, and scaled using the HKL2000 program suite (HKL Research Inc.) or XDS (48).

Initial phases were determined by the single-wavelength anomalous dispersion (SAD) method using the SeMet dataset, practically by using the phenix.autosol module implemented in Phenix program suite (49). The resultant model was built mostly via phenix.autobuild and extended to the whole of the molecule manually, with viewing using the program Coot (50). Initial models corresponding to the other datasets were determined by molecular replacement, using the program Molrep (51) if needed. Models were refined iteratively using REFMAC5 in the CCP4 program suite (52, 53), and heteroatoms found were added according to $m|F_o| - D|F_c|$ maps at the posterior rounds. The crystals originally belong to the $P2_12_12_1$ space group, but are prone to being adapted as $P4_1$ during pretreatment processes. The twin refinement option implemented in

REFMAC5 was applied for the datasets indexed as the $P4_1$ space group since nearly perfect twinning was found by phenix.xtriage. In addition, the occupancies of all of the atoms were set to 0.8 in the REFMAC5 calculations to reduce the phase bias-conducted lack of the resulting map densities around a region of one of the eight molecules in an asymmetric unit. The final model was verified by respective composite omit maps. Crystallographic and refinement statistics are listed in Table S1. Molecular graphics were prepared using PyMOL (Schrödinger LLC). Structure similarity searches were queued on Dali (54). Superposition and alignments of models were carried out on Coot. Electrostatic potentials were calculated using APBS (55) after charge state assignments with PROPKA (56). Coordinates and structure factors were deposited in the Protein Data Bank (PDB) with the accession codes of 5GTE (solute-free), 5GTF (with glycerol), and 5GTG (with 1,2-propanediol).

Product assay

Liberated PTSO was isolated by reverse-phase chromatography at 25°C, as described previously (11). Briefly, a 3-min-reaction mixture was subjected to

378 Pegasil ODS 4.5×250 (Senshu Kagaku) equilibrated with 0.01% trifluoroacetic
379 acid in 30% methanol solution (pH~3.3) and then developed for 12 min with a
380 flow rate of 0.6 mL/min. Absorbance at 254 nm was monitored using an
381 Alliance instrument (Waters). A total batch of 25 μ L initially consisted of 5 μ L
382 of 0.01 mg/mL AcLFS, 5 μ L of respective inhibitor in 5-fold concentrate and 10
383 μ L of garlic alliinase (250 U/mL), then 5 μ L of 20 mg/mL *trans*-1-PRENCISO
384 was added to start the reaction. Cysteine thiols on AcLFS were inactivated by
385 adding iodoacetamide at a final concentration of 1 mM in 0.01 mg/mL AcLFS
386 and incubating the mixture at 4°C for several hours in the dark. Chemical
387 reagents were purchased from Wako Pure Chemicals and Tokyo Chemical
388 Industry.
389

Footnotes

¹ T. A., Y. S. and J. T. contributed equally to this work.

² To whom correspondence should be addressed. Email: arakawa@mail.ecc.u-tokyo.ac.jp.

Author contributions: S.F. conceived the research and T.A. designed experiments; J.T., Y.S., and T.A. performed research; N.M., M.K., M.A, T.K., N.T., and S.I. supplied *AcLFS* gene and materials; All the authors discussed data; T.A. wrote the paper with assistance from all other authors.

The authors declare no conflict of interest.

403 **Acknowledgments**

404 The authors would thank to the beamline scientists of KEK and SPring-8. Also
 405 thank to Drs. Yoshitaka Moriwaki and Kentaro Shimizu for discussion and
 406 preliminary simulations. This work was supported by JSPS KAKENHI Grant
 407 Numbers JP26660289 (to T.A.), JP15H02443, and JP26660083 (to S.F.).

408

References

1. Kris-Etherton PM, *et al.* (2002) Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. *Am J Med* 113 Suppl 9B:71S-88S.
2. Lanzotti V (2006) The analysis of onion and garlic. *J Chromatogr A* 1112(1-2):3-22.
3. Breme K, Fernandez X, Meierhenrich UJ, Brevard H, & Joulain D (2007) Identification of new, odor-active thiocarbamates in cress extracts and structure-activity studies on synthesized homologues. *J Agric Food Chem* 55(5):1932-1938.
4. Naithani R, *et al.* (2008) Antiviral activity of phytochemicals: a comprehensive review. *Mini Rev Med Chem* 8(11):1106-1133.
5. Vasanthi HR, Mukherjee S, & Das DK (2009) Potential health benefits of broccoli- a chemico-biological overview. *Mini Rev Med Chem* 9(6):749-759.
6. Jones MG, *et al.* (2004) Biosynthesis of the flavour precursors of onion and garlic. *J Exp Bot* 55(404):1903-1918.

- 425 7. Block E (1992) The Organosulfur Chemistry of the Genus *Allium* --
426 Implications for the Organic Chemistry of Sulfur. *Angew Chem Int Ed Engl*
427 31(9):1135-1178.
- 428 8. Aoyagi M, *et al.* (2011) Structure and bioactivity of thiosulfinates resulting from
429 suppression of lachrymatory factor synthase in onion. *J Agric Food Chem*
430 59(20):10893-10900.
- 431 9. Brodnitz MH & Pascale JV (1971) Thiopropanal S-oxide: a lachrymatory
432 factor in onions. *J Agric Food Chem* 19(2):269-272.
- 433 10. Imai S, *et al.* (2002) Plant biochemistry: an onion enzyme that makes the eyes
434 water. *Nature* 419(6908):685.
- 435 11. Masamura N, *et al.* (2012) Identification of amino acid residues essential for
436 onion lachrymatory factor synthase activity. *Biosci Biotechnol Biochem*
437 76(3):447-453.
- 438 12. Block E, *et al.* (1996) *Allium* Chemistry: Microwave Spectroscopic
439 Identification, Mechanism of Formation, Synthesis, and Reactions of
440 (E,Z)-Propanethial S-Oxide, the Lachrymatory Factor of the Onion (*Allium*
441 *cepa*). *J Am Chem Soc* 118(32):7492-7501.

- 442 13. Gupta V & Carroll KS (2014) Sulfenic acid chemistry, detection and cellular
443 lifetime. *Biochim Biophys Acta* 1840(2):847-875.
- 444 14. Nagy P & Ashby MT (2007) Reactive sulfur species: kinetics and mechanisms
445 of the oxidation of cysteine by hypohalous acid to give cysteine sulfenic acid. *J*
446 *Am Chem Soc* 129(45):14082-14091.
- 447 15. Yang J, Gupta V, Carroll KS, & Liebler DC (2014) Site-specific mapping and
448 quantification of protein S-sulphenylation in cells. *Nat Commun* 5:4776.
- 449 16. Yeh JI, Claiborne A, & Hol WG (1996) Structure of the native cysteine-sulfenic
450 acid redox center of enterococcal NADH peroxidase refined at 2.8 Å resolution.
451 *Biochemistry* 35(31):9951-9957.
- 452 17. Arakawa T, *et al.* (2007) Structure of thiocyanate hydrolase: a new nitrile
453 hydratase family protein with a novel five-coordinate cobalt(III) center. *J Mol*
454 *Biol* 366(5):1497-1509.
- 455 18. Devarie-Baez NO, Silva Lopez EI, & Furdul CM (2016) Biological chemistry
456 and functionality of protein sulfenic acids and related thiol modifications. *Free*
457 *Radic Res* 50(2):172-194.
- 458 19. Parsonage D, *et al.* (2015) Dissecting peroxiredoxin catalysis: separating

- 459 binding, peroxidation, and resolution for a bacterial AhpC. *Biochemistry*
460 54(7):1567-1575.
- 461 20. Salmeen A, *et al.* (2003) Redox regulation of protein tyrosine phosphatase 1B
462 involves a sulphenyl-amide intermediate. *Nature* 423(6941):769-773.
- 463 21. Wood ZA, Poole LB, & Karplus PA (2003) Peroxiredoxin evolution and the
464 regulation of hydrogen peroxide signaling. *Science* 300(5619):650-653.
- 465 22. Nakamura T, *et al.* (2008) Oxidation of archaeal peroxiredoxin involves a
466 hypervalent sulfur intermediate. *Proc Natl Acad Sci U S A* 105(17):6238-6242.
- 467 23. Arakawa T, *et al.* (2009) Structural basis for catalytic activation of thiocyanate
468 hydrolase involving metal-ligated cysteine modification. *J Am Chem Soc*
469 131(41):14838-14843.
- 470 24. Yamanaka Y, *et al.* (2015) Time-Resolved Crystallography of the Reaction
471 Intermediate of Nitrile Hydratase: Revealing a Role for the Cysteinesulfenic
472 Acid Ligand as a Catalytic Nucleophile. *Angew Chem Int Ed Engl*
473 54(37):10763-10767.
- 474 25. Block E, Dane AJ, Thomas S, & Cody RB (2010) Applications of direct analysis
475 in real time mass spectrometry (DART-MS) in Allium chemistry.

- 476 2-propenesulfenic and 2-propenesulfinic acids, diallyl trisulfane S-oxide, and
477 other reactive sulfur compounds from crushed garlic and other Alliums. *J Agric*
478 *Food Chem* 58(8):4617-4625.
- 479 26. Penn RE, Block E, & Revelle LK (1978) Flash vacuum pyrolysis studies. 5.
480 Methanesulfenic acid. *J Am Chem Soc* 100(11):3622-3623.
- 481 27. Eady CC, *et al.* (2008) Silencing onion lachrymatory factor synthase causes a
482 significant change in the sulfur secondary metabolite profile. *Plant Physiol*
483 147(4):2096-2106.
- 484 28. Iyer LM, Koonin EV, & Aravind L (2001) Adaptations of the helix-grip fold for
485 ligand binding and catalysis in the START domain superfamily. *Proteins*
486 43(2):134-144.
- 487 29. Gutteridge A & Thornton JM (2005) Understanding nature's catalytic toolkit.
488 *Trends Biochem Sci* 30(11):622-629.
- 489 30. Marchler-Bauer A, *et al.* (2015) CDD: NCBI's conserved domain database.
490 *Nucleic Acids Res* 43(Database issue):D222-226.
- 491 31. Hao Q, *et al.* (2011) The molecular basis of ABA-independent inhibition of
492 PP2Cs by a subclass of PYL proteins. *Mol Cell* 42(5):662-672.

- 493 32. Choi SH, *et al.* (2015) Structure of ginseng major latex-like protein 151 and its
494 proposed lysophosphatidic acid-binding mechanism. *Acta Crystallogr D Biol*
495 *Crystallogr* 71(Pt 5):1039-1050.
- 496 33. Sinha M, *et al.* (2014) Current overview of allergens of plant pathogenesis
497 related protein families. *ScientificWorldJournal* 2014:543195.
- 498 34. Ames BD, *et al.* (2011) Structural and biochemical characterization of ZhuI
499 aromatase/cyclase from the R1128 polyketide pathway. *Biochemistry*
500 50(39):8392-8406.
- 501 35. Hertweck C, Luzhetskyy A, Rebets Y, & Bechthold A (2007) Type II
502 polyketide synthases: gaining a deeper insight into enzymatic teamwork. *Nat*
503 *Prod Rep* 24(1):162-190.
- 504 36. Miyazono K-I, *et al.* (2009) Structural basis of abscisic acid signalling. *Nature*
505 462(7273):609-614.
- 506 37. Sultana A, *et al.* (2004) Structure of the polyketide cyclase SnoaL reveals a
507 novel mechanism for enzymatic aldol condensation. *EMBO J* 23(9):1911-1921.
- 508 38. Marković-Housley Z, *et al.* (2003) Crystal structure of a hypoallergenic isoform
509 of the major birch pollen allergen Bet v 1 and its likely biological function as a

- 510 plant steroid carrier. *J Mol Biol* 325(1):123-133.
- 511 39. Chruszcz M, *et al.* (2013) Structural and bioinformatic analysis of the kiwifruit
512 allergen Act d 11, a member of the family of ripening-related proteins. *Mol*
513 *Immunol* 56(4):794-803.
- 514 40. Nishimura N, *et al.* (2009) Structural mechanism of abscisic acid binding and
515 signaling by dimeric PYR1. *Science* 326(5958):1373-1379.
- 516 41. Lee M-Y, Ames BD, & Tsai S-C (2012) Insight into the molecular basis of
517 aromatic polyketide cyclization: crystal structure and in vitro characterization of
518 WhiE-ORFVI. *Biochemistry* 51(14):3079-3091.
- 519 42. Ames BD, *et al.* (2008) Crystal structure and functional analysis of
520 tetracenomycin ARO/CYC: implications for cyclization specificity of aromatic
521 polyketides. *Proc Natl Acad Sci U S A* 105(14):5349-5354.
- 522 43. Okuyama T, *et al.* (1992) Equilibrium and kinetic studies of reactions of
523 2-methyl-2-propanesulfenic acid. *Heteroatom Chem* 3(5-6):577-583.
- 524 44. Masamura N, Aoyagi M, Tsuge N, Kamoi T, & Imai S (2012) Proton transfer in
525 a reaction catalyzed by onion lachrymatory factor synthase. *Biosci Biotechnol*
526 *Biochem* 76(9):1799-1801.

- 527 45. Goto K, Holler M, & Okazaki R (1997) Synthesis, Structure, and Reactions of a
528 Sulfenic Acid Bearing a Novel Bowl-Type Substituent: The First Synthesis of a
529 Stable Sulfenic Acid by Direct Oxidation of a Thiol. *J Am Chem Soc*
530 119(6):1460-1461.
- 531 46. Yoshimura T, *et al.* (1992) Synthesis of a stable sulfenic acid,
532 trans-decalin-9-sulfenic acid. *J Chem Soc, Chem Commun*:1337-1338.
- 533 47. Hiraki M, *et al.* (2008) High-throughput operation of sample-exchange robots
534 with double tongs at the Photon Factory beamlines. *Journal of Synchrotron*
535 *Radiation* 15(3):300-303.
- 536 48. Kabsch W (2010) XDS. *Acta Crystallogr D Biol Crystallogr* 66(Pt 2):125-132.
- 537 49. Adams PD, *et al.* (2010) PHENIX: a comprehensive Python-based system for
538 macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* 66(Pt
539 2):213-221.
- 540 50. Emsley P, Lohkamp B, Scott WG, & Cowtan K (2010) Features and
541 development of Coot. *Acta Crystallogr D Biol Crystallogr* 66(Pt 4):486-501.
- 542 51. Vagin A & Teplyakov A (2000) An approach to multi-copy search in molecular
543 replacement. *Acta Crystallogr D Biol Crystallogr* 56(Pt 12):1622-1624.

52. Murshudov GN, *et al.* (2011) REFMAC5 for the refinement of macromolecular crystal structures. *Acta Crystallogr D Biol Crystallogr* 67(Pt 4):355-367.
53. Winn MD, *et al.* (2011) Overview of the CCP4 suite and current developments. *Acta Crystallogr D Biol Crystallogr* 67(Pt 4):235-242.
54. Holm L & Rosenström P (2010) Dali server: conservation mapping in 3D. *Nucleic Acids Res* 38(Web Server issue):W545-549.
55. Baker NA, Sept D, Joseph S, Holst MJ, & McCammon JA (2001) Electrostatics of nanosystems: application to microtubules and the ribosome. *Proc Natl Acad Sci U S A* 98(18):10037-10041.
56. Olsson MHM, Søndergaard CR, Rostkowski M, & Jensen JH (2011) PROPKA3: Consistent Treatment of Internal and Surface Residues in Empirical pKa Predictions. *J Chem Theory Comput* 7(2):525-537.

Figure legends

Fig. 1. Chemical utilization pathways of γ -glutamyl-*S*-alkenyl cysteines via *trans*-1-PRENCISO in crushed onion. Solid and dashed arrows represent enzymatic and non-enzymatic processes, respectively. Brackets indicate undetectable materials. Numbered chemical names are noted in the main text, except for pyruvic acid (4).

Fig. 2. Structure and the active site of AcLFS. (A) Overall structure displayed by schematic representation in spectral colors. (B) Positions of mutated residues colored by impacts on the LFS activity. Spheres are γ -atoms colored in cyan (none) to red (significant), based on the activities of respective alanine variants, R71K and E112Q. (C) Pocket-lining residues. Views from the entrance (top) and the direction of the parallel to longitudinal axis (bottom, stereoview). The side chains of Wall and Floor layers (sticks) are shown with the cartoon drawing of the main chain. The carbon atoms of Wall are colored as in (A) and those of Floor are in gray. (D) List of variants and relative productivities. Error bars

represent standard deviations of at least triplicate measurements.

Fig. 3. Solute molecules inside the pocket. (A) Non-protein electron densities found at the active site. $m|F_o| - D|F_c|$ omit maps (red, 3σ) are displayed. Surfaces colored in gray are provided by the hydrophobic residues. Atoms suggested to form hydrogen bonds are connected with dashes. (B) Superposition of a protomer of each model. Traces of C α atoms are taken from the data of +glycerol (white), +1,2-propanediol (pink), and solute-free (cyan). Sticks represent exogenous compounds observed on the crystals. (C) Inhibition curves of AcLFS in the presence of glycerol, 1,2-propanediol, and crotyl alcohol.

Fig. 4. Comparisons with structural homologs. Close-up views of (A) PYL10-ABA (PDB 3r6p) and (B) ZhuI-CAPS (3tfz) complexes. ABA-contacting residues of PYL10 are colored cyan. Suggested primal catalytic residues of ZhuI and WhiE aro/cyc (PDB 3tvr) are colored magenta and pink. Corresponding residues of AcLFS are overlaid as green sticks. (C) Superposition around the R71-E88 dyad motif of AcLFS. Structure-based sequence alignment

592 is depicted below. Arrowheads indicate residues whose side chains point inward.

593 (D) Cutaway views of *AcLFS*, MLP151 (PDB 4reh), *PYL10*, and *ZhuI* from left

594 to right. Calculated electrostatic potentials are indicated on the surfaces

595 following the order of blue to red.

596

597 Fig. 5. PTSO generation principles (A) under FVP condition and (B) with the

598 assistance of *AcLFS*.

Figure 2

Figure 3

Figure 4

Supplemental information

Supplemental tables and figure legends

Table S1 Data collection and refinement statistics

	SeMet (peak)	Solute-free	+ glycerol	+1,2-propanediol
PDB ID		5GTE	5GTF	5GTG
Data collection				
Beamline	PF NE3A	PF NE3A	PF BL5A	PF NW12A
Wavelength (Å)	0.9787	1.0000	1.0000	1.0000
Space group	$P4_1$	$P2_12_12_1$	$P2_12_12_1$	$P4_1$
Cell dimensions				
a, b, c (Å)	173.1, 173.1, 64.9	65.1, 118.7, 124.5	65.0, 119.9, 125.6	172.9, 172.9, 65.0
Molecules per ASU	8	4	4	8
Resolution (Å)	50.00-2.10 (2.14-2.10) ¹	50.00-2.00 (2.07-2.00)	50.00-1.80 (1.83-1.80)	50.00-1.70 (1.76-1.70)
Unique reflections	138,410 (5,543)	66,057 (6,398)	91,533 (8,871)	204,662 (16,550)
Completeness (%)	99.8 (99.3)	99.6 (98.0)	99.7 (98.6)	97.3 (79.1)
Multiplicity	6.1 (5.5)	6.7 (6.2)	5.8 (5.2)	6.1 (4.9)
Mean $I/\sigma I$	24.6 (3.7)	15.2 (2.8)	18.9 (2.2)	19.8 (4.2)
R_{merge}	0.079 (0.459)	0.100(0.676)	0.053 (0.600)	0.062 (0.378)
$CC_{1/2}$	(0.98)	(0.80)	(0.93)	(0.86)
Refinement				
Resolution (Å)		50.00-2.00	50.00-1.80	50.00-1.70
No. of reflections		62,884	86,948	194,424
$R_{\text{work}} / R_{\text{free}}$		0.191/0.233	0.170/0.199	0.230/0.260 ²
No. of atoms		5271	5626	10295
Protein		4873	4893	9646
Solute		43	85	65
Water		352	648	584
B-factors				
Protein (Å ²)		29.7	20.3	20.1
Solute (Å ²)		55.9	40.7	22.0
Water (Å ²)		37.4	34.8	25.9

RMS deviation			
Bond lengths (Å)	0.019	0.020	0.031
Bond angles (°)	1.87	1.93	2.83
Ramachandran plot			
Favored (%)	99.0	98.7	95.0
Allowed (%)	0.8	1.2	4.1
Outlier (%)	0.17	0.17	0.85

¹ Values in parentheses are for the highest resolution shell. $R_{merge} = \sum_h \sum_i |I_{h,i} - I_h| / \sum_h \sum_i I_{h,i}$, where I_h is the mean intensity of the i observations of symmetry-related reflections of h . $R = \sum |F_{obs} - F_{calc}| / \sum F_{obs}$, where F_{calc} is the calculated protein structure factor from the atomic model (R_{free} was calculated with 5% of the reflections selected randomly).

² High R factor values because of twinning (see Materials and Methods).

Table S2 Proteins of similar structure to AcLFS ¹

Name	Source	Z _{max}	Identity ²	Similarity ²	Domain family ³	PDB ID ⁴
PYL10	<i>Arabidopsis thaliana</i>	19.0	16.7% (31/186)	32.3% (60/186)	PYR_PYL_RCAR_like (cd07821)	3R6P
PYL9	<i>A. thaliana</i>	17.3	17.3% (33/191)	30.4% (58/191)	PYR_PYL_RCAR_like	3OQU
MLP151	<i>Panax ginseng</i>	17.2	18.8% (32/170)	34.1% (58/170)	Bet_v_1_like (cd07816/smart01037)	4REH
PYL2	<i>A. thaliana</i>	17.1	16.7% (32/192)	32.3% (62/192)	PYR_PYL_RCAR_like	3KB0
PYL1	<i>A. thaliana</i>	16.9	16.8% (30/179)	31.8% (57/179)	PYR_PYL_RCAR_like	3JRS
ZhuI	<i>Streptomyces</i> sp. R1128	16.6	12.1% (21/174)	24.7% (43/174)	OtcD1_ARO-CYC_like (cd08861)	3TFZ
PYL3	<i>A. thaliana</i>	16.5	17.2% (36/209)	31.6% (66/209)	PYR_PYL_RCAR_like	4DSC
WhiE aro/cyc	<i>S. coelicolor</i> A3(2)	15.7	14.1% (25/177)	27.7% (49/177)	OtcD1_ARO-CYC_like	3TVQ

¹ Sorted by a descending order of the maximum Z-scores of DaliLite v. 3.

² Taken from the output of EMBOSS Stretcher.

³ Acquired from Conserved Domain Database (CDD).

⁴ Representative entry (ligand complex if available) is shown.

Figure S1

Figure S2

Figure S3

Figure S4

Figure legends

Figure S1 (A) Distribution of cysteine residues on AcLFS structure. (B) chromatograms of the selected measurements of reaction products. Arrowheads indicate the retention times of PTSO.

Figure S2 Inhibitions by organic hydroxide compounds. (A) Inhibition curves in the presence of normal alcohols (top), unsaturated alcohols (bottom) and diols (right). (B) Array according to the estimated IC_{50} values of solute molecules tested in this study. Structural formulae stand for 1-propanol (**10**), 1-butanol (**11**), 1-pentanol (**12**), 1-hexanol (**13**), crotyl alcohol (**14**), 4-pentene-1-ol (**15**), *trans*-2-pentene-1-ol (**16**), *cis*-2-pentene-1-ol (**17**), *cis*-2-hexene-1-ol (**18**), ethylene glycol (**19**), 1,2-propanediol (**20**), 2-methyl-2,4-pentanediol (**21**) and glycerol (**22**).

Figure S3 (A) Comparison of structurally relative helix-grip fold proteins. Primary substrate or ligand of each protein is displayed at the bottom. Structural formulae stand for abscisic acid (**23**) and octaketide–acyl carrier protein conjugate (**24**). (B) Superposition around the entrance. Cartoon representations of AcLFS (green), MLP15 (orange), PYL10 (cyan) and ZhuI (magenta). Superposed view from Floor to Gate of the pocket. (C) Cluster of hydrophobic residues on Gate layer. Spheres are drawn as respective van der Waals radii of carbon (yellow) and sulfur (orange).

Figure S4 Structure-based amino acid sequence alignment of AcLFS and structural homologs. Conserved residues represent blue characters. Residues are highlighted for Wall and Floor (green), Gate (yellow), direct association with ABA (orange in PYL10) and CAPS (orange in ZhuI). The key residues of AcLFS are marked with magenta dots and secondary structure elements drawn upside are for AcLFS. Arrowheads indicate residues pointing inward.