1 **Genotype, nitrogen and herbivory shape plant defense: the case of a vitamin-**
2 **enriched maize**

3 **Agnès Ardanuy¹, Victoria Pastor², Gaëtan Glauser³, Paul Christou^{1 4}, Ted C. J. Turlings⁵,**
4 **Ramon Albajes¹**

5

6 ¹Departament de Producció i Protecció Vegetal, Agrotecnio Center, Universitat de Lleida,
7 Rovira Roure 191, E-25198 Lleida, Spain. email: agnes.ardanuy@gmail.com

8 ²Metabolic Integration and Cell Signaling Group, Plant Physiology Section, Department of
9 CAMN, Universitat Jaume I, Castelló, Spain. email: pastorm@uji.es

10 ³Neuchâtel Platform of Analytical Chemistry, University of Neuchâtel, Rue Emile-Argand 11,
11 2000 Neuchâtel, Switzerland. email: gaetan.glauser@unine.ch

12 ⁴ICREA, Catalan Institute for Research and Advanced Studies, Passeig Lluís Companys 23,
13 08010 Barcelona, Spain. email: christou@pvcf.udl.cat

14 ⁵Laboratory for Fundamental and Applied Research in Chemical Ecology (FARCE), University
15 of Neuchâtel, Rue Emile-Argand 11, 2000 Neuchâtel, Switzerland. email:
16 ted.turlings@unine.ch

17

18 **HIGHLIGHT**

19 We show the separate and interactive effects of nitrogen availability and genotype on the
20 performance and behavior of a herbivore, and related these changes to constitutive and inducible
21 maize defenses.

22

23 **ABSTRACT**

24 The cultivation of crops with novel traits could interfere with ecosystem services delivered by
25 arthropods through bottom-up effects. Here we tested the hypothesis that a vitamin enriched
26 maize (Carolight^R) is similar in terms of plant-arthropod interactions to its wild type when
27 compared in controlled environment and under field conditions. In order to assess the
28 robustness of their equivalence we tested two nitrogen availability regimes. We used arthropod
29 field abundance, the behavior and fitness of a keystone maize herbivore - the leafhopper
30 *Zyginidia scutellaris* - and above ground chemistry of maize plants (volatile, hormone and
31 metabolite profiling) as indicators of potential changes in plant-insect interactions. Nitrogen
32 availability was the key driver of herbivore abundance and behavior, and determined direct and
33 indirect chemical defense in maize plants. Both genotypes presented similar constitutive and
34 inducible phytohormone profiles independently of the nitrogen regime. However, feeding by the
35 herbivore suppressed the levels of JA-Ile and JA, without impairing the release of induced plant
36 volatiles. Carolight^R and M37W differed to some degree in the concentrations of phenolics
37 (hydroxycinnamic acids and lignans) and in the abundance of a volatile compound. Overall the
38 effect of maize genotype on the herbivores was smaller than the effect of nitrogen fertilization.

39 **Key words:** hormone suppression, maize, metabolomics, modified metabolism, nitrogen
40 fertilization, plant defense, plant-insect interactions, *Zyginidia scutellaris*.

41

42

43 INTRODUCTION

44 One of the issues regarding the cultivation of novel crops (genetically modified or
45 otherwise) is their possible effect on insect biodiversity and associated ecosystem
46 services in agriculture. The mandatory environmental risk assessment for cultivation of
47 novel crops addresses the hypothesis that the traits introduced into the novel crops do
48 not adversely affect the non-target arthropods. Risk assessments are comparative in the
49 sense that novel crops are screened for phenotypic and compositional equivalence to
50 their wild type counterpart, and the biologically meaningful differences observed
51 between them are a consequence of the novel trait (Wolt *et al.*, 2010) and are
52 subsequently evaluated. In addition to risk assessment purposes, studying the herbivore
53 community responses to novel varieties, and in turn the biochemical responses of those
54 novel plant varieties to insect herbivory and nutrient availability may help improve our
55 understanding of plant chemical profiles and their role in plant–herbivore interactions.

56 An elite South African maize inbred was engineered to deliver pro-vitamin A (and other
57 nutritionally important carotenoids such as lutein, zeaxanthin and lycopene) in the diet
58 and thus address vitamin A and other nutritional deficiencies in at-risk populations in
59 developing countries. The kernels of this novel maize (Carolight^R) accumulate higher
60 levels of 3 vitamins in the endosperm through the simultaneous engineering of 3
61 separate metabolic pathways: 169-fold the normal amount of beta-carotene (provitamin
62 A), 6-fold the normal amount of ascorbate (vitamin C), and double the normal amount
63 of folate (vitamin B9) (Naqvi *et al.*, 2009). Molecular and biochemical characterization
64 of Carolight^R seeds (transcript, proteome, and metabolite profiles) indicated changes in
65 sugar and lipid metabolism in the endosperm with respect to the wild type due to the
66 higher up-stream metabolite demand by the extended biosynthesis capacities for
67 terpenoids and fatty acids (Decourcelle *et al.*, 2015). Nevertheless under field conditions
68 the metabolic phenotype of vitamin-enriched maize kernels under contrasting soil
69 nitrogen conditions was indistinguishable from the wild type in terms of carotenoid
70 accumulation in leaves, photosynthetic activity, sensitivity to source limitation and yield
71 (Zanga *et al.*, 2016). Authors concluded that the additional metabolic requirements of
72 Carolight^R endosperm did not affect agronomic performance. Interestingly gravid
73 females of the key Mediterranean maize pest *Sesamia nonagrioides* preferred the
74 volatiles of the wild type to Carolight^R in an olfactometer setting (Cruz and Eizaguirre,

2015), which led to the notion that vitamin enriched maize might modify the outcome of plant-insect interactions.

The strong influence of plant chemical traits on food webs has been demonstrated experimentally both above and below ground (e.g. van der Putten et al. 2001, Ode 2006). As it is not possible to measure all ecological interactions between a plant and its associated insect species, we used the arthropod field abundance, the behaviour and fitness of an herbivore keystone species (Albajes *et al.*, 2011) and above ground chemistry of maize plants as indicators of possible modifications in plant-insect interactions. We therefore tested the hypothesis that Carolight^R is similar in terms of plant-arthropod interactions to its wild type line (M37W) when compared in a controlled environment and in the field. The over-arching objectives of the current study were to: (i) determine if Carolight^R and M37W influence abundance and dynamics of herbivores and natural enemies in the field; (ii) determine potential impact of both genotypes on herbivore choice and performance under controlled conditions; (iii) characterize the chemical profiles of leaves usually consumed by most herbivores (and thus involved in plant-insect interactions) in both genotypes. Characterization was carried out through volatile, hormone and metabolite profiling. In order to broaden the range of environments in the study and to test the consistency of performance between Carolight^R and M37W, we compared both genotypes under different substrate nitrogen availability regimes. The data and conclusions from our studies not only validate the use of plant-insect interactions in the environmental risk assessment of crops with novel traits, but importantly also shed light into the biochemical and metabolic components that underpin the mechanisms involved in maize-insect interactions.

MATERIAL AND METHODS

Plants and nitrogen treatments

Seeds of the elite South African maize (*Zea mays* L.) inbred cv. M37W and its vitamin enriched derived line Carolight^R were obtained from the Applied Plant Biotechnology Group at Universitat de Lleida- Agrotecnio Center.

A field experiment was carried out in order to evaluate the performance of Carolight^R and M37W in terms of arthropod community composition and dynamics. The

experimental design encompassed a factorial combination of the two maize genotypes and two nitrogen treatments. Plots were randomized with four replicates per genotype-nitrogen combination, each consisting of 6 rows, 70 cm apart and 6.47 m in length (approximately 4 plants per meter). Maize was planted on 5 May 2013. Two different fertilization regimes were applied on 9 July 2013: Control = 0 kg ha⁻¹ and +N = 200 kg ha⁻¹ as urea at the V6 stage (six fully expanded leaves). Each plot was fully irrigated.

For laboratory experiments, seeds from each line were sown in plastic pots (10 cm high, 5 cm diameter) in vermiculite, and germinated in the greenhouse. Forty maize plants (seven to ten days old) were placed in plastic containers and provided 2.5 l of hydroponic solution for 10-12 days. Two hydroponic solutions were tested: a control solution and a solution with an increased content in nitrogen (+N). The control solution consisted of a half-strength modified Hoagland solution with micro-nutrients provided at full strength. The solution with nitrogen (+N) consisted of a control solution in which 8 mM of NH₄NO₃ was added. The hydroponic solutions were adjusted to pH 5.9, and were buffered with MES tampon. The solution was replaced every 3-4 days.

Insects and herbivory treatments

A colony of the leafhopper *Z. scutellaris* was established from small grain cereal and maize fields at the Universitat de Lleida (Spain). The colony was reared under controlled conditions (16:8 h L:D, 24±5 °C) on maize plants (var. Delprim).

Plants were transferred to an experimental chamber equipped with full spectrum light benches (24±2 °C, 40±10% r.h., 16:8 h L/ D, and 8000 lm m⁻²) the day prior the experiments started. Plants used for volatile collection were enclosed in custom made Nalophan bags (Omya AG, Oftringen, Switzerland, 150 mm diameter) closed with a parafilm seal at the top of the plastic pot. Plants used for hormone and non-targeted metabolome profiling were enclosed in bottom cut PET plastic bottles covered with muslin cloth. Herbivore treatment was initiated on the following day by exposing plants to ten *Z. scutellaris* adults for 24 h in the case of volatile analysis and non-targeted metabolome profiling, and 24, 48 and 96h for hormone profiling. The timing was chosen based on a previous study which indicated a strong induction of plant volatiles at 24h after the start of leafhopper feeding (Ardanuy *et al.*, 2016).

135 *Field herbivore and natural enemy abundance*

136 Visual sampling of arthropod fauna was conducted on whole plants from the 9th of July
137 (V6-7 stage) to the 16th of September, 2013 every other week (5 samplings in total)
138 according to Albajes et al. (2011). We sampled four plants from each plot randomly,
139 and we recorded the number of herbivores and their natural enemies per plant.
140 Herbivore counts were grouped in five taxonomic units: Thysanoptera (thrips),
141 Hemiptera\Aphididae (aphids), Hemiptera\Cicadellidae (leafhoppers, mainly *Zyginidia*
142 *scutellaris*), and Hemiptera\ Delphacidae (planthoppers, *Laodelphax striatellus*) and
143 Lepidoptera (*Spodoptera spp.*, *Helicoverpa armigera*, corn borers). Later we
144 transformed aphid counts into an abundance scale (0, no aphids; 1, isolated aphids; 2,
145 small colony; 3, medium colony; 4, large colony). Natural enemy counts were grouped
146 in Hemiptera\Anthocoridae, Hemiptera\Miridae, Neuroptera, Coccinellidae,
147 Thysanoptera (thrips) and Arachnida.

148 We calculated the sum of abundances per plot and sampling date for all taxonomic
149 units. We tested the effects of genotype, nitrogen, and sampling date on herbivore and
150 natural enemy community with a permutational MANOVA using the Adonis function in
151 the package vegan in R (Oksanen *et al.*, 2013). We then performed univariate analysis
152 at the species level for herbivore abundance data with a generalized linear model
153 following a Negative Binomial distribution in which sampling date, nitrogen treatment
154 and genotype and their interactions were used as fixed factors. Aphid abundance was
155 analyzed with an ordinal logistic regression. All statistical analyses were performed
156 using R (R Development Core Team) unless otherwise indicated.

157 *Herbivore performance and plant choice*

158 Leafhopper performance was tested by transferring 1-day old leafhopper nymphs from
159 the colony to maize plants and letting them develop until adult stage. Plant treatments
160 consisted of a factorial combination of the two maize genotypes and two N treatments
161 (control and +N) (n=13-15 plants per treatment). Plants were enclosed in plastic bottles
162 with their bottom open, covered by cloth to prevent leafhoppers from escaping; each
163 plant contained 3 leafhoppers. Plants were monitored daily until leafhoppers reached
164 adult stage. Leafhoppers were then removed and placed in 0.5 mm eppendorfs and
165 frozen at -20°C until sexed and weighed. When there was more than one leafhopper per
166 sex in a plant we averaged final weight and developmental time. Final weight of

167 leafhopper individuals and developmental time was analyzed with a GLM following a
168 Gaussian distribution using the variables insect sex, nitrogen regime and genotype and
169 their interactions as factors.

170 The effects of plant volatiles emitted by the different combination of varieties and
171 nitrogen treatments on the behavior of the leafhopper were investigated in a six-arm-
172 olfactometer (for details see Turlings, Davison, & Tamò, 2004). A plant from each
173 genotype-nitrogen treatment was placed in glass vessels one hour before the assay
174 began. Two empty vessels were used as blanks. Purified and humidified air entered each
175 odor source bottle at 0.8 l/min via Teflon tubing (adjusted by a manifold with four flow-
176 meters; Analytical Research System, Gainesville, FL, USA) and carried the volatiles
177 through to the olfactometer compartment. The position of the odor sources in the
178 olfactometer was randomly assigned each experimental day to avoid position-bias.

179 At least half an hour before the experiment started groups of six *Z. scutellaris* females
180 were isolated in pipette tips by means of a manual aspirator, and covered in parafilm.
181 Twelve leafhoppers were freed at the base of the olfactometer and left for 45 minutes.
182 Only when an insect entered an arm and passed the screw cap fitting or was recovered
183 in the bulb we considered it had made a choice. Three times twelve females were tested
184 per experiment per day. All olfactometer tests were conducted between 10 am and 4 pm
185 under light benches (24 ± 2 °C). Each experiment was performed 7 times on different
186 days. This resulted in 7 independent replicates for each olfactometer setup.

187 Olfactometer choice counts were analyzed with a GLM following a Poisson
188 distribution, with nitrogen regime and genotype and their interactions as factors. Pair-
189 wise comparisons were performed with using Tukey's HSD.

190 *Analysis of volatile profiles*

191 VOCs were collected simultaneously from herbivore-damaged plants and from control
192 non-damaged plants for all the treatments consisting of the factorial combination of
193 genotype and nitrogen treatments. Two tubular glass outlets (23x17x12 mm) with a
194 screw cap were attached to the bottom and top of the bag respectively (as described by
195 Turlings et al. 1998). Clean air was supplied to the system through the top outlet via
196 Tygon tubing connected to a flowmeter (Analytical Research Systems) and through the
197 bottom device air was pulled through a volatile adsorbent trap at a rate of 1 l/min using

198 a vacuum pump. We collected volatiles of each odor source for 5h using adsorbent traps
199 consisting of a glass tube (4 mm ID) packed with 25 mg Super-Q polymer (80–100
200 mesh) (Alltech Associates, Deerfield, Illinois, USA). We performed seven experimental
201 replicates for all treatments on different days.

202 The traps were then extracted with 150 µl dichloromethane (Suprasolv, Merck,
203 Dietikon, Switzerland), and 200 ng of n-octane and n-nonyl acetate (Sigma, Buchs,
204 Switzerland) in 10 µl dichloromethane were added to the samples as internal standards.
205 Samples were analyzed with a GC-MS as described in Ardanuy et al., (2016). The
206 detected volatiles were identified by comparison of their mass spectra with those of the
207 NIST 05 library and by comparison of retention times with those from a library from
208 earlier assays.

209 Permutational MANOVA was used to evaluate whether the VOC blend varied between
210 herbivore treatments, nitrogen availability regimes and among genotypes. The
211 abundance of the components of the volatile blend was used as the response variable,
212 while herbivore treatment, nitrogen regime, plant genotype, and their double
213 interactions were used as independent variables. In addition the amount of each
214 individual compound was compared among treatments using a non-parametric Kruskal-
215 Wallis test followed by Dunn's test.

216

217 *Plant hormones and hydroxycinnamic acid analysis*

218 A targeted analysis of plant hormones and phenolic compounds of herbivore-damaged
219 plants (n=3) and control plants (n=3) was performed for each combination of genotype-
220 nitrogen levels at three time points (24, 48 and 96h) after the experiment started. The
221 aboveground part of the plants was flash frozen with liquid nitrogen and stored at -80°C
222 until freeze dried. The experiment was repeated three times. The hormones jasmonoyl-
223 L-isoleucine (JA-Ile), 12-oxo-phytodienoic acid (OPDA), jasmonic acid (JA), salicylic
224 acid (SA), abscisic acid (ABA) and indole-3-acetic acid (IAA), and the
225 hydroxycinnamic acids caffeic acid, chlorogenic acid and ferulic acid were analyzed by
226 ultra-performance liquid chromatography coupled to mass spectrometry (UPLC-MS), as
227 described by Camañes et al. (2012). Data from the three experiments were log-
228 transformed and analyzed by a linear model with nitrogen regime and genotype and
229 their interactions as factors, and experiment as a block. Within an experiment pair-wise
230 comparisons were performed using the Klustal-Wallis test.

231 *Metabolite fingerprinting*

232 Non-targeted metabolite profiling of herbivore damaged (n=5) and control plants (n=5)
 233 was performed for each combination genotype-nitrogen levels. The aboveground part of
 234 the plants was flash frozen with liquid nitrogen 24h after the experiment started and
 235 stored at -80°C. Each sample was ground to powder using a mortar previously frozen in
 236 liquid nitrogen. The frozen powder was weighed (100 mg $\pm$ 1mg) in an Eppendorf tube,
 237 and 500 μ l of extraction solvent (MeOH:H₂O:formic acid 80:20:0.5) and a few glass
 238 beads were added. Samples were briefly vortexed and then extracted in a bead mill for
 239 three minutes at 30Hz. After centrifugation at 10,000 rpm for 10 min (Hettich
 240 mikrolitter D 7200, Buford, GA, USA) the supernatant was transferred to a new
 241 Eppendorf tube, to which 350 μ l of dichloromethane was added. Samples were vortexed
 242 and centrifuged again to separate the two phases. The upper phase was recovered (150
 243 μ l) and transferred to HPLC vials.

244 Metabolite analysis was performed using an Acquity UPLCTM system (Waters) coupled
 245 to Synapt G2 QTOF mass spectrometer (Waters) through an electrospray interface
 246 (ESI). The separation was performed on an Acquity BEH C18 column (50 $\times$ 2.1 mm
 247 i.d., 1.7 μ m particle size) at a flow rate of 0.6 mL min⁻¹. The injection volume was 3 μ l
 248 and the autosampler and column temperatures were kept at 15 and 40 °C, respectively.
 249 The mobile phase consisted of 0.05% formic acid (FA) in water (phase A) and 0.05%
 250 FA in acetonitrile (phase B). The segmented gradient program was as follows: 2% B to
 251 35% B in 3.0 min, 35% B to 100% B in 3.0 min, held at 100% B for 1.5 min, re-
 252 equilibrated to initial conditions (2% B) for 1.5 min. Data acquisitions was performed in
 253 ESI-negative and ESI-positive modes over a mass range of 100–1000 Da. The MSe
 254 mode, in which the instrument alternatively acquires data at low (4 eV; 0.15 s scan
 255 time) and high (10-30 eV ramp; 0.15 s scan time) collision energies, was used. The
 256 mass spectrometer was internally calibrated by infusing a 500 ng/mL solution of
 257 leucine-enkephalin at a flow rate of 15 μ l/min through the LockSprayTM probe. The
 258 system was controlled by Masslynx v4.1.

259 Metabolite raw data was transformed to CDF using Databridge provided by the
 260 Masslynx package. The CDF data was processed with R for statistical computing using
 261 XCMS package for relative quantification (Smith *et al.*, 2006). ESI-negative and ESI-
 262 positive data were combined, log-transformed and Pareto scaled prior to analysis. Pareto
 263 scaling gives each variable a variance equal to the square root of its standard deviation.

264 The advantage of using this technique rather than scaling to unit variance is that the
 265 former reduces the relative importance of large values but keeps data structure partially
 266 intact (van den Berg *et al.*, 2006). First a permutational MANOVA was used to evaluate
 267 whether the metabolite fingerprint consistently varied among genotypes, nitrogen
 268 availability regimes and herbivore treatments and the influences of the interactions of
 269 the factors (permutations=999). Next, a principal component analysis (PCA) was
 270 conducted as an unsupervised method to visualize variability and clustering in the data
 271 set.

272 Partial least squares–discriminant analyses (PLS–DA) were performed to identify
 273 differently detected ions between plant experimental factors - wild type vs. Carolight^R,
 274 control nitrogen vs. nitrogen treatment, and controls vs. leafhopper-induced plants -
 275 given that interactions between factors were non-significant in the perMANOVA. PLS–
 276 DA is a supervised multivariate analysis technique, which maximizes the covariance
 277 between the X–(spectral intensities) and the Y–matrix (group information). We assessed
 278 model reliability using CV-ANOVA. New components were only added to the model
 279 when significant according to the cross–validation. R^2X and R^2Y represent the fraction
 280 of the variance of X and Y matrix, respectively, while Q^2Y suggests the predictive
 281 accuracy of the model. Variable influence on projection (VIP) was used to select the
 282 most influential metabolites to group separation in the validated PLS-DA models. The
 283 VIP values summarize the overall contribution of each X-variable to the model,
 284 summed over all components and weighted according to the Y variation accounted for
 285 by each component. The Sum of squares of all VIP's is equal to the number of terms in
 286 the model - the average VIP is equal to 1- and thus terms with large VIP are the most
 287 important for explaining Y. We considered that metabolites with a $VIP > 2$ were
 288 extremely influential for treatment separation. The ions with $VIP > 2$ for each
 289 experimental factor (genotype, nitrogen and herbivory) were screened for putative
 290 identification using the pathway tool from MarVis (Kaeffer *et al.*, 2014). The MS/MS
 291 fragmentation of the metabolites was compared with candidate compounds identified in
 292 databases or earlier publications, especially when the metabolites were already reported
 293 in maize. Metabolite multivariate analysis (PCA, PLS-DA) was performed with
 294 SIMCA–P software (v. 11.0, Umetrics, Umeå, Sweden).

295

296 RESULTS

297 *Effects of genotype and nitrogen on arthropod communities in the field*

298 The most prominent source of variation in insect abundances in the field was plant
299 developmental stage, which reflects seasonal insect dynamics in the plot (Table 1).
300 Thus, abundance of maize herbivores was mainly influenced by the developmental stage
301 of the plant (perMANOVA $R^2 = 0.62$, $p < 0.001$) and to a minor extent by nitrogen
302 regime ($R^2 = 0.13$, $p = 0.085$) while no effects were attributable to genotype ($R^2 = 0.07$,
303 $p = 0.213$) or genotype x nitrogen interaction ($R^2 = 0.02$, $p = 0.684$). Similarly, maize
304 developmental stage was the main factor explaining the variation in the abundance of
305 the natural enemies recorded in the study ($R^2 = 0.28$, $p < 0.001$) whilst genotype and
306 nitrogen were not significant for determining community composition.

307 Leafhoppers and thrips were the most abundant herbivore taxa in the field, and
308 Anthocoridae and spiders the most abundant natural enemy taxa (Supplementary
309 material, Fig. S.2, S.3, S.4). Univariate analysis revealed that Hemipteran herbivores
310 (leafhoppers, planthoppers and aphids) were more abundant in the higher nitrogen
311 treatments independently of population dynamics (Table 1). Only leafhopper
312 populations were influenced by plant genotype: Carolight^R plots supported lower
313 populations of leafhopper nymphs than the wild type (Table 1). Levels of other
314 herbivores such as thrips and Lepidoptera were not influenced by nitrogen treatment or
315 genotype (Table 1). Overall the variation of natural enemy taxa was attributable to
316 population dynamics, and no differences were detected between any of the treatments
317 (Table 1).

318 *Effects of plant variety and nitrogen levels on herbivore choice and performance*

319 Plants from both genotypes in the high nitrogen hydroponic treatments (+N) were taller
320 and shoots were more robust than plants grown under control nitrogen conditions
321 (Supplementary material Fig. S.1). Genotype and nitrogen factors did not impact
322 herbivore performance as sex was the only significant predictor of final weight
323 ($F_{1,62} = 121.40$, $p < 0.001$) and developmental time ($F_{1,62} = 8.71$, $p = 0.032$). Overall, plants
324 from both genotypes grown under high nitrogen attracted more female leafhoppers than
325 plants grown with no additional nitrogen ($\chi^2_1 = 25.22$, $p < 0.001$); however, when
326 considering only the high nitrogen treatment Carolight^R was preferred ($\chi^2_1 = 4.19$,
327 $p = 0.04$) (Fig. 2). Leafhoppers chose maize plants over empty bottle control treatments
328 ($\chi^2_5 = 30.70$, $p < 0.001$) a result that validates the experimental setup.

329

330 *Effects of plant variety, nitrogen levels and herbivory on volatile compounds*

331 Seven volatile compounds were quantified in our study (Table 2) and all seven had been
 332 previously reported in maize (Degen *et al.*, 2004). We expected a small number and
 333 amount of volatile compounds in control and herbivore induced plants given that (i) the
 334 wild type line M37W produces low amounts of volatile inducible terpenes (Richter *et al.*,
 335 2016) and that (ii) *Z. scutellaris* induced plants do not emit the green leaf volatiles
 336 (Z)-3-hexenal and (E)-2-hexenal (Ardanuy *et al.*, 2016). Herbivory explained the most
 337 variability in volatile blends (perMANOVA $R^2=0.647$, $p=0.001$), and a clear separation
 338 between control and herbivore induced plants was observed in PC1 (Supplementary
 339 material, Fig. S.5). Herbivore damaged plants emitted DMNT, indole, E- β -farnesene
 340 and (E)- β -bergamotene in addition to α -copaene, E- β -caryophyllene and β -
 341 sesquiphellandrene. However, a significant genotype per nitrogen interaction was
 342 detected (perMANOVA $R^2=0.036$, $p=0.007$). In particular, individual differences in
 343 volatile emission between nitrogen regimes could be attributed for α -copaene and E- β -
 344 caryophyllene (Table 2), while differences between genotypes were only detected for β -
 345 sesquiphellandrene in the high nitrogen treatment consistent with the preference of *Z.*
 346 *scutellaris* females for Carolight^R +N in the olfactometer assay. An effect of the
 347 experimental day of volatile collection was detected on the volatile blend ($R^2=0.028$,
 348 $p=0.011$).

349 *Effects of plant variety, nitrogen availability and herbivory on phytohormone and* 350 *hydroxycinnamic acid accumulation*

351 To further investigate the effect of genotype, nitrogen and herbivore attack on plant
 352 defenses, the concentrations of the phytohormones JA, OPDA, JA-Ile, SA, ABA and
 353 IAA were measured together with the hydroxycinnamic acids caffeic, ferulic and
 354 chlorogenic acid. The concentration of JA-Ile, JA, and SA, was significantly influenced
 355 by herbivory and time point (Fig. 3, models in Supplementary material Table S.1).
 356 Interestingly, feeding by the herbivore *Z. scutellaris* significantly repressed JA-Ile and
 357 JA, as mean levels of JA-Ile and JA in herbivore-damaged plants was lower than in
 358 their respective undamaged controls (Fig 3). This trend was also significant but not as
 359 clear for SA and ABA accumulation after herbivory by maize leafhoppers (Fig. 3,
 360 models in Supplementary material Table S.1). Hormone concentrations were similar

among genotype per nitrogen treatments at all time points with the exception of (i) SA levels that were lower in Carolight^R relatively to M37W (Fig 3) and (ii) OPDA accumulated in higher concentrations in plants when grown under high nitrogen (Fig 3). Overall, caffeic and chlorogenic acid concentrations were up to 2-fold lower in Carolight^R than in the wild type (Fig 4). Caffeic acid concentration also depended on herbivory, time point and time point per nitrogen interaction (Fig. 4 Supplementary material Table S.1), whereas chlorogenic acid accumulation varied greatly between nitrogen regimes with its concentration practically doubling under control versus high nitrogen treatments (Fig 4). No consistent differences were detected for ferulic acid accumulation for any of the factors (Supplementary material, Table S.1).

Effects of plant variety, nitrogen availability and herbivory on the metabolite fingerprint

In total 4271 and 2002 markers were detected in ESI (+) and ESI (-) mode, respectively. Overall, nitrogen availability was the main factor contributing to the observed chemotypes (perMANOVA, $R^2=0.124$, $p=0.001$), followed by genotype ($R^2=0.038$, $p=0.030$) and herbivory ($R^2=0.034$, $p=0.048$) while interactions of the experimental factors were non-significant. An unsupervised approach (PCA) showed that nitrogen metabolites from plants subjected to control and high nitrogen treatments clearly grouped in the first two PCs (Fig. 5), independently of the plant genotype and herbivore treatment. In contrast, genotype and herbivory related profiles could not be separated by PCA. However, a supervised partial least squares discriminant analysis (PLS-DA) model separated (i) nitrogen regimes (ii) maize genotypes, and (iii) healthy and herbivore damaged plants (Table 3, validated through CV-ANOVA). These PLS-DA models were used to identify the metabolites showing the maximum difference between treatments with VIP values >2 (Table 3), and subsequently the selected metabolites for each experimental factor (variety, nitrogen and herbivory) were screened for putative identification using the pathway tool from MarVis 2.0 software (Kaeffer et al 2014) (Table 3, Table 4). Mean intensities of the markers plant genotype, nitrogen and herbivory by *Z. scutellaris* are represented in Supplementary material (Fig. S6, S.7 and S.8).

393 DISCUSSION

394 We addressed the hypothesis that a nutritionally enhanced maize (Carolight^R), similar in
 395 terms of biomass and yield to its wild type line M37W (Zanga *et al.*, 2016), will also be
 396 equivalent in terms of plant-insect interactions. Evaluating Carolight^R and wild type
 397 genotypes under contrasting nitrogen levels allowed for (i) a broader characterization of
 398 the resulting chemotypes and their impact on insect behavior/performance; and (ii) a
 399 comparative analysis of the impact of experimental factors (nitrogen, genotype,
 400 herbivory) on the final chemotypes. We demonstrated that nitrogen availability is the
 401 main factor determining herbivore preference and the metabolite fingerprint in
 402 Carolight^R and M37W, followed by the introduced traits and herbivory. There were no
 403 significant effects of nitrogen x genotype or herbivory x genotype interactions,
 404 suggesting that both genotypes behaved similarly when grown under the same nitrogen
 405 conditions.

406 *Insect abundance and performance on Carolight^R in contrasting nitrogen availability* 407 *conditions*

408 Overall, the community of herbivores was similar for both Carolight^R and M37W
 409 genotypes. Yet in the case of Hemiptera (leafhoppers, planthoppers and aphids) higher
 410 abundances were detected in plots with high nitrogen while only the leafhopper *Z.*
 411 *scutellaris* nymph abundances were significantly higher for M37W. Nitrogen is one of
 412 the most frequently used fertilizers in agricultural production and is known to exert a
 413 variety of bottom-up effects and potentially alter tritrophic interactions through various
 414 mechanisms (Chen *et al.*, 2010), especially for herbivorous Hemiptera (Butler *et al.*,
 415 2012). Hemipterans are insects with a high potential sensitivity to plant quality as they
 416 have been reported to prefer and perform better on some genotypes or on plants that
 417 differ in quality in terms of nutritional requirements (e.g. nitrogen content), physical or
 418 chemical plant defense (e.g. Kallenbach *et al.*, 2011; Zytynska and Preziosi, 2011). A
 419 number of reports suggest that herbivore Hemiptera (especially leafhoppers and aphids)
 420 are more abundant and/or perform better on Bt maize lines compared to their
 421 corresponding near isogenic counterparts (Lumbierres *et al.*, 2004, 2010; Pons *et al.*,
 422 2005; Obrist *et al.*, 2006; Virla *et al.*, 2010; Rauschen *et al.*, 2011). The underlying
 423 mechanism(s) responsible for such differences have not been attributed to specific
 424 factors, rather to pleiotropic effects. Pleiotropic effects reported for Bt maize that might

influence Hemipteran densities are higher lignin content in the stem of Bt plants (Saxena and Totzky, 2001), reduced amount of VOC emission in a Bt line (Turlings *et al.*, 2005) and sap amino acid content (Faria *et al.*, 2007).

Insect herbivores are limited by low nitrogen concentrations in food plants, and therefore herbivore performance is generally thought to be positively related to increases in nitrogen content in plants (Awmack and Leather, 2002; Behmer, 2009; Butler *et al.*, 2012). The performance of *Z. scutellaris* nymphs was similar when fed on Carolight^R and M37W grown under control and high nitrogen levels. This result was unexpected as we hypothesized that nitrogen availability would be the main factor contributing to adult final weight as a proxy for reproductive fitness. However, female leafhoppers preferred maize plants grown under high nitrogen in the olfactometer test, and even preferred Carolight^R over M37W when plants were grown under high nitrogen. This fact - together with field data on leafhopper abundance - supports the notion that host plant quality (resulting from enhanced nitrogen fertilization) might indeed offer other advantages to the species, such as reproductive success, that are not reflected by adult body weight or duration of nymphal development. Prestidge (1982) reported an increasing oviposition of *Z. scutellaris* as the nitrogen fertilization increased in the grass *Holcus lanatus*. Therefore the lack of differences in adult weight and developmental time for maize leafhoppers in our experiments could be a product of a mismatch between adult size and fecundity in *Z. scutellaris* as it has been previously described for grasshoppers (Joern and Behmer, 1998). Several features including field abundance, plant preference and fecundity could provide the best measures of performance for *Z. scutellaris* in general.

Maize defense responses to a mesophyll-feeding leafhopper

Plant damage together with salivary secretions of phytophagous arthropods are known to trigger plant inducible defense responses (Alborn *et al.* 1997, Musser *et al.* 2002). In turn inducible plant defenses can be major determinants of ecological interactions, and in particular defenses depending on JA and SA pathways appear to play important roles in determining community composition (Thaler *et al.*, 2001; Thaler, 2002; Kallenbach *et al.*, 2011). Hence it was vital to determine whether Carolight^R and its wild type parent (M37W) behave similarly in terms of constitutive profiles of JA and SA and in hormonal response when facing herbivory. Cell content feeders, such as the spider mite

457 *Tetranychus urticae* Koch (Acari: Prostigmata) and the thrips *Frankliniella occidentalis*
 458 (Pergande) (Thysanoptera: Thripidae) usually stimulate JA-inducible genes upon attack
 459 (Vos *et al.*, 2005), although there are reports that confirm the activation of both SA- and
 460 JA-inducible genes (Kant *et al.*, 2004; Kawazu *et al.*, 2012). Given that typhlocybine
 461 leafhoppers such as *Z. scutellaris* feed on the mesophyll using a sawing laceration
 462 strategy (Marion-Poll *et al.* 1987, Backus *et al.* 2005) we predicted that feeding by the
 463 leafhopper would activate either the JA and/or SA pathways. Interestingly feeding by
 464 this herbivore appears to decrease the constitutive levels of JA and SA on maize plants,
 465 and this was reflected in that phytohormone levels in leafhopper damaged plants were
 466 similar or even lower than the constitutive levels in healthy control plants, in particular
 467 those of JA-Ile.

468 Suppression of plant defenses is a well-known phenomenon in plant pathogens such as
 469 pathogenic bacteria, rust fungi, oomycetes, viruses, and herbivores such as nematodes
 470 and spider mites (reviewed by Kant *et al.* 2015 and Zhang *et al.* 2017). Spider mite
 471 *Tetranychus evansi* suppresses both JA and SA dependent defenses in tomato enhancing
 472 their performance (Sarmiento *et al.*, 2011; Alba *et al.*, 2015). In the case of insects the
 473 majority of cases of plant defense suppression has been attributed to JA-SA hormonal
 474 crosstalk (Walling, 2000; Zhang *et al.*, 2017) and not to a direct blocking of JA or SA
 475 defenses. However, recently aphids and mites have been reported to deliver effectors
 476 when feeding as a strategy to overcome host-plant defenses and improve their fitness
 477 (Hogenhout and Bos, 2011; Kant *et al.*, 2015; Mugford *et al.*, 2016; Villarroel *et al.*,
 478 2016). In our experimental system, *Z. scutellaris* - by feeding and oviposition -
 479 suppresses JA and does not induce SA in maize plants, and hence hormonal suppression
 480 appears to occur independently of SA-JA cross talk (as was the case for *T. evansi*). Non-
 481 targeted metabolomics fingerprinting allowed the identification of markers of herbivory
 482 by *Z. scutellaris*, which opens a door to further research on the potential effectors
 483 delivered by the leafhopper and the mechanism of defense suppression.

484 Defense manipulation by maize leafhopper impaired phytohormone accumulation in the
 485 plant without disturbing plant indirect defense by means of herbivore induced plant
 486 volatile emission. Previous work showed that maize plants damaged by ten *Z.*
 487 *scutellaris* adults emitted a similar amount of volatiles than plants damaged by the five
 488 2nd instar *Spodoptera littoralis*, and that the predatory anthocorid *Orius majusculus* was
 489 innately attracted towards the volatile blend (Ardanuy *et al.*, 2016). We hypothesize that

the suppression of JA defenses ultimately benefits leafhopper reproduction and nymphal performance, but natural enemies will still protect the plant through top-down control. However, defense manipulation by maize leafhoppers might also have consequences for subsequent colonizing herbivores since maize plants with suppressed defenses might promote the performance of co-occurring herbivores (Stam *et al.*, 2014; Kant *et al.*, 2015).

Nitrogen determines the chemical defense attributes of Carolight and M37W

Evaluating Carolight^R and wild type genotypes in contrasting nitrogen conditions allowed for a comparative analysis of the impact of experimental factors (nitrogen, genotype, herbivory) on the final maize chemotypes. We demonstrated that nitrogen is the main factor determining the metabolite fingerprint in Carolight^R and M37W, followed by the introduced traits and herbivory. Our results corroborate the work of Coll *et al.* (2010) where transcript analysis in two maize Bt(Cry1Ab trait)/wild type pairs in the field indicated that differences between lines (genetic background) exerted the highest impact on gene expression patterns, followed by nitrogen availability, while the Cry1Ab trait had the lowest impact. Barros *et al.* (2010) compared two GM maize pairs - Bt (Cry1Ab) and glyphosate tolerant - using transcriptome, proteome, and metabolome profiling and reported that the environment affected gene expression, protein distribution, and metabolite content more strongly than the introduced traits. Our results are therefore consistent with the literature and show that environmental factors (e.g. field location, sampling time during the season or at different seasons, mineral nutrition) consistently exert a greater influence on crop lines than the genetic modification itself (reviewed by Riccroch *et al.* 2011).

In general, nitrogen fertilization increases plant growth and reproduction, decreases concentrations of carbon-based secondary compounds (e.g. phenolics and terpenoids), and increases nitrogenous compounds (Koricheva *et al.*, 1998; Lou and Baldwin, 2004; Scheible *et al.*, 2004; Hermans *et al.*, 2006; Kusano *et al.*, 2011). Nitrogen levels influenced Carolight^R and M37W phenotypes at the metabolite level substantially, including compounds involved in direct and indirect plant defenses. Of the potential 405 markers with a VIP>2 only few were putatively identified. Some of these are secondary metabolites and contribute to the plant's constitutive defense as flavonoids or hydroxamic acids (benzoxazinoids) (Table 4, Supplementary material Fig. S.6).

522 Targeted analysis of defense metabolites showed that chlorogenic acid greatly varied
 523 with nitrogen treatments - at higher concentrations in plant tissues when nitrogen was
 524 limiting - but also with the plant genotype - M37W had higher levels of both
 525 chlorogenic and caffeic acids. Higher concentration of constitutive phenolics in plants
 526 under low nitrogen is consistent with results in *Nicotiana attenuata* (Lou and Baldwin,
 527 2004) and tomato (Stout *et al.*, 1998). Carolight^R accumulated up to 2-fold lower
 528 amounts of plant hydroxycinnamic acids (caffeic and chlorogenic acids) depending on
 529 the nitrogen treatment and time-point, and higher amounts of lignans (especially at low
 530 nitrogen) than the wild type. This suggests an effect of the genotype on the
 531 phenylpropanoid biosynthetic pathway. In addition, phenolics in the form of
 532 unidentified flavonoids were more abundant in control nitrogen maize plants (Table 4,
 533 Supplementary material Fig. S.6).

534 Metabolite fingerprinting showed that nitrogen surplus increased the accumulation of
 535 tryptophan in plants, which we identified as a marker of high nitrogen treatment.
 536 Tryptophan serves as precursor of a broad variety of nitrogen-containing aromatic
 537 secondary metabolites, such as hydroxamic acids (Fig. S.6), which play crucial roles in
 538 plant defense against herbivore feeding (Niemeyer, 2009; Balmer *et al.*, 2013). Higher
 539 levels of constitutive phenolics and hydroxamic acids would theoretically increase plant
 540 tolerance towards herbivores, as increased levels of these secondary compounds have
 541 been associated to reduced herbivory (Mithöfer and Boland, 2012; Balmer *et al.*, 2013).
 542 Olfactometer plant choice might indicate a preference towards plants with lower
 543 concentration of phenolics and higher concentration of hydroxamic acids in the plant;
 544 however, it fails to explain higher abundance of maize leafhopper nymphs in wild type
 545 plots in the field.

546 The VOC blend was also modified by nitrogen availability: a higher concentration of
 547 the sesquiterpenes α -copaene and E- β -caryophyllene was detected for Carolight^R and
 548 M37W plants under higher nitrogen. This might explain leafhopper preference towards
 549 plants grown under higher nitrogen levels. These results contrast with previous findings
 550 (Schmelz *et al.* 2003) which reported higher VOC emission in maize with limited
 551 nitrogen availability, though differences could be explained by maize varieties or by the
 552 source of nitrogen used in each study. While we applied nitrogen as both nitrate and
 553 ammonium, in the later study nitrogen was applied as nitrate. Nitrogen deficient
 554 soybean plants emitted the same range of herbivore-induced VOCs as control plants, but

quantitative changes occurred in the release of the main compound β -farnesene and two other volatiles ((Z)-3-hexenyl- α -methylbutyrate and β -bergamotene) (Winter and Rostás, 2010) and no differences were detected in *Nicotiana attenuata* (Lou and Baldwin, 2004).

Carolight^R damaged plants grown under high nitrogen emitted a larger amount of β -sesquiphellandrene; however the change in the volatile blend did not influence the community of natural enemies in the field. A blend of VOCs that varies in the composition or quantity of its components may constitute a signal with altered information content and may potentially modify the host finding behavior of herbivores and natural enemies, as it is the case for the maize leafhopper *Z. scutellaris*, which prefers Carolight^R to the wild type when grown under high nitrogen. It remains unclear whether leafhoppers respond in a dose-dependent manner to the total blend of VOCs or if other compounds at doses too small to detect (D'Alessandro et al. 2006) triggered leafhopper preference in the olfactometer.

Conclusion

We show the separate and interactive effects of nitrogen availability and genotype on the arthropod community and on the performance and behavior of a herbivore, and correlated these changes to constitutive and inducible maize defenses. We conclude that: (i) nitrogen availability greatly shapes maize metabolism, and the resulting plant chemotypes, and promotes *Z. scutellaris* preferences through the emission of a more attractive blend of VOCs; (ii) feeding by *Z. scutellaris* suppresses the accumulation of JA-Ile, JA and SA, while triggering the emission of herbivore-induced plant volatiles; and (iii) that the minor differences detected among Carolight^R and its wild-type counterpart in the phenylpropanoid pathway do not substantially alter aboveground plant-arthropod interactions.

ACKNOWLEDGMENTS

We would like to thank Michael Gaillard and Daniel Maag for help in sample preparation, Gemma Camañes for advice on nitrogen treatments, Diana LaForgia for help in arthropod sampling, Joan Safont, Carmen López and Matilde Eizaguirre for

586 providing *Z. scutellaris*, Daniela Zanga for discussions on Carolight^R agronomic
 587 performance, Victor Flors and the Metabolic Integration and Cell Signaling Group for
 588 discussion on results and data analysis, and all the members of the FARCE and E-Vol
 589 groups in Neuchâtel for their collaboration. A.A. was funded with an FPU scholarship
 590 and a visiting grant to FARCE laboratory from the Ministerio de Educación, Spain. The
 591 work was partially funded by the Spanish Government projects AGL2011-23996 and
 592 AGL2014-53970-C2-1-R, Agrotecnio (Fundació Centre de Recerca en Agrotecnologia,
 593 VitaMaize 01 project) and Fundació la Caixa (Grant P13005).
 594

REFERENCES

- Alba JM, Schimmel BCJ, Glas JJ, Ataíde LMS, Pappas ML, Villarroel CA, Schuurink RC, Sabelis MW, Kant MR. 2015. Spider mites suppress tomato defenses downstream of jasmonate and salicylate independently of hormonal crosstalk. *New Phytologist* 205, 828–840.
- Albajes R, Lumbierres B, Pons X. 2011. Two heteropteran predators in relation to weed management in herbicide-tolerant corn. *Biological Control* 59, 30–36.
- Alborn HT, Turlings TCJ, Jones TH, Stenhagen G, Loughrin JH, Tumlinson JH. 1997. An elicitor of plant volatiles from beet armyworm oral secretion. *Science* 276, 945–949.
- Ardanuy A, Albajes R, Turlings TCJ. 2016. Innate and Learned Prey-Searching Behavior in a Generalist Predator. *Journal of Chemical Ecology* 42, 497–507.
- Awmack CS, Leather SR. 2002. Host plant quality and fecundity in herbivorous insects. *Annual Review of Entomology* 47, 817–844.
- Backus EA, Serrano MS, Ranger CM. 2005. Mechanisms of hopperburn: an overview of insect taxonomy, behavior, and physiology. *Annual Review of Entomology* 50, 125–151.
- Balmer D, Flors V, Glauser G, Mauch-Mani B. 2013. Metabolomics of cereals under biotic stress: current knowledge and techniques. *Frontiers in Plant Science* 4, 82.
- Barros E, Lezar S, Anttonen MJ, Dijk JP Van, Ro RM, Kok EJ, Engel K. 2010. Comparison of two GM maize varieties with a near- isogenic non-GM variety using transcriptomics , proteomics and metabolomics. *Plant Biotechnology Journal* 8, 436–451.
- Behmer ST. 2009. Insect herbivore nutrient regulation. *Annual Review of Entomology* 54, 165–87.
- van den Berg RA, Hoefsloot HCJ, Westerhuis JA, Smilde AK, van der Werf MJ. 2006. Centering, scaling, and transformations: improving the biological information content of metabolomics data. *BMC genomics* 7, 142.
- Butler J, Garratt MPD, Leather SR. 2012. Fertilisers and insect herbivores: A meta-analysis. *Annals of Applied Biology* 161, 223–233.
- Camañes G, Pastor V, Cerezo M, Garcia-Andrade J, Vicedo B, Garcia-Agustin P, Flors V. 2012. A Deletion in NRT2.1 Attenuates *Pseudomonas syringae*-Induced Hormonal Perturbation, Resulting in Primed Plant Defenses. *Plant Physiology* 158, 1054–1066.
- Chen Y, Olson DM, Ruberson JR. 2010. Effects of nitrogen fertilization on tritrophic interactions. *Arthropod-Plant Interactions* 4, 81–94.
- Coll A, Nadal A, Collado R, Capellades G, Kubista M, Messeguer J, Pla M. 2010. Natural variation explains most transcriptomic changes among maize plants of MON810 and comparable non-GM varieties subjected to two N-fertilization farming practices. *Plant Molecular Biology* 73, 349–362.

- Cruz D, Eizaguirre M. 2015. Do *Sesamia nonagrioides* (Lepidoptera; Noctuidae) Gravid Females Discriminate Between Bt or Multivitamin Corn Varieties? Role of Olfactory and Visual Cues. *Journal of Insect Science* 15, 33–33.
- Decourcelle M, Perez-Fons L, Baulande S, et al. 2015. Combined transcript, proteome, and metabolite analysis of transgenic maize seeds engineered for enhanced carotenoid synthesis reveals pleiotropic effects in core metabolism. *Journal of Experimental Botany* 66, 3141–3150.
- Degen T, Dillmann C, Marion-Poll F, Turlings TCJ. 2004. High Genetic Variability of Herbivore-Induced Volatile Emission within a Broad Range of Maize Inbred Lines. *Plant Physiology* 135, 1928–1938.
- Faria CA, Wäckers FL, Pritchard J, Barrett D a, Turlings TCJ. 2007. High susceptibility of Bt maize to aphids enhances the performance of parasitoids of lepidopteran pests. *PloS one* 2, e600.
- Hermans C, Hammond JP, White PJ, Verbruggen N. 2006. How do plants respond to nutrient shortage by biomass allocation? *Trends in Plant Science* 11, 610–617.
- Hogenhout SA, Bos JIB. 2011. Effector proteins that modulate plant-insect interactions. *Current Opinion in Plant Biology* 14, 422–428.
- Joern A, Behmer ST. 1998. Impact of diet quality on demographic attributes in adult grasshoppers and the nitrogen limitation hypothesis. *Ecological Entomology* 23, 174–184.
- Kaever A, Landesfeind M, Feussner K, Mosblech A, Heilmann I, Morgenstern B, Feussner I, Meinicke P. 2014. MarVis-Pathway: integrative and exploratory pathway analysis of non-targeted metabolomics data. *Metabolomics*, 764–777.
- Kallenbach M, Bonaventure G, Gilardoni P a, Wissgott A, Baldwin IT. 2011. Empoasca leafhoppers attack wild tobacco plants in a jasmonate-dependent manner and identify jasmonate mutants in natural populations. *Proceedings of the National Academy of Sciences of the United States of America* 109, E1548-57.
- Kant MR, Ament K, Sabelis MW, Haring M a, Schuurink RC. 2004. Differential timing of spider mite-induced direct and indirect defenses in tomato plants. *Plant Physiology* 135, 483–495.
- Kant MR, Jonckheere W, Knecht B, et al. 2015. Mechanisms and ecological consequences of plant defence induction and suppression in herbivore communities. *Annals of Botany* 115, 1015–1051.
- Kawazu K, Mochizuki A, Sato Y, Sugeno W, Murata M, Seo S, Mitsuhashi I. 2012. Different expression profiles of jasmonic acid and salicylic acid inducible genes in the tomato plant against herbivores with various feeding modes. *Arthropod-Plant Interactions* 6, 221–230.
- Koricheva J, Larsson S, Haukioja E, Keinänen M. 1998. Regulation of woody plant secondary metabolism by resource availability: hypothesis testing by means of meta-analysis. *Oikos* 83, 212–226.
- Kusano M, Fukushima A, Redestig H, Saito K. 2011. Metabolomic approaches toward understanding nitrogen metabolism in plants. *Journal of Experimental Botany* 62, 1439–1453.

- Lou Y, Baldwin IT. 2004. Nitrogen Supply Influences Herbivore-Induced Direct and Indirect Defenses and Transcriptional Responses in *Nicotiana attenuata*. *Plant Physiology* 135, 496–506.
- Lumbierres B, Albajes R, Pons X. 2004. Transgenic Bt maize and *Rhopalosiphum padi* (Hom., Aphididae) performance. *Ecological Entomology* 29, 309–317.
- Lumbierres B, Starý P, Pons X. 2010. Effect of Bt maize on the plant-aphid-parasitoid tritrophic relationships. *BioControl* 56, 133–143.
- Mithöfer A, Boland W. 2012. Plant defense against herbivores: Chemical aspects. *Annual Review of Plant Biology* 63, 431–450.
- Mugford S, Barclay E, Drurey C, Findley K, Hogenhout S. 2016. An immuno-suppressive aphid saliva protein is delivered into the cytosol of plant mesophyll cells during feeding. *bioRxiv* X.
- Musser RO, Hum-Musser SM, Eichenseer H, Peiffer M, Ervin G, Murphy JB, Felton GW. 2002. Herbivory: caterpillar saliva beats plant defences. *Nature* 416, 599–600.
- Naqvi S, Zhu C, Farre G, Ramessar K, Bassie L, Perez D, Ros G, Sandmann G, Capell T, Christou P. 2009. Transgenic multivitamin corn through biofortification of endosperm with three vitamins representing three. *PNAS* 106, 7762–7767.
- Niemeyer H. 2009. Hydroxamic Acids Derived from 2-Hydroxy-2 H-1, 4-Benzoxazin-3 (4 H)-one: Key Defense Chemicals of Cereals. *Journal of Agricultural and Food Chemistry* 57, 1677–1696.
- Obrist LB, Dutton A, Albajes R, Bigler F. 2006. Exposure of arthropod predators to Cry1Ab toxin in Bt maize fields. *Ecological Entomology* 31, 143–154.
- Ode PJ. 2006. Plant chemistry and natural enemy fitness: effects on herbivore and natural enemy interactions. *Annual Review Entomology* 51, 163–185.
- Oksanen J, Blanchet FG, Friendly M, et al. 2013. Package ‘vegan’ | Community Ecology Package. R package.
- Pons X, Lumbierres B, López C, Albajes R. 2005. Abundance of non-target pests in transgenic Bt-maize: A farm scale study. *European Journal of Entomology* 102, 73–79.
- Prestidge RA. 1982. Instar duration, adult consumption, oviposition and nitrogen utilization efficiencies of leafhoppers feeding on different quality food (Auchenorrhyncha: Homoptera). *Ecological Entomology* 7, 91–101.
- Van Der Putten WH, Vet LEM, Harvey JA, Wäckers FL. 2001. Linking above- and belowground multitrophic interactions of plants, herbivores, pathogens, and their antagonists. *Trends in Ecology and Evolution* 16, 547–554.
- Rauschen S, Schultheis E, Hunfeld H, Schaarschmidt F, Schuphan I, Eber S. 2011. Diabrotica-resistant Bt-maize DKc5143 event MON88017 has no impact on the field densities of the leafhopper *Zyginiidia scutellaris*. *Environmental Biosafety Research* 9, 87–99.

- Richter A, Schaff C, Zhang Z, et al. 2016. Characterization of the herbivore-induced biosynthetic pathways of the volatile homoterpenes 3,8-dimethyl-1,4,7-nonatriene (DMNT) and 4,8,12-trimethyltrideca-1,3,7,11-tetraene, in *Zea mays*. Plant Cell accept wit.
- Ricroch E, Berge JB, Kuntz M, Ricroch AE, Bergé JB, Kuntz M. 2011. Evaluation of genetically engineered crops using transcriptomic, proteomic, and metabolomic profiling techniques. Plant Physiology 155, 1752–1761.
- Sarmiento RA, Lemos F, Bleeker PM, Schuurink RC, Pallini A, Oliveira MGA, Lima ER, Kant M, Sabelis MW, Janssen A. 2011. A herbivore that manipulates plant defence. Ecology Letters 14, 229–236.
- Saxena D, Totzky GS. 2001. Corn has a higher lignin content than non-Bt corn. American Journal of Botany 88, 1704–1706.
- Scheible W-R, Morcuende R, Czechowski T, Fritz C, Osuna D, Palacios-Rojas N, Schindelasch D, Thimm O, Udvardi MK, Stitt M. 2004. Genome-wide reprogramming of primary and secondary metabolism, protein synthesis, cellular growth processes, and the regulatory infrastructure of *Arabidopsis* in response to nitrogen. Plant Physiology 136, 2483–2499.
- Schmelz EA, Alborn HT, Engelberth J, Tumlinson JH. 2003. Nitrogen deficiency increases volicitin-induced volatile emission, jasmonic acid accumulation, and ethylene sensitivity in maize. Plant Physiology 133, 295–306.
- Smith CA, Want EJ, O'Maille G, Abagyan R, Siuzdak G, CA Smith, J Elizabeth, G O'Maille, Ruben Abagyan, and Gray S. 2006. XCMS: processing mass spectrometry data for metabolite profiling using Nonlinear Peak Alignment, Matching, and Identification. Analytical Chemistry 78, 779–87.
- Stam JM, Kroes A, Li Y, Gols R, van Loon JJAA, Poelman EH, Dicke M. 2014. Plant interactions with multiple insect herbivores: from community to genes. Annual Review of Plant Biology 65, 689–713.
- Stout MJ, Brovont RA, Duffey SS. 1998. Effect of nitrogen availability on expression of constitutive and inducible chemical defenses in tomato, *Lycopersicon esculentum*. Journal of Chemical Ecology 24, 945–963.
- Thaler JS. 2002. Effect of jasmonate-induced plant responses on the natural enemies of herbivores. Journal of Animal Ecology 71, 141–150.
- Thaler JS, Stout MJ, Karban R, Duffey SS. 2001. Jasmonate-mediated induced plant resistance affects a community of herbivores. Ecological Entomology 26, 312–324.
- Turlings TCJ, Davison AC, Tamò C. 2004. A six-arm olfactometer permitting simultaneous observation of insect attraction and odour trapping. Physiological Entomology 29, 45–55.
- Turlings TCJ, Jeanbourquin PM, Held M, Degen T. 2005. Evaluating the induced-odour emission of a Bt maize and its attractiveness to parasitic wasps. Transgenic Research 14, 807–16.
- Turlings TCJ, Lengwiler UB, Bernasconi ML, Wechsler D. 1998. Timing of induced volatile emissions in maize seedlings. Planta 207, 146–152.

- Villarroel CA, Jonckheere W, Alba JM, Glas JJ, Dermauw W, Haring MA, Van Leeuwen T, Schuurink RC, Kant MR. 2016. Salivary proteins of spider mites suppress defenses in *Nicotiana benthamiana* and promote mite reproduction. *Plant Journal* 86, 119–131.
- Virla EG, Casuso M, Frias E A. 2010. A preliminary study on the effects of a transgenic corn event on the non-target pest *Dalbulus maidis* (Hemiptera: Cicadellidae). *Crop Protection* 29, 635–638.
- De Vos M, Van Oosten VR, Van Poecke RMP, et al. 2005. Signal Signature and Transcriptome changes of *Arabidopsis* during pathogen and insect attack. *Molecular Plant-Microbe Interactions* 18, 923–937.
- Walling L. 2000. The Myriad Plant Responses to Herbivores. *Journal of Plant Growth Regulation* 19, 195–216.
- Winter TR, Rostás M. 2010. Nitrogen deficiency affects bottom-up cascade without disrupting indirect plant defense. *Journal of Chemical Ecology* 36, 642–651.
- Wolt JD, Keese P, Raybould A, Fitzpatrick JW, Burachik M, Gray A, Olin SS, Schiemann J, Sears M, Wu F. 2010. Problem formulation in the environmental risk assessment for genetically modified plants. *Transgenic Research* 19, 425–36.
- Zanga D, Capell T, Slafer GA, Christou P, Savin R. 2016. A carotenogenic mini-pathway introduced into white corn does not affect development or agronomic performance. *Scientific Reports* 6, 38288.
- Zhang L, Zhang F, Melotto M, Yao J, He SY. 2017. Jasmonate signaling and manipulation by pathogens and insects. *Journal of Experimental Botany*, erw478.
- Zytynska SE, Preziosi RF. 2011. Genetic interactions influence host preference and performance in a plant-insect system. *Evolutionary Ecology* 25, 1321–1333.

1 Table 1. Effects of maize genotype, nitrogen treatment, their interaction and maize's developmental stage on field abundances of herbivores and natural enemies. Arthropod
2 abundance was determined by visual sampling on five maize developmental stages. Significant effects ($\alpha=0.05$) appear in bold.

3

Herbivores		Leafhoppers		Planthoppers		Thrips		Lepidoptera		Aphids	
<i>Factors</i>	df	χ^2	p	χ^2	p	χ^2	p	χ^2	p	χ^2	p
genotype	1	4.09	0.046	0.09	0.779	1.41	0.233	2.19	0.076	0.64	0.423
nitrogen	1	5.28	0.023	11.9	<0.002	0.15	0.693	0.18	0.613	10.91	0.001
genotype $\times$ nitrogen	1	2.1	0.150	0.43	0.547	0.45	0.501	1.38	0.161	2.57	0.109
develop. stage	4	223.14	<0.001	32.98	<0.001	157.96	<0.001	181.53	<0.001	45.65	<0.001

Natural enemies		Anthocoridae		Chrysopidae		Thrips		Coccinellidae		Miridae		Arachnida	
<i>Factors</i>	df	χ^2	p	χ^2	p	χ^2	p	χ^2	p	χ^2	p	χ^2	p
genotype	1	0.012	0.914	0.29	0.564	0.161	0.686	0.05	0.83	1.17	0.286	1.14	0.306
nitrogen	1	0.49	0.49	0.29	0.564	0.875	0.346	0.21	0.67	0.16	0.69	1.14	0.306
genotype $\times$ nitrogen	1	1.95	0.168	0.02	0.881	0.24	0.619	0.15	0.72	1.90	0.175	0.00	0.992
develop. stage	4	52.41	<0.001	16.45	<0.001	45.08	<0.001	22.68	<0.001	57.08	<0.001	16.93	0.004

4 Fig 1. Choice of maize volatiles by leafhopper *Z. scutellaris* on the olfactometer. Tested plants consisted
5 of Wild type and Carolight^R plants grown under control or surplus nitrogen (+N) conditions. Different
6 letters indicate differences between treatments ($\alpha=0.05$).

7

8

9

Table 2. Volatile emissions (ng/h) from control and herbivore-induced maize plants (Carolight^R, Wild type) at two different N availability treatments (control, +N) (n=7). Amounts of each compound were compared among treatments using a non-parametric Kruskal-Wallis test followed by Dunn's test (*p<0.05, **p<0.01, ***p<0.001). Compounds denoted with "N" were only tentatively identified by comparison of their MS to that reported in libraries.

	Wild type		Carolight		Wild type+N		Carolight+N		χ^2	P
	mean	±SE	mean	±SE	mean	±SE	mean	±SE		
<i>Control</i>										
α-copaene	0.59	0.13	0.67	0.17	0.51	0.09	1.21	0.22	6.11	0.11
E-β-caryophyllene	0.44	0.06	0.35	0.06	0.32	0.05	0.78	0.15	6.31	0.10
β-sesquiphellandrene^N	1.36	0.26	1.16	0.34	1.01	0.19	2.96	0.69	5.53	0.14
<i>Induced</i>										
DMNT	3.74	0.79	3.44	0.80	3.51	1.46	4.27	0.94	0.98	0.80
Indole	7.79	2.55	5.29	1.40	7.64	4.00	6.85	1.68	1.13	0.77
α-copaene	0.75ab	0.16	0.60a	0.11	1.19bc	0.25	1.35c	0.12	9.10	0.03
E-β-caryophyllene	0.45a	0.08	0.40a	0.07	0.79b	0.18	0.92b	0.13	9.00	0.03
(E)-β-bergamotene	1.33	0.34	1.01	0.20	1.54	0.49	1.37	0.22	1.97	0.58
E-β-farnesene	4.97	1.45	2.81	0.85	5.37	1.90	4.32	1.03	1.48	0.69
β-sesquiphellandrene^N	2.08b	0.44	1.57ab	0.32	0.80a	0.14	3.68c	0.45	13.90	<0.001

Fig 3. Hormonal content (ng/g dry weight) in Carolight^R and Wild type plants grown under two different nitrogen regimes (control and +N) upon *Z. scutellaris* feeding. Control and herbivore-damaged plants (+H) were collected at different time points (24h, 48h and 96h after herbivore feeding), and phytohormone levels were determined in freeze-dried material by UPLC-MS. The experiment was replicated 3 times with similar results. Full factorial models combining data of the three experiments are available in Supplementary material Table S.1. The results shown are mean ($\pm$ SE) hormone levels of one experiment. Asterisks indicate differences among treatments (non-parametric Kruskal-Wallis test).

Fig 4. Caffeic acid (A) and chlorogenic acid (B) content (ng/g dry weight) of Carolight^R and Wild type plants grown under two different nitrogen regimes (control and +N). Plants were collected at different time-points (24h, 48h and 96h after the start of the experiment), and caffeic and chlorogenic acid levels were determined in freeze-dried material by UPLC-MS. The experiment was replicated 3 times with similar results. Full factorial models combining data of the three experiments are available in Supplementary material Table S.1. The results shown are mean ($\pm$ SE) hormone levels of one experiment. Asterisks indicate differences among treatments (non-parametric Kruskal-Wallis test).

34 Fig 5. Principal component analysis (PCA) showing groups generated from signals obtained in ESI+ and
 35 ESI- by non-targeted analysis. Carolight^R and Wild type (Wt) plants were grown under two different
 36 nitrogen regimes (control and +N) upon *Z. scutellaris* feeding (control and +H).

41 Table 3. PLS-DA models of the factors nitrogen, genotype and herbivory. For each factor, the number of
42 ions with a VIP>2 for the best model and the number of MarVis Pathway IDs are specified.

Factor	Components	R ²	Q ²	CV-ANOVA		VIP>2	MarVis ID
Nitrogen	1	0.875	0.744	F_{2,35}=50.89	p=4.38x10⁻¹¹	405	48
	1+1	0.968	0.999	F _{4,33} =34.80	p=2.08x10 ⁻¹¹		
Genotype	1+1	0.956	0.581	F_{4,33}=7.87	p=0.00014	69	19
	1+1+1	0.993	0.803	F _{6,31} =6.25	p=0.00022		
Herbivory	1+1	0.967	0.486	F_{4,33}=3.97	p=0.0097	163	15
	1+1+1	0.99	0.634	F _{6,31} =2.79	p=0.027		

43

44

Table 4. Metabolites with higher loadings on the best PLS-DA models (VIP>2) for the factors Nitrogen, Genotype and Herbivory that could be tentatively identified. Metabolites in A were identified by comparison of the MS/MS to online databases, while metabolites in B were assigned after comparing the accurate mass to reference compound databases. Mean abundances of the metabolites can be found in Supplementary material Fig. S.6, S.7 and S.8. Pathways were assigned using the pathway tool from MarVis 2.0 (Kaeffer et al., 2014).

A. Identified	Mass neutral	ESI	RT (min)	Factor	Pathway
Kynurenic acid	189.0431	+	0.99	Herbivory (H↑)	Tryptophan-kynurenine pathway
11-trans-LTD4	496.2592	-	3.43	Herbivory (H↑)	Leukotriene- Arachidonic acid metabolism
19-HETE/ 20-HETE	320.236	+	2.06	Genotype (Wt↑)	Arachidonic acid metabolism
Thiamine	265.1153	+	0.86	Genotype (Wt↑)	Vitamin and cofactor - Thiamine metabolism (primary metabolism)
Phytosphingosine	317.2932	+	4.41	Genotype (Wt↑)	Sphingolipid metabolism (primary metabolism)
L-Tryptophan	204.0897	-	0.93	Nitrogen (+N↑)	Tryptophan pathway (primary metabolism)
DIMBOA-Glu	373.1009	-	1.46	Nitrogen (+N↑)	Benzoxazinoid biosynthesis
B. Assigned to	Mass neutral	ESI	RT (min)	Factor	Pathway
Sinapoyl malate	340.0791	-	2.73	Herbivory (H↓)	Biosynthesis of phenylpropanoids
Porphobilinogen	226.0965	-	0.76	Herbivory (H↑)	Porphyrin and chlorophyll metabolism
(-)-Jasmonoyl-L-isoleucine	323.2097	+	4.97	Herbivory (H↓)	Biosynthesis of plant hormones
A-tocopherol	430.3777	+	5.92	Herbivory (H↑)	Biosynthesis of plant secondary metabolites
2,3-Dihydroxybenzoate	154.0263	-	1.44	Herbivory (H↑)	Biosynthesis of phenylpropanoids
Coniferol	180.0788	+	0.84	Genotype (Wt↑)	Biosynthesis of phenylpropanoids
Cis-hinokiresinol	252.1195	-	0.84	Genotype (Wt↑)	Biosynthesis of phenylpropanoids
Unknown flavonoid	306.0775	-	1.36	Genotype (Wt↓)	Flavonoid biosynthesis
Unknown flavonoid	578.1634	+	1.78	Genotype (Wt↓) Nitrogen (+N↓)	Flavone and flavonol biosynthesis
Unknown flavonoid	610.1544	+	1.79	Genotype (Wt↓) Nitrogen (+N↓)	Flavone and flavonol biosynthesis
Unknown flavonoid	448.1016	+/-	1.59/2.05	Nitrogen (+N↓)	Flavone and flavonol biosynthesis
Unknown flavonoid	594.1579	-	1.61	Nitrogen (+N↓)	Flavone and flavonol biosynthesis
Unknown flavonoid	286.0483	+	1.99	Nitrogen (+N↓)	Flavone and flavonol biosynthesis
Unknown flavonoid	464.0949	-	1.86	Nitrogen (+N↓)	Flavone and flavonol biosynthesis
Unknown flavonoid	592.1802	+	2.73	Nitrogen (+N↓)	Flavone and flavonol biosynthesis
Zeatin/Pantothenate	219.1113	+	0.8	Nitrogen (+N↓)	Biosynthesis of plant hormones
HBOA	149.0471	-	0.73	Nitrogen (+N↑)	Benzoxazinoid biosynthesis
DHBOA/ DIBOA-Glc	343.0891	-	1.27	Nitrogen (+N↑)	Benzoxazinoid biosynthesis
Dhurrin	311.1002	-	0.73	Nitrogen (+N↑)	Cyanoamino acid metabolism