1 **Title**

2 In vitro evaluation of antagonism by *Trichoderma* spp. towards *Phellinus noxius* associated with rain
3 tree (*Samanea saman*) and Senegal mahogany (*Khaya senegalensis*) in Singapore

4

5 **Authors**

6 Daniel C. Burcham^{1*}; Wong Jia Yih²; Nelson V. Abarrientos, Jr.¹; Mohamed Ismail Mohamed Ali²;
7 Fong Yok King¹; Francis W.M.R. Schwarze³

8

9 ¹Centre for Urban Greenery and Ecology

10 National Parks Board

11 Singapore 259601

12

13 ²Phytopathology and Integrated Pest Management Section

14 Plant Health Laboratory Department

15 Agri-Food & Veterinary Authority

16 Singapore 718827

17

18 ³Applied Wood Materials

19 Empa Swiss Federal Laboratories for Science and Technology

20 St. Gallen, Switzerland CH-9014

21

22 *Corresponding author: daniel_christopher_burcham@nparks.gov.sg

23

1 **Abstract** (250 words)

2 A series of laboratory tests was conducted to screen and identify *Trichoderma* spp. for the biological
 3 control of *P. noxius* on pruning wounds. In total, 7 *Trichoderma* isolates were evaluated for their
 4 biological fitness and antagonism towards 2 different *P. noxius* isolates associated with Senegal
 5 mahogany and rain tree in Singapore. The competitive ability of various *Trichoderma* isolates was
 6 assessed by their germination rates, growth rates, and chlamydospore production; and the antagonism
 7 of *P. noxius* by *Trichoderma* was assessed by the interaction of these fungi in dual culture and on
 8 wood blocks. In this study, the *Trichoderma* isolates showed greater competitiveness and antagonism
 9 towards *P. noxius* compared to similar reports in the literature. A majority of the *Trichoderma* isolates
 10 germinated and grew at consistently high rates, but there was considerable variability in the
 11 production of chlamydospores among isolates. All *Trichoderma* isolates routinely antagonized *P.*
 12 *noxius* in the different bioassays, but there was significant variability in the antagonistic capacity of
 13 various isolates and in the susceptibility of the *P. noxius* isolates to confrontation with *Trichoderma*.
 14 Relative to the controls, *P. noxius* cultures grew significantly slower in the presence of volatile
 15 organic compounds emitted by most *Trichoderma* isolates, and a large majority of *Trichoderma*
 16 isolates caused a significant reduction to the dry weight loss of wood blocks inoculated with *P.*
 17 *noxius*. Although no single *Trichoderma* isolate consistently antagonized *P. noxius* better than all
 18 others in every test, *T. harzianum* 9132 and *T. virens* W23 notably did so more regularly than others.

19

20 **Keywords**

21 Biological control; Wood decay; Pruning wounds

22

23 **Introduction**

24 Trees are pruned for many reasons, including aesthetics, clearance, and safety. However, the exposed
 25 wood on pruning wounds is susceptible to wood decay infection (Barry et al., 2000; Deflorio et al.,
 26 2007), and these infections are a concern to people interested in preserving the environmental benefits
 27 and commercial value of trees. Historically, arborists often applied various treatments to pruning
 28 wounds to create a physical barrier or inhibit wood decay fungi (Lonsdale, 1984). Although some of

1 these treatments improved wound occlusion (Mercer, 1983) and temporarily prevented infection
 2 (Mercer et al., 1983), the physical barriers frequently deteriorated over time due to weathering, insect
 3 damage, and growth stress that, in some cases, exacerbated decay compared to untreated wounds
 4 (Mercer et al., 1983). Similarly, some fungicides were reported as effective at limiting infection but
 5 their phytotoxicity often caused harmful cambial dieback (Mercer et al., 1983; Mercer, 1979) that
 6 inhibited wound occlusion and increased wood decay by expanding the exposed wound surface. In
 7 addition to their limited efficacy, the use of synthetic chemicals, especially on trees planted in densely
 8 populated urban areas, creates environmental and human health risks that many consider
 9 unacceptable. As a result, most arboriculture industry standards discourage wound treatment
 10 (Anonymous 2008).

11
 12 Biological control agents, however, offer a promising alternative to traditional wound treatments. In
 13 particular, many fungal antagonists have been selected from the genus *Trichoderma* for biological
 14 control of plant diseases, including economically important fungal pathogens affecting fruit trees
 15 (Ricard and Highley, 1988) and conifers (Kallio and Hallaksela, 1979). Notably, Pottle and Shigo
 16 (1975) reported that basidiomycetes were absent from 1-year-old red maple [*Acer rubrum* L.
 17 (Sapindaceae)] wounds preventatively treated with *T. viride*, but decay fungi were present in 80% of
 18 untreated control wounds after the same period. Mercer and Kirk (1984b) similarly reported a
 19 significant reduction in wood decay fungi infection of European beech [*Fagus sylvatica* L.
 20 (Fagaceae)] wounds treated with a different isolate of *T. viride*, compared to control wounds. Notably,
 21 *T. viride* was more effective and persisted longer than other biological controls (Mercer and Kirk,
 22 1984b). More recently, Schubert et al. (2008a) demonstrated the ability of a *T. atroviride* isolate to
 23 significantly reduce wood decay infection rates among more than 1,400 wounds treated from 6
 24 different broadleaf tree species.

25
 26 *Trichoderma* spp. (Samuels, 1996) are ubiquitous, globally distributed saprophytes that occur
 27 primarily in highly organic soils (Papavizas, 1985), but they are capable of colonizing a wide range of
 28 natural substrates (Klein and Eveleigh, 1998). Many *Trichoderma* spp. are avirulent plant symbionts

1 (Harman et al., 2004) that antagonize phytopathogenic fungi through mycoparasitism (Lorito et al.,
2 1996a), antibiosis (Ghisalberti and Sivasithamparam, 1991), enzyme production (Markovich and
3 Kononova, 2003), and competition for resources (Sivan and Chet, 1989). In most cases, a single
4 *Trichoderma* spp. isolate simultaneously employs more than one of these antagonistic mechanisms to
5 suppress disease-causing fungi; there are several reports of synergistic enhancement between multiple
6 antagonistic processes and substances (Di Pietro et al., 1993; Lorito et al., 1994a; Lorito et al., 1994b;
7 Lorito et al., 1996b). In addition to suppressing harmful microbes, soil application of *Trichoderma*
8 spp. enhances root growth and nutrient uptake efficiency, resulting in increased overall plant growth
9 and reproductive yield (Harman et al., 2004). Several authors have also demonstrated that some
10 *Trichoderma* spp. cause systemic induced resistance (Pieterse and van Loon, 1999) in plants by
11 observing physical separation between the introduced *Trichoderma* spp. and area of disease reduction
12 (Bigirimana et al., 1997; Djonovic et al., 2006; Yedidia et al., 1999).

13
14 Similar to mycorrhizae, many benefits accrue to plants after *Trichoderma* spp. colonize the outermost
15 epidermal layers of their roots (Yedidia et al., 2000). Although most *Trichoderma* spp. are avirulent, a
16 few isolates are pathogenic or phytotoxic (Koike et al., 2001), and it is important to screen for
17 potential biological control agents with representative laboratory tests that mimic the temperature,
18 moisture, and nutrient availability conditions expected for the desired field application (Howell,
19 2003). Moreover, many authors caution that individual *Trichoderma* spp. isolates usually do not have
20 the intrinsic ability to exert all of the reported biocontrol mechanisms (Harman, 2000; Howell, 2003),
21 and hybridization, genetic modification, or mixing may be needed to combine the desirable
22 characteristics of multiple isolates (Brunner et al., 2005; Djonovic et al., 2007).

23
24 However, most studies addressing biological control with *Trichoderma* have been conducted in
25 temperate climates with associated species. *Phellinus noxius* (Corner) G. Cunn (Hymenochaetaceae)
26 is an aggressive, often lethal wood decay fungus that infects the root systems of many tree species
27 throughout the global tropics and subtropics (Rossman and Farr, 2014). Macroscopically, the wood of
28 infected trees appears brown and the adjacent root flare is often covered by a dark brown mycelial mat

(Chang, 1995). Although the disease is commonly called brown root rot, *P. noxius* actually causes a simultaneous rot by degrading cellulose and lignin at the same time (Nicole et al., 1995). A few recent studies have evaluated the ability of *Trichoderma* to antagonize *P. noxius* obtained from root system infections in Australia (Schwarze et al., 2012) and Hong Kong (Ribera et al., 2016). Recent work in Singapore, however, demonstrated the ability of *P. noxius* to colonize pruning wounds, invade host defensive responses, and cause severe wood decay in the aboveground parts of Senegal mahogany [*Khaya senegalensis* (Desr.) A. Juss. (Meliaceae)]; this work revealed that *P. noxius* occupies a broader ecological niche than previously understood with physiological capacity for multiple trophic strategies (Burcham et al., 2015). At the same time, diagnostic efforts have shown that *P. noxius* similarly infects other trees in Singapore, including rain tree [*Samanea saman* (Jacq.) Merr. (Fabaceae)].

Collectively, Senegal mahogany and rain tree comprise more than 10% of Singapore's urban forest and generate a significant part of the total benefits rendered to the surrounding community. In general, the ubiquity of pruning wounds creates considerable opportunity for *P. noxius* infections, and limiting these infections is an important strategy to protect susceptible trees and, by extension, the health of the urban forest. As a result, the objectives of this study were (1) to screen *Trichoderma* spp. isolates for antagonistic potential against *P. noxius* associated with pruning wounds on Senegal mahogany and rain tree and (2) to identify the most competitive isolate(s) for field testing. To this end, a series of bioassays were designed to evaluate antagonism by *Trichoderma* spp. isolates towards *P. noxius* in laboratory conditions that resembled situations expected for landscape application in Singapore.

Material and methods

Origin of fungal isolates

P. noxius isolates were obtained from decayed wood columns that were excised near existing branch pruning wounds on Senegal mahogany and rain tree in Singapore. Among several obtained from these tree species, the most aggressive *P. noxius* isolate was selected for screening biological control agents based on its superior ability to cause dry weight loss to species-specific sapwood blocks in laboratory

1 tests (data not shown). In total, two *P. noxi* isolates were selected—one caused maximum dry weight
2 loss to Senegal mahogany and the other to rain tree wood blocks. Separately, 7 *Trichoderma* spp.
3 isolates were obtained from the fruiting bodies of basidiomycetes colonizing various plant species and
4 organic substrates in Singapore. The identity and origin of fungal isolates used in this study is given in
5 Table 1.

6 *Cultural media and growth conditions*

8 Three cultural media were used, including a malt extract agar (MEA, Thermo Fisher Scientific,
9 Waltham, MA, USA); a low nutrient agar (LNA) prepared according to Huttermann and Volger
10 (1973), with each liter containing 0.013 g H₂O: L-asparagine, 1 g KH₂PO₄, 0.3 g MgSO₄, 0.5 g KCL,
11 0.01 g FeSO₄, 0.008 g MnSO₄ 4H₂O, 0.002 g ZnSO₄ 6H₂O, 0.05 g CaNO₃ 4H₂O, 0.002 g CuSO₄,
12 0.008 g NH₄NO₃, 5 g D-glucose, and 10 g agar; and a basidiomycete selective medium (BSM)
13 modified from Sieber (1995), with each liter of media containing 50 g MEA, 105.75 mg
14 thiabendazole dissolved in 2 ml of concentrated lactic acid, 200 mg chloramphenicol, and 300 mg
15 streptomycin sulfate. Unless indicated otherwise, cultures were incubated consistently in the dark at
16 28 °C and 50–70% RH on 90 or 50 mm Petri dishes containing approximately 20 or 11 ml,
17 respectively, of media. All plates were inoculated with 8 mm mycelial discs extracted from the
18 actively growing margins of pure cultures on MEA and sealed with plastic paraffin film (Parafilm[®],
19 Pechiney Plastic Packaging, Chicago, IL, USA).

20 *Germination and growth rates*

22 Growth rates of the *Trichoderma* isolates were evaluated on MEA and LNA. The LNA was chosen
23 because its higher C:N ratio better replicated the nutritional status of wood (Srinivasan et al., 1992)
24 than MEA. To assess growth, 8 mm mycelial discs of *Trichoderma* were inoculated centrally onto
25 MEA or LNA in a 90 mm Petri dish. Growth rates were determined by recording the average of 2
26 orthogonal fungal culture diameter measurements at regular intervals over a 36-hour period.

27

1 To determine germination rates, a conidial suspension of each *Trichoderma* isolate was prepared by
 2 flooding mature cultures with sterile water, dislodging conidia by physical agitation, and skimming
 3 buoyant conidia from the surface. The concentration of suspensions was assessed using a
 4 haemocytometer and adjusted to obtain approximately 10^5 colony forming units (CFU) $\cdot$ ml⁻¹.
 5 Subsequently, 10 μ l of each suspension was placed in 3 separate locations on a single Petri dish
 6 containing LNA. Test plates were covered, sealed with paraffin tape, and incubated at constant
 7 conditions for 18 hours. After incubation, conidia were observed at 10 $\times$ magnification with a light
 8 microscope (Nikon Eclipse E200, Tokyo, Japan) in 5 randomly selected locations in each Petri dish.
 9 The germination rate was calculated for each plate as the average percent of germinated conidia
 10 contained in these fields of observation.

11

12 *Chlamydospore production*

13 To assess the production of thick-walled resting spores during adverse environmental conditions,
 14 chlamydospores were counted on 14-day old mature cultures of *Trichoderma* growing on LNA in a 50
 15 mm Petri dish. The number of chlamydospores was determined by counting those observed at 10 $\times$
 16 magnification in 5 randomly selected locations on each Petri dish using a digital camera system
 17 (Nikon Digital Sight DS-Fi1, Tokyo, Japan) attached to a light microscope. The number of
 18 chlamydospores on each plate was computed as the average number observed across the different
 19 locations.

20

21 *VOC inhibition*

22 Inhibition of wood decay fungi by volatile organic compounds (VOCs) produced by *Trichoderma*
 23 isolates was evaluated using tests described by Dennis and Webster (1971). Initially, 8 mm mycelial
 24 discs of *Trichoderma* isolates were placed in the center of a Petri dish containing MEA. At the same
 25 time, each of the *P. noxius* isolates was inoculated identically onto a separate Petri dish. Each
 26 *Trichoderma* isolate was then paired with a *P. noxius* isolate by inverting the bottom of one Petri dish
 27 containing *Trichoderma* and replacing the top of another containing *P. noxius*. The rims of the paired
 28 Petri dishes were aligned and sealed. The diameter of *P. noxius* cultures was recorded as the average

1 of 2 orthogonal measurements at regular 24-hour intervals over a 5-day period. Similar Petri dishes
2 without *Trichoderma* isolates served as a control.

3

4 *Interaction tests*

5 Mycoparasitism was assessed in dual culture and on wood blocks broadly according to Highley et al.
6 (1997) and Schubert et al. (2008a), respectively. For dual culture tests, 8 mm mycelial discs were
7 removed from 14-day old, actively growing pure cultures of each *P. noxius* isolate and placed
8 individually at a regular distance (~ 10 mm) from the edge of a 90 mm Petri dish containing MEA.
9 After incubation for 72 hours, equivalent mycelial discs were removed from 7-day old actively
10 growing pure cultures of each *Trichoderma* isolate, inserted on a Petri dish opposite the growing
11 wood decay fungus, covered, and sealed. The Petri dishes were incubated under constant conditions
12 for 4 weeks. The ability of *Trichoderma* isolates to parasitize *P. noxius* isolates was assessed by
13 removing 6 mycelial discs from the final extent of the wood decay fungus culture and placing discs on
14 BSM. The media's inhibition of *Trichoderma* allowed a selective determination of the lethal effect of
15 *Trichoderma* isolates, determined as one minus the percent of these 6 discs yielding *P. noxius*.

16

17 For wood block interaction tests, wood blocks (10 × 8 × 30 mm) were removed from healthy sapwood
18 in Senegal mahogany and rain tree. Individual wood blocks were oven-dried at 100 °C for 48 hours,
19 cooled in a vacuum desiccator, and weighed using a precision balance. The blocks were subsequently
20 autoclaved twice at 121 °C for 30 min. Each wood block was inoculated by immersion for 10 minutes
21 in a conidial suspension containing 10⁵ CFU · ml⁻¹ of a *Trichoderma* isolate, 0.2% D-glucose, and
22 0.1% urea. The inoculated wood blocks were incubated at constant conditions in a 50 ml sterile
23 centrifuge tube for 4 weeks. Afterwards, the Senegal mahogany and rain tree wood blocks pre-treated
24 with *Trichoderma* were placed on a 14-day old pure culture of one *P. noxius* isolate growing on MEA
25 in a 50 mm Petri dish. Test plates were covered, sealed, and incubated under constant conditions for
26 12 weeks. Untreated, sterilized wood blocks were similarly placed on 14-day old actively growing
27 pure cultures of *P. noxius* and served as controls. After incubation, the wood blocks were cleaned by

1 removing surface mycelia, oven-dried, cooled, and reweighed. The dry-weight loss was then
2 determined for each wood block on a percentage basis.

3

4 *Statistical analysis*

5 For all experiments, 5 replicates were maintained of each treatment combination. Since germination
6 rates and chlamydospore counts were not measured under experimentally manipulated conditions, the
7 results were not statistically analyzed but instead presented as descriptive data. The effect of
8 experimental treatments on the measured response was determined using analysis of variance in SAS
9 9.4 (SAS Institute, Inc., Cary, NC, USA). For growth rate bioassays, model fixed effects included
10 media type, *Trichoderma* isolate, and their interaction. Since the rate of change in culture diameter
11 with respect to time was constant, growth rates were analyzed as the slope of a linear function (Gams
12 and Bissett, 1998) fit to paired observations of culture diameter and time using least squares
13 regression. For the remaining bioassays, model fixed effects included *Trichoderma* isolates, *P. noxius*
14 isolates, and their interaction. Given our primary interest in antagonism by *Trichoderma* isolates,
15 significant interactions were separated to determine differences among *Trichoderma* isolates within
16 specific levels of the other fixed effect. Mean separation was performed using Dunnett's one-way test
17 and Tukey's honestly significant difference for experiments with and without controls, respectively.

18

19 **Results**

20 Except for *T. reesei* 1056TD1, all *Trichoderma* isolates germinated, on average, at nearly maximum
21 rates between 99 and 100%. Still, light microscopy revealed that a considerable majority (87%) of *T.*
22 *reesei* 1056TD1 conidia germinated on LNA. In contrast, there was sizeable variability in the
23 production of chlamydospores among *Trichoderma* isolates. The average number of chlamydospores
24 observed among 5 randomly-selected visual fields at 10× magnification on each Petri dish ranged
25 between 6 (SD 2) for *T. saturnisporum* SPH20109132 and 189 (SD 30) for *T. virens* W23 (Table 2).

26

27 Overall, there were significant differences in growth rates among *Trichoderma* isolates ($F = 103.46$;
28 $df = 6, 24$; $p < 0.001$), and these growth rates differed significantly between the 2 media types ($F =$

1 107.83; $df = 1, 24$; $p < 0.001$). Specifically, *Trichoderma* isolates grew significantly faster on MEA
2 than on LNA (Table 3). However, *Trichoderma* isolates and media types interacted significantly to
3 affect growth rates ($F = 34.55$; $df = 6, 24$; $p < 0.001$). There were significant differences in growth
4 rates among *Trichoderma* isolates growing on both LNA and MEA, but the relative ranking of
5 *Trichoderma* isolates within each media type was not consistent (Table 3). Notably, the growth rate of
6 *T. virens* W23 was significantly slower than all other *Trichoderma* isolates, except *T. virens* W26, on
7 MEA, but its growth rate actually increased by 8.0% on LNA to a value not significantly different
8 from the 4 other fastest growing isolates. The growth rate of *T. harzianum* 9132 decreased marginally
9 by only 2.9% on LNA compared to MEA; the other *Trichoderma* isolates grew between 11.6 and
10 40.2% slower on LNA (Table 3).

11
12 Compared to the controls, all *P. noxius* cultures grew slower, in absolute terms, in the presence of
13 VOCs emitted by *Trichoderma*. Overall, growth rates of *P. noxius* varied significantly in the presence
14 of VOCs produced by *Trichoderma* isolates ($F = 5.40$; $df = 7, 28$; $p < 0.001$), and the growth of *P.*
15 *noxius* SSN3P cultures was inhibited significantly more by *Trichoderma* VOC emission than *P.*
16 *noxius* KSK6 ($F = 24.14$; $df = 1, 28$; $P = 0.008$). However, *Trichoderma* and *P. noxius* isolates
17 interacted significantly to affect growth rates because VOCs emitted by *Trichoderma* isolates did not
18 inhibit the 2 *P. noxius* isolates in a proportionally consistent way; *Trichoderma* isolates significantly
19 inhibiting one *P. noxius* isolate did not necessarily inhibit the other equally well. Compared to the
20 controls, 6 *Trichoderma* isolates caused a significant reduction to the growth rate of *P. noxius* SSN3P,
21 but only 4 *Trichoderma* isolates caused similar reductions to *P. noxius* KSK6 (Table 4). Separately, *T.*
22 *harzianum* 9132 and *T. virens* W23 caused the greatest inhibition to *P. noxius* KSK6 and *P. noxius*
23 SSN3P, respectively (Table 4).

24
25 In dual culture tests on MEA, most of the *Trichoderma* isolates exhibited comparably high
26 antagonistic potential with percent mortality to *P. noxius* ranging between 80 and 100%. All
27 *Trichoderma* isolates overspread and colonized the *P. noxius* cultures; none were deadlocked in
28 distant confrontation or evaded by the wood decay fungus. Overall, there were no significant

1 differences in the percent mortality of *P. noxius* challenged by various *Trichoderma* isolates in dual
2 culture tests ($F = 0.98$; $df = 6, 12$; $p = 0.479$), and there were no significant differences in the
3 susceptibility of the 2 *P. noxius* isolates to antagonism by *Trichoderma* ($F = 0.61$; $df = 1, 12$; $p =$
4 0.517). However, *Trichoderma* and *P. noxius* isolates interacted significantly to affect percent
5 mortality ($F = 3.25$; $df = 6, 12$; $p = 0.039$). Although the lethal effect of various *Trichoderma* isolates
6 towards *P. noxius* SSN3P in dual culture did not vary significantly, the same was not true for *P.*
7 *noxius* KSK6. Specifically, the percent mortality of *P. noxius* KSK6 confronted by *T. harzianum* 9132
8 was significantly less than that caused by the other *Trichoderma* isolates (Table 5).

9
10 Overall, dry weight loss of wood blocks varied significantly according to their pre-treatment with
11 various *Trichoderma* isolates ($F = 7.65$; $df = 7, 28$; $p < 0.001$), and pre-treatment with *Trichoderma*
12 isolates gave significantly better protection against decomposition by *P. noxius* SSN3P than *P. noxius*
13 KSK6 ($F = 20.52$; $df = 1, 28$; $p = 0.011$). The interaction between *Trichoderma* and *P. noxius* isolates,
14 however, was not significant ($F = 1.11$; $df = 7, 28$; $p = 0.386$) because *Trichoderma* isolates inhibited
15 decomposition by the 2 *P. noxius* isolates similarly. Compared to the controls, pre-treatment with 6
16 different *Trichoderma* isolates caused a significant reduction to the dry weight loss of wood blocks
17 inoculated with *P. noxius*. Among all isolates, *T. harzianum* 9132 offered the greatest protection to
18 wood blocks against decomposition by *P. noxius*, and *T. virens* W26 was the only isolate that did not
19 significantly inhibit wood block decomposition compared to the controls (Table 6).

20

21 **Discussion**

22 Despite apparent variability, all of the *Trichoderma* isolates tested in this study antagonized both *P.*
23 *noxius* isolates in the different bioassays. In contrast with similar studies (Mercer and Kirk, 1984a;
24 Ribera et al., 2016; Schubert et al., 2008a; Schwarze et al., 2012), none of the *Trichoderma* isolates
25 enhanced the activity of *P. noxius* isolates in culture. Still, no single *Trichoderma* isolate was
26 consistently superior among all others in its ability to antagonize *P. noxius* across every bioassay. In
27 addition, the presence of *Trichoderma* caused a greater reduction to the growth (Table 4) and wood
28 decay rates (Table 6) of *P. noxius* SSN3P than *P. noxius* KSK6, indicating the greater susceptibility of

1 the former to biological control by *Trichoderma*. Although the wood block interaction tests were
 2 performed on different wood substrates corresponding to the tree species from which each *P. noxius*
 3 isolate was originally obtained (Table 1), the equally greater antagonistic effect of *Trichoderma* VOC
 4 emission on *P. noxius* SSN3P suggests that the disparity is not entirely caused by differences in test
 5 conditions associated with wood anatomy. In agreement with existing reports (Schwarze et al., 2012;
 6 Wells and Bell, 1979), this intraspecific variability and target specificity affirms the importance of
 7 screening potential biological control agents against the desired pathogen in laboratory tests.

8
 9 Germination rates were nearly complete for the conidia of *Trichoderma* isolates cultured on LNA
 10 (Table 2). Although exogenous nutrients are required for germination, existing reports indicate that
 11 leached nutrients and secondary metabolites can provide adequate resources to support germination
 12 (Danielson and Davey, 1973; Naar and Kecskes, 1998). Since conidia were unwashed, germination
 13 rates reported in this study may not reflect those for *Trichoderma* in sterile environments. Practically,
 14 these results suggest that conidial preparations for biological control can be amended with modest
 15 supplemental nutrients to support high germination rates. Analogously, Schubert et al. (2008a)
 16 reported that an unmodified conidial suspension and another amended with glucose and urea persisted
 17 equally well on tree wounds over a 30-month period. In addition, high germination rates for
 18 introduced conidia valuably limits the availability of dead propagules as a nutrient source to other
 19 deleterious microbes that might interfere with biological control by competing with *Trichoderma* or
 20 infecting the plant host (Harman et al., 1991).

21
 22 In contrast, however, there was considerable variability in the production of chlamydospores among
 23 *Trichoderma* isolates (Table 2). Relatively few reports exist on *Trichoderma* chlamydospore biology;
 24 Lewis and Papavizas (1984) reported that their production rates similarly varied among *Trichoderma*
 25 isolates. Although it is generally accepted that these resting spores enhance survival during
 26 unfavorable conditions in the soil, it is not clear to what extent they might serve a similar purpose on
 27 other substrates, such as wood. Practically, chlamydospores are valuable for commercial *Trichoderma*

1 formulations since these propagules better tolerate desiccation and storage than conidia or mycelial
2 fragments (Li et al., 2016).
3
4 Growth rates varied significantly among *Trichoderma* isolates, but all isolates grew at faster rates than
5 reported for others selected from temperate or sub-tropical climates for similar biological control
6 applications (Ribera et al., 2016; Schubert et al., 2008a; Schwarze et al., 2012). In agreement with
7 existing work (Schubert et al., 2008b), growth rates decreased for isolates cultured under nutrient
8 limiting conditions on LNA. However, the increased growth of *T. virens* W23 under these conditions
9 is a notable exception (Table 3). Although difficult to demonstrate experimentally (Harman, 2000),
10 competition for space and nutrients is likely an important mechanism for biological control by
11 *Trichoderma*.
12
13 It is useful to note that germination and growth rates were determined under constant environmental
14 conditions chosen to replicate typical values for Singapore. At this location, the climate is
15 representative of the equatorial tropics with stable temperatures ranging daily between 24° and 32° C,
16 abundant rainfall approaching 250 cm annually, and elevated humidity between 60 and 90%
17 (Micheline and Ng, 2012). In general, the optimum temperature for the growth of many *Trichoderma*
18 spp. is between 25° and 30° C (Klein and Eveleigh, 1998). There is considerable experimental
19 evidence that *Trichoderma* germination and growth rates are proportional to water activity and
20 temperature (Kredics et al., 2003), up to some limiting value that depends on the climatic origin of a
21 given *Trichoderma* isolate (Ribera et al., 2016; Schubert et al., 2008b; Schwarze et al., 2012); but it
22 was assumed that tolerance to a wide range of unfavorable temperature and moisture conditions is not
23 critical for biological control by *Trichoderma* in Singapore.
24
25 Compared to similar work on wood decay fungi (Mercer and Kirk, 1984a; Schubert et al., 2008b),
26 including *P. noxius* (Ribera et al., 2016; Schwarze et al., 2012), VOC emission by *Trichoderma*
27 isolates had a comparatively large effect on the growth of *P. noxius* cultures in this study. Ribera et al.
28 (2016) reported that less than half of all tested *Trichoderma* isolates inhibited the radial growth rates

1 of *P. noxius* isolates associated with landscape trees in Hong Kong. Similarly, Schwarze et al. (2012)
 2 reported that only one out of 16 possible combinations of *P. noxius* and *Trichoderma* isolates resulted
 3 in a significant decrease to the radial growth rate of the wood decay fungus compared to their
 4 respective controls. Based on similar observations, Schubert et al. (2008b) postulated that antibiosis
 5 might not be an important mechanism by which *Trichoderma* antagonizes wood decay fungi, but
 6 these results provide contrasting evidence that VOCs can significantly inhibit the growth of *P. noxius*
 7 cultures. Notably, *T. harzianum* 9132 and *T. virens* W23 caused a 68.6% and 81.3% reduction to the
 8 growth rates, respectively, of *P. noxius* KSK6 and *P. noxius* SSN3P.

9

10 The lethal effect of *Trichoderma* on *P. noxius* in dual culture tests was consistently high, except for
 11 the interaction between *P. noxius* KSK6 and *T. harzianum* 9132 (Table 5). The uniformly high lethal
 12 effect might be partially explained by the favorable habitability of MEA; other authors have reported
 13 that inhibition of wood decay fungi by *Trichoderma* in dual culture tests was significantly greater on
 14 MEA than LNA (Schubert et al., 2008b; Srinivasan et al., 1992). Specifically, Srinivasan et al. (1992)
 15 reported that soluble metabolites and volatile antibiotics were produced in significantly greater
 16 quantities by *Trichoderma* growing on MEA. As a result, the results from this test preclude
 17 discrimination among candidate *Trichoderma* isolates for biological control, but there was much
 18 greater variability in the protection afforded by various *Trichoderma* isolates to wood blocks against
 19 decay by *P. noxius*.

20

21 In this study, almost all *Trichoderma* isolates significantly inhibited wood decay by *P. noxius*
 22 compared to the controls (Table 6). Although the *Trichoderma* isolates failed to completely prevent
 23 wood decay, the sizeable reduction is in broad agreement with existing studies (Ribera et al., 2016;
 24 Schubert et al., 2008a; Schubert et al., 2008b; Schwarze et al., 2012). There are 2 possible
 25 explanations for the incomplete reduction of dry weight loss to wood blocks by *Trichoderma*. First,
 26 the wood blocks were inoculated by introducing an artificially large amount of *P. noxius* inoculum,
 27 and the disproportionate size of *P. noxius* and *Trichoderma* cultures may have favored the former.
 28 Second, a fraction of the total dry weight loss to wood blocks was probably caused by the

1 decomposition of non-structural carbohydrates by *Trichoderma* (Klein and Eveleigh, 1998). Still, the
2 significant reductions to the dry weight loss of wood blocks shows the strong antagonistic potential of
3 several *Trichoderma* isolates, especially *T. harzianum* 9132 (Table 6).

4

5 After incubation for 12 weeks, the percent dry weight loss to control wood blocks inoculated with *P.*
6 *noxius* KSK6 and SSN3P was much greater than reported for balsa [*Ochroma pyramidale* (Cav. ex
7 Lam.) Urb. (Malvaceae)] wood blocks inoculated with different *P. noxius* isolates obtained from
8 landscape trees in Hong Kong (Ribera et al., 2016), but the same values in this study are more
9 comparable to dry weight losses reported for wood blocks removed from several landscape tree
10 species associated with *P. noxius* in Brisbane, Australia (Schwarze et al., 2012). Collectively, these
11 results demonstrate the importance of using species-specific wood substrates to represent the host-
12 fungus interaction in which *Trichoderma* must intercede as a biological control agent, and this
13 distinction gives confidence to the observations made on Senegal mahogany and rain tree wood
14 blocks in this study.

15

16 In this study, the systematic investigation of 7 *Trichoderma* isolates demonstrated their promising
17 antagonism towards *P. noxius*. Although the 2 *P. noxius* isolates differed in their susceptibility,
18 several *Trichoderma* isolates showed a significant ability to inhibit growth and reduce decomposition
19 by this aggressive wood decay fungus. None of the *Trichoderma* isolates increased the activity of *P.*
20 *noxius*, and the deliberate collection of *Trichoderma* isolates from an ecological niche that resembled
21 the desired target specificity, i.e., from basidiomycete fruiting bodies, may have been an effective
22 selection strategy for biological control agents. In agreement with existing reports, there was no clear
23 evidence of a primary mechanism by which *Trichoderma* antagonized *P. noxius*, and the results from
24 different bioassays should be given equal weight in the selection of *Trichoderma* isolates. Although
25 no single *Trichoderma* isolate outperformed all others in every test, there were a few that did so more
26 than others. In particular, *T. virens* W23 produced the most chlamydospores, best inhibited the growth
27 of *P. noxius* SSN3P, and uniquely grew faster on LNA. On the other hand, *T. harzianum* 9132 best
28 inhibited the growth of *P. noxius* KSK6 and prevented dry weight loss to wood blocks. Still, these

1 results are only indicative of the ability of *Trichoderma* to antagonize *P. noxius* on tree wounds, and
2 these competitive *Trichoderma* isolates should be further examined under field conditions to confirm
3 their efficacy. Laboratory conditions were more favorable and simplified than those governing host-
4 fungus interaction in dynamic natural systems.

6 **Acknowledgments**

7 The Agri-Food & Veterinary Authority and National Parks Board of Singapore funded this research.
8 The contribution of laboratory assistance by Yushiella binte Ahmad, Choy May Yee, and Doris Tee
9 Hui Ying is gratefully acknowledged.

11 **Bibliography**

- 12 Anonymous. American National Standard for Tree Care Operations - Tree, Shrub, and Other Woody
13 Plant Management - Standard Practices (Pruning), 2008 Tree Care Industry Association, Inc,
14 Londonderry, NH, USA.
- 15 Barry, K.M., Pearce, R.B., Mohammed, C.L., 2000. Properties of reaction zones associated with
16 decay from pruning wounds in plantation-grown *Eucalyptus nitens*. Forest Pathol. 30, 233-245.
- 17 Benson, D.A., Boguski, M.S., Lipman, D.J., Ostell, J., 1997. GenBank. Nucleic Acids Res. 25, 1-6.
- 18 Bigirimana, J., de Meyer, G., Poppe, J., Elad, Y., Hoefte, M., 1997. Induction of systemic resistance
19 on bean (*Phaseolus vulgaris*) by *Trichoderma harzianum*. Int S Crop. 62, 1001-1008.
- 20 Brunner, K., Zeilinger, S., Ciliento, R., Woo, S.L., Lorito, M., Kubicek, C.P., et al., 2005.
21 Improvement of the fungal biocontrol agent *Trichoderma atroviride* to enhance both antagonism and
22 induction of plant systemic disease resistance. Appl Environ Microb. 71, 3959-3965.
- 23 Burcham, D.C., Wong, J.Y., Ali, M.I.M., Abarrientos, N.V., Fong, Y.K., Schwarze, F.W.M.R., 2015.
24 Characterization of host-fungus interactions among wood decay fungi associated with *Khaya*
25 *senegalensis* (Desr.) A. Juss (Meliaceae) in Singapore. Forest Pathol. 45, 492-504.
- 26 Chang, T.-., 1995. Decline of nine tree species associated with brown root rot caused by *Phellinus*
27 *noxius*. Plant Dis. 79, 962-965.
- 28 Danielson, R.M., Davey, C.B., 1973. Effects of nutrients and acidity on phialospore germination of
29 *Trichoderma* in vitro. Soil Biol Biochem. 5, 517-524.
- 30 Deflorio, G., Barry, K.M., Johnson, C., Mohammed, C.L., 2007. The influence of wound location on
31 decay extent in plantation-grown *Eucalyptus globulus* and *Eucalyptus nitens*. Forest Ecol Manag. 242,
32 353-362.
- 33 Dennis, C., Webster, J., 1971. Antagonistic properties of species-groups of *Trichoderma* II.
34 Production of volatile antibiotics. T Brit Mycol Soc. 57, 41-48.

- 1 Di Pietro, A., Lorito, M., Hayes, C.K., Broadway, R.M., Harman, G.E., 1993. Endochitinase from
2 *Gliocladium virens*: Isolation, characterization, and synergistic antifungal activity in combination with
3 gliotoxin. Mol Plant Pathol. 83, 308-313.
- 4 Djonovic, S., Pozo, M.J., Dangott, L.J., Howell, C.R., Kenerley, C.M., 2006. Sm1, a proteinaceous
5 elicitor secreted by the biocontrol fungus *Trichoderma virens* induces plant defense responses and
6 systemic resistance. Mol Plant Microbe In. 19, 838-853.
- 7 Djonovic, S., Vittone, G., Mendoza-Herrera, A., Kenerley, C.M., 2007. Enhanced biocontrol activity
8 of *Trichoderma virens* transformants constitutively coexpressing β -1,3- and β -1,6-glucanase genes.
9 Mol Plant Pathol. 8, 469-480.
- 10 Gams, W., Bissett, J., 1998. Morphology and identification of *Trichoderma*, in: Kubicek, C.P.,
11 Harman, G.E. (Eds.), *Trichoderma and Gliocladium* Volume 1. Taylor & Francis, London, England,
12 UK, pp. 3-31.
- 13 Ghisalberti, E.L., Sivasithamparam, K., 1991. Antifungal antibiotics produced by *Trichoderma* spp.
14 Soil Biol Biochem. 23, 1011-1020.
- 15 Harman, G.E., 2000. Myths and dogmas of biocontrol: Changes in perceptions derived from research
16 on *Trichoderma harzianum* T-22. Plant Dis. 84, 377-393.
- 17 Harman, G.E., Howell, C.R., Viterbo, A., Chet, I., Lorito, M., 2004. *Trichoderma* species-
18 opportunistic, avirulent plant symbionts. Nat Rev Microbiol. 2, 43-56.
- 19 Harman, G.E., Jin, X., Stasz, T.E., Peruzzotti, G., Leopold, A.C., Taylor, A.G., 1991. Production of
20 conidial biomass of *Trichoderma harzianum* for biological control. Biol Control. 1, 23-28.
- 21 Highley, T.L., Padmanabha, H.S.A., Howell, C.R., 1997. Control of wood decay by *Trichoderma*
22 (*Gliocladium*) *virens* II. Antibiosis. Mater Organismen. 31, 157-166.
- 23 Howell, C.R., 2003. Mechanisms employed by *Trichoderma* species in the biological control of plant
24 diseases: The history and evolution of current concepts. Plant Dis. 87, 4-10.
- 25 Huttermann, A., Volger, C., 1973. Induction of aryl- β -glucosidase in *Fomes annosus* by cellobiose.
26 Archiv Mikrobiol. 93, 195-204.
- 27 Kallio, T., Hallaksela, A.M., 1979. Biological control of *Heterobasidion annosum* (Fr.) Bref. (*Fomes*
28 *annosus*) in Finland. Eur J Forest Pathol. 9, 298-308.
- 29 Klein, D., Eveleigh, D.E., 1998. Ecology of *Trichoderma*, in: Kubicek, C.P., Harman, G.E. (Eds.),
30 *Trichoderma and Gliocladium* Volume 1. Taylor & Francis, London, England, UK, pp. 57-74.
- 31 Koike, N., Kyakumachi, M., Kageyama, K., Tsuyumu, S., Doke, N., 2001. Induction of systemic
32 resistance in cucumber against several diseases by plant growth-promoting fungi: Lignification and
33 superoxide generation. Eur J Plant Pathol. 107, 523-533.
- 34 Kredics, L., Antal, Z., Manczinger, L., Szekeres, A., Kevei, F., Nagy, E., 2003. Influence of
35 environmental parameters on *Trichoderma* strains with biocontrol potential. Food Technol
36 Biotechnol. 41, 37-42.

- 1 Kuhls, K., Lieckfeldt, E., Samuels, G.J., Kovacs, W., Meyer, W., Petrini, O., et al., 1996. Molecular
2 evidence that the asexual industrial fungus *Trichoderma reesei* is a clonal derivative of the
3 ascomycete *Hypocrea jecorina*. Proc Natl Acad USA. 93, 7755-7760.
- 4 Kuhls, K., Lieckfeldt, E., Samuels, G.J., Meyer, W., Kubicek, C.P., Borner, T., 1997. Revision of
5 *Trichoderma* sect. *Longibrachiatum* including related teleomorphs based on analysis of ribosomal
6 DNA internal transcribed spacer sequences. Mycologia. 89, 442-460.
- 7 Lewis, J.A., Papavizas, G.C., 1984. Chlamydospore formation by *Trichoderma* spp. in natural
8 substrates. Can J Microbiol. 30, 1-7.
- 9 Li, Y.Q., Song, K., Li, Y.C., Chen, J., 2016. Statistical culture-based strategies to enhance
10 chlamydospore production by *Trichoderma harzianum* SH2303 in liquid fermentation. J Zhejiang
11 Univ-Sc B. 17, 619-627.
- 12 Lieckfeldt, E., Kullnig, C.M., Kubicek, C.P., Samuels, G.J., Borner, T., 2001. *Trichoderma*
13 *aureoviride*: Phylogenetic position and characterization. Mycol Res. 105, 313-322.
- 14 Lieckfeldt, E., Samuels, G.J., Nirenberg, H.I., Petrini, O., 1999. A morphological and molecular
15 perspective of *Trichoderma viride*: Is it one or two species?. Appl Env Microbiol. 65, 2418-2428.
- 16 Lonsdale, D., 1984. Available treatments for tree wounds: An assessment of their value. Arboric J. 8,
17 99-107.
- 18 Lorito, M., Farkas, V., Rebuffat, S., Bodo, B., Kubicek, C.P., 1996a. Cell wall synthesis is a major
19 target of mycoparasitic antagonism by *Trichoderma harzianum*. J Bacteriol. 178, 6382-6385.
- 20 Lorito, M., Hayes, C.K., Di Pietro, A., Woo, S.L., Harman, G.E., 1994a. Purification,
21 characterization, and synergistic activity of a glucan 1,3- β -glucosidase and an *N*-acetyl- β -
22 glucosaminidase from *Trichoderma harzianum*. Phytopathology. 84, 398-405.
- 23 Lorito, M., Peterbauer, C., Hayes, C.K., Harman, G.E., 1994b. Synergistic interaction between fungal
24 cell wall degrading enzymes and different antifungal compounds enhances inhibition of spore
25 germination. Microbiology. 140, 623-629.
- 26 Lorito, M., Woo, S.L., D'Ambrosio, M., Harman, G.E., Hayes, C.K., Kubicek, C.P., et al., 1996b.
27 Synergistic interaction between cell wall degrading enzymes and membrane affecting compounds.
28 Mol Plant Microbe In. 9, 206-213.
- 29 Markovich, N.A., Kononova, G.L., 2003. Lytic enzymes of *Trichoderma* and their role in plant
30 defense from fungal diseases: A review. Appl Biochem Microbiol. 39, 341-351.
- 31 Mercer, P.C., Kirk, S.A., Gendle, P., Clifford, D.R., 1983. Chemical treatments for control of decay in
32 pruning wounds. Ann Appl Biol. 102, 435-453.
- 33 Mercer, P.C., 1983. Callus growth and the effect of wound dressings. Ann Appl Biol. 103, 527-540.
- 34 Mercer, P.C., 1979. Phytotoxicity and fungitoxicity tests for tree wound paints. Ann Appl Biol. 91,
35 199-202.
- 36 Mercer, P.C., Kirk, S.A., 1984a. Biological treatments for the control of decay in tree wounds I.
37 Laboratory tests. Ann Appl Biol. 104, 211-219.

- 1 Mercer, P.C., Kirk, S.A., 1984b. Biological treatments for the control of decay in tree wounds II.
2 Field tests. *Ann Appl Biol.* 104, 221-229.
- 3 Micheline, F., Ng, L.K., 2012. The Weather and Climate of Singapore, Meteorological Service
4 Singapore, National Environment Agency, Singapore.
- 5 Naar, Z., Kecskes, M., 1998. Factors influencing the competitive saprophytic ability of *Trichoderma*
6 species. *Microbiol Res.* 153, 119-129.
- 7 Nicole, M., Chamberland, H., Rioux, D., Xixuan, X., Blanchette, R.A., Geiger, J.P., et al., 1995.
8 Wood degradation by *Phellinus noxius*: Ultrastructure and cytochemistry. *Can J Microbiol.* 41, 253-
9 265.
- 10 Ottenheim, C., Meier, K., Zimmermann, W., Wu, J.C., 2015. Isolation of filamentous fungi exhibiting
11 high endoxylanase activity in lignocellulose hydrolysate. *Appl Biochem Biotech.* 175, 2066-2074.
- 12 Papavizas, G.C., 1985. *Trichoderma* and *Gliocladium*: Biology, ecology, and potential for biocontrol.
13 *Annu Rev Phytopathol.* 23, 23-54.
- 14 Pieterse, C.M.J., van Loon, L.C., 1999. Salicylic acid-independent plant defence pathways. *Trends*
15 *Plant Sci.* 4, 52-58.
- 16 Pottle, H.W., Shigo, A.L., 1975. Treatment of wounds on *Acer rubrum* with *Trichoderma viride*. *Eur J*
17 *Forest Pathol.* 5, 274-279.
- 18 Ribera, J., Tang, A.M.C., Schubert, M., Lam, R.Y.C., Chu, L.M., Leung, M.W.K., et al., 2016. In-
19 vitro evaluation of antagonistic *Trichoderma* strains for eradicating *Phellinus noxius* in colonised
20 wood. *J Trop For Sci.* 28, 457-468.
- 21 Ricard, J.L., Highley, T.L., 1988. Biocontrol of pathogenic fungi in wood and trees, with particular
22 emphasis on the use of *Trichoderma*. *Biocontrol News and Information*, 133-142.
- 23 Rossman, A.Y., Farr, D.F., 2014. *Fungal Databases*. 2014.
- 24 Samuels, G.J., 1996. *Trichoderma*: A review of biology and systematics of the genus. *Mycol Res.*
25 100, 923-935.
- 26 Schubert, M., Fink, S., Schwarze, F.W.M.R., 2008a. Evaluation of *Trichoderma* spp. as a biocontrol
27 agent against wood decay fungi in urban trees. *Biol Control.* 45, 111-123.
- 28 Schubert, M., Fink, S., Schwarze, F.W.M.R., 2008b. In vitro screening of an antagonistic
29 *Trichoderma* strain against wood decay fungi. *Arboric J.* 31, 227-248.
- 30 Schwarze, F.W.M.R., Jauss, F., Spencer, C., Hallam, C., Schubert, M., 2012. Evaluation of an
31 antagonistic *Trichoderma* strain for reducing the rate of wood decomposition by the white rot fungus
32 *Phellinus noxius*. *Biol Control.* 61, 160-168.
- 33 Sieber, T.N., 1995. *Pyrenochaeta ligni-putridi* sp. nov., a new coelomycete associated with butt rot of
34 *Picea abies* in Switzerland. *Mycol Res.* 99, 274-276.
- 35 Sivan, A., Chet, I., 1989. The possible role of competition between *Trichoderma harzianum* and
36 *Fusarium oxysporum* on rhizosphere colonization. *Phytopathology.* 79, 198-203.

- 1 Srinivasan, U., Staines, H.J., Bruce, A., 1992. Influence of media type on antagonistic modes of
2 *Trichoderma* spp. against wood decay basidiomycetes. Mater Organismen. 21, 302-321.
- 3 Wells, H.D., Bell, D.K., 1979. Variable antagonistic reaction in vitro of *Trichoderma harzianum*
4 against several pathogens. Phytopathology. 69, 1048-1049.
- 5 Yedidia, I., Benahmou, N., Kapulnik, Y., Chet, I., 2000. Induction and accumulation of PR proteins
6 activity during early stages of root colonization by the mycoparasite *Trichoderma harzianum* strain T-
7 203. Plant Physiol Bioch. 38, 863-873.
- 8 Yedidia, I., Benhamou, N., Chet, I., 1999. Induction of defense responses in cucumber plants
9 (*Cucumis sativus* L.) by the biocontrol agent *Trichoderma harzianum*. Appl Environ Microb. 65,
10 1061-1070.

11

Table 1: Identity, origin, and accession numbers of the fungal isolates collected for use in this study

Species	Isolate	Accession No.	Substrate	Origin	Reference	Identity (%)
a. Wood decay fungi						
<i>P. noxius</i>	KSK6	KF233592	ex. <i>Khaya senegalensis</i>	Bugis, Singapore	FR821770 (Schwarze et al., 2012)	99
<i>P. noxius</i>	SSN3P	KP202302	ex. <i>Samanea saman</i>	Clementi, Singapore	FR821769 (Schwarze et al., 2012)	98
b. <i>Trichoderma</i> spp.						
<i>T. reesei</i>	1056TD1	KY025557	<i>Ganoderma boninense</i> on <i>Cocos nucifera</i>	Singapore	JQ979435.1 (Kuhls et al., 1996)	99
<i>T. asperellum</i>	3	KY025555	<i>Ganoderma</i> sp. on coarse woody debris	Bukit Batok, Singapore	AJ230669.1 (Lieckfeldt et al., 1999)	100
<i>T. harzianum</i>	9132	KY025556	<i>Ganoderma boninense</i> on <i>Cyrtostachys renda</i>	Singapore	KR856210.1 (Lieckfeldt et al., 2001)	99
<i>T. saturnisporum</i>	SPH20109132	KY025558	<i>Ganoderma boninense</i> on <i>Cyrtostachys renda</i>	Singapore	NR_103704.1 (Kuhls et al., 1997)	99
<i>T. harzianum</i>	W21	KY025559	<i>Ganoderma boninense</i> on <i>Ptychosperma macarthurii</i>	Woodlands, Singapore	KJ000324.1 (Lieckfeldt et al., 2001)	99
<i>T. virens</i>	W23	KY025560	<i>Ganoderma boninense</i> on <i>Ptychosperma macarthurii</i>	Woodlands, Singapore	KP009289.1 (Ottenheim et al., 2015)	100
<i>T. virens</i>	W26	KY025561	<i>Ganoderma boninense</i> on <i>Ptychosperma macarthurii</i>	Woodlands, Singapore	KR296867.1 (Ottenheim et al., 2015)	100

Note: Fungal ITS sequences were deposited in GenBank (Benson et al., 1997).

Table 2: Germination rates (%) and chlamydospore counts (*n*) for various *Trichoderma* spp. isolates cultured on a low nutrient agar (LNA)

<i>Trichoderma</i> spp.	Germination rate (%)	Chlamydospore count (<i>n</i>)
1056TD1	87 (21)	9 (4)
3	100 (0)	20 (15)
9132	100 (0)	20 (13)
SPH20109132	100 (0)	6 (2)
W21	100 (0)	39 (8)
W23	99 (1)	189 (30)
W26	99 (1)	165 (44)

Note: Values listed are mean (SD). See Table 1 for the taxonomic identity of fungal isolates.

Table 3: Growth rates (cm·day⁻¹) of *Trichoderma* spp. isolates cultured on 2 different media, including a malt extract agar (MEA) and low nutrient agar (LNA)

<i>Trichoderma</i> spp.	Media type	
	MEA	LNA
1056TD1	7.96 (0.11) a	4.76 (0.04) a
3	4.61 (0.01) c	3.12 (0.27) c
9132	4.74 (0.02) c	4.60 (0.02) ab
SPH20109132	8.04 (0.10) a	5.28 (0.48) a
W21	5.07 (0.11) b	4.48 (0.76) ab
W23	4.38 (0.10) d	4.73 (0.07) a
W26	4.41 (0.08) d	3.62 (0.97) bc
Mean	5.61 (1.56)	4.37 (0.84)

Note: Values listed are mean (SD). Means followed by the same letter within each column are not significantly different at the $\alpha = 0.05$ level. See Table 1 for the taxonomic identity of fungal isolates.

Table 4: *Phellinus noxius* KSK6 and *P. noxius* SSN3P growth rates (cm·day⁻¹) in the presence of VOCs emitted by various *Trichoderma* spp. isolates

<i>Trichoderma</i> spp.	<i>P. noxius</i>	
	KSK6	SSN3P
Control	1.75 (0.01)	1.34 (0.08)
1056TD1	1.48 (0.19)	0.42 (0.47)*
3	1.46 (0.09)	0.68 (0.49)*
9132	0.55 (0.45)*	0.69 (0.29)*
SPH20109132	0.90 (0.32)*	0.86 (0.21)*
W21	1.18 (0.36)*	1.04 (0.25)
W23	1.52 (0.23)	0.25 (0.33)*
W26	0.94 (0.59)*	0.39 (0.55)*
Mean	1.22 (0.48)	0.71 (0.48)

Note: Values listed are mean (SD). Within each column, asterisks (*) indicate a significant reduction in growth rates compared to the control at the $\alpha = 0.05$ level. See Table 1 for the taxonomic identity of fungal isolates.

Table 5: Percent mortality (%) to *Phellinus noxius* KSK6 and *P. noxius* SSN3P due to mycoparasitism by various *Trichoderma* spp. isolates in dual culture

<i>Trichoderma</i> spp.	<i>P. noxius</i>	
	KSK6	SSN3P
1056TD1	89 (19) a	89 (10)
3	83 (17) a	94 (10)
9132	33 (58) b	100 (0)
SPH20109132	94 (10) a	89 (10)
W21	78 (10) a	94 (10)
W23	89 (10) a	67 (58)
W26	100 (0) a	100 (0)
Mean	81 (29)	90 (22)

Note: Values listed are mean (SD). Within left-hand column, means followed by the same letter are not significantly different at the $\alpha = 0.05$ level. See Table 1 for the taxonomic identity of fungal isolates.

Table 6: Dry weight loss (%) caused to Senegal mahogany (*Khaya senegalensis*) and rain tree (*Samanea saman*) wood blocks by *Phellinus noxius* KSK6 and *P. noxius* SSN3P, respectively, after pre-treatment with various *Trichoderma* spp. isolates

<i>Trichoderma</i> spp.	<i>P. noxius</i>		
	KSK6	SSN3P	Mean
Control	12.7 (4.4)	10.8 (5.0)	11.7
1056TD1	8.0 (2.6)	3.6 (5.0)	5.8*
3	4.6 (0.5)	2.8 (0.8)	3.7*
9132	5.0 (0.8)	1.8 (1.3)	3.4*
SPH20109132	7.7 (2.4)	4.0 (4.2)	5.7*
W21	8.7 (0.8)	3.6 (2.9)	6.2*
W23	7.2 (1.2)	5.8 (4.2)	6.5*
W26	8.2 (2.1)	8.6 (1.7)	8.4
Mean	7.8 (3.1) a	5.1 (4.3) b	

Note: Values listed are mean (SD). Within right-hand column, asterisks (*) indicate a significant reduction in dry weight loss compared to the control at the $\alpha = 0.05$ level. Means followed by the same letter within the bottom row are not significantly different at the $\alpha = 0.05$ level. See Table 1 for the taxonomic identity of fungal isolates.

Figure 1: In one dual culture test, *T. harzianum* W21 (bottom) overgrew *P. noxius* KSK6 (top) during confrontation. The distribution of *T. harzianum* W21 over the wood decay fungus is evident by its green conidia, likely formed in response to its antagonistic interaction with *P. noxius* KSK6.