A genetic-based algorithm for recovery: A pilot study.

Craig Pickering^{1,2}, John Kiely¹, Bruce Suraci³, Charlie Quigley³, Jake Watson⁴.

Institutional Affiliations:

1. Institute of Coaching and Performance, School of Sport and Wellbeing, University of Central Lancashire, Preston, UK
2. Exercise and Nutritional Genomics Research Centre, DNAFit Ltd, London, UK
3. Suraci Consultancy, Portsmouth, UK
4. University of Portsmouth, University House, Winston Churchill Avenue, Portsmouth, UK.

Abstract

Exercise training creates a number of physical challenges to the body, the overcoming of which drives exercise adaptation. The balance between sufficient stress and recovery is a crucial, but often under-explored, area within exercise training. Genetic variation can also predispose some individuals to a greater need for recovery after exercise. In this pilot study, 18 male soccer players underwent a repeated sprint training session. Countermovement jump (CMJ) heights were recorded immediately pre- and post-training, and at 24- and 48-hours post-training. The reduction in CMJ height was greatest at all post-training time points in subjects with a larger number of gene variants associated with a reduced exercise recovery. This suggests that knowledge of genetic information can be important in individualizing recovery timings and modalities in athletes following training.

Introduction

Exercise training produces a variety of acute physiological challenges to the body, perturbing homeostasis and inducing a stress response, the overcoming of which leads to exercise adaptation. Successful adaptation is comprised of the accumulation of periods of stress-induced response (training), and periods of recovery, which takes place away from exercise (Bishop et al., 2008). This relationship is finely balanced, and if there is insufficient time between training sessions for recovery to occur, the athlete increases their risk of undue accumulation of fatigue, potentially resulting in acute underperformance, injury (Mair et al., 1996), illness (Schwellnus et al., 2016), and eventually non-functional overreaching and unexplained underperformance syndrome (UPS) (Soligard et al., 2016). These indicators of maladaptation are common in athletes, with 10-20% of endurance athletes suffering from UPS each year (Budgett, 2000). Overuse injury incidence is also frequent, with rates between 37% and 85% reported, depending on the sport (DiFiori et al., 2013; Wilber et al., 1995). The prevalence of these symptoms of stress-recovery imbalance at epidemic proportions indicates that there is a mismatch in our knowledge of how to create stress and how to recover from it. Indeed, a search on PubMed yields over 37,000 papers with “exercise training” in the title and abstract. A similar result with “exercise recovery” as the search field only results in 13,000 papers.

At the cellular level, the physiological challenges induced by exercise include increased oxidative metabolism within the mitochondria, leading to increased formation of reactive oxygen species (ROS) (Fisher-Wellman & Bloomer, 2009). Under normal, non-exercise conditions, the body can neutralise ROS through its endogenous antioxidant defence system, which is comprised of enzymes such as superoxide dismutase (Belviranli & Gokbel,

2006). However, when ROS production is increased through exercise, an imbalance between ROS production and neutralisation occurs, leading to elevated oxidative stress (Fisher-Wellman & Bloomer, 2009). In turn, this elevates lipid peroxidation and tissue damage. Mechanical load also increases muscle damage (Baumert et al., 2016), initiating an inflammatory response partially driven by pro-inflammatory cytokines such as interleukin-6 (IL-6) and tumour necrosis factor (TNF) (Yamin et al., 2008).

These changes at the molecular and cellular level drive the whole-body symptoms of under-recovery that athletes and coaches are well aware of. Increases in plasma IL-6 levels occur following exercise (Robson-Ansley et al., 2007), and administration of exogenous IL-6 into athletes induces feelings of fatigue and impairs performance (Robson-Ansley et al., 2004). Increased IL-6 is also a risk factor for the development of UPS (Robson, 2003). Both TNF and IL-6 can act on the central nervous system (Ament & Verkerke, 2009), potentially decreasing the drive to exercise. Increased ROS and oxidative stress are also associated with a decrease in physical performance (Powers & Jackson, 2007), and increased feelings of muscle soreness following exercise (Konig et al., 2001).

Recently, a number of best practices for recovery have been put forward (Soligard et al., 2016; Schwellnus et al., 2016; Leeder et al., 2011; Bishop et al., 2008). However, there is likely considerable inter-individual variation in the time course of exercise recovery (Nosaka et al., 1996). This is partially governed by genetic variation between individuals, with several single nucleotide polymorphisms (SNPs) already identified as potentially impacting the speed of post-exercise recovery (Baumert et al., 2016). Knowledge of this variation may enable athletes and support staff to create individualised recovery interventions based on the identification of individuals at increased risk of exercise induced muscle damage or oxidative stress (Del Coso et al., 2017).

Whilst this emerging research of the impact of genotype on exercise recovery is interesting, the translation of these findings to the field are currently under-explored. Given the proposed impact of genetic polymorphisms on exercise recovery, the purpose of this study was to attempt to bridge this gap, by determining whether a seven SNP algorithm successfully differentiated between the recovery speed of male soccer players. It is believed that individuals possessing a greater number of alleles associated with increased oxidative stress, muscle damage or inflammation would see a greater reduction in neuromuscular function post-training, and that this reduction would take longer to abate relative to those individuals in possession of a more favourable genetic profile.

Methods

Subjects

18 male soccer players aged between 16-19 years of age from a college soccer academy volunteered to participate in this study. Each player had an average of 11 years' football training experience, and was actively competing in the English College Football Association Premier League.

Methodology

As part of their normal soccer training, and following 24-hours rest, the subjects took part in a repeated sprints testing session. Both before, and immediately upon completion of this session, the subjects underwent Countermovement Jump (CMJ) testing. This test was repeated at 24- and 48-hours post-training to monitor their recovery status. Subjects were

familiarised to all tests as they are regularly used during their normal soccer training. Within the CMJ trials, subjects undertook three trials at each time point, which were averaged to give a mean score. Subjects had two minutes' recovery between trials. Prior to the initial exercise bout and subsequent testing, players carried out a standardised 15-minute warm up, consisting of pre-activation exercises and dynamic drills. The initial exercise bout was comprised of two sets of seven 25m sprints undertaken outdoors on a soccer pitch. The recovery period was 30 seconds between sprint reps, and 5 minutes between sets.

The CMJ was chosen as it is a reliable and valid measure of neuromuscular fatigue (Cormack et al., 2008a, 2008b, 2008c; McLean et al., 2010; Gathercole et al., 2015) that is widely used in sporting settings (Taylor et al., 2012). The CMJ was measured using Optojump (Microgate, Italy), a valid and reliable tool for the assessment of vertical jump height in the field (Glatthorn et al., 2011). Prior to undertaking each CMJ, subjects were instructed to keep their hands on their hips throughout the jump to eliminate any influence of arm swing. If the arms lost contact with the hips, the jump was classed as a no-jump, and an additional jump was performed following two-minutes recovery. In each CMJ trial, subjects began standing upright, then performed a fast downwards eccentric action followed immediately by a jump for maximal height. Individual results were expressed as height jumped in centimetres (cm).

Genetic Testing

Alongside the training programme, subjects underwent genetic testing by DNAFit Ltd; this occurred via a sterile buccal swab. The samples were sent to IDna Genetics Laboratory (Norwich, UK), where DNA was extracted and purified using the Isohelix Buccalyse DNA extraction kit BEK-50 (Kent, UK), and amplified through PCR on an ABI7900 real-time thermocycler (Applied Biosystem, Waltham, USA). Through this process, genetic information regarding SNPs believed to affect post-exercise recovery speed (*CRP* rs1205, *GSTM1* & *GSTT1* INDEL, *IL-6* -G174C rs1800795, *IL-6R* rs2228145, *SOD2* rs4880, *TNF* G-308A rs1800629) was determined. The DNAFit test uses an algorithm to stratify subjects into "slow", "medium" or "fast" recovery speed by utilising a Total Genotype Score (TGS) method. Each allele is given a score of between 0 and 4 points depending on the expected magnitude of its impact on post-exercise recovery speed. The strength of the rating was based on the evidence from cumulative literature results averaged over time. The sum of these points was combined to give an overall score. This method is identical to Jones et al. (2016), and similar to the methods used in other studies utilising genetic algorithms (Ruiz et al., 2009; Meckel, Ben-Zaken, Nemet, Dror & Eliakim, 2014). An overall score of 40% or less is classed as a fast genetic recovery speed. Scores of 41-60% are classed as a medium genetic recovery speed. A score of >60% is classed as a slow genetic recovery speed. The athletes were blinded to their genetic results until completion of the final testing.

Statistical Analysis

As this is not a genetic association study, but an observational study into the impact of TGS on exercise recovery, gene-by-gene analysis was not carried out. Instead, we focused on data pertaining to exercise recovery, and compared this to TGS group. Means and standard deviations were calculated for whole group and sub-groups for both pre- and post-training (0h, 24h and 48h) test scores. CMJ height at the three post-training time points was converted to a percentage of pre-training height. Given the small sample size, and the fact that significance testing is both sensitive to low sample sizes, and doesn't inform as to the magnitude (Buccheit 2016), we instead calculated effect sizes (Cohen's d) between the groups at the three post-training time points. The thresholds used were 0.2 (trivial), 0.5

(small), 0.8 (moderate), >0.8 (large) (Cohen 1988). Data were analysed using Microsoft Excel 15.29 (Microsoft Corporation, Redmond, WA, USA). All data are reported as mean $\pm$ SD.

Results

Overall, 12 subjects were classed as having a fast recovery speed, with 6 subjects having a medium recovery speed. No subjects were found to have a slow recovery speed; based on an analysis of 17,000 samples tested by DNAFit Ltd (Pickering, unpublished data), approximately 6% of all individuals within a population would be expected to be in the slow group. Within this sample population, we would expect one subject to be in this group; the lack of this TGS is therefore not unusual. Table 1 shows the absolute CMJ results for the fast and medium genetic recovery speed groups. Figure 1 illustrates the between group differences as a percentage over the 48-hour period following the exercise bout, along with effect sizes.

Group	CMJ Pre-Training	CMJ Post-Training	CMJ 24h Post-Training	CMJ 48h Post-Training
Fast (n=12)	37.6 $\pm$ 6.2 cm	37.0 $\pm$ 6.2 cm	36.0 $\pm$ 6.9 cm	37.0 $\pm$ 6.1 cm
Slow (n=6)	35.7 $\pm$ 5.6 cm	34.1 $\pm$ 6.3 cm	33.3 $\pm$ 5.5 cm	33.7 $\pm$ 5.6 cm

Table 1 – CMJ values for both groups across all time points.

Figure 1 – Percentage change in CMJ height immediately post-training, 24h post-training, and 48h post-training as a percentage relative to pre-training values. Effect sizes are # = 0.7 (medium), * = 0.5 (medium), § = 1.0 (large).

Discussion

The results of this study indicate that rate of recovery, as measured by CMJ, is modified by a total genotype score comprised of seven SNPs thought to impact exercise recovery. Overall, players in the “fast” genetic recovery speed group saw a smaller reduction in CMJ height relative to those players in the “medium” genetic recovery speed group, and

were closer to baseline score following 48-hours of recovery. Immediately upon completion of the exercise bout, and 24 hours later, the magnitude of this effect was medium. Forty-eight hours after training, this effect had grown in magnitude to large. This suggests that these seven SNPs impact recovery speed, such that individuals with more favourable alleles suffer a smaller percentage loss in CMJ height after repeated sprints, and have regained a greater percentage of their pre-training CMJ height 48 hours post-training.

The ability to predict the recovery time needed following intense exercise may be useful for several reasons. Firstly, it ensures that optimal recovery time can be given to individual athletes, reducing the fatigue that will accumulate across a training programme. This will ensure that the athlete is not placed at undue risk of suffering from injuries, of which the risk increases under fatigue, and may guard against the developing of unexplained underperformance syndrome. It may also be useful when planning the final physical conditioning session before a competition, with players genetically predisposed to slower recovery speeds having a longer rest period pre-competition. Finally, it may increase the motivation of players to carry out the correct recovery modalities post-training or post-competition, particularly if they are shown to have a slower recovery speed. However, the results from nutrigenetic research indicate that, at present, individuals rarely make behavioural changes based on genetic information (McBride et al., 2010); whether this is the case in highly motivated sports people is unclear.

The SNPs that comprise the genetic algorithm used appear to influence both the inflammatory response to exercise and the ability to tolerate oxidative stress. *IL-6* -G174C (rs1800795) has been shown to influence creatine kinase (CK) levels following eccentric exercise, with the C allele of *IL-6* associated with higher levels (Yamin et al., 2008; Lappalainen 2009). The *IL-6* C allele is also associated with increased post-exercise plasma IL-6 (Huuskonen et al., 2009), which is associated with increased fatigue in athletes (Robson-Ansley et al., 2007). *IL-6R* (rs2228145) is also associated with post-exercise IL-6 levels, with the C allele associated with higher concentrations (Reich et al., 2007). Increases in plasma IL-6R concentrations are associated with increased C-Reactive Protein (CRP) levels post-exercise, as well as increased feelings of fatigue in athletes (Robson-Ansley et al., 2009). The *TNF* G-308A (rs1800629) polymorphism is associated with plasma CRP levels following aerobic exercise, with AA genotypes having higher CRP levels (Lakka et al., 2006). *CRP* (rs1205) alters plasma CRP concentrations, a common exercise recovery marker (Miles et al., 2007; Ingram et al., 2009), with G allele carriers having significantly higher CRP levels than AA genotypes (Eiriksdottir et al., 2009).

SOD2 (rs4880) encodes for manganese superoxide dismutase (MnSOD), which supports the dismutation of mitochondrial superoxide radicals into hydrogen peroxide and oxygen (Li et al., 2005). The T allele of this SNP is associated with increased CK post-exercise (Akimoto et al., 2010; Ahmetov et al., 2014), although this relationship is complex and potentially modified by a subject's habitual antioxidant nutrient intake (Li et al., 2005). *GSTM1* (glutathione S-transferase M1) and *GSTT1* (glutathione S-transferase T1) are insertion/deletion polymorphisms, with deleted genotypes having poor activity of the enzymes. Whilst these polymorphisms are well studied with regards to health and diet interactions (Palli et al., 2004), only one study has examined their impact on post-exercise muscle damage markers, with no significant differences between the inserted or deleted genotypes (Akimoto et al., 2010). These SNPs were included in the algorithm based on their theoretical impact on exercise recovery (Evelo et al., 1992; Vani et al., 1990).

The identification of athletes who may be genetically predisposed to increased recovery times can lead to the use of targeted recovery modalities. These include nutritional interventions. Phillips et al. (2003) reported that 14 days of supplementation with vitamin E, omega-3 and flavonoids blunted the release of IL-6 and CRP following eccentric exercise. Similar results have been reported from other studies (Satoshi et al., 1989; Jouris et al., 2011; Satchek et al., 2003). Other interventions that may enhance recovery include cold water immersion (CWI) and the use of compression garments, although the results are currently equivocal (Leeder et al., 2011; Bleakley & Davison, 2009; Ascensao et al., 2010; Jakeman et al., 2009; Duffield et al., 2010; Duffield et al., 2008; Jakesment et al., 2010). It should be noted that exercise adaptation relies on the application of stress to the body, and the use of antioxidant supplements and CWI may blunt this adaptation (Draeger et al., 2014; Yamane et al., 2015), or in some cases reduce speed of recovery (Close et al., 2006).

There are some potential limitations within this study. The population used is very modest (n=18). A standard soccer team is often comprised of 20-25 players; although at any given time some players may be injured and unable to train. The sample size in this study therefore represents a realistic size that coaches and practitioners may encounter in the real world. It is also representative of sample sizes often used in genetic pilot studies (Loy et al., 2015). However, we recognise that this research requires replication in a larger cohort, which we intend to carry out. In addition, this pilot study had no subjects with a TGS that suggested they had a slow genetic recovery speed, only medium and fast. The result of having a slow recovery speed is uncommon, with approximately only one subject for every sixteen expected to be in this category. This requires very large sample sizes in order to recruit sufficient subjects into this category. The subjects were all male, so it is unclear whether the results would be the same in females. Finally, it must be recognised that the use of this algorithm represents a crude measure, as many other SNPs doubtless impact exercise recovery. However, the results of this pilot study are both novel and promising, such that further research in this field should enable the development of personalised recovery guidelines.

Conclusion

It can be seen that there is considerable variation in the rate of recovery from a repeated sprint exercise in a group of well-trained male soccer players. The use of a seven SNP genetic algorithm aids in the identification of those players who may require longer recovery times between intense exercise bouts, or who may most benefit from targeted recovery interventions. These findings replicate those of Del Coso et al. (2017), although the SNPs utilised vary. The implications of these findings suggest that knowledge of genetic information may be important in individualising recovery timings and modalities in athletes. Future research is required to replicate these findings in a larger cohort, as well as in females, but nevertheless the results potentially herald a further step towards an individualised training process, potentially revolutionising the field of exercise recovery.

Disclosure Statement

Craig Pickering is an employee of DNAFit Ltd. He received no financial incentive for the production of this manuscript, and had full control over the data used. Bruce Suraci is a freelance consultant to DNAFit. John Kiely, Charlie Quigley and Jake Watson have no interests to declare.

References

- Ahmetov II, Naumov VA, Donnikov AE, Maciejewska-Karłowska A, Kostryukova ES, Larin AK, Maykova EV, Alexeev DG, Fedotovskaya ON, Generozov EV, Jastrzębski Z. SOD2 gene polymorphism and muscle damage markers in elite athletes. *Free Radical Research*. 2014;48(8):948-55.
- Akimoto AK, Miranda-Vilela AL, Alves PC, Pereira LC, Lordelo GS, Hiragi CD, da Silva IC, Grisolia CK, Klautau-Guimarães MD. Evaluation of gene polymorphisms in exercise-induced oxidative stress and damage. *Free Radical Research*. 2010;44(3):322-31.
- Ament W, Verkerke GJ. Exercise and fatigue. *Sports Medicine*. 2009;39(5):389.
- Ascensao A, Leite M, Rebelo AN, Magalhães S, Magalhães J. Effects of cold water immersion on the recovery of physical performance and muscle damage following a one-off soccer match. *Journal of Sports Sciences*. 2011;29(3):217-25.
- Baumert P, Lake MJ, Stewart CE, Drust B, Erskine RM. Genetic variation and exercise-induced muscle damage: implications for athletic performance, injury and ageing. *European Journal of Applied Physiology*. 2016;116(9):1595-625.
- Belviranlı M, Gökbel H. Acute exercise induced oxidative stress and antioxidant changes. *Eur J Gen Med*. 2006;3(3):126-31.
- Bleakley CM, Davison GW. What is the biochemical and physiological rationale for using Cold Water Immersion in Sports Recovery? A Systematic Review. *British Journal of Sports Medicine*. 2010;44(3):179-87.
- Bishop PA, Jones E, Woods AK. Recovery from training: a brief review: brief review. *The Journal of Strength & Conditioning Research*. 2008;22(3):1015-24.
- Budgett R. Overtraining and chronic fatigue: The unexplained underperformance syndrome (UPS). *International Sport Med Journal*. 2000;1(3).
- Buchheit M. The numbers will love you back in return—I promise. *International Journal of Sports Physiology and Performance*. 2016;11(4):551-4.
- Close GL, Ashton T, Cable T, Doran D, Holloway C, McArdle F, MacLaren DP. Ascorbic acid supplementation does not attenuate post-exercise muscle soreness following muscle-damaging exercise but may delay the recovery process. *British Journal of Nutrition*. 2006;95(05):976-81.
- Cohen, J. (1988). *Statistical Power Analysis for the Behavioural Sciences*. 2nd ed. Hillsdale, NJ: Lawrence Erlbaum.
- Cormack SJ, Newton RU, McGuigan MR, Doyle TL. Reliability of measures obtained during single and repeated countermovement jumps. *International Journal of Sports Physiology and Performance*. 2008a;3(2):131-44.

Cormack SJ, Newton RU, McGuigan MR. Neuromuscular and endocrine responses of elite players to an Australian rules football match. *International Journal of Sports Physiology and Performance*. 2008b;3(3):359-74.

Cormack SJ, Newton RU, McGuigan MR, Cormie P. Neuromuscular and endocrine responses of elite players during an Australian rules football season. *International Journal of Sports Physiology and Performance*. 2008c;3(4):439-53.

Del Coso J, Valero M, Salinero JJ, Lara B, Gallo-Salazar C, Areces F. Optimum polygenic profile to resist exertional rhabdomyolysis during a marathon. *PloS One*. 2017;12(3):e0172965.

DiFiori JP, Benjamin HJ, Brenner JS, Gregory A, Jayanthi N, Landry GL, Luke A. Overuse injuries and burnout in youth sports: a position statement from the American Medical Society for Sports Medicine. *Br J Sports Med*. 2014;48(4):287-8.

Draeger CL, Naves A, Marques N, Baptistella AB, Carnauba RA, Paschoal V, Nicastro H. Controversies of antioxidant vitamins supplementation in exercise: ergogenic or ergolytic effects in humans? *Journal of the International Society of Sports Nutrition*. 2014;11(1):4.

Duffield R, Cannon J, King M. The effects of compression garments on recovery of muscle performance following high-intensity sprint and plyometric exercise. *Journal of Science and Medicine in Sport*. 2010;13(1):136-40.

Duffield R, Edge J, Merrells R, Hawke E, Barnes M, Simcock D, Gill N. The effects of compression garments on intermittent exercise performance and recovery on consecutive days. *International Journal of Sports Physiology and Performance*. 2008;3(4):454-68.

Eiriksdottir G, Smith AV, Aspelund T, Hafsteinsdottir SH, Olafsdottir E, Launer LJ, Harris TB, Gudnason V. The interaction of adiposity with the CRP gene affects CRP levels: age, gene/environment susceptibility-Reykjavik study. *International Journal of Obesity*. 2009;33(2):267-72.

Evelo CT, Palmen NG, Artur Y, Janssen GM. Changes in blood glutathione concentrations, and in erythrocyte glutathione reductase and glutathione S-transferase activity after running training and after participation in contests. *European Journal of Applied Physiology and Occupational Physiology*. 1992;64(4):354-8.

Fisher-Wellman K, Bloomer RJ. Acute exercise and oxidative stress: a 30 year history. *Dynamic Medicine*. 2009;8(1):1.

Gathercole R, Sporer B, Stellingwerff T, Sleivert G. Alternative countermovement-jump analysis to quantify acute neuromuscular fatigue. *International Journal of Sports Physiology and Performance*. 2015;10(1):84-92.

Glatthorn JF, Gouge S, Nussbaumer S, Stauffacher S, Impellizzeri FM, Maffiuletti NA. Validity and reliability of Optojump photoelectric cells for estimating vertical jump height. *The Journal of Strength & Conditioning Research*. 2011;25(2):556-60.

Huuskonen A, Tanskanen M, Lappalainen J, Oksala N, Kyröläinen H, Atalay M. A common variation in the promoter region of interleukin-6 gene shows association with exercise performance. *Journal of Sports Science & Medicine*. 2009;8(2):271.

Ingram J, Dawson B, Goodman C, Wallman K, Beilby J. Effect of water immersion methods on post-exercise recovery from simulated team sport exercise. *Journal of Science and Medicine in Sport*. 2009;12(3):417-21.

Jakeman JR, Macrae R, Eston R. A single 10-min bout of cold-water immersion therapy after strenuous plyometric exercise has no beneficial effect on recovery from the symptoms of exercise-induced muscle damage. *Ergonomics*. 2009;52(4):456-60.

Jakeman JR, Byrne C, Eston RG. Lower limb compression garment improves recovery from exercise-induced muscle damage in young, active females. *European Journal of Applied Physiology*. 2010;109(6):1137-44.

Jouris KB, McDaniel JL, Weiss EP. The effect of omega-3 fatty acid supplementation on the inflammatory response to eccentric strength exercise. *Journal of Sports Science and Medicine*. 2011;10(3):432-8.

König D, Wagner KH, Elmadfa I, Berg A. Exercise and oxidative stress: significance of antioxidants with reference to inflammatory, muscular, and systemic stress. *Exercise Immunology rReview*. 2001;7:108-33.

Lakka HM, Lakka TA, Rankinen T, Rice T, Rao DC, Leon AS, Skinner JS, Bouchard C. The TNF- α G-308A polymorphism is associated with C-reactive protein levels: the HERITAGE Family Study. *Vascular Pharmacology*. 2006;44(5):377-83.

Lappalainen J. IL6 genotype and creatine kinase response to exercise. *European Journal of Applied Physiology*. 2009;106(2):315-.

Leeder J, Gissane C, van Someren K, Gregson W, Howatson G. Cold water immersion and recovery from strenuous exercise: a meta-analysis. *British Journal of Sports Medicine*. 2012;46(4):233-40.

Li H, Kantoff PW, Giovannucci E, Leitzmann MF, Gaziano JM, Stampfer MJ, Ma J. Manganese superoxide dismutase polymorphism, prediagnostic antioxidant status, and risk of clinical significant prostate cancer. *Cancer Research*. 2005;65(6):2498-504.

Loy BD, O'Connor PJ, Lindheimer JB, Covert SF. Caffeine is ergogenic for adenosine A2A receptor gene (ADORA2A) T allele homozygotes: a pilot study. *Journal of Caffeine Research*. 2015;5(2):73-81.

Mair SD, Seaber AV, Glisson RR, Garrett JR WE. The role of fatigue in susceptibility to acute muscle strain injury. *The American Journal of Sports Medicine*. 1996;24(2):137-43.

McBride CM, Koehly LM, Sanderson SC, Kaphingst KA. The behavioral response to personalized genetic information: will genetic risk profiles motivate individuals and families to choose more healthful behaviors?. *Annual Review of Public Health*. 2010;31:89-103.

McLean BD, Coutts AJ, Kelly V, McGuigan MR, Cormack SJ. Neuromuscular, endocrine, and perceptual fatigue responses during different length between-match microcycles in professional rugby league players. *International Journal of Sports Physiology and Performance*. 2010;5(3):367-83.

Miles MP, Pearson SD, Andring JM, Kidd JR, Volpe SL. Effect of carbohydrate intake during recovery from eccentric exercise on interleukin-6 and muscle-damage markers. *International Journal of Sport Nutrition and Exercise Metabolism*. 2007;17(6):507-20.

Nosaka K, Clarkson PM. Variability in serum creatine kinase response after eccentric exercise of the elbow flexors. *International Journal of Sports Medicine*. 1996;17(02):120-7.

Palli D, Masala G, Peluso M, Gaspari L, Krogh V, Munnia A, Panico S, Saieva C, Tumino R, Vineis P, Garte S. The effects of diet on DNA bulky adduct levels are strongly modified by GSTM1 genotype: a study on 634 subjects. *Carcinogenesis*. 2004;25(4):577-84.

Powers SK, Jackson MJ. Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiological Reviews*. 2008;88(4):1243-76.

Reich D, Patterson N, Ramesh V, De Jager PL, McDonald GJ, Tandon A, Choy E, Hu D, Tamraz B, Pawlikowska L, Wassel-Fyr C. Admixture mapping of an allele affecting interleukin 6 soluble receptor and interleukin 6 levels. *The American Journal of Human Genetics*. 2007;80(4):716-26.

Robson PJ. Elucidating the unexplained underperformance syndrome in endurance athletes: the interleukin-6 hypothesis. *Sports Medicine*. 2003;33(10):771-82.

Robson-Ansley PJ, Blannin A, Gleeson M. Elevated plasma interleukin-6 levels in trained male triathletes following an acute period of intense interval training. *European Journal of Applied Physiology*. 2007;99(4):353-60.

Robson-Ansley P, Barwood M, Canavan J, Hack S, Eglin C, Davey S, Hewitt J, Hull J, Ansley L. The effect of repeated endurance exercise on IL-6 and sIL-6R and their relationship with sensations of fatigue at rest. *Cytokine*. 2009;45(2):111-6.

Sacheck JM, Milbury PE, Cannon JG, Roubenoff R, Blumberg JB. Effect of vitamin E and eccentric exercise on selected biomarkers of oxidative stress in young and elderly men. *Free Radical Biology and Medicine*. 2003;34(12):1575-88.

Satoshi S, Kiyoji T, Hiroyo K, Fumio N. Exercise-induced lipid peroxidation and leakage of enzymes before and after vitamin E supplementation. *International Journal of Biochemistry*. 1989;21(8):835-8.

Schwellnus M, Soligard T, Alonso JM, Bahr R, Clarsen B, Dijkstra HP, Gabbett TJ, Gleeson M, Häggglund M, Hutchinson MR, Van Rensburg CJ. How much is too much? (Part 2) International Olympic Committee consensus statement on load in sport and risk of illness. *British Journal of Sports Medicine*. 2016;50(17):1043-52.

Soligard T, Schwellnus M, Alonso JM, Bahr R, Clarsen B, Dijkstra HP, Gabbett T, Gleeson M, Häggglund M, Hutchinson MR, Van Rensburg CJ. How much is too much? (Part 1)

International Olympic Committee consensus statement on load in sport and risk of injury. British Journal of Sports Medicine. 2016;50(17):1030-41.

Taylor K, Chapman D, Cronin J, Newton MJ, Gill N. Fatigue monitoring in high performance sport: a survey of current trends. J Aust Strength Cond. 2012;20(1):12-23.

Vani M, Reddy GP, Reddy GR, Thyagaraju K, Reddanna P. Glutathione-S-transferase, superoxide dismutase, xanthine oxidase, catalase, glutathione peroxidase and lipid peroxidation in the liver of exercised rats. Biochemistry International. 1989;21(1):17-26.

Wilber CA, Holland GJ, Madison RE, Loy SF. An epidemiological analysis of overuse injuries among recreational cyclists. International Journal of Sports Medicine. 1995;16(03):201-6.

Yamane M, Ohnishi N, Matsumoto T. Does regular post-exercise cold application attenuate trained muscle adaptation? International Journal of Sports Medicine. 2015;36(08):647-53.

Yamin C, Duarte JA, Oliveira JM, Amir O, Sagiv M, Eynon N, Sagiv M, Amir RE. IL6 (-174) and TNFA (-308) promoter polymorphisms are associated with systemic creatine kinase response to eccentric exercise. European Journal of Applied Physiology. 2008;104(3):579.