**Nonconsumptive predator effects on prey demography: Dogwhelk
cues decrease benthic mussel recruitment**

Sonja M. Ehlers, Ricardo A. Scrosati, and Julius A. Ellrich

*St. Francis Xavier University, Department of Biology, Antigonish, Nova Scotia B2G 2W5,
Canada*

Corresponding author: R A. Scrosati. Email: rscrosat@stfx.ca

Abstract

Predators have often been shown to have nonconsumptive effects (NCEs) on prey behaviour, but the demographic consequences for prey remain poorly known. This is important to understand because demography influences the impact of a species in its community. We used an intertidal predator–prey system to investigate predator NCEs on prey recruitment, a key demographic process for population persistence. Pelagic mussel larvae are known to avoid waterborne cues from dogwhelks, which prey on intertidal mussels. Through a field experiment done in Atlantic Canada, we manipulated the presence of dogwhelks in intertidal habitats during the mussel recruitment season. We measured mussel recruitment in collectors that could be reached by waterborne dogwhelk cues but not by dogwhelks themselves. We found that the nearby presence of dogwhelks significantly decreased mussel recruit density. A previous study done in the same habitats under the same experimental conditions showed that dogwhelk cues also limit the recruitment of barnacles, another prey item for dogwhelks. However, such NCEs were four times stronger than those observed for mussel recruitment. This difference relates well to the higher ability of mussels to escape predation, as mussels can relocate while barnacles cannot. Therefore, basic features of natural history may be useful to predict predator NCEs on prey recruitment.

Keywords: nonconsumptive effect; mussel; predator; prey; recruitment; snail.

Introduction

Nonconsumptive effects (NCEs) of predators on prey are ubiquitous in nature. When organisms of a prey species detect cues from nearby predators, a variety of responses are often triggered to limit predation risk (Ferrari *et al.*, 2010; Brönmark & Hansson, 2012). As cues from a predator can reach many prey organisms at the same time, NCEs can be extensive in prey populations (Preisser *et al.*, 2005; Peacor *et al.*, 2013). Thus, understanding what prey traits are affected and how has become an important research line in ecology (Weissburg *et al.*, 2014).

Immediate prey responses are typically behavioural. They include moving away to minimize the chance of being reached by predators or limiting movements to avoid being detected by predators (Keppel & Scrosati, 2004; Molis *et al.*, 2011; Johnston *et al.*, 2012; Matassa *et al.*, 2016, Johnson *et al.*, 2017). The consequences of such behavioural responses for prey demography have received, however, little attention (Creel *et al.*, 2007; Schoener & Spiller, 2012; Ellrich *et al.*, 2016a). This is important to understand because demography ultimately determines to a large extent the function of a species in its community. This paper focuses on predator NCEs on prey recruitment, which is a key demographic process for population persistence (Caley *et al.*, 1996; Palumbi & Pinsky, 2014).

Benthic invertebrates with pelagic larvae are useful model organisms for this kind of research. For instance, a laboratory experiment has shown that larvae of blue mussels (*Mytilus edulis*) avoid waterborne chemical cues from predatory dogwhelks (*Nucella lapillus*; Morello & Yund, 2016). Dogwhelks feed on benthic mussel stages, not on their pelagic larvae (Hunt & Scheibling, 1998). However, larval avoidance of dogwhelk cues may have evolved to aid settlement-seeking larvae to find habitats with a reduced predation pressure for juveniles and adults. Such an avoidance behaviour might ultimately decrease benthic recruitment (the addition

of new organisms to a benthic population after larval settlement and metamorphosis). In fact, field experiments in intertidal habitats have shown that cues from *N. lapillus* limit barnacle (*Semibalanus balanoides*) recruitment (Ellrich *et al.*, 2015a,b). This barnacle is another important prey for *N. lapillus* and it also has pelagic larvae, which settle elsewhere when dogwhelk cues are detected (Ellrich *et al.*, 2016a). Thus, the mussel–dogwhelk system offers the opportunity to start evaluating how broadly predator NCEs can limit the recruitment of benthic invertebrate prey. Through a field experiment, the present study tests the hypothesis that dogwhelk cues limit intertidal mussel recruitment.

Basic differences in natural history between mussels and barnacles may influence the intensity of such NCEs, however. The location of a barnacle is fixed for life after a larva settles and metamorphoses into a recruit (Jenkins *et al.*, 2000). However, mussel recruits can detach themselves from the substrate and relocate (Bayne, 1964; Le Corre *et al.*, 2013). Additionally, older mussels can immobilize dogwhelks through the production of byssus threads (Farrell & Crowe, 2007). These processes provide mussels with opportunities to escape predation that barnacles lack. Thus, we also predict that the expected dogwhelk NCEs on mussel recruitment are weaker than the NCEs recently reported for barnacles.

Materials and Methods

We did the experiment in rocky intertidal habitats from Deming Island (45° 12' 45" N, 61° 10' 26" W), on the Atlantic coast of Nova Scotia (Canada), between May–July 2016. These habitats are constituted by stable bedrock and are protected from direct oceanic swell by rocky formations. Maximum water velocity measured with dynamometers (see design in Bell & Denny, 1994) during the study period was $6.0 \pm 0.3 \text{ m s}^{-1}$ (mean $\pm$ SE, $n = 48$). These wave-sheltered habitats were used in previous years to demonstrate that dogwhelk cues limit barnacle

recruitment (Ellrich & Scrosati, 2016; Ellrich *et al.*, 2015b, 2016b). In-situ temperature measured every 30 minutes during the study period using submersible loggers (HOBO Pendant Logger, Onset Computer Corp., Pocasset, MA, USA) was 12.8 ± 0.1 °C (mean $\pm$ SE, $n = 7$ loggers). Coastal seawater salinity measured on 21 May 2016 with a refractometer was 35 ‰.

The dogwhelk used for this study was *Nucella lapillus*, which is the only dogwhelk species on the studied coast (Scrosati & Heaven, 2007). On the Atlantic coast of Nova Scotia, two blue mussel congeners, *Mytilus edulis* and *M. trossulus*, co-occur (Tam & Scrosati, 2011, 2014) and are preyed upon by *N. lapillus* (Hunt & Scheibling, 1998). These mussel species show only subtle morphological differences (Innes & Bates, 1999) and can form hybrids (Riginos & Cunningham, 2005). Thus, their visual identification is very difficult, especially at the recruit stage. Therefore, recruit counts in this study were done as *Mytilus* spp., as commonly done in ecological field studies involving these species (Cusson & Bourget, 2005; Le Corre *et al.*, 2013).

We evaluated dogwhelk cue effects on mussel recruitment by manipulating dogwhelk presence in cages attached to the intertidal substrate. Each cage (Fig. 1) was made using a PVC ring (25 cm in diameter and 2.5 cm tall) and plastic mesh (0.5 cm x 0.5 cm of opening size). Each cage was divided by mesh into a central compartment (area = 144 cm²) and a peripheral compartment (area = 347 cm²). The peripheral compartment was used to create two dogwhelk treatments (presence vs. absence) by enclosing either 10 dogwhelks (2.23 ± 0.02 cm in shell length, mean $\pm$ SE, $n = 104$) or none. The used dogwhelk density (ca. 3 individuals dm⁻²) was representative of the studied coast (Ellrich & Scrosati, 2016). The central compartment held a plastic mesh scourer (Our Compliments Poly Pot Scrubbers, Mississauga, ON, Canada) attached with cable ties (Fig. 1). Mesh scourers have often been used to measure intertidal mussel recruitment (Menge & Menge, 2013; South, 2016), as scourers resemble habitats where mussel

larvae preferentially settle (filamentous algae or byssal threads of established mussels; Menge, 1992; Le Corre *et al.*, 2013). For *Mytilus edulis* and *M. trossulus*, pelagic pediveliger larvae of at least approximately 0.25 mm in shell length settle in those habitats and, then, undergo metamorphosis, becoming recruits (Bayne, 1965; Menge *et al.*, 2009; Martel *et al.*, 2014). After growing to a shell length of about 0.5 mm (Hunt & Scheibling, 1996; Le Corre *et al.*, 2013), such recruits may enter a second pelagic dispersal phase (Bayne, 1964). For instance, recruits of *M. edulis* up to 2.5 mm long can passively drift in the water aided by a byssus thread (Sigurdsson *et al.*, 1976). In our study, observations under a stereomicroscope indicated that 70-80 % of the recruits found in the scourers belonged to the first phase, the remaining organisms belonging to the second phase. Precise counts are unavailable because the threshold size between both phases is not accurately known (Le Corre *et al.*, 2013). As all of those organisms ultimately contribute to mussel recruitment (Le Corre *et al.*, 2013), at the end of the experiment we counted the recruits of both phases together to determine recruit density for each scourer, as often done in field studies of this kind (Menge & Menge, 2013).

We set up the experiment on 21 May 2016 following a randomized complete block design with replicated treatments within blocks (Quinn & Keough, 2002). We established 12 blocks, each one including two replicate cages of each of the two dogwhelk treatments, thus yielding 24 replicates for each dogwhelk treatment. Block size was $7.7 \pm 0.4 \text{ m}^2$ (mean $\pm$ SE, $n = 12$ blocks) and the distance between cages within blocks was at least 0.5 m. We established the blocks at an intertidal elevation of $0.9 \pm 0.1 \text{ m}$ (mean $\pm$ SE, $n = 12$ blocks) above chart datum (the full vertical intertidal range is 1.8 m). We attached the cages to the substrate using PVC plates and screws (Fig. 1). Before installing the cages, we removed all seaweeds (mainly *Ascophyllum nodosum* and *Fucus vesiculosus*) and benthic invertebrates from the substrate to avoid chemical and

physical influences from those organisms (Johnson & Strathmann, 1989; Jenkins *et al.*, 1999; Beermann *et al.*, 2013). During the experiment, we kept these areas devoid of free-living dogwhelks. We did not feed the caged dogwhelks during the experiment but, to prevent their starvation, we replaced them every 10-14 days with mussel-fed dogwhelks that were kept in separate cages tens of meters away from the blocks. We used mussel-fed dogwhelks because prey reacts strongly to chemical cues from predators fed conspecific prey (Cheung *et al.*, 2006; Weissburg & Beauvais, 2015; Scherer & Smee, 2016). We ended the experiment on 29 July 2016, when we took all of the scourers to the laboratory to measure mussel recruit density.

In the laboratory, we stored the scourers in a freezer to preserve the integrity of the recruits until each scourer was analyzed. To count the recruits in a scourer, we unrolled the scourer and manually rinsed it in tap water to separate the recruits from the mesh. The recruits were retained in a sieve (0.212 mm x 0.212 mm of opening size) and then transferred to a Petri dish. We subsequently counted the recruits under a stereomicroscope. For each scourer, we calculated mussel recruit density by dividing the encountered number of recruits by the total area of the scourer. This standardization was necessary because small area differences could exist among the replicate scourers provided by the vendor. To calculate the total area of a scourer, we first unrolled the scourer. Then, we used scissors to cut alongside the resulting cylindrical mesh to produce a two-dimensional mesh, which we extended flat on a table. As mussel recruits occurred on both sides of this surface, we calculated the total area of the scourer as the area of that two-dimensional mesh viewed from the top multiplied by two. We evaluated the effect of dogwhelk cues (fixed factor with two levels: dogwhelk presence and absence) on mussel recruit density through an analysis of variance (ANOVA) that was appropriate for a randomized complete block design with replicated treatments within blocks (random factor with 12 levels). We confirmed

the homoscedasticity and normality assumptions using Cochran's *C*-test and the Kolmogorov-Smirnov test, respectively.

We also conducted a side experiment to verify that the presence of dogwhelks in a cage did not alter water motion at the place of attachment of the mesh scourer. For this purpose, we established 24 different cages on the shore on 1 June 2016. Each of those cages held a gypsum piece (Jonsson *et al.*, 2006; Beermann *et al.*, 2013) in the same place in which the cages used for the main experiment held a mesh scourer. We prepared the gypsum pieces following Howerton & Boyd (1992). We determined the initial dry mass of each gypsum piece to the nearest 0.1 mg. Twelve randomly selected cages each contained 10 dogwhelks in the peripheral compartment, whereas the other 12 cages lacked dogwhelks. On 2 June 2016, we collected the gypsum pieces, dried them at 60 °C for 24 h, and then measured the percent loss of mass for each piece. We compared percent loss of gypsum mass between both treatments with a *t*-test. We conducted all of the data analyses with STATISTICA 13.5 (Statsoft, Tulsa, OK, USA).

Results

The ANOVA for the field experiment revealed that the presence of dogwhelks decreased intertidal mussel recruitment (Table 1). On average, mussel recruit density was 13 % lower with nearby dogwhelks than in their absence (Fig. 2). Blocks had a significant effect on mussel recruit density (Table 1), but that result merely indicates that mussel recruitment differed among blocks. The important result is that the interaction between the dogwhelks factor and the blocking factor was not significant (Table 1), indicating that the negative dogwhelk NCEs on mussel recruitment were spatially consistent on the shore. The side field experiment revealed that the presence of dogwhelks in the cages did not affect water motion ($t_{22} = 1.14$, $P = 0.267$) in the place in which the cages used for the main experiment held a mesh scourer.

Discussion

This study has experimentally demonstrated that cues from predatory dogwhelks decrease mussel recruitment in intertidal habitats. This is a valuable contribution because it adds to the growing literature that is revealing predator NCEs on prey demography. Other studies have shown negative NCEs on prey reproduction (Selden *et al.*, 2009; Zanette *et al.*, 2011; Ellrich *et al.*, 2016a) and also recruitment (Creel *et al.*, 2011; Ellrich *et al.*, 2015a; Benkwitt, 2017). These studies are important because most NCE research to date has focused on behavioural responses in prey (Ferrari *et al.*, 2010; Brönmark & Hansson, 2012; Schoener & Spiller, 2012), likely because of the short times required to document such responses. Evaluating the demographic consequences requires more time, but this knowledge is necessary to better understand predator NCEs on prey population dynamics (Weissburg *et al.*, 2014).

Nucella lapillus preys on blue mussels (Crothers, 1985). Young *N. lapillus* consume juvenile mussels and even recently hatched *N. lapillus* prey on young mussels by drilling a hole through their shells (Largen, 1967). Hence, dogwhelks are a threat to various age classes of mussels. Such an extended predation pressure is, therefore, what may have selected for the larval avoidance behaviour (Morello & Yund, 2016) that can ultimately decrease recruitment. In intertidal habitats, dogwhelks are patchily distributed (Johnson *et al.*, 1998) and have a restricted activity range (Crothers, 1985; Fretter & Graham, 1994; Carro *et al.*, 2012). Thus, by avoiding dogwhelk cues, young mussels likely contribute to limiting future predation risk.

This study has also revealed that the recruitment limitation caused by dogwhelk cues is weaker for mussels than for barnacles. Barnacles cannot change their location once recruited (Anderson, 1994) and cues from *Nucella lapillus* were found to limit barnacle (*Semibalanus balanoides*) recruitment by 50 % in the same habitats where we conducted the present study and

under the same dogwhelk density (Ellrich *et al.*, 2016b). Mussels are also sessile, but to some extent they can relocate across the substrate throughout their benthic existence (Bayne, 1964; Hunt & Scheibling, 2002; de Vooy, 2003; van de Koppel *et al.*, 2008). Older mussels can also immobilize dogwhelks using byssus threads (Farrell & Crowe, 2007). Therefore, mussels have more opportunities to escape predation than barnacles, which might explain why mussel recruitment is less responsive to dogwhelk cues than barnacle recruitment.

Pre-recruitment avoidance of predator cues has been found not only for mussels and barnacles, but also for lobsters (Boudreau *et al.*, 1993), crabs (Welch *et al.*, 1997; Banks & Dinnel, 2000; Tapia-Lewin & Pardo, 2014), and sea urchins (Metaxas & Burdett-Coutts, 2006). Thus, negative predator NCEs on prey recruitment might be common in benthic invertebrates with pelagic dispersal stages. The intensity of such NCEs may depend, as discussed above and among other factors, on the capacity of benthic prey stages to relocate across the substrate.

Indirect NCEs of predators on third species mediated by the direct NCEs on the predator's prey have often been reported (Molis *et al.*, 2011; Schoener & Spiller, 2012; Matassa *et al.*, 2016). Those studies have generally evaluated effects on only one or a few of such third species (although exceptions evaluating responses on entire assemblages exist; Hammill *et al.*, 2015). Intertidal mussels are foundation species (Altieri & van de Koppel, 2014), because they often occur in extensive patches that host several small species among the mussels (Valdivia & Thiel, 2006; O'Connor & Crowe, 2007; Arribas *et al.*, 2014). Therefore, by nonconsumptively limiting mussel recruitment, dogwhelks have the potential to alter intertidal species composition. Evaluating this possibility would enrich models of community organization that currently consider only the consumptive effects of predators on foundation species and its associated biodiversity (Bruno *et al.*, 2003; Scrosati *et al.*, 2011).

Overall, the present study shows that predator NCEs limit intertidal mussel recruitment, potentially affecting mussel population dynamics. Moreover, the study has linked behavioural observations obtained in the laboratory (Morello & Yund, 2016) to population processes occurring under natural conditions. The field nature of our experiment is important because the complexity of intertidal environments cannot be replicated in laboratory settings. Thus, our approach agrees with recent calls to study predator NCEs under realistic conditions in order to advance NCE theory further (Weissburg *et al.*, 2014; Babarro *et al.*, 2016).

Acknowledgements

We thank Jadine Krist and Zachary Sherker for field assistance. This project was funded by a Discovery Grant from the Natural Sciences and Engineering Research Council (NSERC) and a Leaders Opportunity Grant from the Canada Foundation for Innovation (CFI) awarded to R.A.S. and by a postdoctoral fellowship from the German Academic Exchange Service (DAAD) awarded to J.A.E.

References

- Altieri, A.H. & van de Koppel, J. (2014). Foundation species in marine ecosystems. In *Marine community ecology and conservation*: 37–56. Bertness, M.D., Bruno, J.F., Silliman, B.R. & Stachowicz, J.J. (Eds). Sunderland: Sinauer.
- Anderson, D.T. (1994). *Barnacles. Structure, function, development, and evolution*. London: Chapman & Hall.
- Arribas, L.P., Donnarumma, L., Palomo, M.G. & Scrosati, R.A. (2014). Intertidal mussels as ecosystem engineers: their associated invertebrate biodiversity under contrasting wave exposures. *Mar. Biodiv.* 44: 203–211.

230 Babarro, J.M.F., Vázquez, E. & Olabarria, C. (2016). Importance of phenotypic plastic traits on
 231 invasive success: response of *Xenostrobus securis* to the predatory dogwhelk *Nucella lapillus*.
 232 *Mar. Ecol. Prog. Ser.* 560: 185–198.

233 Banks, J. & Dinnel, P. (2000). Settlement behavior of Dungeness crab (*Cancer magister* Dana,
 234 1852) megalopae in the presence of the shore crab, *Hemigrapsus* (Decapoda, Brachyura).
 235 *Crustaceana* 73: 223–234.

236 Bayne, B.L. (1964). Primary and secondary settlement in *Mytilus edulis* L. (Mollusca). *J. Anim.*
 237 *Ecol.* 33: 513–523.

238 Bayne, B.L. (1965). Growth and the delay of metamorphosis of the larvae of *Mytilus edulis* (L.)
 239 *Ophelia* 2: 1-47.

240 Beermann, A.J., Ellrich, J.A., Molis, M. & Scrosati, R.A. (2013). Effects of seaweed canopies
 241 and adult barnacles on barnacle recruitment: the interplay of positive and negative influences. *J.*
 242 *Exp. Mar. Biol. Ecol.* 448: 162–170.

243 Bell, E.C. & Denny, M.W. (1994). Quantifying "wave exposure": a simple device for recording
 244 maximum velocity and results of its use at several sites. *J. Exp. Mar. Biol. Ecol.* 156: 199–215.

245 Benkwitt, C.E. (2017). Predator effects on reef fish settlement depend on predator origin and
 246 recruit density. *Ecology* 98: 896–902.

247 Boudreau, B., Bourget, E. & Simard, Y. (1993). Behavioural responses of competent lobster
 248 postlarvae to odor plumes. *Mar. Biol.* 117: 63–69.

249 Brönmark, C. & Hansson, L.A. (2012). *Chemical ecology in aquatic systems*. Oxford: Oxford
 250 University Press.

251 Bruno, J.F., Stachowicz, J.J. & Bertness, M.D. (2003). Inclusion of facilitation into ecological
252 theory. *Trends Ecol. Evol.* 18: 119–125.

253 Caley, M.J., Carr, M.H., Hixon, M.A., Hughes, T.P., Jones, G.P. & Menge, B.A. (1996).
254 Recruitment and the local dynamics of open marine populations. *Annu. Rev. Ecol. Syst.* 27:
255 477–500.

256 Carro, B., Quintela, M., Ruiz, J.M. & Barreiro, R. (2012). AFLPs reveal different population
257 genetic structure under contrasting environments in the marine snail *Nucella lapillus*. *PLoS*
258 *ONE* 7: e49776.

259 Cheung, S.G., Luk, K.C. & Shin, P.K.S. (2006). Predator-labeling effect on byssus production in
260 marine mussels *Perna viridis* (L.) and *Brachidontes variabilis* (Krauss). *J. Chem. Ecol.* 32:
261 1501–1512.

262 Creel, A., Christianson, D., Liley, S. & Winney, J.A. (2007). Predation risk affects reproductive
263 physiology and demography of elk. *Science* 315: 960.

264 Creel, A., Christianson, D. & Winney, J.A. (2011). A survey of the effects of wolf predation risk
265 on pregnancy rates and calf recruitment in elk. *Ecol. Appl.* 21: 2847–2853.

266 Crothers, J.H. (1985). Dog-whelks: an introduction to the biology of *Nucella lapillus* (L.). *Field*
267 *Studies* 6: 291–360.

268 Cusson, M. & Bourget, E. (2005). Small-scale variations in mussel (*Mytilus* spp.) dynamics and
269 local production. *J. Sea Res.* 53: 255–268.

270 de Vooy, C.G.N (2003). Effect of a tripeptide on the aggregational behaviour of the blue mussel
271 *Mytilus edulis*. *Mar. Biol.* 142: 1119–1123.

272 Ellrich, J.A. & Scrosati, R.A. (2016). Water motion modulates predator nonconsumptive
273 limitation of prey recruitment. *Ecosphere* 7: e01402.

274 Ellrich, J.A., Scrosati, R.A., Bertolini, C. & Molis, M. (2016a). A predator has nonconsumptive
275 effects on different life-history stages of a prey. *Mar. Biol.* 163: article 5.

276 Ellrich, J.A., Scrosati, R.A. & Molis, M. (2015a). Predator nonconsumptive effects on prey
277 recruitment weaken with recruit density. *Ecology* 96: 611–616.

278 Ellrich, J.A., Scrosati, R.A. & Petzold, W. (2015b). Predator density affects nonconsumptive
279 predator limitation of prey recruitment: field experimental evidence. *J. Exp. Mar. Biol. Ecol.*
280 472: 72–76.

281 Ellrich, J.A., Scrosati, R.A., Romoth, K. & Molis, M. (2016b). Adult prey neutralizes predator
282 nonconsumptive limitation of prey recruitment. *PLoS ONE* 11: e0154572.

283 Farrell, E.D. & Crowe, T.P. (2007). The use of byssus threads by *Mytilus edulis* as an active
284 defense against *Nucella lapillus*. *J. Mar. Biol. Assoc. U. K.* 87: 559–564.

285 Ferrari, M.C.O., Wisenden, B.D. & Chivers, D.P. (2010). Chemical ecology of predator–prey
286 interactions in aquatic ecosystems: a review and prospectus. *Can. J. Zool.* 88: 698–724.

287 Fretter, V. & Graham, A. (1994). *British prosobranch molluscs. Their functional anatomy and*
288 *ecology*. London: The Ray Society.

289 Hammill, E., Atwood, T.B. & Srivastava, D.S. (2015). Predation threat alters composition and
290 functioning of bromeliad ecosystems. *Ecosystems* 18: 857–866.

291 Howerton, R.D. & Boyd, C.E. (1992). Measurement of water circulation in ponds with gypsum
292 blocks. *Aquacult. Eng.* 11: 141–155.

293 Hunt, H.L. & Scheibling, R.E. (1996). Physical and biological factors influencing mussel
294 (*Mytilus trossulus*, *M. edulis*) settlement on a wave-exposed rocky shore. *Mar. Ecol. Prog. Ser.*
295 142: 135–145.

296 Hunt, H.L. & Scheibling, R.E. (1998). Effects of whelk (*Nucella lapillus* (L.)) predation on
297 mussel (*Mytilus trossulus* (Gould), *M. edulis* (L.)) assemblages in tidepools and on emergent
298 rock on a wave-exposed rocky shore in Nova Scotia, Canada. *J. Exp. Mar. Biol. Ecol.* 226: 87–
299 113.

300 Hunt, H.L. & Scheibling, R.E. (2002). Movement and wave dislodgment of mussels on a wave-
301 exposed rocky shore. *Veliger* 45: 273–277.

302 Innes, D.J. & Bates, J.A. (1999). Morphological variation of *Mytilus edulis* and *Mytilus trossulus*
303 in eastern Newfoundland. *Mar. Biol.* 133: 691–699.

304 Jenkins, S.R., Åberg, P., Cervin, G., Coleman, R.A., Delany, J., Della Santina, P., Hawkins, S.J.,
305 LaCroix, E., Myers, A.A., Lindegarth, M., Power, A.M., Roberts, M.F. & Hartnoll, R.G.
306 (2000). Spatial and temporal variation in settlement and recruitment of the intertidal barnacle
307 *Semibalanus balanoides* (L.) (Crustacea: Cirripedia) over a European scale. *J. Exp. Mar. Biol.*
308 *Ecol.* 243: 209–225.

309 Jenkins, S.R., Norton, T.A. & Hawkins, S.J. (1999). Settlement and post-settlement interactions
310 between *Semibalanus balanoides* (L.) (Crustacea: Cirripedia) and three species of fucoid
311 canopy algae. *J. Exp. Mar. Biol. Ecol.* 236: 49–67.

312 Johnson, G.C., Karajah, M.T., Mayo, K., Armenta, T.C. & Blumstein, D.T. (2017). The bigger
313 they are the better they taste: size predicts predation risk and anti-predator behavior in giant
314 clams. *J. Zool.* 301: 102–107.

- 315 Johnson, L.E. & Strathmann, R.R. (1989). Settling barnacle larvae avoid substrata previously
316 occupied by a mobile predator. *J. Exp. Mar. Biol. Ecol.* 128: 7–103.
- 317 Johnson, M.P., Hughes, R.N., Burrows, M.T. & Hawkins, S.J. (1998). Beyond the predation
318 halo: small scale gradients in barnacle populations affected by the relative refuge value of
319 crevices. *J. Exp. Mar. Biol. Ecol.* 231: 163–170.
- 320 Johnston, B.R., Molis, M. & Scrosati, R.A. (2012). Predator chemical cues affect prey feeding
321 activity differently in juveniles and adults. *Can. J. Zool.* 90: 128–132.
- 322 Jonsson, P.R., Granhag, L., Moschella, P.S., Åberg, P., Hawkins, S.J. & Thompson, R.C. (2006).
323 Interactions between wave action and grazing control the distribution of intertidal macroalgae.
324 *Ecology* 87: 1169–1178.
- 325 Keppel, E. & Scrosati, R. (2004). Chemically mediated avoidance of *Hemigrapsus nudus*
326 (Crustacea) by *Littorina scutulata* (Gastropoda): effects of species coexistence and variable
327 cues. *Anim. Behav.* 68: 915–920.
- 328 Largen, M.J. (1967). The diet of the dog-whelk, *Nucella lapillus* (Gastropoda Prosobranchia). *J.*
329 *Zool.* 151: 123–127.
- 330 Le Corre, N., Martel, A.L., Guichard, F. & Johnson, L.E. (2013). Variation in recruitment:
331 differentiating the roles of primary and secondary settlement of blue mussels *Mytilus* spp. *Mar.*
332 *Ecol. Prog. Ser.* 481: 133–146.
- 333 Martel, A.L., Tremblay, R., Toupont, N., Olivier, F. & Myrand, B. (2014). Veliger size at
334 metamorphosis and temporal variability in prodissococonch II morphometry in the blue mussel
335 (*Mytilus edulis*): potential impact on recruitment. *J. Shellfish Res.* 33: 443–455.

- 336 Matassa, C.M., Donelan, S.C., Luttbeg, B. & Trussell, G.C. (2016). Resource levels and prey
337 state influence antipredator behavior and the strength of nonconsumptive predator effects.
338 *Oikos* 125: 1478–1488.
- 339 Menge, B.A. (1992). Community regulation: under what conditions are bottom-up factors
340 important on rocky shores? *Ecology* 73: 755–765.
- 341 Menge, B.A., Chan, F., Nielsen, K.J., Di Lorenzo, E. & Lubchenco, J. (2009). Climatic variation
342 alters supply-side ecology: impact of climate patterns on phytoplankton and mussel
343 recruitment. *Ecol. Monogr.* 79: 379–395.
- 344 Menge, B.A. & Menge, D.N.L. (2013). Dynamics of coastal meta-ecosystems: the intermittent
345 upwelling hypothesis and a test in rocky intertidal regions. *Ecol. Monogr.* 83: 283–310.
- 346 Metaxas, A. & Burdett-Coutts, V. (2006). Response of invertebrate larvae to the presence of the
347 ctenophore *Bolinopsis infundibulum*, a potential predator. *J. Exp. Mar. Biol. Ecol.* 334: 187–
348 195.
- 349 Molis, M., Preuss, I., Firmenich, A. & Ellrich, J. (2011). Predation risk indirectly enhances
350 survival of seaweed recruits but not intraspecific competition in an intermediate herbivore
351 species. *J. Ecol.* 99: 807–817.
- 352 Morello, S.L. & Yund, P.O. (2016). Response of competent blue mussel (*Mytilus edulis*) larvae
353 to positive and negative settlement cues. *J. Exp. Mar. Biol. Ecol.* 480: 8–16.
- 354 O'Connor, N.E. & Crowe, T.P. (2007). Biodiversity among mussels: separating the influence of
355 sizes of mussels from the ages of patches. *J. Mar. Biol. Assoc. U. K.* 87: 551–557.

- 356 Palumbi, S.R. & Pinsky, M.L. (2014). Marine dispersal, ecology, and conservation. In *Marine*
357 *community ecology and conservation*: 57–83. Bertness, M.D., Bruno, J.F., Silliman, B.R. &
358 Stachowicz, J.J. (Eds). Sunderland: Sinauer.
- 359 Peacor, S.D., Peckarsky, B.L., Trussell, G.C. & Vonesh, J.R. (2013). Costs of predator-induced
360 phenotypic plasticity: a graphical model for predicting the contribution of nonconsumptive and
361 consumptive effects of predators on prey. *Oecologia* 171: 1–10.
- 362 Preisser, E.L., Bolnick, D.I. & Benard, M.F. (2005). Scared to death? The effects of intimidation
363 and consumption in predator–prey interactions. *Ecology* 86: 501–509.
- 364 Quinn, G.P. & Keough, M.J. (2002). *Experimental design and data analysis for biologists*.
365 Cambridge: Cambridge University Press.
- 366 Riginos, C. & Cunningham, C.W. (2005). Local adaptation and species segregation in two
367 mussel (*Mytilus edulis* x *Mytilus trossulus*) hybrid zones. *Mol. Ecol.* 14: 381–400.
- 368 Scherer, A.E. & Smee, D.L. (2016). A review of predator diet effects on prey defensive
369 responses. *Chemoecology* 26: 83–100.
- 370 Schoener, T.W. & Spiller, D.A. (2012). Perspective: kinds of trait-mediated indirect effects in
371 ecological communities. A synthesis. In *Trait-mediated indirect interactions: ecological and*
372 *evolutionary perspectives*: 9–27. Ohgushi, T., Schmitz, O.J. & Holt, R.D. (Eds) Cambridge:
373 Cambridge University Press.
- 374 Scrosati, R. & Heaven, C. (2007). Spatial trends in community richness, diversity, and evenness
375 across rocky intertidal environmental stress gradients in eastern Canada. *Mar. Ecol. Prog. Ser.*
376 342: 1–14.

- 377 Scrosati, R.A., van Genne, B., Heaven, C.S. & Watt, C.A. (2011). Species richness and diversity
378 in different functional groups across environmental stress gradients: a model for marine rocky
379 shores. *Ecography* 34: 151–161.
- 380 Selden, R., Johnson, A.S. & Ellers, O. (2009). Waterborne cues from crabs induce thicker
381 skeletons, smaller gonads, and size-specific changes in growth rate in sea urchins. *Mar. Biol.*
382 156: 1057–1071.
- 383 Sigurdsson, J.B., Titman, C.W. & Davies, P.A. (1976). The dispersal of young post-larval
384 bivalve molluscs by byssus threads. *Nature* 262: 386–387.
- 385 South, P.M. (2016). An experimental assessment of measures of mussel settlement: effects of
386 temporal, procedural, and spatial variations. *J. Exp. Mar. Biol. Ecol.* 482: 64–74.
- 387 Tam, J.C. & Scrosati, R.A. (2011). Mussel and dogwhelk distribution along the north-west
388 Atlantic coast: testing predictions derived from the abundant-centre model. *J. Biogeogr.* 38:
389 1536–1545.
- 390 Tam, J.C. & Scrosati, R.A. (2014). Distribution of cryptic mussel species (*Mytilus edulis* and *M.*
391 *trossulus*) along wave exposure gradients on northwest Atlantic rocky shores. *Mar. Biol. Res.*
392 10: 51–60.
- 393 Tapia-Lewin, S. & Pardo, L.M. (2014). Field assessment of the predation risk–food availability
394 trade-off in crab megalopae settlement. *PLoS ONE* 9: e95335.
- 395 Valdivia, N. & Thiel, M. (2006). Effects of point-source nutrient addition and mussel removal on
396 epibiotic assemblages in *Perumytilus purpuratus* beds. *J. Sea Res.* 56: 271–283.

- 397 van de Koppel, J., Gascoigne, J.C., Theraulaz, G., Rietkerk, M., Mooij, W.M. & Herman, P.M.J.
 398 (2008). Experimental evidence for spatial self-organization and its emergent effects in mussel
 399 bed ecosystems. *Science* 322: 739–742.
- 400 Weissburg, M. & Beauvais, J. (2015). The smell of success: the amount of prey consumed by
 401 predators determines the strength and range of cascading non-consumptive effects. *PeerJ* 3:
 402 e1426.
- 403 Weissburg, M., Smee, D.L. & Ferner, M.C. (2014). The sensory ecology of nonconsumptive
 404 predator effects. *Am. Nat.* 184: 141–157.
- 405 Welch, J.M., Rittschof, D., Bullock, T.M. & Fordward, R.B. (1997). Effects of chemical cues on
 406 settlement behaviour of blue crab *Callinectes sapidus* postlarvae. *Mar. Ecol. Prog. Ser.* 154:
 407 143–153.
- 408 Zanette, L.Y., White, A.F., Allen, M.C. & Clinchy, M. (2011). Perceived predation risk reduces
 409 the number of offspring songbirds produce per year. *Science* 334: 1398–1401.

Table 1. Results of the ANOVA conducted to evaluate dogwhelk NCEs on mussel recruitment.

Source of variation	df	SS	MS	<i>F</i>	<i>P</i>
Dogwhelks	1	43.86	43.86	7.37	0.020
Blocks	11	401.36	36.49	2.29	0.044
Dogwhelks x Blocks	11	65.50	5.96	0.37	0.954
Residual	24	382.70	15.95

Figure 1. (a) Top view of a cage, showing: (i) the PVC ring determining the cage's shape (25 cm in diameter and 2.5 cm tall), (ii) the peripheral compartment (which had 10 dogwhelks or none, depending on the treatment), (iii) the central compartment with (iv) the mesh scourer to collect mussel recruits, and (v) the four plates used to secure the cage to the intertidal substrate.

(b) Simplified diagram of a cage, showing: (ii) its peripheral compartment, (iii) its central compartment, and (iv) the mesh scourer.

Figure 2. Mussel recruit density (recruits dm⁻², mean ± SE, $n = 24$) depending on the presence or absence of dogwhelks during the field experiment.