**Comparison of gene expression patterns of *Kappaphycus alvarezii* (Rhodophyta, Solieriaceae)
under different light wavelengths and CO₂ enrichment**

Thien Yun Yee, Kenneth Francis Rodrigues, Clemente Michael Wong Vui Ling, Wilson Yong Thau
Lym

Biotechnology Research Institute, Universiti Malaysia Sabah, Jalan UMS, 88400 Kota Kinabalu,
Sabah Malaysia.

E-mail: kennethr@ums.edu.my

Abstract

Transcriptomes associated with the process of photosynthesis and carbon fixation have offered insights into the mechanism of gene regulation in terrestrial plants, however limited information is available as far as macroalgae are concerned. Intertidal red alga, *Kappaphycus alvarezii* is exposed to different wavelengths of light in their lives as light quantity and quality changes at different depths in seawater. This investigation aims to study the underlying mechanisms associated with photosynthesis and carbon fixation under specific light qualities and CO₂ enrichment. Light regulation of gene expression has not been previously described for red algae. By using next generation sequencing, transcriptome profiling of *K. alvarezii* generated 76,871 qualified transcripts with a mean length of 979bp and a N50 length of 1,707bp and 55.83% transcripts were annotated on the basis of function. Blue, green and red light all have demonstrated roles in modulating light responses, such as changes in gene expression. Here we analysed the effects of light regulation on four selected photosynthesis aspects (light-harvesting complex, phycobilisomes, photosystems and photoreceptors). We observed that light-regulated gene expression in this species is not a single light response and different light qualities are transduced to regulate the same metabolic pattern. The carbon fixation pathway was analysed and key genes encoding enzymes involved in the carbon fixation pathway such as ppc, pepc, prk, pgk, ppdk, provided that unequivocal molecular evidence that most of the C₃ and C₄ pathway genes were actively transcribed in *K. alvarezii*. In addition to this the CO₂ induced transcriptome suggested the possibility of shifting carbon metabolism pathway after acclimation to increased level of CO₂. Impact of CO₂ enrichment on the cultures has provided new insight into the response to rising CO₂.

Keywords: carbon fixation, *Kappaphycus alvarezii*, transcriptome sequencing

Introduction

Transcriptome shifts associated by the light signals perceived by photoreceptors have offered insights into the gene regulation in higher plants, however limited information is available as far as macroalgae under light treatment are concerned. Red alga, *Kappaphycus alvarezii* grows in intertidal zone where they may face irradiance environments depleted in both light quantity and quality as the light spectral distribution changes at different depths in seawater. Intertidal seaweeds exposed to light spectrum which is similar as in air during low tide and blue light becomes predominant where the seaweeds grow under seawater of 2-5 m in depth at high tide (Dring, 1981). Seaweeds can acclimate to the changes of light quantity and qualities by employing efficient light-harvesting mechanisms, such as increase the quantity of photosynthetic pigments, change the ratio of accessory pigments to chlorophyll *a* (Lobban and Harrison, 1994). Red alga, *Chondrus crispus* have been reported to have increased phycoerythrin content and the efficiencies of photosynthesis increased with depth (Sagert *et al.*, 1997).

Recently, insights in the light-mediated physiological responses and molecular mechanisms have been gained in brown alga, *Saccharina japonica* (Deng *et al.*, 2012; Wang *et al.*, 2013). Expression profiling of their researches indicate that light induces profound gene expression changes in *S. japonica*. These light responsive genes include many transcription factors and fall into various functional categories mainly involved in photomorphogenesis, circadian clock function, photoreactivation, photosynthetic carbon, metabolism and biosynthesis. Wang *et al.* (2013) suggested that promotion of metabolism and growth in kelps under blue light predominating environment was attributed to both reduction of red light and increase of blue light illumination. Thus, it suggested that different light qualities may play significant roles in the lives of red algae.

The transcriptome of *K. alvarezii* was first sequenced by Wu *et al.* (2014) together with 19 Phaeophyceae (brown algae) and other 20 Rhodophyceae (red algae), provide a broad range of algal transcriptome information that could contribute to algal genetic and biological study. Then, Song *et*

al. (2014) performed de novo transcriptome sequencing of *K. alvarezii* to investigate the mechanisms underlying the biosynthesis of carrageenan in Family Solieriaceae (*Betaphycus gelatinus*, *K. alvarezii* and *Eucheuma denticulatum*). They have identified 861 KEGG orthologs which might contain the main genes regulating the biosynthesis of carrageenan with different types and possessions. More recently, the transcriptome of *K. alvarezii* was profiled by Zhang *et al.* (2015). The results elucidate some genetic information and reveal many important metabolic pathways in *K. alvarezii*. For instance, the significantly enriched pathways include selenocompound metabolism (ko00450) and sulphur metabolism (ko00920) which were probably related to the alga's resistance to grazers and expelling of toxins (Lobanov *et al.*, 2007).

No clear evidence for C₃ or C₄ photosynthetic pathways have been found in the *K. alvarezii*. The occurrence photosynthetic pathways in *K. alvarezii* are therefore still in question and await further research. Many studies reported that the operation of C₃ pathway is predominant in algae (Beer *et al.*, 1986; Tsuji *et al.*, 2009); however, it has been demonstrated that C₄ photosynthesis is found in some eukaryotic algae (Derelle *et al.*, 2006; Leliaert *et al.*, 2012). The study of *Ulva prolifera* and *Chara contraria* on their primary photosynthetic carbon metabolism revealed that some algae possess both C₃ and C₄ pathways and may alter their carbon metabolism pursuant to the environment (Keeley, 1999; Xu *et al.*, 2012). The alterations of photosynthetic pathways under environmental changes probably contributing to the adaptation of plants to environmental stress (Ehleringer *et al.*, 1997). A submerged aquatic plant, *Hydrilla verticillata*, was reported to change its photosynthetic pathways from C₃ to C₄ under conditions of CO₂ deficiency (Reiskind *et al.*, 1997).

In this study, we used next generation sequencing technology to profile the transcriptome of red alga, *K. alvarezii*, with the aim to characterise its functional genome under specific light qualities and CO₂ enrichment. This investigation will enhance our understanding of molecular mechanisms underlying light-induced responses in lower plants as well as facilitate our understanding in inorganic carbon fixation in red algae. In addition, this study would improve our understanding of the impact of

changing light conditions on *K. alvarezii* and the potential mechanism of light adaptation of this red alga.

Materials and methods

Seaweed materials and culture conditions

Seedlings of *Kappaphycus alvarezii* (var. *tambalang* ‘giant’) were obtained from Biotechnology Research Institute (BRI), Universiti Malaysia Sabah. The seaweed was originally collected from Pulau Sebangkat (4°33′ 31″N, 118°39′49″E), Semporna, Sabah. The young seedlings were cultured under laboratory conditions (Yong *et al.* 2014). The experimental samples were selected from healthy and disease-free explants. Three replicates consisted of five algal tips were cut to a total length of 2 cm and used for each experimental condition. The algal tips were cultured in the Fernbach culture vessel with 800 mL artificial seawater (Fluval marine salt, 36 g L⁻¹, salinity 31.4 ppt) enriched with 50 % Provasoli's enriched seawater (PES) media. For light treatment, the cultures grown under 75 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ white light (WL), blue light (BL) (wavelength = 492-455 nm), green light (GL) (wavelength = 577-492 nm) and red light (RL) (wavelength = 780-622 nm), respectively. Cultures were carried out at 25±1.0°C with 18 h light and 6 h dark cycle. Light-emitting diodes (LEDs) were used as light sources. Detected irradiances were measured with a digital light meter (Kyoritsu, Japan). For CO₂ treatment, the cultures were illuminated by white LED, providing an irradiance of 75 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$, on a 18 h light and 6 h dark cycle. CO₂ was provided 500 mL per day and controlled by supplying chambers with air/CO₂ mix. Provision of 500 mL of CO₂ per day was proved to achieve the highest growth rate of *K. alvarezii* (Barat, 2011). To avoid overly acidifying cultures, supplemental CO₂ was only supplied during photoperiod. Temperature was maintained at 25±1.0°C. The cultures from both treatments were collected for RNA extraction after 14 days experimental period.

RNA isolation and preparation of cDNA library

Total RNA was extracted using QIAGEN RNeasy Plant Mini Kit (QIAGEN, Germany) according to manufacturer's protocol. The experimental samples were wash with DEPC-treated water and then quickly frozen with liquid nitrogen and ground to a fine powder using RNase-free, chilled mortar and pestle. Typically, RNA samples with an A260/A280 ratio between 1.8 – 2.0 and A260/A230 ratio between 1.8 – 2.1 and RIN number of 8 and above were recommended to be sufficiently pure for further library construction.

Five cDNA libraries were generated with messenger RNA (mRNA) isolation and cDNA synthesis from five RNA samples were performed using NEBNext Ultra RNA Library Prep Kit for Illumina according to manufacturer's protocol. In brief, the mRNA was purified from total RNA using poly-T oligo-attached magnetic beads and sheared into short segments of about 200 bp. Using the cleaved short RNA fragments as templates, first strand cDNAs were synthesised by random hexamer-primers and reverse transcriptase. The second strand cDNA was synthesised using DNA polymerase I and RNase H. The purified double strand cDNAs were subjected to end repair and NEBNext adaptor ligation. The resulting libraries were then loaded onto a single lane of the flow cell, and 209 cycles on the Illumina HiSeq 2000 platform (Illumina, USA) performed by Malaysian Genomics Resource Centre (MGRC). The high quality reads were deposited at the NCBI Short Read Archive (SRA) with the accession number: SRR2757332 (Green), SRR2757333 (Blue), SRR2757334 (CO₂), SRR2757335 (White) and SRR2757337 (Red).

Quality control and de novo transcriptome assembly

Sequencing reads from the Illumina sequencer were exported in FASTQ format with the corresponding Phred quality scores. First, the quality of the sequencing raw reads was evaluated with FastQC v0.11.2 (www.bioinformatics.babraham.ac.uk/projects/fastqc). Strict reads filtering was performed before assembly: (i) removing sequencing adaptor sequences; (ii) filtering the reads

containing unknown nucleotides (Ns); (iii) removing the reads with more than 10% bases below Q20 sequencing quality.

The obtained clean reads were subjected to transcriptome de novo assembly using Trinity (Grabherr *et al.*, 2011). Briefly, Inchworm first assembles reads by searching for paths in a 25 bp K-mer graph, resulting in a collection of linear contigs. Next, Chrysalis clusters the reads with certain length of overlap and constructs a De Bruijn graph for each cluster. Finally Butterfly reconstructs plausible full-length, linear transcripts by reconciling the individual de Bruijn graphs generated by Chrysalis. After generating final assembly, TGIR Gene Indices clustering tools (TGICL) software (Pertea *et al.*, 2003) was used to remove redundant sequences.

Transcriptome analysis

The assembled sequences were blastx searched against the UniProt protein databases (The UniProt Consortium, 2015) and Kyoto Encyclopaedia of Genes and Genomes (KEGG) database (Kanehisa *et al.*, 2008) with 10^{-5} *E*-value cutoff. The sequences were then functionally clustered using Blast2GO (Conesa *et al.*, 2005) to get GO annotation and further classification by WEGO (Ye *et al.*, 2006) for all assembled sequences.

Identification of differentially expressed genes (DEGs)

RSEM was used to estimate transcripts abundance (Li and Dewey, 2011) and differential expression between two groups was determined by software edgeR (Robinson *et al.*, 2010). In order to analysing differential expression, the transcripts with adjusted *p*-value < 1e-3 and log₂ fold change of two were extracted and clustered according to their patterns of differential expression across the sample. All the DEGs were mapped to the terms in KEGG database and searched for significantly enriched KEGG pathways compared to the whole transcriptome background.

Quantitative real-time PCR analysis

Total RNA was extracted with QIAGEN RNeasy Plant Mini Kit (QIAGEN, Germany). The cDNA synthesis was carried out with QuantiTect Reverse Transcription kit (QIAGEN, Germany). The 18S rDNA gene (U25437) was chosen as internal control for normalisation of real-time PCR data. Primers for amplification of targeted genes (Supplementary Table S1) and 18S rDNA gene were designed using Primer3 software (<http://frodo.wi.mit.edu/primer3/>) aiming for an amplicon length of 80 – 200 bp, a GC content of 40-60%, a primer length of 18 – 30 bp and a primer melting temperature of 60 – 65°C. Primers were assessed for melting temperature, hairpins and primer dimers using the web-based tool Beacon Designer (<http://free.premierbiosoft.com>). The real-time quantitative PCR was performed with QuantiNova SYBR green master mix (QIAGEN, Germany) on the Eco Real-Time PCR System (Illumina). Real-time PCR was performed in volume of 20 µL, and cycling conditions were 95°C for 2 min; 40 cycles of 95°C for 5 s and 60°C for 10 s. The PCR reactions were repeated in triplicates together with negative control and data were analysed using the $2^{-\Delta\Delta CT}$ method.

Results

Transcriptome sequencing and de novo assembly

The sequencing run generated a total of 31.89 Gb raw paired-end reads. After trimming adapters and filtering out low quality reads, over 25 Gb clean reads (~79%) were retained for assembly and further analysis. We generated 76,871 assembled transcripts by Trinity, which have lengths ranging from 300 to 17,257 bp (Fig. 1) and an N50 of 1,707 bp (Table 1). There were no ambiguous bases within the assembled sequences. A large number of the reads (93.51%) aligned back to the transcripts and average read depth of 256.47. The reads that did not map back to the assembled transcripts corresponded to either shorter than 300 bp or failed to match with any clean read.

Figure 1: Graphical length distribution summary of transcripts identified in transcriptome data.

Table 1: Assembly results of *K. alvarezii* transcriptome generated from Trinity.

	Trinity
Total transcripts	76871
total length (bp)	75,239,915
Mean length (bp)	979
N50 length (bp)	1,707
The shortest transcripts (bp)	300
The longest transcripts (bp)	17,257
Transcripts ≥ 500 bp (%)	50.63%
Transcripts ≥ 1000 bp (%)	24.44%
GC (%)	51.17
Annotation rate (%)	55.83

Functional annotation of *K. alvarezii* transcriptome

Blastx was used to map assembled transcripts against the protein databases UniProt, KEGG and COG with a cutoff *E*-value of 10^{-5} . The top hit from each of these assembled transcripts comparisons was used as the annotation reference for the respective transcripts. To sum up, 42,915 transcripts were assigned to putative functions, accounting 55.83% of the total assembled sequences (Supplementary Table S2). With UniProt annotation, 23,701 sequences had perfect matches with *E*-value $<10^{-45}$ and 21.42% of the matches had similarity over 95% (Fig. 2a, b). The species distribution showed that only 13.45% had top matches to *Chondrus crispus* and next top matching species was *Phaeodactylum tricornutum*, accounting for 3.61% (Fig. 2c).

Figure 2: Characteristics of similarity search of the assembled sequences against the Uniprot databases. **a** E value distribution of blast hits for each unique sequence with E value $\leq 10^{-5}$. **b** Similarity distribution of the top blast hits for each sequence. **c** Species distribution of the total homologous sequences with E value $\leq 10^{-5}$. The first hit of each sequence was used for statistics.

There were 45 transcripts encoding light harvesting proteins and 72 gene tags belonging to photosynthetic pigments including chlorophyll, carotenoids (beta- and zeta-carotene), phycobilisome, phycocyanin and phycoerythrin. Three types photoreceptors were identified including one cryptochrome, two blue-light receptor (PHR2) and four phytochrome protein. One phytochrome protein had significant homology to the phytochrome-like protein 2, a photoreceptor which exists in two forms that are reversibly interconvertible by light (Kaneko *et al.*, 1995). In addition, a total of 38 gene tags belonging to photosystem proteins (I and II). The main subunits of PS I, PsaA, PsaB and PsaC, PS I chlorophyll A apoprotein, the Ycf3 and Ycf4 protein domain are identified. Out of 38 photosystem proteins, 23 of them are photosystem II proteins including D1, D2, CP43, CP47, PsbP, PS II reaction center protein M). There were 307 gene tags belonging to protein kinase (PK) family, including serine/threonine PK, calcium-dependent PK and receptor-like PK. A total of 59.93% PKs in *K. alvarezii* were highly homologous to those in *Chondrus crispus*.

To facilitate the organization of the *K. alvarezii* transcripts into putative functional groups, Gene Ontology (GO) terms were assigned using WEGO software. A total of 28,079 annotated transcripts were further categorized to the three main GO domains: biological processes, cellular components and molecular function, including 20,847 sequences at term “biological process”, 14,291 sequences at term “cellular component” and 41,698 sequences at term “molecular function” (Supplementary Fig. S3).

Functional and pathway analysis of the transcripts of *K. alvarezii* were carried out using the KEGG pathway database (Kanehisa *et al.*, 2004). A total of 10,460 transcripts had assigned to 278 KEGG pathways (Table 2). Category “metabolic pathways” had the largest numbers of transcripts among all the categories, followed by the categories of “biosynthesis of secondary metabolites”, “biosynthesis of amino acids” and “carbon metabolism”.

250 Table 2: The top 10 pathways identified in the *Kappaphycus* transcriptome.

Pathway	Number of transcripts	Pathway ID
Metabolic pathways	814	ko01100
Biosynthesis of secondary metabolites	337	ko01110
Biosynthesis of amino acids	131	ko01230
Carbon metabolism	117	ko01200
Ribosome	111	ko03010
Purine metabolism	95	ko00230
Oxidative phosphorylation	94	ko00190
Pyrimidine metabolism	73	ko00240
RNA transport	70	ko03013
Spliceosome	65	ko03040

251

252

253 Both C3 and C4 photosynthesis genes were found in *K. alvarezii* by transcriptome sequencing. In
254 total, 97 transcripts assigned to all the essential enzymes in both C3 and C4 carbon fixations were
255 identified (Supplementary Table S4). Some key genes of enzymes involved in the carbon fixation
256 pathway in *K. alvarezii* were discovered, such as aspartate aminotransferase (ast),
257 phosphoribulokinase (prk), pyruvate orthophosphate dikinase (ppdk), phosphoglycerate kinase (pgk),
258 malate dehydrogenase (mdh), phosphoenolpyruvate carboxylase (ppc) and phosphoenolpyruvate
259 carboxykinase (pck), which provided unequivocal molecular evidence that most of the C₃ and C₄
260 pathway genes were actively transcribed in *K. alvarezii*.

261

262 Identification of DEGs

263 The high quality reads from individual samples were mapped to the transcriptome database with the
264 mapping percentage ranged from 84 to 86% (Table 3). The lower mapping rate reflects the low
265 transcriptome sequencing depth and the high heterozygosity found in *K. alvarezii*.

Table 3: Paired-end read mapping statistics of transcripts identified in transcriptome data.

Sample	WL	BL	GL	RL	C02E*
Total reads pair	30,115,700	22,796,939	27,106,635	23,380,754	24,122,534
Total mapped reads pair	25,435,206	19,493,673	23,083,936	19,978,487	20,784,761
Percentage of mapped reads pair (%)	84.46	85.51	85.16	85.45	86.16

*CO2E indicated CO₂ enriched cultures.

The read normalisation was performed using TMM methods (Robinson and Oshlack, 2011) which implemented in the edgeR Bioconductor package. Setting FDR $\leq 10^{-3}$ and log2 fold change of two as the cutoff, 11,769 DEGs were obtained (Supplementary Table S5). The results were validated by qPCR (data not shown), confirming the accuracy of RNA-Seq in mRNA quantification. A total of 7,962 DEGs (67.65%) were successfully annotated through the sequence similarity searching against the UniProt databases with a significance of E-value $\leq 10^{-5}$.

The most DEGs were found to be down-regulated between every two samples. For the light treatment, the most DEGs were detected between GL and WL, with 390 DEGs upregulated and 908 DEGs downregulated (Supplementary Fig. S5). If mRNA abundance of a gene was similar between BL, GL and WL while higher than RL, the DEG was interpreted to be RL-downregulated. If the expression level of a gene was higher or lower under WL than under both BL, GL and RL, the DEG was suggested to be either BL-, GL- or RL-regulated. Based on the gene expression patterns among the four lights (Table 4), a total of 5,967 DEGs were designated to four categories: BL-regulated, GL-regulated, RL-regulated and either BL- or GL- or RL-regulated (Supplementary Fig. S5). For CO₂ enrichment experiment, a total of 54,938 transcripts were differentially expressed between WL

and CO₂ enrichment. Based on adjusted p -value $<10^{-3}$ and log2 fold change of two, 2,519 DEGs were identified, including 1,125 upregulated and 1,394 downregulated transcripts (Supplementary Fig. S5).

Table 4: Number of transcripts significantly differentially expressed in the four samples.

Condition 1	Condition 2	Up-regulated	Down-regulated	Total	Category
BL	GL+RL+WL	498	54	552	BL-regulated
GL	BL+RL+WL	320	1,656	1,976	GL-regulated
RL	BL+GL+WL	601	177	778	RL-regulated
WL	BL+GL+RL	127	2,534	2,661	Either BL- or GL- or RL-regulated

To better understand the functions of DEGs, all the DEGs were mapped to the terms in KEGG database and compared with whole transcriptome background to search for genes involved in significantly pathways. Pathway enrichment analysis revealed that the annotated DEGs were mainly involved in ‘metabolic pathways’, ‘biosynthesis of secondary metabolites’, ‘oxidative phosphorylation’, ‘carbon metabolism’ and ‘porphyrin and chlorophyll metabolism’. The annotation shed light on the regulatory functions of light qualities on the specific processes, functions and pathways in *K. alvarezii*.

Discussion

Intertidal seaweeds are subjected to rapidly changing environmental conditions. As a result, they have developed mechanisms that help them to cope with unfavourable disturbances or environmental stresses. These adaptation and acclimation responses contribute significantly to their survival and fitness, providing them with the plasticity necessary for a stationary life. This investigation highlights the understanding of gene expressions and responses of red alga *K. alvarezii* towards their light environment (BL, GL, RL and WL) and increasing CO₂ level.

Photosynthesis begins with light absorbing, thus light harvesting is the first step in the photosynthetic process which mediated by pigment-binding proteins forming light-harvesting antenna systems. Light-harvesting complex (LHC) proteins constitute a large family of proteins, which includes chlorophyll *a/b*-binding proteins, fucoxanthin chlorophyll *a/c*-binding proteins, high light-induced proteins and early light-induced proteins (Green and Kuhlbrandt, 1995; Caron *et al.*, 2001). Chlorophyll *a/b*-binding and fucoxanthin chlorophyll *a/c*-binding proteins were frequently reported to be transcriptionally repressed in response to light stress (Tonkyn *et al.*, 1992; Maxwell *et al.*, 1995; Teramoto *et al.*, 2002). In this study, the amount of genes encoding light-harvesting proteins transcripts were significantly downregulated under different light qualities (BL, GL and RL) (Supplementary S5) and no significance difference between control and CO₂ enrichment. Fucoxanthin chlorophyll *a/c*-binding and early light-induced proteins were upregulated under GL. Meanwhile, similar responses of *K. alvarezii* were found under BL and RL that both fucoxanthin chlorophyll *a/c*-binding and high light-induced proteins were downregulated. Recent transcriptomic studies of stress responses highlighted fucoxanthin chlorophyll *a/c*-binding proteins were upregulated in responses to heat-, salt-, oxidative-, or light stress in both brown algae (Hwang *et al.*, 2008; Dittami *et al.*, 2009; Pearson *et al.*, 2010). Similar observations were made concerning chlorophyll *a/b*-binding and fucoxanthin chlorophyll *a/c* -binding proteins in *Chlamydomonas reinhardtii* after high light treatment (Savard *et al.*, 1996; Miura *et al.*, 2004). It is reported that there is possible additional functions of these proteins. For example, fucoxanthin chlorophyll *a/c*-binding proteins were observed to be downregulated in a developmental mutant of the brown alga *Ectocarpus siliculosus* (Peters *et al.*, 2008). The recently discovered “red lineage chlorophyll *a/b*-binding-like proteins” (RedCAPs) by Sturm *et al.* (2013) which was found to be restricted to the red algal lineage, reported to participate in the light (intensity- and quality-) dependent structural remodelling of light-harvesting antennae in the red lineage. However, the existence and expression of RedCAPs in response to different light qualities and CO₂ level in *K. alvarezii* remains unknown.

Chlorophyll and carotenoids in red algae is not that of a primary light absorber, this role is taken over by the phycobilisomes. Phycobilisome functions as primary light absorber in red algae and funnel the energy to the reaction center of PS II for conversion into chemical forms (Gantt *et al.*, 2003). This complex contains two or three types of pigment-proteins known as biliproteins: phycoerythrin, phycocyanin and allophycocyanin. They differ in protein identity, chromophore type and attachment and their relative location in the phycobilisome complex (Blankenship, 2002). The phycobilisome absorbs light across the between 590-650 nm region of the solar spectrum that neither chlorophyll nor carotenoids are capable of absorbing light. Thus, organisms with phycobilisomes have greater accessibility to usable light within the visible spectrum and hence greater adaptability and light capturing capacity (Blankenship, 2002). Although RL had the significant impact on the growth rate of *K. alvarezii* as compared to those treated with BL and GL, however, the phycobilisomes showed no significant different expressions among the four light qualities. Different light spectral seemed to induce the same effect on phycobilisome contents in *K. alvarezii*. In contrast, many experiments (Figueroa and Niell 1990; Talarico and Cortese, 1993; Franklin *et al.*, 2002; Godinez-Ortega *et al.*, 2008) observed that the synthesis of chlorophyll, phycoerythrin, phycocyanin in the red algae have been shown to be influence by both irradiance and spectral composition. Such changes do not always imply statistical significance in differential expression analysis in transcriptome data as the differential analysis takes into account of overall depth of sequencing, read and fragment length, gene density in the genome, transcriptome splicing complexity and transcript abundance that could be vary for each sample (Trapnell *et al.*, 2012). On the other hand, three DEGs tag to phycobilisome were identified between control and cultures treated with CO₂ enrichment. Phycobilisome and phycoerythrin were upregulated while phycocyanin was downregulated under CO₂ enrichment. The upregulation of photosystem proteins and phycobilisomes suggested that photosynthesis efficiency of *K. alvarezii* was enhanced by raising CO₂ level.

The presence study has identified 38 gene tags of photosystems proteins (15 PS I and 23 PS II). Photosystems are functional and structural units of protein complexes involved in photosynthesis.

The photosystem proteins showed no significant different expressions among the four light qualities. The most had the highest mRNA levels under RL (33) and the lowest under BL (18). It can be said that RL is more efficient than BL, GL and WL in term of photosynthetic efficiency in *K. alvarezii*. Meanwhile, out of 38 transcripts of photosystem proteins, 9 genes encoding of photosystem proteins were upregulated by raising CO₂ concentrations, including 4 PS I proteins (ycf4, subunit III, reaction center subunit XI and chlorophyll A apoprotein) and 5 PS II proteins (D1, CP43 and CP47). These findings exhibited the importance of CO₂ enhancing the quantum efficiency of photosystem I and II. Interaction between elevated CO₂ levels and responses of seaweeds are still largely unknown. The contradictory results were obtained on growth and photosynthetic rates under CO₂ enrichment (Gao *et al.*, 1991, 1993; Garcia-Sanchez *et al.*, 1994; Israel and Hophy, 2002; Zou, 2005). It is suggested that the functional PS II is increased under CO₂ enrichment, therefore, in order to balance the stoichiometry of PS I and PS II, genes encoding subunits of PS I are simultaneously highly expressed to increase amount of PS I, improving the tolerance of plant to high CO₂ levels.

Light-regulated gene expression mediated by photoreceptors has been study in plants, including algae (Ma *et al.*, 2001; Jiao *et al.*, 2005; see review Kianianmomeni and Hallman, 2014). We have identified two types of photoreceptors in *K. alvarezii*: phytochromes and cryptochromes. Phytochromes are red/far-red photoreceptors, measure the changes in light quality in the red and far-red regions of the visible spectrum, allowing plants to assess the quantity of photosynthetically active light (Franklin and Whitelam, 2005); cryptochromes are flavoproteins, function as blue light receptors that share sequence similarity to DNA photolyases, DNA enzymes that use blue light to repair UV-induced (Sancar, 2003). In the present study, phytochrome-like protein and cryptochrome transcripts were observed expressed under WL. The phytochrome gene is not expressed under BL, GL and RL. Thus we suggest that the expression of phytochrome genes in *K. alvarezii* seems to be triggered by others light qualities. In *Arabidopsis thaliana*, the expression of all photoreceptors genes has been demonstrated to be regulated by light or circadian clock (Reka, 2003). Similarly in brown alga *Saccharina japonica*, expression of photoreceptor genes seems to be triggered by all light

qualities (BL, RL and WL) (Wang *et al.*, 2013). In contrast, no phytochrome genes could be identified in the genomes of green alga *Volvox carteri* and *Chlamydomonas reinhardtii*, even though red- and far-red-regulated gene expression has been observed in these algae (Alizadeh and Cohen, 2010; Beel *et al.*, 2012). The absence of phytochromes might indicate that other phytochrome photoreceptors such as animal-like cryptochromes, which absorb both blue and red light (Beel *et al.*, 2012). On the other hand, the expression of cryptochrome genes was highly expressed in all light qualities and CO₂ enrichment except under RL. Danon *et al.* (2006) and Lopez *et al.* (2011) reported that cryptochrome in plants are involved in the light-dependent gene expression, the light dependence mainly affect genes involved in the response to biotic/abiotic stress and regulation of photosynthesis. The presence of cryptochrome under GL is not surprising as cryptochromes are able to process GL, however, less effectively compared to BL (Folta and Maruhnich, 2007).

All the genes necessary to encode the enzyme involved in photosynthetic inorganic carbon fixation were identified (Supplementary Table S4). The data allow the identification of all enzymes necessary for the reductive pentose phosphate cycle, glyceraldehyde-3P => ribulose-5P (C₃) and the C₄-dicarboxylic acid cycle, phosphoenolpyruvate carboxykinase type (PCK) carbon fixation pathway, indicated the possibility of the existence of two photosynthetic pathways (C₃ and C₄) in *K. alvarezii*. Some experimental results suggest that the PCK-type is maximal in biomass production and CO₂ fixation (Fravolini *et al.*, 2002; Wang *et al.*, 2012). Recent papers have reported the evidence for the operation of C₄ photosynthesis as an alternative inorganic carbon-concentrating mechanism (CCM) in marine organisms (Tachibana *et al.*, 2011). The expression levels of C₄ enzymes, such as mdh, ast, ppc, pck and ppdk were higher than C₃ enzymes, indicate C₄ photosynthesis may function under atmospheric CO₂ level. Most of the key enzymes of C₃ such as glyceraldehyde 3-phosphate dehydrogenase (gapdh), pgk, and fructose-bisphosphate aldolase (fba) were expressed higher under CO₂ enrichment (Supplementary Table S6), however the difference is not significant. These results suggested CO₂ enhancement may alter carbon metabolism and lead to C₃-type carbon metabolism in *K. alvarezii*. As CO₂ level increase, the stomata plants increasingly close up and thus reduce the

amount of water lost. In other words, plants significantly improve the water use efficiency as less water is transpired. Increase in the level of CO₂, C₄ plants show drastically reduced rates of photorespiration because CO₂ is concentrated at the carboxylation site (RuBisCo) and is able to out compete molecular oxygen (Weber and von Caemmerer, 2010). The elimination of oxygenation reaction to a great extent and the loss of energy connected with the photorespiratory pathway is largely decreased (Heldt and Piechulla, 2011). For these reasons, the photosynthetic efficiency gap between C₃ and C₄ rapidly closes. As competing for the resources (water, nutrients and light) on the same patch, C₃ is increasing outcompete the C₄ metabolism. However, current understanding of the underlying mechanisms of the response of red algae to elevated CO₂ concentrations is largely unknown; the demand of response of red algae to increasing CO₂ warrants further research.

Conclusion

To summarise, a total of 76,871 sequences were assembled using Trinity and 42,915 transcripts were assigned to functional annotation. 55.83% of transcripts were annotated through UniProt, provided important clues to the functions of many unknown sequences that could not be annotated using sequence comparison. Differential expression analysis revealed that only a small portion of genes (15.31%) were significantly expressed. The most DEGs were found under CO₂ enrichment. Three different wavelengths were used: BL, GL and RL and compared to full spectrum (WL), similar gene expression patterns were found in term of photosynthesis aspect. The data presented suggesting light-regulated gene expression in *K. alvarezii* is not a single light response. On the other hand, CO₂ enrichment leads to changes of photosynthetic pathway from C₄ to C₃ may contribute to the adaptation of this species to global CO₂ changes. The alteration of photosynthetic pathways under CO₂ enrichment highlights new questions to be addressed in subsequent work. This transcriptome study resulted in a number of new insights regarding the regulation of genes respond to different wavelengths of light and addition of CO₂.

Acknowledgements

The authors would like to thank the Ministry of Higher Education for granting a Fundamental Research Grant Scheme (FRGS) entitled “**Transcriptomic and carrageenan analyses of micropropagated *Kappaphycus* and *Eucheuma* seaweeds under different light regimes and carbon dioxide concentrations**”.

References

- Blankenship, R.E. 2002. Molecular mechanisms of photosynthesis. Blackwell Science, Oxford.
- Heldt, H.W., Piechulla, B. 2011. Plant biochemistry. Academic Press, Elsevier.
- Barat, W. O. B. (2011). Utilization of carbon dioxide for the optimization of the growth of seaweed *kappaphycus alvarezii*. UT-Aquatic product technology.
<http://repository.ipb.ac.id/handle/123456789/52986>
- Conesa, A., Gotz, S., Garcia-Gomez, J.M., Terol, J., Talon, M., Robles, M. 2005. Blast2GO: a universal tool for annotation, visualisation and analysis in functional genomics research. *Bioinformatics* **21**: 3674-3676.
- Weber, A.P.M., von Caemmerer, S. 2010. Plastid transport and metabolism of C3 and C4 plants – comparative analysis and possible biotechnological exploitation. *Current Opinion Plant Biology* 13:257-265.
- Franklin, K.A., Whitelam, G.C. 2005. Phytochromes and shade-avoidance responses in plants. *Annals of Botany* 96:169-175.
- Folta, K.M., Maruhnich, S.A. 2007. Green light: a signal to slow down or stop. *Journal of Experimental Botany* 58:3099-3111.

468

469 Lopez L, Carbone F, Bianco L, Giuliano G, Facella P, Perrotta G (2011) Tomato plants

470 overexpressing cryptochrome 2 reveals altered expression of energy and stress related gene products

471 in response to diurnal cues. *Plant, Cell Environ* 35:994–1012

472

473 Danon A, Coll NS, Apel K (2006) Cryptochrome-1-dependent execution of programmed cell death

474 induced by singlet oxygen in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 103:17036–17041

475

476 Sancar A. Structure and function of DNA photolyase and cryptochrome blue-light

477 photoreceptors. *Chem Rev.* 2003;103:2203–2237. doi: 10.1021/cr0204348.

478

479 Ma, L., Li, J., Qu, L., Hager, J., Chen, Z., Zhao, H., Deng, X.W. 2001. Light control of Arabidopsis

480 development entails coordinated regulation of genome expression and cellular pathway. *Plant Cell*

481 13:2589-2608.

482

483 Jiao, Y., Ma, L., Strickland, E., Deng, X.W. 2005. Conservation and divergence of light-regulated

484 genome expression patterns during seedling development in rice and Arabidopsis. *Plant Cell*

485 17:3239-3256.

486

487 Kianianmomeni, A., Hallman, A. 2014. Algal photoreceptors: in vivo functions and potential

488 applications. *Planta* 239:1-26.

489

490 Alizadeh D, Cohen A (2010) Red light and calmodulin regulate the expression of the *psbA* binding

491 protein genes in *Chlamydomonas reinhardtii*. *Plant Cell Physiol* 51:312–322

492

Beel B, Prager K, Spexard M, Sasso S, Weiss D, Müller N, Heinnickel M, Dewez D, Ikoma D,
Grossman AR, Kottke T, Mittag M (2012) A flavin binding cryptochrome photoreceptor responds to
both blue and red light in *Chlamydomonas reinhardtii*. *Plant Cell* 24:2992–3008

Reka, T. 2003. Analysis of regulatory function of circadian clock on photoreceptor gene expression.
Ph. D thesis. Biological Research Center of the Hungarian Academy Sciences, Hungary.

Fravolini A, Williams DG, Thompson TL (2002) Carbon isotope discrimination and bundle sheath
leakiness in three C(4) subtypes grown under variable nitrogen, water and atmospheric CO(2) supply. *J*
Exp Bot 53(378): 2261–2269.

Wang C, Guo L, Li Y, Wang Z (2012) Systematic comparison of c3 and c4 plants based on metabolic
network analysis. *BMC Syst Biol* 2: S9.

Grabherr, M.G., Haas, B.J., Yassour, M., et al. 2011. Full length transcriptome assembly from
RNA-Seq data without a reference genome. *Nature Biotechnology* **29**: 644-652.

Kanehisa, M., Araki, M., Goto, S., et al. 2008. KEGG for linking genomes to life and the environment.
Nucleic Acids Research **36** (database issue): D480-D484.

Tachibana M, Allen AE, Kikutani S, Endo Y, Bowler C (2011) Localization of putative carbonic
anhydrases in two marine diatoms, *Phaeodactylum tricornutum* and *Thalassiosira*
pseudonana. *Photosynth Res* 109: 205–221.

Ma, L., Li, J., Qu, L., Hager, J., Chen, Z., Zhao, H., Deng, X.W. 2001. Light control of *Arabidopsis*
development entails coordinated regulation of genome expression and cellular pathways. *Plant Cell* **13**:
2589-2608.

Robinson, M.D., Oshlack, A. 2010. A scaling normalisation method for differential expression analysis of RNA-seq data. *Genome Biology* 11:R25.

Sun, L., Wang, S., Zhao, M., Fu, X., Gong, X., et al. 2009 Phycobilisomes from cyanobacteria. In: Gault, P.M., Maler, H.J. (editors) *Handbook on Cyanobacteria: Biochemistry, Biotechnology and Application*. New York: Nova Science Publishers. 105–160.

The UniProt Consortium. 2015. UniProt: a hub for protein information. *Nuclei Acids Research* **43**: D204-D212.

Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D. R., et al. 2012. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nature Protocols* 7: 562-578.

Ye, J., Ye, J., Fang, L., Zheng, H., Zheng, H., et al. 2006: WEGO: a web tool for plotting GO annotations. *Nucleic Acids Research* **34** (web server issue): W293-W297.

Yong, W.T.L., Ting, S.H., Yong, Y.S., Thien, V.Y., Wong, S.H., Chin, W.L., Rodrigues, K.F. and Anton, A. 2014. Optimization of culture conditions for the direct regeneration of *Kappaphycus alvarezii* (Rhodophyta, Solieriaceae). *Journal of Applied Phycology* **26**: 1597-1606.

Tonkyn, J.C., Deng, X.W., Grussem, W. 1992. Regulation of plastid gene expression during photooxidative stress. *Plant Physiology* 99:1406-1415.

Maxwell, D.P., Laudenbach, D.E., Huner, N. 1995. Redox regulation of light-harvesting complex II and CAB mRNA abundance in *Dunaliella salina*. *Plant Physiology* 109:787-795.

547

548 Teramoto, H., Nakamori, A., Minagawa, J., T-aki Ono. 2002. Light-intensity-dependent expression of

549 Lhc gene family encoding light-harvesting chlorophyll a/b proteins of photosystem II in

550 *Chlamydomonas reinhardtii*. *Plant Physiology* 130:325-333.

551

552 Peters, A.F., Scornet, D., Ratin, M., Charrier, B., Monnier, A., Merrien, Y., Corre, E., Coelho, S.M.,

553 Cock, J.M. 2008. Life-cycle-generation –specific developmental processes are modified in the

554 immediate upright mutant of the brown alga *Ectocarpus siliculosus*. *Development* 135:1503-1512.

555

556 Hwang, Y.S., Jung, G., Jin, E. 2008. Transcriptomic analysis of acclamatory responses to thermal

557 stress in Antarctic algae. *Biochem Biophys Res Comm* 367:635-641.

558

559 Dittami, S.M., Scornet, D., Petit, J.L., Ségurens, B., Da Silva C, Corre E, Dondrup M, Glatting K-H,

560 König R, Sterck L, *et al.* 2009. Global expression analysis of the brown alga *Ectocarpus*

561 *siliculosus* (Phaeophyceae) reveals large-scale reprogramming of the transcriptome in response to

562 abiotic stress. *Genome* 10:R66.

563

564 Pearson GA, Hoarau G, Lago-Leston A, Coyer James A, Kube M, Reinhardt R, Henckel K, Serrão

565 ETA, Corre E, Olsen JL: An expressed sequence tag analysis of the intertidal brown seaweeds *Fucus*

566 *serratus* (L.) and *F. vesiculosus* (L.) (Heterokontophyta, Phaeophyceae) in response to abiotic

567 stressors. *Mar Biotechnol* 2010, **12**:195–213.

568

569 Savard F, Richard C, Guertin M: The *Chlamydomonas reinhardtii* LI818 gene represents a distant

570 relative of the cabI/II genes that is regulated during the cell cycle and in response to illumination. *Plant*

571 *Mol Biol* 1996, **32**:461–473.

572

Miura K, Yamano T, Yoshioka S, Kohinata T, Inoue Y, Taniguchi F, Asamizu E, Nakamura Y, Tabata SY, Katsuyuki T, *et al.*: Expression profiling-based identification of CO₂-responsive genes regulated by CCM1 controlling a carbon-concentrating mechanism in *Chlamydomonas reinhardtii*. *Plant Physiol* 2004, **135**:1595–1607.

Green BR, Kühlbrandt W: Sequence conservation of light-harvesting and stress-response proteins in relation to the three-dimensional molecular structure of LHCII. *Photosynth Res* 1995, **44**:139–148.

Caron L, Douady D, de Martino A, Quinet M: Light harvesting in brown algae. *Cah Biol Mar* 2001, **42**:109–124.

Dring, M.J. 1981. Chromatic adaptation of photosynthesis in benthic marine algae: an examination of its ecological significance using a theoretical model. *Limnol Oceanography* 26:271-284.

Lobban C.S. & Harrison P.J. (1994) *Seaweed ecology and physiology*. Cambridge University Press, Cambridge.

Sagert, S., Forster, R.M., Feuerpfeil, P., Schubert, H. 1997. Daily course of photosynthesis and photoinhibition in *Chondrus crispus* (Rhodophyta) from different shore levels. *European Journal of Phycology* 32:363-371.

Deng, Y., Yao, J., Wang, X., Guo, H. & Duan, D. 2012. Transcriptome sequencing and comparative analysis of *Saccharina japonica* (Laminariales, Phaeophyceae) under blue light induction. *PLoS ONE* 7:e39704.

Wang, W.J., Wang, F.J., Sun, X.T., Liu, F.L. & Liang, Z.R. 2013. Comparison of transcriptome under red and blue light culture of *Saccharina japonica* (Phaeophyceae). *Planta* 237:1123-1133.

600

601 Wu, S.X., Sun, J., Chi, S., Wang, L., Wang, X.M., Liu, C., Li, X., Yin, L., Liu, T., Yu, J. 2014.

602 Transcriptome sequencing of essential marine brown and red algal species in China and its significance

603 in algal biology and phylogeny. Acta Oceanologica Sinica 33:1-12.

604

605 Song, L., Wu, S.X., Sun, J., Wang, L., Liu, T., Chi, S., Liu, C., Li, X., Yin, L., Wang, X., Yu, J. 2014.

606 De novo sequencing and comparative analysis of three red algal species of family Solieriaceae to

607 discover putative genes associated with carrageenan biosynthesis. Acta Oceanologica Sinica 33:45-53.

608

609 Zhen Zhang, Tong Pang, Qianqian Li, Litao Zhang, Ling Li & Jianguo Liu (2015) Transcriptome

610 sequencing and characterization for *Kappaphycus alvarezii*, European Journal of Phycology, 50:4,

611 400-407

612

613 Lobanov, A., Fomenko, D., Zhang, Y., Sengupta, A., Hatfield, D. & Gladyshev, V. (2007).

614 Evolutionary dynamics of eukaryotic selenoproteomes: large selenoproteomes may associate with

615 aquatic life and small with terrestrial life. Genome Biology, 8:1 –16.

616

617 Tsuji Y, Suzuki I, Shiraiwa Y (2009) Photosynthetic carbon assimilation in the

618 coccolithophorid *Emiliania huxleyi* (Haptophyta): evidence for the predominant operation of the

619 C₃ cycle and the contribution of β -carboxylases to the active anaplerotic reaction. Plant Cell Physiol

620 50: 318–329.

621

622 Beer S, Israel A (1986) Photosynthesis of *Ulva* sp. III. O₂ effects, carboxylase activities, and the

623 CO₂ incorporation pattern. Plant Physiol 81: 937–938.

624

Derelle E, Ferraz C, Rombauts S, Rouze P, Worden AZ, et al. (2006) Genome analysis of the smallest free-living eukaryote *Ostreococcus tauri* unveils many unique features. Proc Natl Acad Sci USA 103: 11647–11652.

Leliaert F, Smith DR, Moreau H, Herron MD, Verbruggen H, et al. (2012) Phylogeny and molecular evolution of the green algae. Crit Rev Plant Sci 31: 1–46.

Xu, J., Fan, X, Zhang, X., Xu, D., Mou, S., Cao, S., Zheng, Z, Miao, J., Ye, N. 2012. Evidence of coexistence of C3 and C4 photosynthetic pathways in green-tide-forming alga, *Ulva prolifera*. PLoS ONE 7(5):e37438.

Keeley, J.E. 1999. Photosynthetic pathway diversity in a seasonal pool community. Funct Ecol 13:106-118.

Ehleringer, J.R., Cerling, T.E., Helliker, B.R. 1997. C4 photosynthesis, atmospheric CO2 and climate. Oecologia 112:285-299.

Reiskind, J.B., Madsen, T.V., Vanginkel, L.C., Bowes, G. 1997. Evidence that inducible C4 type photosynthesis is a chloroplastic CO2-concentrating mechanism in *Hydrilla*, a submersed monocot. Plant, Cell and Environment 20:211-220.

Pertea, G., Huang, X., Liang, F., Antonescu, V., Sultana, R., Karamycheva, S., et al. 2003. TIGR Gene Indices clustering tools (TGICL): a software system for fast clustering of large EST datasets. Bioinformatics 19:651-652.

Li B, Dewey CN (2011) RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics 12: 1–16. doi: 10.1186/1471-2105-12-323

652

653 Sturm, S., Engelken, J., Gruber, A., Vugrinec, S., Kroth, P.G, Adamska, I., Lavaud, J. 2013. A novel

654 type of light-harvesting antenna protein of red algal origin in algae with secondary plastids. BMC

655 Evolutionary Biology 13:159.

656

657 Gantt, E., Grabowski, B., Cunningham Jr., F.X. 2003. Antenna systems of red algae: phycobilisomes

658 with photosystem II and chlorophyll complexes with photosystem I. In Green, B.R., Parson, W.W.

659 (editors). Light harvesting antennas in photosynthesis. Springer Netherlands.

660

661 F. Lopez-Figueroa and X. Niell, Effects of light quality of chlorophyll and biliprotein accumulation

662 on seaweeds, *Mar. Biol.*, 104 (1990) 321-327.

663

664 Godinez-Ortega, J.L., Snoeijs, P., Freile-Pelegrin, Y., Pedersen, M. 2008. Growth and pigment

665 composition in the red alga *Halymenia floresii* cultured under different light qualities. Journal of

666 Applied Phycology 20:253-260.

667

668 Talarico L, Cortese A (1993) Response of *Audouinella-saviana* (Meneghini) Woelkerling

669 (Nemaliales, Rhodophyta) cultures to monochromatic light. Hydrobiologia 261:477–484

670

671 Franklin LA, Krâbs G, Kuhlenkamp R (2002) Blue light and UV radiation control the synthesis of

672 mycosporine-like amino acids in *Chondrus crispus* (Florideophyceae). J Phycol 37:257–270.

673

674 Zou D (2005) Effects of elevated atmospheric CO₂ on growth, photosynthesis and nitrogen

675 metabolism in the economic brown seaweed, *Hizikia fusiforme* (Sargassaceae, Phaeophyta).

676 Aquaculture 250:726–735

677

678 Israel A, Hophy M (2002) Growth, photosynthetic properties and Rubisco activities and amounts of
679 marine macroalgae grown under current and elevated seawater CO₂ concentrations. Glob Chang Biol
680 8:831–840
681
682 Garcia-Sanchez MJ, Fernandez JA, Niell FX (1994) Effect of inorganic carbon supply on the
683 photosynthetic physiology of *Gracilaria tenuistipitata*. Planta 194:55–61
684
685 Gao K, Aruga Y, Asada K, Ishihara T, Akano T, Kiyohara M (1991) Enhanced growth of the red alga
686 *Porphyra yezoensis* Ueda in high CO₂ concentrations. J Appl Phycol 3:355–362
687
688 Gao K, Aruga Y, Asada K, Kiyohara M (1993) Influence of enhanced CO₂ on growth and
689 photosynthesis of the red algae *Gracilaria* sp. and *Gracilaria chilensis*. J Appl Phycol 5:563–571.
690
691 Kaneko T., Tanaka A., Sato S., Kotani H., Sazuka T., Miyajima N., Sugiura M., Tabata S. 1995.
692 Sequence analysis of the genome of the unicellular cyanobacterium *Synechocystis* sp. strain PCC6803.
693 I. Sequence features in the 1 Mb region from map positions 64% to 92% of the genome. DNA Res
694 2:153-166.