1 ***NAC-NOR* mutations in tomato Penjar accessions attenuate multiple**
2 **metabolic processes and prolong the fruit shelf life**

3 Rakesh Kumar, Vajir Tamboli, Rameshwar Sharma, Yellamaraju Sreelakshmi*
4 Repository of Tomato Genomics Resources, Department of Plant Sciences, University of
5 Hyderabad, Hyderabad-500046, India

6 ***Corresponding author**

7
8 **Email addresses of authors:**

9 Rakesh Kumar: rakeshgupta.hcu@gmail.com

10 Vajir Tamboli: vajirchem@gmail.com

11 Rameshwar Sharma: rameshwar.sharma@gmail.com

12 Yellamaraju Sreelakshmi: syellamaraju@gmail.com; Phone: 0091-40-23134771

13

14 **Running head:** Metabolite analysis of Penjar accessions

15 Word count: 5741

16 No. of Tables: 1

17 No. of Figures: 6

18 Supplementary Information: Figures S1-S16; Supplementary Tables S1-S6.

19

20 **Highlight**

21 The prolonged shelf life of tomato Penjar accessions bearing mutations in NAC-NOR
22 transcription factor appears to be regulated by a combined effect of attenuation of respiration,
23 altered cuticle composition, enhanced ABA and sucrose levels in fruits and downregulation
24 of cell wall modification.

25 **Abstract**

26 Several Penjar accessions of tomato (*Solanum lycopersicum*), widely grown in the
27 Mediterranean region, exhibit prolonged shelf life, and harbor *alcobaca* mutation with valine-
28 106-aspartic acid substitution in the NAC-NOR protein. To uncover the metabolic basis
29 underlying the prolonged shelf life, we compared four Penjar accessions to Ailsa Craig (AC).
30 Three accessions bore *alcobaca* mutation, whereas fourth was a novel *NAC-NOR* allele with
31 only six amino acids in the encoded protein. The cuticle composition among Penjars varied
32 widely during the progression of fruit ripening. All Penjars exhibited delayed ripening,
33 prolonged on-vine and off-vine shelf life, low ethylene emission and carotenoid levels albeit
34 with accession-specific differences. Metabolic profiling revealed shifts in Krebs cycle
35 intermediates, amino acids, and γ -aminobutyric acid levels indicating the attenuation of
36 respiration in Penjars during post-harvest storage. The prolonged shelf life of Penjar fruits
37 was associated with a concerted downregulation of a number of cell-wall modifying genes
38 and cell-wall-related metabolites. The accumulation of higher ABA and sucrose levels at the
39 onset of senescence in Penjar fruits likely contribute to reduced water loss. Our analyses
40 reveal that in addition to specialized cuticle composition, the attenuation of various metabolic
41 processes by *NAC-NOR* mutation likely prolongs the shelf life of Penjar fruits.

42

43 **Key words:** Tomato (*Solanum lycopersicum*), fruit shelf life, NAC-NOR, metabolite
44 analysis, cuticle composition, hormones, cell wall modification.

45

46

Introduction

Among the factors that influence the economic value of fruits, the post-harvest shelf life is the foremost, as shorter shelf life causes losses during transportation, distribution, and storage. Consistent efforts have been made over last few decades to decipher the molecular-genetic basis underlying the process of fruit ripening and spoilage (Klee and Giovannoni, 2011; Pirrello, 2009; Seymour et al. 2013). Genetic regulation of fruit ripening is mainly deciphered in tomato, where spontaneous mutants such as *rin*, *nor* and *Cnr*, fail to undergo characteristic accumulation of lycopene in the fruits and remain firm for a long time (Giovannoni, 2007). The *RIN*, *CNR* and *NOR* genes encode for transcription factors belonging to MADS, SBP box, and NAC family respectively (Seymour et al. 2013).

The *nor* mutant family has two alleles; *nor* with truncated protein, and *alcobaca* (*alc*) with a single amino acid substitution (Casals et al. 2012; Giovannoni, 2004). The cultivars harboring either of these alleles show prolonged shelf life of fruits, albeit duration depends on the fruit size and genetic background (Garg et al. 2008a,b; Casals et al. 2012; Mutschler et al. 1988). The biochemical characterization of *nor/alc* mutants revealed that compared to normal ripening cultivars, the *nor/alc* fruits had altered cuticle composition with denser cutin matrix (Kosma et al. 2010; Saladié et al. 2007), were deficient in carotenoids accumulation, and were resistant to infections (Garg et al. 2008a,b). As *NOR* mutation is strongly associated with long shelf life, search for additional NAC family transcription factors revealed a role for *NAC1* and *NAC4* in tomato ripening (Ma et al. 2014; Zhu et al. 2014).

The onset of ripening in tomato initiates several irreversible processes such as changes in metabolite composition, loss of chlorophyll, accumulation of carotenoids and cell wall softening (Carrari et al. 2006; Fraser et al. 2007; Giovannoni, 2007). These responses are triggered by the climacteric rise of ethylene in the fruits, a process that is subdued in the *nor* mutants. Current evidences indicate that in addition to ethylene, other hormones like abscisic acid (ABA), jasmonic acid (JA), methyl jasmonate (MeJA), auxin, etc. also affect fruit ripening and quality (Kumar et al. 2014; McAtee et al. 2013). It is believed that *NOR* acts upstream to ethylene in regulating the ripening of tomato (Giovannoni, 2007). However, it is not known whether *NOR* influences the levels of other hormones during ripening.

The studies using transgenic manipulation of genes revealed that softening of tomato fruits can be prevented by silencing of ethylene biosynthesis (Oeller et al. 1991; Xie et al. 2006) and perception (Tieman et al. 2000), increasing polyamine biosynthesis (Nambeesan et al. 2010), suppression of ABA biosynthesis (Sun et al. 2012), and downregulation of enzymes participating in cell wall degradation (Brummell et al. 2002; Vicente et al. 2007;

81 Uluisik et al. 2016). The transpirational water loss through cuticle also affects the fruit
82 firmness. Thus fruits that retain cellular turgidity remain firm for longer periods (Saladié et
83 al. 2007). From the foregoing, it is apparent that a combination of multiple processes likely
84 regulates prolonged shelf life of *nor/alc* mutants. However, only limited information is
85 available about modulation of these processes in the *nor/alc* mutants (Casals et al. 2015;
86 Saladie et al. 2007, Osorio et al. 2011).

87 Breeders have used *alc* for better fruit quality, though hybrid fruits have a slightly
88 shorter shelf life than *rin* or *nor* hybrids (Garg et al. 2008a,b). Among the landraces grown in
89 Mediterranean, the Penjars are popular for long post-harvest shelf life. Several Penjar
90 accessions bear the *alc* mutation (Casals et al. 2012), however, little is known about the
91 metabolic profiles of fruits during postharvest storage. In the current study, we analyzed four
92 Penjar accessions during ripening and post-harvest storage to decipher the metabolic basis for
93 the prolonged shelf life.

94

95 **Materials and Methods**

96 **Plant material and post-harvest analysis**

97 Penjar accessions (a gift from RF Muñoz) and tomato (*Solanum lycopersicum*)
 98 cultivar Ailsa Craig were grown in pots between October to February (day 30±2°C, night
 99 18±2°C) season, which is similar to Mediterranean summer. The fruits were collected at
 100 mature green (MG), breaker (BR) and red ripe (RR) stages. For Penjars, attainment of
 101 uniform fruit coloration was considered equivalent to RR stage. The pericarps were snap-
 102 frozen in liquid nitrogen and stored at -80°C until analysis. Freshly harvested RR fruits were
 103 incubated at 24±2°C under natural day/night conditions. The appearance of wrinkling was
 104 considered as the onset of senescence (SEN). On post-harvest storage, the Penjar-2 fruit was
 105 first to show wrinkling at 65-days, which was considered as SEN. The weight loss during
 106 post-harvest storage was monitored by periodic weighing of fruits.

107 **Sequence, SNP, promoter and SIFT analysis**

108 The *NOR* promoter (1834 bp) and cDNA were amplified from AC and Penjars
 109 (Primer details in Table S1), and PCR products were sequenced (Macrogen, South Korea).
 110 The SNPs were identified using Multialin Interface page (Corpet et al. 1988). The amino acid
 111 sequences were aligned using Multialin or PRALINE software
 112 (<http://www.ibi.vu.nl/programs/pralinewww/>) (Bawono and Heringa, 2014). The promoter
 113 sequences (3 Kb) of genes were retrieved from SOL genomics network
 114 (<https://solgenomics.net/>) and analyzed for NAC transcription factor binding sites using
 115 PlantPAN 2.0 (<http://PlantPAN2.itsps.ncku.edu.tw>) (Chow et al. 2016). The deleterious effect
 116 of non-synonymous polymorphisms was calculated by SIFT (www.sift.dna.org; version
 117 4.0.5) (Sim et al. 2012). For SIFT score ≥0.05, SNPs were predicted as tolerated, whereas
 118 score ≤ 0.05 was predicted to be deleterious for the protein function.

119 **Carotenoids, hormones, transcripts and fruit metabolites analysis**

120 The carotenoid content was estimated using Gupta et al. (2015) protocol. The ethylene
 121 emission and hormone analysis were carried out as described in Kilambi et al. (2013) and
 122 Bodanapu et al. (2016) respectively. RNA was extracted from fruits in three biological
 123 replicates using hot phenol method (Verwoerd et al. 1989) and q-PCR was carried out as
 124 described earlier (Kilambi et al. 2013) (Primer details in Table S1). Tomato fruit metabolites
 125 were extracted, analyzed and identified using GC-MS as described in Bodanapu et al. 2016.

126 **Analysis of cuticle components**

For cuticular wax and cutin monomers analysis, five biological replicates at MG and RR were used. The total cuticular wax was isolated as described by Leide et al. (2007). Intact fruits were dipped for 2 min in chloroform and 5-Pregnen-3 β -ol-20-one was added as an internal standard. The solvent was evaporated under nitrogen to 1 mL followed by drying in a Speed Vac before GC-MS. The cuticle membrane (CM) of fruits was isolated using pectinase and cellulase in 50 mM citrate buffer (pH 4.5, supplemented with NaN₃). Isolated 1 cm² size CMs were repeatedly washed with MilliQ water and air dried. CMs were dewaxed as described by Kosma et al. (2010). Dried CMs were treated with chloroform: methanol (1:1; v/v) for 24 h followed by washing with methanol (4-5 times) to remove chloroform. The dewaxed CMs were depolymerized by alkaline hydrolysis (Osman et al. 1995) and derivatized for 30 min with 80 μ L MSTFA at 37°C. 10 μ L of 5-Pregnen-3 β -ol-20-one (1 mg/mL) was used as internal standard for cuticular wax and cutin analysis.

For GC-MS, 1 μ L sample was injected to RXi column as described for fruit metabolites in Bodanapu et al. 2016 in a splitless mode. Cutin and wax components were separated using the following program: 1 min at 50°C with a linear ramp of 10°C/min to 180°C and held at 180°C for 2 min, again a linear ramp of 3°C/min to 300°C and then held at 300°C for 18 min. Both ion source and injector temperature were 250°C and helium was used as carrier gas at a flow rate of 1.5 mL/min. The mass spectra were recorded at a scan rate of 2 scans/sec with a scanning range of 40 to 850 m/z. The raw data were processed, and metabolite identity was assigned as described above for GC-MS analysis.

Statistical analysis

For all experiments, a minimum of 3-5 biological replicates was used and mean with the standard error was calculated. The **StatisticalAnalysisOnMicrosoft-Excel** software (http://prime.psc.riken.jp-/Metabolomics_Software/StatisticalAnalysisOn-MicrosoftExcel/) was used to obtain significant differences between data points using Student's test ($P \leq 0.05$). PCA was carried out using metaboanalyst 3.0.

Network analysis

The carotenoid, hormone, *PSYI* and *CYCB* abundances for Penjars and AC were pairwise correlated at MG, BR and RR stages ($n=3$, $P\text{-value} \leq 0.05$) using the Pearson's Correlation coefficient (PCC). The associations with a PCC value (r) ≥ 0.8 (+/-) were used to create a network, and were visualized using Cytoscape (<http://www.cytoscape.org/>) (Shannon et al. 2003). The red and gray edges indicate negative and positive correlations respectively.

Accession Numbers

160 The accession numbers of genes examined in this study are given in Table S1 along
161 with the primer sequences used for amplification.

162

Results

Penjar tomatoes cultivated in northeastern Spain are harvested during July to September for consumption in winter owing to their longer shelf life. In this study, we compared the cuticle composition, primary metabolites, hormones, carotenoids composition, and cell wall modification in four Penjar accessions with a normal ripening cultivar, Alisa Craig (AC).

Penjar-1 has two novel *NOR* mutations

To uncover the genetic basis of prolonged shelf life, we amplified and sequenced the full-length cDNA of *NOR* gene from all four Penjars (Figure 1; Fig. S1A; Table S1). Notably, two mutations present in Penjar-1 were novel, the C to A transversion at 20th position created a stop codon resulting in a truncated *NOR* protein with only six amino acids (Figure 1) and downstream C to A transversion at 37th position led to glutamine to lysine substitution at the 13th position of the protein. Excepting Penjar-1, other three accessions possessed a single base change similar to *alc* mutant (T to A at 316th position of cDNA; substitution of valine to aspartic acid at 106th position of *NOR* protein, Fig. S1B) akin to the mutation present in 27 Penjar accessions (Casals et al. 2012). The low SIFT score (<0.05) (Ng and Henikoff, 2003; Sim et al. 2012) predicted that the mutations in all four accessions are deleterious for *NOR* protein function. The examination of *NOR* gene promoter (1834 bp) did not reveal any additional SNPs in above accessions (Fig. S1C).

Penjars exhibit delayed ripening and prolonged shelf life

The tomato fruits are considered fully ripe when the fruits acquire uniform red coloration, albeit ripe Penjar fruits exhibited different colors. At the ripe stage, fruits of Penjar-2 were light red, Penjar-1 were yellow-orange and Penjar-3 and -4 were orange colored (Fig. S2A). The attainment of uniform coloration by Penjar fruits was considered equivalent to the red ripe stage (RR) of AC.

The fruit growth of Penjars was nearly similar to AC up to mature green stage (MG). Post-MG stage, the transition period to breaker stage (BR) in Penjars and AC was nearly similar and also akin to *nor*, *alc* and DFD (Mutschler et al. 1992; Saladié et al. 2007; Simpson et al. 1976) (Fig. S2B). However, BR to RR transition (8-15 days) was more prolonged in Penjars than AC, with Penjar-1 being slowest (15 days). Apart from delayed ripening, all Penjars exhibited extended shelf life both on-vine and off-vine. The on-vine ripened Penjar fruits manifested no visible signs of shriveling even after 130 days post-anthesis (DPA) (Simpson et al. 1976), while AC fruits shriveled at 70-80 DPA (Fig. S3).

We next examined in detail the off-vine shelf life of Penjars and AC. The fruits harvested at RR were monitored for wrinkling at regular intervals. In AC, wrinkling appeared on 10th-day post-harvest (DPH), became prominent by 20 DPH followed by shrinking of fruits due to water loss (Figure 2A). At 65 DPH, the shrinkage was markedly apparent, and fruits lost 40% of initial weight. Contrastingly, all Penjar fruits even after 65 DPH retained 80-90% of initial weight and displayed no wrinkling (Figure 2B). By 80 DPH, AC fruits were completely shriveled, whereas, Penjar-1, -3 and -4 showed no signs of wrinkling even at 150 DPH. Among these, Penjar-2 fruits manifested wrinkling by 65 DPH; therefore 65 DPH was selected as the onset of senescence (SEN) for the subsequent post-harvest studies.

***NOR* and *RIN* expression is altered in Penjars**

In tomato, *RIN* and *NOR* are two key regulators of ripening and shelf life (Giovannoni, 2001). To ascertain the role of these regulators in ripening and shelf life of Penjar fruits, their transcript levels were analyzed by qRT-PCR (Fig. S4). Consistent with the long shelf life, the *NOR* expression in Penjar fruits at RR was significantly lower than AC. The *NOR* expression declined in AC and Penjars (except Penjar-1, -2) at SEN. Interestingly, *RIN* transcript was below the detection limit at MG in AC and thereafter increased progressively. The reduced expression of *RIN* in Penjar-1 at RR and SEN is in consonance with it being a knockout mutant. Other Penjars showed higher *RIN* expression at RR and the levels were similar to AC at SEN.

Penjars show wide variation in cuticle composition

The cuticle protects fruit firmness by preventing transpirational water loss. Consistent with this, stripping of the cuticular wax (CW) by chloroform accelerated shrinkage (Figure 3A) and water loss of Penjar fruits akin to AC (Fig. S5A). The analysis of CW revealed the presence of nearly 100 compounds that were classified into five categories: hydrocarbons (alkanes, alkenes), fatty acids, fatty acid alcohols, aromatic and miscellaneous compounds (sterols and triterpenoids) (Fig. S5B, Table S6). At MG, the CW levels were very high in Penjars than AC barring Penjar-1. At RR though CW content increased in all Penjars and AC, only Penjar-3, -4 had levels higher than AC (Figure 3B).

The relative abundance of different cuticle components distinctly differed between four Penjars as well as with AC (Figure 3B-F; Fig. S5). The abundance of nine detected alkanes in RR fruits widely differed in Penjars and AC, excepting C29 alkane that was 1.5-2 fold higher in Penjars than AC. While few fatty acids classes at MG were more abundant in Penjars than AC, the abundance of these fatty acids distinctly changed at RR. Several fatty acid alcohols were most abundant in Penjars particularly at MG, and their levels significantly

increased during ripening. Compared to stigmasterol, the levels of all triterpenes changed during the transition to RR, albeit variably among different Penjars and AC.

Several cutin components such as glycerol, hydroxy hexadecanoic acid, octadecanoic acid, 18-OH octadecanoic and 9,18-diOH octadecanoic were higher in all four Penjars at MG compared to AC (Table 1). However, at RR only glycerol was higher in all four Penjars than AC. The other cutin components in Penjars though varied did not show any concerted increase or decrease like glycerol on comparison to AC. Analogously, the analysis of transcript levels of fourteen genes putatively regulating cuticle biogenesis showed upregulation of *cutin deficient 2 (CD2)*, *MIXTA*, *ABC transporter G family member 11 (WBC11)*, *glycerol-3-phosphate acyltransferase 6 (GPAT6)*, *3-ketoacyl-CoA reductase 1 (KCR1)*, *enoyl-CoA reductase (ECR)*, and downregulation of *fatty acyl-ACP thioesterase (FATB)* in MG fruits of all Penjars compared to AC (Fig. S6; Table S1). However, at RR only *FATB* transcript was commonly altered reinforcing that the cuticle biogenesis is more specifically modulated in Penjars at MG than at RR.

Cell wall modifying genes are downregulated in Penjars

In tomato, fruit ripening is accompanied by progressive softening of cell walls by a battery of cell-wall specific enzymes, whose suppression extends fruit shelf life (Brummell et al. 2002; Cantu et al. 2007; Meli et al. 2010; Uluisik et al. 2016). To ascertain whether the extended shelf life of Penjar fruits is linked with the reduced expression of genes encoding cell wall modifying enzymes, expression of *polygalacturonase (PG-2A, PG-β)*, *expansin (EXP)*, *pectin methylesterase (PME)*, *galactosidase (α-GAL, β-GAL)*, *mannosidase (α-MAN, β-MAN)*, *deoxyhypusine synthase (DHS)* and *hexosaminidase (HEX)* was examined. Consistent with the above notion, expression of at least eight cell wall modifying genes- *β-GAL*, *α-GAL*, *α-MAN*, *β-MAN*, *PG-β*, *PME*, *EXP*, and *DHS* was downregulated at RR of Penjar fruits (Figure 4). Only *PG2A* and *HEX* genes showed higher expression at RR in Penjar-1, and -2 than AC. At SEN, Penjars showed accession-specific variation in gene expressions. All Penjars showed higher expression of *α-MAN*, *β-MAN*, *PG-β* genes than AC. Among different Penjars, expression of *α-GAL* and *PG2A* was greater in Penjar-1, and -3, *β-GAL* was elevated in Penjar 2, *PME* and *EXP* was higher in Penjar-3, and *DHS*, *HEX* was enhanced in Penjar-3, -4 than AC. The analysis of promoters of above genes revealed the presence of multiple NAC binding sites (Fig. S7; Table S4).

Hormone profiling

In tomato, a climacteric fruit, ethylene production is coupled to ripening and the associated fruit coloration and softening (Alexander and Grierson, 2002). Though the overall

pattern of ethylene emission was similar in AC and Penjars, consistent with delayed ripening and lower carotenoid content, the Penjar fruits emitted significantly less ethylene than AC (Fig. S8A), excepting Penjar-2 that emitted slightly higher ethylene at BR. In consonance with reduced ethylene emission, at RR, the transcript levels of key genes of system II ethylene biosynthesis - *1-aminocyclopropane-1-carboxylate synthase 2 and 4 (ACS2, ACS4)*, and at BR of *1-aminocyclopropane-1-carboxylic acid oxidase 1 (ACO1)* were significantly lower in Penjars than in AC (Fig. S8B-D). In contrast, system I ethylene biosynthesis gene-*ACO3* levels (Fig. S8E) were similar in Penjars and AC at BR (except Penjar-3 and -4) and RR (except Penjar-2 and -4). Nearly equal gene expression of *ACS2, 4* and *ACO1* at BR in Penjar-2 and AC may be related to slightly higher ethylene emission from Penjar-2 fruits.

In tomato, other hormones like jasmonate (JA), methyl jasmonate (MeJA) and salicylic acid (SA) also play important roles in lycopene accumulation (Kumar et al. 2014; Liu et al. 2012). In both AC and Penjars, the transition to RR upregulated JA levels. However, the stimulation was substantially lower in Penjars (Figure 5). The upregulation of JA was sustained at SEN in AC, and light-red fruited Penjar-2, while it declined in yellow/orange-fruited Penjars. Contrastingly, Penjar accessions showed changes in MeJA levels that were opposite to AC during the transition from MG to BR with a steady level at RR (except Penjar-2) and a decline at SEN. In tomato, lycopene accumulation in fruits requires higher SA levels during early stages of ripening (Ding and Wang, 2003). Consistent with this, while SA level in MG/BR fruits of AC was significantly higher than RR, it was much lower in Penjars at MG and BR.

Also, the levels of abscisic acid (ABA), indole-3-acetic acid (IAA), indole-3-butyric acid (IBA) and zeatin were altered in Penjar fruits (Figure 5). In AC, ABA level increased during ripening and then declined at SEN, whereas Penjars showed only modest variations during ripening, but with upregulation at SEN. Both Penjars and AC did not show a consistent pattern for IAA levels, except that Penjars had significantly low IAA levels (barring Penjar-1 at MG). Conversely, the IBA levels gradually declined in AC from MG to SEN, and Penjar-4 retained consistently steady IBA levels at all stages. In higher plants leaves, cytokinin reportedly inhibits the senescence process (Lim et al. 2007), however, in both AC and Penjars, zeatin levels did not correlate with fruit senescence.

Penjars show reduced *PSYI* expression

Since ripe Penjar fruits do not acquire typical deep red coloration, it was assumed that *NOR* mutations might have influenced the carotenogenesis in fruits. Profiling of carotenoids at different stages revealed substantially low carotenoid levels in Penjars (Fig. S9), akin to

nor and *alc* (Kopeliovitch et al. 1979; Sink Jr. et al. 1974). Consistent with light-red color, Penjar-2 fruits accumulated higher levels of phytoene, phytofluene, and lycopene than other three Penjars. In tomato, *phytoene synthase 1* (*PSY1*) and chromoplast-specific *lycopene β -cyclase* (*CYCB*) genes are two key regulators of carotenogenesis in fruits, and their increased expression from MG to RR is closely associated with carotenoid accumulation (Hirschberg, 2001). The reduced *PSY1* expression at RR is consistent with lower carotenoid levels in respective Penjars, with the least reduction in Penjar-2 (Fig. S10). Interestingly, Penjar-1 fruits showed 2-fold higher *CYCB* expression at RR than AC, which is reflected as high β -carotene/lycopene ratio than other Penjars (Table S2A). The transition from RR to SEN in Penjar fruits was marked by an accelerated loss in carotenoids levels (~50-70%) than AC (26%) (Table S2B). Among *PSY1* and *CYCB*, at SEN, the *PSY1* expression declined more severely in Penjars than in AC. In contrast, *CYCB* expression though decreased in Penjars yet was 2-3 folds higher than AC. The correlation networks ($r = 0.8$) consisting of most abundant carotenoids, *PSY1* and *CYCB* expression and the hormones during ripening revealed interesting patterns (Fig. S11; Table S3). Penjar-1, -2 -3, and AC showed a strong positive correlation between JA and carotenoid levels, and ethylene and *PSY1* expression levels. In contrast, Penjar-4 showed very few interactions compared to other Penjars and AC. Though other hormones also interacted with different carotenoids and transcripts, these varied between different Penjars.

Metabolite analysis in Penjars

Using GC-MS, we identified ~110 primary metabolites in the MG, BR, RR and SEN fruits of AC and Penjars (Table S5). Notwithstanding the diverse metabolite composition in AC and different Penjars, principal component analysis (PCA) revealed that the metabolite profiles of ripening Penjars were closer and overlapped with each other (Fig. S12A). At SEN, while the profiles were closer in PC1, they showed accession-specific differences in PC2 (Fig. S12B). Based on functional groups, the identified metabolites were classified as organic acids, amino acids, amines, fatty acids, and sugars. On the metabolic pathway, only those metabolites with ≥ 1.5 fold (Log_2 Penjar/AC value of 0.58; $P \leq 0.05$) upregulation or downregulation in Penjar fruits compared to AC were mapped (Figure 6).

Penjars show differential accumulation of Krebs cycle intermediates

The fruits of Penjars displayed differential accumulation of Krebs cycle intermediates during ripening and postharvest storage (Fig. S13). Particularly in Penjars, organic acids levels were lower than AC during ripening. Citrate, the most abundant organic acid in ripe fruits, was significantly lower in Penjar fruits at SEN, whereas, at RR, the citrate level was

lower in Penjar-1 and -3 than AC. Likewise, the acotinate levels in RR fruits were also highly reduced than AC, albeit at SEN, the levels were nearly similar. Though the levels were low at RR, malate, fumarate and methyl maleate were significantly upregulated at SEN in Penjars. The levels of succinate in Penjar fruits decreased during ripening and SEN.

Penjars exhibit low levels of amino acids during ripening and post-harvest storage

During tomato ripening, suppression of climacteric rise of ethylene reportedly reduces the amino acid content (Gao et al. 2007). Consistent with reduced ethylene emission in Penjar fruits, the levels of several amino acids (12) were lower than in AC. At MG, free amino acids such as alanine, leucine, isoleucine, valine, glycine, serine, alanine-3-cyano, threonine, aspartic acid, beta-alanine, glutamine, GABA, etc., were relatively low in both AC and Penjar (except Penjar-3) fruits (Fig. S14). The onset of ripening stimulated a modest increase in the amino acids and amines in AC and Penjar-2 fruits at BR. Interestingly, the GABA levels declined in all Penjars by 2.5-5 fold at BR. At RR, in most Penjars, the amino acids such as beta-alanine, GABA, isoleucine, glycine, serine, alanine-cyano, threonine, aspartic acid, asparagine, and hydroxylamine were much lower than AC. At SEN, reduction in the amino acids was most severe in Penjars with ~10-30 fold decrease in alanine, valine, leucine, and isoleucine. The levels of only a few amino acids such as glutamate at RR of Penjar-2, phenylalanine at RR and SEN of Penjar-3 and -4, and aspartate at SEN of Penjar-2 and -3 were higher than AC at respective stages. All Penjar fruits had reduced level of serotonin (5-hydroxytryptamine) at SEN while serotonin levels were high in AC fruits throughout ripening and SEN.

Penjars exhibit differential accumulation of sugars and cell wall-related metabolites

During fruit ripening, glucose and fructose, cell wall-derived sugars, and polyols increase, while sucrose and sugar phosphates and fatty acids decrease (Fraser et al. 2007). Ripening Penjar fruits characteristically exhibited substantially higher levels of glucose, fructose, glucose-6-P, sucrose, etc. than AC (Fig. S15). At SEN glucose and fructose levels declined in Penjars but increased in AC. The Penjar fruits at SEN also showed increased levels of glucose-6-P, fructose-6-P, and mannose-6-P. At SEN, high levels of cell wall-related metabolites like xylose and cellobiose in AC fruits and reduced levels of galacturonate and galactarate levels was observed in Penjar fruits.

The profiles of other metabolites such as fatty acids, nucleotides, dehydroascorbate, nicotinate, and quinate in Penjar and AC fruits were also distinctly different (Fig. S16). Though the levels of fatty acids such as margaric acid, linolenic acid, stearic acid in Penjar fruits were similar to AC during ripening, at SEN their levels were higher in AC. The

366 metabolites- nonadecyclic acid and heneicosylic acid that are detected only at SEN were
367 much lower in Penjars than AC. In contrast to Penjars, AC at SEN exhibited increased levels
368 of adenosine and guanosine levels. Interestingly, uracil levels were high in Penjars
369 throughout ripening including SEN except for Penjar-2 that had low levels at SEN.

370

Discussion

Tomato being a perishable fruit is the subject of intensive investigations to extend its post-harvest shelf life (Pech et al. 2013). Among the regulatory factors identified, the *nor* and *rin* alleles in tomato prolong the shelf life of fruits, even in heterozygous condition (Garg et al. 2008a; Giovannoni, 2007). The *NOR* gene belongs to NAC transcription factor family (Klee and Giovannoni, 2011; Martel et al. 2011) and has only two reported mutant alleles, *alc* and *nor* in tomato (Casals et al. 2012; Dias et al. 2003; Garg et al. 2008a). In this study, we identified a novel allele in Penjar-1 with two mutations, one with Q13K and second, where a stop codon terminated the NOR protein after six aa, while the other three Penjar accessions had *alc* mutation. All four accessions showed extended fruit shelf-life during post-harvest storage, a phenotype consistent with reported *nor/alc* mutants (Casals et al. 2012). Considering that post-harvest shelf life of Penjars is dependent on genetic background, the same may have contributed to the observed differences in the shelf life of these accessions (Casals et al. 2012; Garg et al. 2008a,b). The 55 days delay in SEN compared to AC strongly indicates that post-harvest shelf life is considerably prolonged in the Penjar fruits.

In *nor/alc* mutant, the delayed onset or total loss of ripening signifies that the *NOR* gene regulates a majority of the ripening triggered processes (Casals et al. 2015; Osorio et al. 2011; Saladié et al. 2007). Interestingly, despite mutations in *NOR* gene, the *RIN* expression was not considerably altered, except in Penjar-1, a total knockout mutant. Such a downregulation of *RIN* expression was also observed in *nor* and *SINAC4* RNAi lines (Martel, 2010; Zhu et al. 2014). Evidently, though *RIN* is a master regulator, *NOR* acts autonomously of *RIN*, and optimal regulation of ripening requires a concerted action between these two regulators (Osorio et al. 2011).

A factor influencing fruit firmness is the cuticle as it prevents water loss and sustains cellular turgidity (Saladié et al. 2007). Consistent with earlier studies (Kosma et al. 2010; Saladié et al. 2007) the overall cuticle composition (wax and cutin) of Penjars is more similar to *nor/alc* mutants than AC. At the same time, the four Penjars widely vary in levels of different constituents of the cuticle. Only at MG, the Penjars share upregulation of few cuticle constituents and gene transcripts, whereas, at RR, such a shared regulation is largely absent. Surprisingly, post-MG despite variations in building blocks of the cuticle, all Penjars share prolonged shelf life and reduced water loss. Seemingly along with the composition, the cuticle architecture is also a key determinant of the prolonged shelf life of Penjars. The endogenous factors regulating cuticle composition/architecture are currently not known (Yeats and Rose, 2013; Fernández et al. 2016). It appears that the cuticle

composition/architecture is not the sole factor, and the regulation of shelf life also involves other cellular processes altered by *NOR* mutation in Penjars.

In addition to the cuticle, the cell wall is also considered a key determinant for retention of fruit firmness. Consistent with this, the silencing of several cell wall modifying enzymes in tomato extends the shelf life of fruits (Vicente et al. 2007). The prolonged post-harvest shelf life indicates that the process of cell wall dissolution is considerably slower in Penjar fruits. The presence of several NAC domains in the promoters of cell wall modifying genes suggests that mutations in *NOR* would lead to lowering of their expression. Consistent with this, the expression of most cell wall modifying genes is downregulated during ripening in Penjars similar to *nor/alc* mutants (Osorio et al. 2011; Saladié et al. 2007). Even at SEN, several of these genes showed reduced expression indicating slower degradation of the cell wall, which is also corroborated by lower levels of cell wall related metabolites in Penjars than in AC.

One of the characteristic features of *nor/alc* fruits is the reduced accumulation of carotenoids compared to normal ripening cultivars (Giovannoni, 2007; Kopeliovitch et al. 1979; Sink Jr. et al. 1974). In Penjars, the reduction in carotenoids was manifested for most fruit-specific carotenoids including lycopene and β -carotene as well as for precursors, phytoene, and phytofluene. The reduced level of precursors is consistent with the lower expression of *PSYI* and *CYCB* in the Penjar fruits during ripening. In tomato fruits, carotenoid accumulation is also strongly influenced by ethylene; consequently *Nr* mutant of tomato shows diminished accumulation of carotenoids, due to a defect in ethylene perception (Barry et al. 2005). Similar to carotenoids, the reduction in ethylene emission from Penjars fruits was associated with lower expression of system II ethylene biosynthesis genes, *ACS2*, *ACS4* and *ACO1* (Barry and Giovannoni, 2007; Pech et al. 2012). The promoter analysis of carotenogenesis and ethylene biosynthesis genes revealed multiple NAC binding sites (Table S4) suggesting the downregulation of above genes may be related to the absence of a functional *NOR* protein in Penjars.

Current evidence indicate that fruit development is regulated by extensive interactions among different plant hormones (Kumar et al. 2014; Liu et al. 2015, Bodanapu et al. 2016). However, the overall hierarchical order of hormonal regulation in tomato development is not fully known. Hormonal profiling of Penjar fruits revealed that the *NOR* mutations influenced several hormones across the different developmental stages of fruits. In tomato, several studies have examined the influence of individual hormones on the accumulation of carotenoids (Kumar et al. 2014; Liu et al. 2015). However, a comprehensive correlation

between hormones and carotenoids is largely missing. Barring Penjar-4, the interactions between hormones and carotenoids were similar in other Penjars albeit with few differences. The similarity in hormone-carotenoids interactions indicates that though *NOR* mutation attenuates carotenoid accumulation in Penjars, it does not influence the process underlying above interactions. The different interaction profiles in Penjar-4 may owe to nearly steady JA levels at all ripening stages. Evidence from JA-deficient mutants has implied that JA influences carotenogenesis in tomato independent of ethylene (Liu et al. 2012). The hormone-carotenoid interactions also indicate that in addition to JA and ethylene, other hormones also influence the regulation of carotenogenesis in tomato.

In tomato, suppression of ethylene biosynthesis delays ripening of fruits and extends the shelf life (Oeller et al. 1991). Consistent with this, lower ethylene emission may have contributed to the prolonged shelf life of Penjar fruits. Also, lower ABA levels at MG and RR in Penjars can lead to slower ripening (Sun et al. 2012). The accumulation of ABA is also characteristic of plant cells that retain water (Iuchi et al. 2000; Wan and Li, 2006). The higher retention of water in Penjar fruits may be assisted by higher ABA levels observed at SEN in all Penjars. Consistent with this view, higher sugar levels in Penjars may help in retention of cellular turgidity as marked by slow water loss (Vicente et al. 2007). A similar correlation between sucrose levels and increased firmness was also observed in *LIN5* suppressed fruits (Vallarino et al. 2017).

The extension of shelf life of fruits demands an optimal utilization of stored resources. During post-harvest storage, the fruits are deprived of support from the mother plant and can stay fresh only by lowering metabolism. Consistent with this, Penjar fruits show reduced metabolic processes such as turnover of proteins, fatty acids and reduced flux via TCA cycle. The lowering of citrate and upregulation of malate indicates an attenuation of respiration that is critical for prolonging the shelf life (Centeno et al. 2011). Likewise, abundances of the aspartate amino acid family (methionine, isoleucine, threonine, and lysine) have been linked to energy metabolism during seed germination (Angelovici et al. 2011; Kirma et al. 2012). The downregulation of isoleucine, threonine observed in Penjars at SEN indicates reduced flux in TCA cycle. Increased lysine levels in Penjar-2 and -4 in SEN fruits was similar to that found in the fruits of *LeACS2* suppressed transgenic line (Gao et al. 2007). Lowering of metabolism was not restricted to respiration, but even protein turnover was reduced, as evident by the decrease in the levels of free amino acids in Penjars. The lowering of metabolism also requires a balance between carbon and nitrogen metabolism, which is consistent with a reduction in GABA levels in Penjars, as GABA acts as a regulatory

molecule to fine tune the cooperation between these two pathways (Takayama and Ezura, 2015). Collectively, above metabolic shifts indicate attenuation of overall metabolism during ripening and post-harvest storage of Penjars, contributing to their long shelf life.

In summary, our study revealed a wide-ranging influence of *NOR* mutations on diverse metabolic processes in four different Penjars. The prolonged shelf life of Penjars appears to involve attenuation of several metabolic processes associated with slower degradation of cell walls. The sustenance of firmness seems to be correlated with higher sucrose and reduced water loss, and hitherto unknown features of cuticle composition/architecture in the Penjars. In future, a better comprehension of the metabolic processes including an understanding of cuticle architecture combined with genome editing of causative genes would facilitate the improvement of the shelf life of perishable fruits.

Acknowledgements

We thank Department of Biotechnology, Government of India (BT/PR11671/PBD/16/828/2008) funding to YS and RS, Council of Scientific and Industrial Research fellowship, New Delhi to RK and DST-Fund for Improvement of Science and Technology to Department of Plant Sciences. We thank Rafael Fernandez Muñoz, Institute for Mediterranean and Subtropical Horticulture, Malaga, Spain for the kind gift of Penjar accessions.

Author contributions

The work was conceptualized by RK and YS. Most experiments were done by RK and VT helped in the wax analysis. RK, RS, and YS were involved in writing the manuscript, and all authors read and approved the manuscript.

Conflict of Interest

The authors declare no conflict of interest.

Supplementary data

The following materials are available in the online version of this article.

Figure S1. Sequence analysis of *NAC-NOR* in Penjars.

Figure S2. Fruit phenotypes (A) and chronological development and ripening (B) in AC and Penjars.

Figure S3. The on-vine shelf life of Penjar and AC fruits.

Figure S4. Relative expression of *RIN* and *NOR* genes in AC and Penjar fruits.

Figure S5. Loss of fruit weight in both AC and Penjars after chloroform treatment (A) and the percent relative abundance of cuticular wax components in MG and RR fruits of AC and Penjars.

Figure S6. Relative expression of genes associated with cuticle biosynthesis in AC and Penjar fruits.

Figure S7. Location of putative NAC motifs in the promoters of cell wall modifying genes.

Figure S8. Ethylene emission and relative expression of ethylene biosynthetic genes in AC and Penjar fruits.

Figure S9. Carotenoid composition in AC and Penjar fruits during ripening and post-harvest storage.

Figure S10. Relative expression of key carotenoid biosynthesis genes in AC and Penjar fruits.

Figure S11. Correlation network of carotenoids, carotenogenic genes, and hormones during fruit ripening.

Figure S12. Principle component analysis (PCA) of metabolic profiles in AC and Penjar fruits.

Figure S13. The relative abundances of organic acids in the AC and Penjar fruits during ripening and post-harvest storage.

Figure S14. The relative abundances of amino acids and amines in the AC and Penjar fruits during ripening and post-harvest storage.

Figure S15. The relative abundances of sugars, sugar alcohols and sugar-derived acids in the AC and Penjar fruits during ripening and post-harvest storage.

Figure S16. The relative abundances of miscellaneous compounds (fatty acids, vitamins and vitamin-derivatives, nucleotides, phenolics, etc.) in the AC and Penjar fruits during ripening and post-harvest storage.

Table S1. Primers used for PCR in the present study.

Table S2. Carotenoid content in AC and Penjar fruits.

Table S3. The interactions between different metabolites and genes in the correlation networks of AC and Penjar fruits.

Table S4. List of NAC binding sites identified in the promoters of various genes.

Table S5. List of metabolites identified and their relative abundances in AC and Penjar fruits during ripening and post-harvest storage using GC-MS.

Table S6. List of cuticular wax components identified and their relative abundances in AC and Penjars during ripening using GC-MS.

References

- Alexander L, Grierson D.** 2002. Ethylene biosynthesis and action in tomato: a model for climacteric fruit ripening. *Journal of Experimental Botany*. 53, 2039–2055.
- Angelovici R, Fait A, Fernie AR, Galili G.** 2011. A seed high-lysine trait is negatively associated with the TCA cycle and slows down Arabidopsis seed germination. *New Phytologist*, 189, 148–159.
- Barry CS, Giovannoni JJ.** 2007. Ethylene and Fruit Ripening. *Journal of Plant Growth and Regulation* 26: 143–159.
- Barry CS, McQuinn RP, Thompson, AJ, Seymour GB, Grierson D, Giovannoni JJ.** 2005. Ethylene insensitivity conferred by the *Green-ripe* and *Never-ripe 2* ripening mutants of tomato. *Plant Physiology* 138, 267–275.
- Bawono P, Heringa J.** 2014. Multiple Sequence Alignment Methods. *Methods in Molecular Biology* 1079, 245–62.
- Bodanapu R, Gupta SK, Basha PO, Sakthivel K, Sadhna S, Sreelakshmi, Y, Sharma R.** 2016. Nitric oxide overproduction in tomato shr mutant shifts metabolic profiles and suppresses fruit growth and ripening. *Frontiers in Plant Science* 7, 1714.
- Brummell DA, Howie WJ, Ma C, Dunsmuir P.** 2002. Postharvest fruit quality of transgenic tomatoes suppressed in expression of a ripening-related *expansin*. *Postharvest Biology and Technology* 25, 209–220.
- Cantu D, Vicente AR, Greve LC, Labavitch JM, Powell ALT.** 2007. Genetic Determinants of Textural Modifications in Fruits and Role of Cell Wall Polysaccharides and Defense Proteins in the Protection Against Pathogens. *Fresh Prod.* 1(2), 101-110
- Carrari F, Baxter C, Usadel B, et al.** 2006. Integrated analysis of metabolite and transcript levels reveals the metabolic shifts that underlie tomato fruit development and highlight regulatory aspects of metabolic network behavior. *Plant Physiology* 142, 1380–96.
- Casals J, Pascual L, Cañizares J, Cebolla-Cornejo J, Casañas F, Nuez F.** 2012. Genetic basis of long shelf life and variability into Penjar tomato. *Genetic Resources and Crop Evolution* 59, 219–229.
- Casals J, Marti Renau R, Casañas Artigas F, Cebolla-Cornejo J.** 2015. Sugar and acid profile of Penjar tomatoes and its evolution during storage. *Scientia Agricola* 72, 314–321.
- Centeno DC, Osorio S, Nunes-Nesi A, et al.** 2011. Malate Plays a Crucial Role in Starch Metabolism, Ripening, and Soluble Solid Content of Tomato Fruit and Affects Postharvest Softening. *Plant Cell* 23, 162–184.

- Chow CN, Zheng HQ, Wu NY, Chien CH, Huang HD, Lee TY, Chiang-Hsieh YF, Hou PF, Yang TY, Chang WC.** 2016. PlantPAN 2.0: an update of plant promoter analysis navigator for reconstructing transcriptional regulatory networks in plants. *Nucleic Acids Research* 44, D1154-60.
- Dias T, Maluf W, Faria M, De Freitas J, Gomes L, Resende J, De Azevedo S.** 2003. *Alcobaça* allele and genotypic backgrounds affect yield and fruit shelf life of tomato hybrids. *Scientia Agricola* 60, 269–275.
- Ding CK, Wang CY.** 2003. The dual effects of methyl salicylate on ripening and expression of ethylene biosynthetic genes in tomato fruit. *Plant Science* 164, 589–596.
- Fernández V, Guzmán-Delgado P, Graça J, Santos S, Gil L.** (2016) Cuticle Structure in Relation to Chemical Composition: Re-assessing the Prevailing Model. *Frontiers in Plant Science* 7, 427.
- Fraser PD, Enfissi EMA, Halket JM, Truesdale MR, Yu D, Gerrish C, Bramley PM.** 2007. Manipulation of Phytoene Levels in Tomato Fruit: Effects on Isoprenoids, Plastids, and Intermediary Metabolism. *Plant Cell* 19, 3194–3211.
- Gao HY, Zhu BZ, Zhu HL, Zhang YL, Xie YH, Li YC, Luo YB.** 2007. Effect of suppression of ethylene biosynthesis on flavor products in tomato fruits. *Russian Journal Plant Physiology* 54, 80–88.
- Garg N, Cheema DS, Dhatt AS.** 2008a. Utilization of *rin*, *nor*, and *alc* alleles to extend tomato fruit availability. *International Journal of Vegetable Science* 14, 41–54.
- Garg N, Cheema DS, Pathak D.** 2008b. Heterosis breeding in tomato involving *rin*, *nor* and *alc* alleles: A review of literature. *Advances in Horticultural Science* 22, 54–62.
- Giovannoni JJ.** 2001. Molecular biology of fruit maturation and ripening. *Annual Review of Plant Physiology and Plant Molecular Biology* 52, 725-749.
- Giovannoni JJ.** 2007. Fruit ripening mutants yield insights into ripening control. *Current Opinion in Plant Biology* 10, 283–289.
- Giovannoni JJ.** 2004. Genetic regulation of fruit development and ripening. *Plant Cell* 16 Suppl, S170–S180.
- Gupta P, Sreelakshmi Y, Sharma R.** 2015. A rapid and sensitive method for determination of carotenoids in plant tissues by high performance liquid chromatography. *Plant Methods* 11, 5.
- Hirschberg J.** 2001. Carotenoid biosynthesis in flowering plants. *Current Opinion in Plant Biology* 4, 210–218.
- Iuchi S, Kobayashi M, Yamaguchi-Shinozaki K, Shinozaki K.** 2000. A stress-inducible

- gene for 9-cis-epoxycarotenoid dioxygenase involved in abscisic acid biosynthesis under water stress in drought-tolerant cowpea. *Plant Physiology* 123, 553–62.
- Kilambi HV, Kumar, R, Sharma R, Sreelakshmi Y.** 2013. Chromoplast-Specific Carotenoid-Associated Protein Appears to Be Important for Enhanced Accumulation of Carotenoids in *hp1* Tomato Fruits. *Plant Physiology* 161:2085–2101.
- Kirma M, Araújo WL, Fernie AR, Galili G.** 2012. The multifaceted role of aspartate-family amino acids in plant metabolism. *Journal of Experimental Botany* 63, 4995–5001.
- Klee HJ, Giovannoni JJ.** 2011. Genetics and Control of Tomato Fruit Ripening and Quality Attributes. *Annual Reviews in Genetics* 45, 41–59.
- Kosma DK, Parsons EP, Isaacson T, Lü S, Rose JKC, Jenks MA.** 2010. Fruit cuticle lipid composition during development in tomato ripening mutants. *Physiology Plantarum* 139, 107–117.
- Kopeliovitch E, Rabinowitch HD, Mizrahi Y, Kedar N.** 1979. The potential of ripening mutants for extending the storage life of the tomato fruit. *Euphytica* 28, 99–104.
- Kumar R, Khurana A, Sharma AK.** 2014. Role of plant hormones and their interplay in development and ripening of fleshy fruits. *Journal of Experimental Botany* 65, 4561–4575.
- Lim PO, Kim HJ, Nam HG.** 2007. Leaf senescence. *Annual Review of Plant Biology* 58, 115–36.
- Leide J, Hildebrandt U, Reussing K, Riederer M, Vogg G.** 2007. The developmental pattern of tomato fruit wax accumulation and its impact on cuticular transpiration barrier properties: effects of a deficiency in a beta-ketoacyl-coenzyme A synthase (*LeCER6*). *Plant Physiology* 144, 1667–1679.
- Liu L, Wei J, Zhang M, Zhang L, Li C, Wang Q.** 2012. Ethylene independent induction of lycopene biosynthesis in tomato fruits by jasmonates. *Journal of Experimental Botany* 63, 5751–5762.
- Liu M, Pirrello J, Chervin C, Roustan JP, Bouzayen M.** 2015. Ethylene Control of Fruit Ripening: Revisiting the Complex Network of Transcriptional Regulation. *Plant Physiology* 169, 2380–90.
- Ma N, Feng H, Meng X, Li D, Yang D, Wu C, Meng Q.** 2014. Overexpression of tomato *SINAC1* transcription factor alters fruit pigmentation and softening. *BMC Plant Biology* 14, 351.
- Martel C.** 2010. Characterization Of The Roles Of *Rin* And *Nor* During Tomato (*Solanum*

- 645 *lycopersicum*) Fruit Ripening. Thesis submitted to Cornell University.
- 646 <https://www.cornell.edu/>
- 647 **Martel C, Vrebalov J, Tafelmeyer P, Giovannoni JJ.** 2011. The Tomato MADS-Box
- 648 Transcription Factor RIPENING INHIBITOR Interacts with Promoters Involved in
- 649 Numerous Ripening Processes in a COLORLESS NONRIPENING-Dependent Manner.
- 650 *Plant Physiology* 157, 1568–1579.
- 651 **McAtee P, Karim S, Schaffer R, David K.** 2013. A dynamic interplay between
- 652 phytohormones is required for fruit development, maturation, and ripening. *Frontiers in*
- 653 *Plant Science* 4, 79.
- 654 **Meli VS, Ghosh S, Prabha TN, Chakraborty N, Chakraborty S, Datta A.** 2010.
- 655 Enhancement of fruit shelf life by suppressing N-glycan processing enzymes.
- 656 *Proceedings of National Academy of Sciences, U. S. A.* 107, 2413–2418.
- 657 **Mutschler M, Guttieri M, Kinzer S, Grierson D, Tucker G.** 1988. Changes in ripening-
- 658 related processes in tomato conditioned by the *alc* mutant. *Theoretical and Applied*
- 659 *Genetics* 76, 285–292.
- 660 **Mutschler M, Wolfe DW, Cobb ED, Yourstone KS.** 1992. Tomato Fruit Quality and Shelf
- 661 Life in Hybrids Heterozygous for the *alc* Ripening Mutant. *Horticultural Science* 27,
- 662 1985–1988.
- 663 **Nambeesan S, Datsenka T, Ferruzzi MG, Malladi A, Mattoo AK, Handa AK.** 2010.
- 664 Overexpression of yeast spermidine synthase impacts ripening, senescence and decay
- 665 symptoms in tomato. *The Plant Journal* 63, 836–847.
- 666 **Ng PC, Henikoff S.** 2003. SIFT: Predicting amino acid changes that affect protein function.
- 667 *Nucleic Acids Research* 31, 3812–3814.
- 668 **Oeller PW, Lu MW, Taylor LP, Pike DA, Theologis A.** 1991. Reversible inhibition of
- 669 tomato fruit senescence by antisense RNA. *Science* 254, 437–439.
- 670 **Osman SF, Gerard HC, Fett WF, Moreau R, Dudley RL.** 1995. Method for the
- 671 Production and Characterization of Tomato Cutin Oligomers. *Journal of Agriculture and*
- 672 *Food Chemistry* 43, 2134–2137.
- 673 **Osorio S, Alba R, Damasceno CMB, et al.** 2011. Systems Biology of Tomato Fruit
- 674 Development: Combined Transcript, Protein, and Metabolite Analysis of Tomato
- 675 Transcription Factor (*nor*, *rin*) and Ethylene Receptor (*Nr*) Mutants Reveals Novel
- 676 Regulatory Interactions. *Plant Physiology* 157, 405–425.
- 677 **Pech JC, Purgatto E, Bouzayen M, Latché A.** 2012. Ethylene and Fruit Ripening, *Annual*
- 678 *Plant Reviews* 44, 275-304.

- 679 **Pech JC, Purgatto E, Girardi CL, Rombaldi CV, Latché A.** 2013. Current challenges in
680 postharvest biology of fruit ripening. *Current Agricultural Science and Technology* 19,
681 1–18.
- 682 **Pirrello J.** 2009. Regulation of tomato fruit ripening. *CAB Reviews Perspectives in*
683 *Agriculture Veterinary Science Nutrition and Natural Resources* 4, 1–14.
- 684 **Saladié M, Matas AJ, Isaacson T, et al.** 2007. A reevaluation of the key factors that
685 influence tomato fruit softening and integrity. *Plant Physiology* 144, 1012–28.
- 686 **Seymour GB, Chapman NH, Chew BL, Rose JKC.** 2013. Regulation of ripening and
687 opportunities for control in tomato and other fruits. *Plant Biotechnology Journal* 11,
688 269–278.
- 689 **Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski**
690 **B, Ideker T.** 2003. Cytoscape: A software Environment for integrated models of
691 biomolecular interaction networks. *Genome Research* 13, 2498–2504.
- 692 **Sim NL, Kumar P, Hu J, Henikoff S, Schneider G, Ng PC.** 2012. SIFT web server:
693 predicting effects of amino acid substitutions on proteins. *Nucleic Acids Research* 40,
694 W452-7.
- 695 **Simpson DJ, Baqar AMR, Mcglasson AWB, Leea TH.** 1976. Changes in Ultrastructure
696 and Pigment Content during Development and Senescence of Fruits of Normal and *rin*
697 and *nor* Mutant Tomatoes. *Australian Journal of Plant Physiology* 3(5), 575-587
- 698 **Sink Jr, KC, Herner RC, Knowlton LL.** 1974. Chlorophyll and carotenoids of the *rin*
699 tomato mutant. *Canadian Journal of Botany* 52, 1657–1660.
- 700 **Sun L, Sun Y, Zhang M, et al.** 2012. Suppression of *9-cis-epoxycarotenoid dioxygenase*,
701 which encodes a key enzyme in abscisic acid biosynthesis, alters fruit texture in
702 transgenic tomato. *Plant Physiology* 158, 283–298.
- 703 **Takayama M, Ezura H.** 2015. How and why does tomato accumulate a large amount of
704 GABA in the fruit? *Frontiers in Plant Science* 6, 1–7.
- 705 **Tieman D, Taylor M, Ciardi J, Klee H.** 2000. The tomato ethylene receptors NR and
706 LeETR4 are negative regulators of ethylene response and exhibit functional
707 compensation within a multigene family. *Proceedings of National Academy of Science,*
708 *U. S. A.* 97, 5663–5668.
- 709 **Uluisek S, Chapman NH, Smith R, Poole M, Adams G, Gillis RB, Besong TMD, et al.**
710 2016. Genetic improvement of tomato by targeted control of fruit softening. *Nature*
711 *Biotechnology* 34, 950–952.
- 712 **Vallarino JG, Yeats TH, Maximova E, Rose JK, Fernie AR, Osorio S.** 2017. Postharvest

- changes in LIN5-down-regulated plants suggest a role for sugar deficiency in cuticle metabolism during ripening. *Phytochemistry* 142, 11-20.
- Verwoerd TC, Dekker BMM, Hoekema A.** 1989. A small-scale procedure for the rapid isolation of plant RNAs. *Nucleic Acids Research* 17, 2362.
- Vicente AR, Saladié M, Rose JKC, Labavitch JM.** 2007. The linkage between cell wall metabolism and fruit softening: Looking to the future. *Journal of the Science of Food and Agriculture* 87, 1435–1448.
- Wan XR, Li L.** 2006. Regulation of ABA level and water-stress tolerance of Arabidopsis by ectopic expression of a peanut *9-cis-epoxycarotenoid dioxygenase* gene. *Biochem. Biophys. Res. Commun.* 347(4), 1030–1038.
- Xie Y, Zhu B, Yang X, Zhang H, Fu D, Zhu H, Shao Y, Li Y, Gao H, Luo Y.** 2006. Delay of postharvest ripening and senescence of tomato fruit through virus-induced *LeACS2* gene silencing. *Postharvest Biology and Technology* 42, 8–15.
- Yeats TH, Rose JKC.** 2013. The formation and function of plant cuticles. *Plant Physiology* 163, 5–20
- Zhu M, Chen G, Zhou S, Tu Y, Wang Y, Dong T, Hu Z.** 2014. A new tomato NAC (NAM ATAF1/2/CUC2) transcription factor, SINAC4, functions as a positive regulator of fruit ripening and carotenoid accumulation. *Plant Cell Physiology* 55, 119–135.

Figure legends

Figure 1. Mutations in *NAC-NOR* gene in *nor*, *alcobaca* and Penjar accessions. AC encodes a full-length NOR protein of 355 aa. *nor* mutant produces a truncated protein of 186 aa. The Penjar-1 encodes a protein of 6 aa length, whereas *alc*, Penjar-2, -3, and -4 encode a full-length protein with a single amino acid change of V to D at 106th position (indicated by an asterisk and red band). NAC indicates NAC domain and TRR indicates transcriptional regulatory region. The details of base changes and the corresponding positions in the cDNA, protein, and SIFT score are indicated in the Table.

Figure 2. Post-harvest shelf life and water loss in AC and Penjar fruits. **A**, The fruits were harvested at ripe stage and incubated under normal day-night conditions as described in methods. By 40 days post-harvest, AC fruits showed prominent wrinkling, and 65 day-old fruits displayed a clear sign of cellular disintegration. **B**, The water loss was measured during post-harvest storage. Data are means of 5 biological replicates $\pm$ SE, ‘*’ indicates $P \leq 0.05$.

Figure 3. The cuticular wax composition of AC and Penjar fruits during ripening. **A**, The fruits were harvested at the ripe stage, and cuticular wax from the surface of both AC and Penjar fruits was removed by using chloroform. This resulted in the severe shrinking of fruits due to loss of moisture. **B**, The total cuticular wax content of AC and Penjar fruits at MG and RR. **C-F**, The relative abundance of cuticular wax components: alkanes (**C**), fatty acids (**D**), fatty alcohols (**E**), sterols and triterpenes (**F**) in MG and RR fruits of AC and Penjars. Data are means of 3 biological replicates $\pm$ SE, ‘*’ indicates $P \leq 0.05$.

Figure 4. Relative expression of cell wall modifying genes in AC and Penjar fruits. The fruits were collected at different ripening and SEN stages, and expression of various cell wall modifying genes was determined by qPCR. The graphs represent the data obtained after normalization with *actin* and *ubiquitin*. **A**, α -Galactosidase; **B**, β -Galactosidase; **C**, α -Mannosidase; **D**, β -Mannosidase; **E**, Polygalacturonase 2A; **F**, Polygalacturonase β subunit; **G**, Pectin methylesterase; **H**, Expansin; **I**, Deoxyhypusine synthase and **J**, β -N-acetyl-D-hexosaminidase. Data are means of 3 biological replicates $\pm$ SE, ‘*’ indicates $P \leq 0.05$.

Figure 5. Hormone profiling in AC and Penjar fruits. The fruits were collected at different ripening and SEN stages, and hormone profiles were determined as described in methods. **A**, IAA; **B**, IBA; **C**, zeatin; **D**, JA; **E**, MeJA; **F**, SA; **G**, ABA. Data are means of 5 biological replicates $\pm$ SE, ‘*’ indicates $P \leq 0.05$.

766

767 **Figure 6. The metabolic shifts in the fruits of Penjars in comparison to AC during**
 768 **ripening and post-harvest storage.** The relative abundances of the metabolites in Penjar
 769 fruits at MG, BR, RR and SEN stages were obtained from the Log₂ Penjar/AC values, and
 770 only those metabolites with ≥ 1.5 fold change ($n=5 \pm SE$, $P \leq 0.05$) were mapped on the
 771 pathway. The scale at the top left corner represents Log₂ fold changes in the range of -3 to +3.
 772 The metabolites indicated in gray letters on the pathway were not detected in the GC-MS
 773 analysis.

774

Table 1. List of compounds identified in the cutin matrix of MG and RR stage fruits of AC and Penjars.

Compound	MG								
	AC	Penjar-1	Penjar-2		Penjar-3		Penjar-4		
	Mean ± SE	Mean ± SE	P-value	Mean ± SE	P-value	Mean ± SE	P-value	Mean ± SE	P-value
Glycerol	1.18 ± 0.28	29.06 ± 1.89	0.0005	6.85 ± 0.50	0.0157	12.41 ± 0.23	0.0001	54.86± 7.72	0.0196
Dodecanoic acid	3.39 ± 0.23	39.10 ± 4.03	0.0045	5.77± 0.40	0.0106	4.87 ± 0.41	0.0759	9.58 ±0.23	0.0002
Hexadecanoic acid	37.63 ± 4.55	264.38 ± 14.89	0.0211	56.61± 4.32	0.0006	3.62± 0.08	0.5516	79.28 ± 3.72	0.0139
Heptadecanoic acid	2.90 ± 0.15	17.72 ± 1.66	0.0547	<i>1.56 ± 0.06</i>	0.0001	11.30 ± 0.25	0.0001	4.48± 0.20	0.0889
Octadecanol	10.90 ± 1.38	<i>5.66 ± 0.81</i>	0.0184	<i>2.45 ± 0.06</i>	0.0184	7.60 ± 0.73	0.0090	39.81 ±1.26	0.0003
hydroxy hexadecanoic acid	25.03 ± 2.01	217.18 ± 3.19	0.0002	47.90 ± 4.67	0.0223	70.95 ± 7.42	0.0326	153.82 ± 2.83	0.0004
9(10),16-diOH hexadecanoic	442.17 ± 42.61	<i>8.31 ± 1.04</i>	0.0086	<i>9.43 ± 0.54</i>	0.0002	847.69 ± 72.10	0.0318	731.33 ± 55.32	0.0006
Octadecanoic acid	57.51 ± 4.66	355.26 ± 19.42	0.0002	88.28±1.81	0.1982	123.74 ± 4.33	0.0264	103.68 ± 1.65	0.0005
18-OH octadecanoic	28.39 ± 4.66	312.40± 9.685	0.0014	90.59± 6.41	0.0094	77.22 ± 5.63	0.0178	167.78± 4.07	0.0004
9,18-diOH octadecanoic	15.09 ± 1.12	168.98 ± 12.04	0.0092	25.96 ± 0.87	0.0210	44.29 ± 3.66	0.7018	54.91 ±2.72	0.0002
Docosanoic acid	0 ± 0	0 ± 0	0.0000	0 ± 0	0.0000	0 ± 0	0.0000	0 ± 0	0.0000
Tetracosanoic acid	0 ± 0	0 ± 0	0.0000	0 ± 0	0.0000	0 ± 0	0.0000	8.79± 0.36	0.0017
TOTAL (µg cm ⁻²)	624.22 ± 59.39	1418.10 ± 68.69		335.43 ± 19.69		1272.24 ± 98.40		1408.37 ± 80.13	
RR									
Compound	AC	Penjar-1	Penjar-2		Penjar-3		Penjar-4		
	Mean ± SE	Mean ± SE	P-value	Mean ± SE	P-value	Mean ± SE	P-value	Mean ± SE	P-value
Glycerol	2.25 ± 0.27	8.55 ± 0.45	0.0017	8.55 ± 0.90	0.0053	9.45 ± 1.35	0.0304	432.45 ± 59.41	0.0030
Dodecanoic acid	6.34 ± 0.45	19.35 ± 1.82	0.0159	6.30 ± 0.45	0.0194	<i>1.80 ± 0.16</i>	0.0004	27.47 ± 2.25	0.0017
Hexadecanoic acid	33.75 ± 2.25	132.75 ± 5.43	0.0005	60.75 ± 6.75	0.0701	30.67 ± 3.24	0.5778	114.35 ± 5.85	0.0006
Heptadecanoic acid	14.62 ± 0.981	<i>4.52 ± 0.45</i>	0.0046	<i>1.80 ± 0.135</i>	0.0047	<i>1.85 ± 0.24</i>	0.0049	13.95 ± 1.84	0.0007
Octadecanol	18.17 ± 0.96	<i>2.70 ± 0.35</i>	0.0028	20.72 ± 1.35	0.0312	36.93 ± 3.6	0.0055	141.75 ± 19.08	0.0077
hydroxy hexadecanoic acid	28.35 ± 2.25	102.15 ± 12.6	0.0010	46.35 ± 5.40	0.5663	27.45 ± 2.25	0.9869	185.85 ± 17.14	0.0041
9(10),16-diOH hexadecanoic	613.35 ± 60.32	<i>45.91 ± 2.25</i>	0.0005	842.85 ± 118.80	0.0654	1683.22 ± 192.15	0.0034	2753.55 ± 239.45	0.0021
Octadecanoic acid	12.61 ± 1.35	54.94 ± 5.71	0.0504	21.62 ± 2.25	0.1170	9.94 ± 0.45	0.1027	119.76 ± 16.65	0.0003
18-OH octadecanoic	54.92 ± 5.85	219.61 ± 28.8	0.0040	91.35 ± 17.59	0.4278	47.73 ± 6.3	0.2023	197.55 ± 32.4	0.0002
9,18-diOH octadecanoic	33.37 ± 2.25	145.87 ± 25.20	0.0027	44.17 ± 3.55	0.0732	40.05 ± 4.5	0.9738	181.86 ± 25.65	0.0019
Docosanoic acid	0 ± 0	0 ± 0	0.0000	0 ± 0	0.0000	1.35 ± 0.20	0.0090	18.91 ± 2.92	0.0007
Tetracosanoic acid	0 ± 0	0 ± 0	0.0000	0 ± 0	0.0000	0 ± 0	0.0000	47.07 ± 19.81	0.1389
TOTAL (µg cm ⁻²)	817.42 ± 76.85	736.21 ± 83.11		1144.35 ± 157.18		1890 ± 214.45		4234.05 ± 442.35	

The values highlighted in bold and italics represent ≥1.5 fold increase or decrease respectively in Penjars compared to AC.

Penjar accessions	SNP position (cDNA)	Base change	SIFT Score (deleterious ≤ 0.05)	Amino acid change and position	Protein length (aa)
Penjar-1	20	C to A	Stop codon, deleterious	Stop codon	6
	37	C to A	0.15, tolerable	Q to K, 13 th	
Penjar-2	316	T to A	0.0, deleterious	V to D, 106 th	355
Penjar-3	316	T to A	0.0, deleterious	V to D, 106 th	355
Penjar-4	316	T to A	0.0, deleterious	V to D, 106 th	355

Figure 1. Mutations in NAC-NOR in *nor*, *alcobaca* and Penjar accessions. AC encodes a full length NOR protein of 355 aa. *nor* mutant produces a truncated protein of 186 aa. The Penjar-1 encodes a protein of 6 aa length, whereas *alcobaca*, Penjar-2, -3, and -4 encode a full length protein with a single amino acid change of V to D at 106th position (indicated by an asterisk and red band). NAC indicates NAC domain and TRR indicates transcriptional regulatory region. The details of base changes and the corresponding positions in the cDNA, protein, and SIFT score are indicated in the Table.

Figure 2. Post-harvest shelf life and water loss in AC and Penjar fruits. **A**, The AC and Penjar fruits were harvested at ripe stage and incubated under normal day-night conditions as described in methods. By 40 days post-harvest, WT fruits showed prominent wrinkling, and 65 day-old fruits displayed a clear sign of cellular disintegration. **B**, The water loss was measured during post-harvest storage. Data are means of 5 biological replicates $\pm$ SE, ‘*’ indicates $P \leq 0.05$.

Figure 3. Cuticular wax composition of AC and Penjar fruits during ripening. **A**, The fruits were harvested at ripe stage and cuticular wax from the surface of both AC and Penjar fruits was removed by using chloroform. This resulted in severe shrinking of fruits due to loss of moisture. **B**, Total cuticular wax content of AC and Penjar fruits at MG and RR. **C-F**, The relative abundance of cuticular wax components: alkanes (**C**), fatty acids (**D**), fatty alcohols (**E**), sterols and triterpenes (**F**) in MG and RR fruits of AC and Penjars. Data are means of 3 biological replicates $\pm$ SE, ‘*’ indicates $P \leq 0.05$.

Figure 4. Relative expression of cell wall modifying genes in AC and Penjar fruits. The fruits of AC and Penjars were collected at different ripening and SEN stages and expression of various cell wall modifying genes was determined by qPCR. The graphs represent the data obtained after normalization with *actin* and *ubiquitin*. **A**, α -Galactosidase, **B**, β -Galactosidase, **C**, α -Mannosidase, **D**, β -Mannosidase, **E**, Polygalacturonase 2A, **F**, Polygalacturonase β subunit, **G**, Pectinesterase, **H**, Expansin, **I**, Deoxyhypusine synthase and **J**, β -N-acetyl-D-hexosaminidase. Data are means of 3 biological replicates $\pm$ SE, ‘*’ indicates $P \leq 0.05$.

Figure 5. Hormone profiling in AC and Penjar fruits. The fruits of AC and Penjars were collected at different ripening and SEN stages and hormone profiles were determined as described in methods. **A**, IAA, indole-3-acetic acid; **B**, IBA, indole-3-butyric acid; **C**, zeatin, **D**, JA, jasmonic acid; **E**, MeJA, methyl jasmonate; **F**, SA, salicylic acid; **G**, ABA, abscisic acid. Data are means of 5 biological replicates $\pm$ SE, ‘*’ indicates $P \leq 0.05$.

Figure 6. The metabolic shifts in the fruits of Penjars in comparison to AC during ripening and post-harvest storage. The relative abundances of the metabolites in Penjar fruits at MG, BR, RR and SEN stages were obtained from the Log₂ Penjar/AC values and only those metabolites with ≥ 1.5 fold change ($n=5 \pm SE$, $P \leq 0.05$) were mapped on the pathway. The scale at the top left corner represents Log₂ fold changes in the range of -3 to +3. The metabolites indicated in gray letters on the pathway were not detected in our GC-MS analysis.